

10-2018

Synthesis of 8-deoxypumiliotoxin 193H and 9-deoxyhomopumiliotoxin 207O

Takuya Okada
University of Toyama

Taiga Yamamoto
University of Toyama

Daiki Kato
University of Toyama

Masashi Kawasaki
Toyama Prefectural University

Ralph Saporito
John Carroll University, rsaporito@jcu.edu

See next page for additional authors

Follow this and additional works at: https://collected.jcu.edu/fac_bib_2018

 Part of the [Biology Commons](https://biologycommons.org/)

Recommended Citation

Okada, Takuya; Yamamoto, Taiga; Kato, Daiki; Kawasaki, Masashi; Saporito, Ralph; and Toyooka, Naoki, "Synthesis of 8-deoxypumiliotoxin 193H and 9-deoxyhomopumiliotoxin 207O" (2018). *2018 Faculty Bibliography*. 73.
https://collected.jcu.edu/fac_bib_2018/73

This Article is brought to you for free and open access by the Faculty Bibliographies Community Homepage at Carroll Collected. It has been accepted for inclusion in 2018 Faculty Bibliography by an authorized administrator of Carroll Collected. For more information, please contact connell@jcu.edu.

Authors

Takuya Okada; Taiga Yamamoto; Daiki Kato; Masashi Kawasaki; Ralph Saporito; and Naoki Toyooka,



Synthesis of 8-deoxypumiliotoxin **193H** and 9-deoxyhomopumiliotoxin **207O**

Takuya Okada^{a,*}, Taiga Yamamoto^b, Daiki Kato^b, Masashi Kawasaki^c, Ralph A. Saporito^d, Naoki Toyooka^{a,b,*}

^a Graduate School of Innovative Life Science, University of Toyama, 3190 Gofuku, Toyama 930-8555, Japan

^b Graduate School of Science and Engineering, University of Toyama, 3190 Gofuku, Toyama 930-8555, Japan

^c Department of Liberal Arts and Sciences, Faculty of Engineering, Toyama Prefectural University, 5180 Kurokawa, Imizu, Toyama 939-0398, Japan

^d Department of Biology, John Carroll University, University Heights, OH 44118, USA

ARTICLE INFO

Article history:

Received 31 July 2018

Revised 28 August 2018

Accepted 6 September 2018

Available online 8 September 2018

Keywords:

Asymmetric synthesis

Neotropical poison frog

Pumiliotoxin

8-Deoxypumiliotoxin **193H**

9-Deoxyhomopumiliotoxin **207O**

ABSTRACT

The asymmetric synthesis of 8-deoxypumiliotoxin **193H** and 9-deoxyhomopumiliotoxin **207O** has been achieved, starting from both enantiomers of (+)- and (–)-**10**. Enantiomerically pure alcohols (+)- and (–)-**10** were obtained by lipase-mediated kinetic resolution of racemic **10**, which was prepared in 3 steps from new lactam-type building block (–)-**8** in a highly stereoselective manner.

© 2018 Published by Elsevier Ltd.

Introduction

To date, over 800 lipophilic alkaloids representing >20 structural classes have been detected from the skin extracts of poison frogs [1]. Pumiliotoxins (PTXs, **1**) are one of the major classes of frog alkaloids, and their structural diversity and pharmacological activity have stimulated considerable interest in their chemical synthesis by several groups. At present, over 30 alkaloids are considered to be PTXs and over 60 alkaloids belong to various congeners, such as homopumiliotoxins (hPTXs, **2**), 8-deoxypumiliotoxins (8-deoxy-PTXs, **3**), 9-deoxy-homopumiliotoxins (9-deoxy-hPTXs, **4**), 8-desmethyl-pumiliotoxins (8-desmethyl-PTXs, **5**), and 9-desmethyl-homopumiliotoxins (9-desmethyl-hPTXs, **6**) (Fig. 1).

Some of the proposed structures for PTXs are tentative, while others have been rigorously established by MS, FT-IR, NMR spectra, and/or synthesis. Among PTXs, over 10 enantioselective syntheses of PTX **251D** have been reported [2]; however, only one synthesis of 8-deoxy-PTX **193H** has been reported to date [3].

As part of a larger research program directed at studying the synthesis of poison frog alkaloids [4], we report here the asymmetric synthesis of 8-deoxy-PTX **193H** and 9-deoxy-hPTX **207O** (Fig. 2).

Results and discussion

The synthesis began with β -keto ester (\pm)-**7**, which was prepared in 3 steps from allylamine [5]. Reduction of (\pm)-**7** with NaBH₄ gave to the racemic alcohol (\pm)-**8** in low yield, while bakers' yeast reduction of (\pm)-**7** under nonfermenting conditions gave rise to the enantiopure alcohol (–)-**8** in good yield and high optical purity (>98% ee) [6]. Treatment of (–)-**8** with bis(2-methoxyethyl) azodicarboxylate and Ph₃P provided the enaminoester (\pm)-**9**, which was subjected to copper-mediated 1,4-addition of methylolithium, followed by reduction of the resulting adduct with Super-Hydride to afford the primary alcohol (\pm)-**10** in high yield as a single isomer (Scheme 1).

To obtain both enantiomers of (+)- and (–)-**10**, lipase-mediated kinetic resolution of (\pm)-**10** with lipase PL and vinyl acetate in *t*-BuOMe was conducted, and resulted in the acetate **11** in 55% yield with 58% ee, and the alcohol (–)-**10** in 36% yield with >98% ee, respectively. Since the ee of the acetate **11** from the initial resolution was not sufficient for further transformation, the acetate **11**

* Corresponding authors at: Graduate School of Innovative Life Science, University of Toyama, 3190 Gofuku, Toyama 930-8555, Japan (N. Toyooka).

E-mail addresses: author@university.edu (T. Okada), author@university.edu (N. Toyooka).




Fig. 1. Structures of pumiliotoxins and various congeners.






Fig. 2. Structures of PTX 251D, 8-deoxy-PTX 193H, and 9-deoxy-hPTX 2070.



Scheme 1. Reagents and conditions: (a) 1.05 eq. Ethyl bromoacetate, 1.05 eq. Et₃N, CH₂Cl₂, r.t.; (b) 1.2 eq. Ethyl succinyl chloride, 1.2 eq. Et₃N, CH₂Cl₂, r.t.; (c) 1.1 eq. NaH, THF, r.t. (70% in 3 steps); (d) bakers' yeast, tap water, r.t. (76%, >98% ee); (e) 1.1 eq. bis(2-methoxyethyl) azodicarboxylate, 1.1 eq. Ph₃P, r.t. (94%); (f) 2.5 eq. Me₂CuLi, Et₂O, -78 to 0 °C; (g) 3.0 eq. Super-Hydride, THF, r.t. (92% in 2 steps).

was hydrolyzed to the corresponding alcohol **10**, which was then subjected to a second kinetic resolution under the same reaction conditions to obtain the acetate (+)-**11**. Hydrolysis of the enantiopure acetate (+)-**11** with NaOH gave the enantiopure alcohol (+)-**10** with 98% ee (Scheme 2).

Once both enantiomers of **10** were obtained, we performed a Swern oxidation of (-)-**10**, which was followed by a Wittig reaction of the resulting aldehyde. The Wittig reaction provided the olefin (-)-**12**, which was converted to the tetrahydroindolizin-5-one (-)-**13** by ring-closing metathesis using a 2nd generation Grubbs catalyst. Hydrogenation of (-)-**13** over 10% Pd/C resulted in the indolizidinone (-)-**14**. The absolute stereochemistry of (-)-**10** was determined to be 5*R*, 6*R* by comparing the optical rotation between our synthetic (-)-**14** with known (+)-**14** [7]. Indolizidinone (-)-**14** was subjected to an aldol condensation with isobutyraldehyde to afford the aldol adduct **15**. Stereospecific *syn*-elimination of **15** using DCC/CuCl [2*b*] afforded the desired (*Z*)-olefin (-)-**16** with no (*E*)-olefin. Finally, a reduction of the lactam moiety of (-)-**16** using AlH₃, resulted in the formation of (-)-8-deoxy-PTX **193H** (Scheme 3). The ¹H and ¹³C NMR spectra of our synthetic (-)-8-deoxy-PTX **193H** were in good accordance




Scheme 2. Reagents and conditions: (a) lipase PL, vinyl acetate, *t*-BuOMe, r.t. ((+)-**11**: 55%, 58% ee, (-)-**10**: 36%, >98% ee); (b) 6 M NaOH (aq.), EtOH, r.t. (84%); (c) lipase PL, vinyl acetate, *t*-BuOMe, r.t. (47%); (d) 6 M NaOH (aq.), EtOH, r.t. (93%, 98% ee).


with data from the literature, however the optical rotation was opposite [3].

The enantiomer (+)-8-deoxy-PTX **193H** was also synthesized from (+)-**10** in the same manner as shown in Scheme 3.

Furthermore, both enantiomers of (+)- and (-)-9-deoxy-hPTX **2070** were also synthesized from the alcohols (-)- and (+)-**10**. The alcohol (-)-**10** was converted to the iodide (+)-**17**, which was subjected to a cross-coupling reaction with vinylmagnesium chloride in the presence of HMPA and P(OEt)₃ [8] to afford the diallylpiperidone (-)-**18** in 62% yield, together with the undesired ring-opening amide **19** in 22% yield, respectively. The ring-closing methathesis reaction of (-)-**18** using a 2nd generation Grubbs catalyst, followed by hydrogenation of the resulting olefin (-)-**20**, provided the quinolizidinone (+)-**21** [9], which was subjected to an aldol condensation to afford the aldol adduct **22**. Stereospecific *syn*-elimination of **22** using DCC/CuCl gave the undesired (*E*)-olefin (+)-**23a** in 81% yield, while anti-elimination [2*b*] by mesylation of **22**, followed by treatment of the resulting meslate with KOH, provided the desired (*Z*)-olefin (+)-**23b** in 70% yield in 2 steps. The double bond geometry in (+)-**23a** and (+)-**23b** was determined based on the chemical shift of the vinyl proton using ¹H NMR spectra. The vinyl proton of the (*Z*)-olefin (+)-**23b** was detected in a high magnetic field (δ 5.45 ppm), while that of the (*E*)-olefin (+)-**23a** was present in a low magnetic field (δ 6.56 ppm) due to the anisotropy of the amide-carbonyl group. These results are very similar to those of previous report [10]. It was noteworthy that the aldol adducts for indolizidinone **14** and quinolizidinone **21** were opposite. Finally, reduction of (+)-**23b** by treatment with AlH₃ resulted in the formation of (+)-deoxy-hPTX **2070** (Scheme 4). The enantiomer (-)-deoxy-hPTX **2070** was also synthesized from (+)-**10** in the same manner as shown in Scheme 4.



Scheme 3. Reagents and conditions: (a) 2.0 eq. $(\text{COCl})_2$, 4.0 eq. DMSO, 6.0 eq. Et_3N , CH_2Cl_2 , -78 to 0 °C; (b) 2.5 eq. $\text{MeP}^+\text{Ph}_3\text{Br}^-$, 2.25 eq. $n\text{-BuLi}$, THF, r.t.; (c) 5 mol% Grubbs catalyst 2nd generation, CH_2Cl_2 , r.t.; (d) H_2 , 10% Pd/C, EtOAc, r.t.; (e) 1.25 eq. isobutyraldehyde, 1.25 eq. lithium diisopropylamide, 1.25 eq. HMPA, THF, -78 to 0 °C; (f) 1.2 eq. DCC, 1.9 eq. CuCl , toluene, reflux; (g) 7.5 eq. LiAlH_4 , 2.5 eq. AlCl_3 , Et_2O , r.t.



Scheme 4. Reagents and conditions: (a) 1.5 eq. I_2 , 1.5 eq. Ph_3P , 1.5 eq. imidazole, CH_2Cl_2 , r.t.; (b) 3.3 eq. vinylmagnesium chloride, 1.1 eq. CuI , 2.2 eq. HMPA, 2.2 eq. $\text{P}(\text{OEt})_3$, THF, -40 to 0 °C; (c) 5 mol% Grubbs catalyst 2nd generation, CH_2Cl_2 , r.t.; (d) H_2 , 10% Pd/C, EtOAc, r.t.; (e) 1.25 eq. isobutyraldehyde, 1.25 eq. lithium diisopropylamide, 1.25 eq. HMPA, THF, -78 to 0 °C; (f) 1.2 eq. DCC, 1.9 eq. CuCl , toluene, reflux; (g) 10.0 eq. MsCl , pyridine, r.t.; (h) 10.0 eq. KOH , MeOH , reflux; (i) 7.5 eq. LiAlH_4 , 2.5 eq. AlCl_3 , Et_2O , r.t.

Conclusion

We have achieved the asymmetric synthesis of both enantiomers of 8-deoxy-PTX **193H** and 9-deoxy-hPTX **2070** from the common enantiopure alcohol **10**. Furthermore, we are trying to synthesize 8-desmethyl-PTXs and 9-desmethyl-hPTXs from the enantiopure alcohol (**-**)-**8**. The antagonistic activities of both enantiomers of these PTX alkaloids against nicotinic acetylcholine receptors are now in progress.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.tetlet.2018.09.015>.

References

- [1] J.W. Daly, T.F. Spande, H.M. Garraffo, *J. Nat. Prod.* **68** (2005) 1556–1575.
- [2] (a) L.E. Overman, K.L. Bell, F. Ito, *J. Am. Chem. Soc.* **106** (1984) 4192–4201; (b) D.N.A. Fox, D. Lathbury, M.F. Mahon, K.C. Molloy, T. Gallagher, *J. Am. Chem. Soc.* **113** (1991) 2652–2656; (c) T. Honda, M. Hoshi, M. Tsubuki, *Heterocycles* **34** (1992) 1515–1518; (d) J. Cossy, M. Cases, D. Gomez Pardo, *Synlett* (1996) 909–910; (e) A.G.M. Barrett, F. Damiani, *J. Org. Chem.* **64** (1999) 1410–1411; (f) S.F. Martin, S.K. Bur, *Tetrahedron* **55** (1999) 8905–8914; (g) A. Sudau, W. Munch, J. Bats, U. Nubbemeyer, *Eur. J. Org. Chem.* (2002) 3304–3314; (h) A. Sudau, W. Munch, J. Bats, U. Nubbemeyer, *Eur. J. Org. Chem.* (2002) 3315–3325; (i) K.S. Woodin, T.F. Jamison, *J. Org. Chem.* **72** (2007) 7451–7454; (j) P.R. Sultane, A.R. Mohite, R.G. Bhat, *Tetrahedron Lett.* **53** (2012) 5856–5858; (k) B. Bernardim, V.D. Pinho, A.C.B. Burtoloso, *J. Org. Chem.* **77** (2012) 9926–9931;

- (l) V.D. Pinho, D.J. Procter, A.C.B. Burtoloso, *Org. Lett.* 15 (2013) 2434–2437;
(m) S.P. Chou, K. Yang, T. Chiu, *Heterocycles* 89 (2014) 679–691.
- [3] G. Smits, R. Zemribo, *Eur. J. Org. Chem.* (2015) 3152–3156.
- [4] (a) N. Toyooka, K. Tanaka, T. Momose, J.W. Daly, H.M. Garraffo, *Tetrahedron* 53 (1997) 9553–9574;
(b) N. Toyooka, A. Fukutome, H. Nemoto, J.W. Daly, T.F. Spande, H.M. Garraffo, T. Kaneko, *Org. Lett.* 4 (2002) 1715–1718;
(c) N. Toyooka, H. Nemoto, *Tetrahedron Lett.* 44 (2003) 569–570;
(d) N. Toyooka, A. Fukutome, H. Shinoda, H. Nemoto, *Angew. Chem. Int. Ed.* 42 (2003) 3808–3810;
(e) H. Tsuneki, Y. You, N. Toyooka, S. Kagawa, S. Kobayashi, T. Sasaoka, H. Nemoto, I. Kimura, J.A. Dani, *Mol. Pharmacol.* 66 (2004) 1061–1069;
(f) N. Toyooka, H. Nemoto, M. Kawasaki, H.M. Garraffo, T.F. Spande, J.W. Daly, *Tetrahedron* 61 (2005) 1187–1198;
(g) N. Toyooka, Z. Dejun, H. Nemoto, H.M. Garraffo, T.F. Spande, J.W. Daly, *Tetrahedron Lett.* 47 (2006) 577–580;
(h) N. Toyooka, Z. Dejun, H. Nemoto, H.M. Garraffo, T.F. Spande, J.W. Daly, *Tetrahedron Lett.* 47 (2006) 581–582;
(i) N. Toyooka, S. Kobayashi, D. Zhou, H. Tsuneki, T. Wada, H. Sakai, H. Nemoto, T. Sasaoka, H.M. Garraffo, T.F. Spande, J.W. Daly, *Bioorg. Med. Chem. Lett.* 17 (2007) 5872–5875;
(j) N. Toyooka, D. Zhou, S. Kobayashi, H. Tsuneki, T. Wada, H. Sakai, H. Nemoto, T. Sasaoka, Y. Tezuka, S. Subehan, S. Kadota, H.M. Garraffo, T.F. Spande, J.W. Daly, *Synlett* (2008) 61–64;
(k) N. Toyooka, D. Zhou, H. Nemoto, Y. Tezuka, S. Kadota, N.R. Andriamaharavo, H.M. Garraffo, T.F. Spande, J.W. Daly, *J. Org. Chem.* 74 (2009) 6784–6791;
(l) X. Wang, H. Tsuneki, N. Urata, Y. Tezuka, T. Wada, T. Sasaoka, H. Sakai, R.A. Saporito, N. Toyooka, *Eur. J. Org. Chem.* (2012) 7082–7092;
(m) X. Wang, J. Li, R.A. Saporito, N. Toyooka, *Tetrahedron* 69 (2013) 10311–10315.
- [5] J. Bonjoch, I. Serret, J. Bosch, *Tetrahedron* 40 (1984) 2505–2511.
- [6] N. Toyooka, Y. Yoshida, Y. Yotsui, T. Momose, *J. Org. Chem.* 64 (1999) 4914–4919.
- [7] (a) F. Abels, C. Lindemann, E. Koch, C. Schneider, *Org. Lett.* 14 (2012) 5972–5975;
(b) F. Abels, C. Lindemann, C. Schneider, *Chem. Eur. J.* 20 (2014) 1964–1979.
- [8] S.A.M.W. van den Broek, J.G.H. Lemmers, F.L. van Delft, F.P.J.T. Rutjes, *Org. Biomol. Chem.* 10 (2012) 945–951.
- [9] C. Lindemann, C. Schneider, *Synthesis* 48 (2016) 828–844.
- [10] L.S. Santos, R.A. Pilli, *Tetrahedron Lett.* 42 (2001) 6999–7001.