



University of Kentucky
UKnowledge

Entomology Faculty Publications

Entomology

4-16-2013

The endosymbiont *Arsenophonus* is widespread in soybean aphid, *Aphis glycines*, but does not provide protection from parasitoids or a fungal pathogen

Jason A. Wulff

University of Kentucky, jasonwulff@uky.edu

Karrie A. Buckman

University of Minnesota - Twin Cities

Kongming Wu

Chinese Academy of Agricultural Sciences, China

George E. Heimpel

University of Minnesota - Twin Cities

Jennifer A. White

University of Kentucky, jenawhite@uky.edu

[Click here to let us know how access to this document benefits you.](#)

Follow this and additional works at: https://uknowledge.uky.edu/entomology_facpub

 Part of the [Entomology Commons](#)

Repository Citation

Wulff, Jason A.; Buckman, Karrie A.; Wu, Kongming; Heimpel, George E.; and White, Jennifer A., "The endosymbiont *Arsenophonus* is widespread in soybean aphid, *Aphis glycines*, but does not provide protection from parasitoids or a fungal pathogen" (2013).

Entomology Faculty Publications. 25.

https://uknowledge.uky.edu/entomology_facpub/25

This Article is brought to you for free and open access by the Entomology at UKnowledge. It has been accepted for inclusion in Entomology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

The endosymbiont *Arsenophonus* is widespread in soybean aphid, *Aphis glycines*, but does not provide protection from parasitoids or a fungal pathogen

Notes/Citation Information

Published in *PLoS ONE*, v. 8, no. 4, e62145.

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

[Minor Correction: Errors in references](#)

Digital Object Identifier (DOI)

<http://dx.doi.org/10.1371/journal.pone.0062145>

The Endosymbiont *Arsenophonus* Is Widespread in Soybean Aphid, *Aphis glycines*, but Does Not Provide Protection from Parasitoids or a Fungal Pathogen

Jason A. Wulff^{1*}, Karrie A. Buckman^{2,3}, Kongming Wu⁴, George E. Heimpel³, Jennifer A. White¹

1 Department of Entomology, University of Kentucky, Lexington, Kentucky, United States of America, **2** USDA, Agricultural Research Service, Center for Grain and Animal Health Research Unit, Manhattan, Kansas, United States of America, **3** Department of Entomology, University of Minnesota, St. Paul, Minnesota, United States of America, **4** State Key Laboratory for Plant Diseases and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing, China

Abstract

Aphids commonly harbor bacterial facultative symbionts that have a variety of effects upon their aphid hosts, including defense against hymenopteran parasitoids and fungal pathogens. The soybean aphid, *Aphis glycines* Matsumura (Hemiptera: Aphididae), is infected with the symbiont *Arsenophonus* sp., which has an unknown role in its aphid host. Our research goals were to document the infection frequency and diversity of the symbiont in field-collected soybean aphids, and to determine whether *Arsenophonus* is defending soybean aphid against natural enemies. We performed diagnostic PCR and sequenced four *Arsenophonus* genes in soybean aphids from their native and introduced range to estimate infection frequency and genetic diversity, and found that *Arsenophonus* infection is highly prevalent and genetically uniform. To evaluate the defensive role of *Arsenophonus*, we cured two aphid genotypes of their natural *Arsenophonus* infection through ampicillin microinjection, resulting in infected and uninfected isolines within the same genetic background. These isolines were subjected to parasitoid assays using a recently introduced biological control agent, *Binodoxys communis* [Braconidae], a naturally recruited parasitoid, *Aphelinus certus* [Aphelinidae], and a commercially available biological control agent, *Aphidius colemani* [Braconidae]. We also assayed the effect of the common aphid fungal pathogen, *Pandora neoaphidis* (Remaudiere & Henebert) Humber (Entomophthorales: Entomophthoraceae), on the same aphid isolines. We did not find differences in successful parasitism for any of the parasitoid species, nor did we find differences in *P. neoaphidis* infection between our treatments. Our conclusion is that *Arsenophonus* does not defend its soybean aphid host against these major parasitoid and fungal natural enemies.

Citation: Wulff JA, Buckman KA, Wu K, Heimpel GE, White JA (2013) The Endosymbiont *Arsenophonus* Is Widespread in Soybean Aphid, *Aphis glycines*, but Does Not Provide Protection from Parasitoids or a Fungal Pathogen. PLoS ONE 8(4): e62145. doi:10.1371/journal.pone.0062145

Editor: Kostas Bourtzis, International Atomic Energy Agency, Austria

Received: December 14, 2012; **Accepted:** March 18, 2013; **Published:** April 16, 2013

This is an open-access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication.

Funding: This research was supported in part by a grant from the Kentucky Science and Engineering Foundation as per Grant Agreement #KSEF-148-502-09-248 with the Kentucky Science and Technology Corporation, USDA-AFRI grant # 2011-67014-3015, and the University of Kentucky. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: jasonwulff@uky.edu

Introduction

Maternally inherited bacterial endosymbionts are common in arthropods [1–5]. Many insects are infected with obligate nutritional endosymbionts that are required for survival, e.g. *Buchnera aphidicola* in aphids [1,2,6]. In contrast, facultative endosymbionts are not strictly required for insect survival, but can provide a selective advantage in certain ecological contexts [7]. For example, facultative endosymbionts have been shown to provide their hosts with heat shock resistance [8], modify host color [9], and potentially facilitate host plant colonization [10]. A subset of these facultative endosymbionts can also defend their insect hosts against natural enemies such as parasitoids, entomopathogenic fungi, viruses, and nematodes [11–14].

Bacterial symbionts in the genus *Arsenophonus* are estimated to infect approximately 5% of arthropods [4,15]. In the parasitoid wasp *Nasonia vitripennis*, *Arsenophonus nasoniae* acts as a male killing reproductive parasite [16–19]. Other strains are thought to be obligate symbionts of triatomine bugs, hippoboscids and streblid

flies, and lice [20–22], and yet others are plant pathogens [23–25]. *Arsenophonus* is also found in multiple whitefly, psyllid, and aphid species [26–30], but its function among these hosts remains uncharacterized. However, there have been suggestions that *Arsenophonus* may play a defensive role. In a geographic survey of the lerp psyllid, *Glycaspis brimblecombei*, Hansen et al. (2007) found a positive correlation between parasitism and the frequency of *Arsenophonus* infection, potentially indicating that *Arsenophonus* provides the psyllid with a selective advantage in populations under heavy parasitism pressure [31].

If *Arsenophonus* provides defense against natural enemies, then it could be an important consideration in biological control programs against *Arsenophonus*-bearing pests. For example, a defensive symbiont that is present at low prevalence within a population could become common under selective pressure provided by a newly released classical biological control agent, thus undercutting the efficacy of the agent [32,33]. Alternatively, laboratory populations, which experience vastly different selective

environments and frequent population bottlenecks [34], might be expected to have a different frequency of symbiont infection than field populations. In such a case, conclusions about natural enemy efficacy drawn from laboratory studies may have little bearing on natural enemy performance in the field.

Multiple important pest species are infected with *Arsenophonus*, including the lespyid, the cotton aphid, *Aphis gossypii*, the sweet potato whitefly, *Bemisia tabaci*, and the soybean aphid, *Aphis glycines* [26,31,35,36]. Soybean aphid is a serious invasive pest of soybeans in North Central United States, causing extensive yield loss and requiring intensive pesticide applications to a crop that required little pesticide input prior to the introduction of the soybean aphid [37]. Early parasitism surveys in North America found that soybean aphids were infrequently parasitized [38–40], leading to ongoing biological control investigations that incorporate augmentation of ambient fungal pathogens and introduction of parasitoids from the aphid's native range [41–43]. The function and prevalence of *Arsenophonus* in field populations of soybean aphid has the potential to affect these pest management tactics.

The goals of this study were 1) to document the frequency and diversity of *Arsenophonus* infection in field-collected soybean aphids from the aphids' native and introduced range and 2) to investigate whether *Arsenophonus* protects soybean aphid against parasitoid wasps or entomopathogenic fungi by assessing natural enemy efficacy against infected versus experimentally cured aphid isolines. For the former goal, we performed *Arsenophonus* diagnostic PCR on six native and seven introduced populations of soybean aphid, followed by multi-locus strain typing (MLST) of *Arsenophonus* using 3 bacterial genes [30,44]. For the latter goal, we assayed three species of parasitoid wasp and one species of fungal pathogen. The first parasitoid species assayed was *Binodoxys communis*, which currently is the only exotic parasitoid to have been intentionally released in the United States to control the soybean aphid as part of a classical biological control program [42]. The second wasp, *Aphelinus certus*, has been identified from parasitized North American soybean aphids, although estimates of parasitism rates are still forthcoming. This parasitoid is native to China, was potentially co-introduced with soybean aphid, and is of interest as a biological control agent [45]. The third wasp, *Aphidius colemani*, is a commercially-available generalist parasitoid of aphids that is known to be susceptible to a defensive symbiont in pea aphid [46]. The aphid fungal pathogen, *Pandora neoaphidis*, is also known to be susceptible to defensive symbionts in pea aphid, and is being investigated for augmentative biological control of the soybean aphid [12,47,48].

Results

Geographic survey

When the prevalence of *Arsenophonus* in native and introduced populations of the soybean aphid was surveyed, we found that the symbiont was very common in all examined populations (Table 1). In the introduced North American range, a mean (\pm S.E.) of $98\pm 1\%$ of aphids were infected, which was slightly, but significantly, higher than the $85\pm 6\%$ infection found in the native Asian range (Wald = 2.128, df = 11, $P = 0.0334$).

Arsenophonus MLST

Arsenophonus fbaA, *ftsK*, *yaeT* genes were sequenced from one aphid from each of our surveyed populations [30,44]. We did not detect any genetic variation among sequences from the native and introduced populations, giving no evidence for multiple strains of *Arsenophonus* within soybean aphid.

Parasitism assays

The influence of *Arsenophonus* on soybean aphid susceptibility to parasitism was assessed using three different parasitoids. Parasitism by the introduced biological control agent *B. communis* did not differ significantly between *Arsenophonus*-infected and experimentally cured aphids of a Kentucky (KY) origin isolate within either a cage assay ($t = 0.88$, $df = 18$, $P = 0.39$), or an observation assay ($t = 0.22$, $df = 22$, $P = 0.83$; Figure 1A). Parasitism of a Minnesota (MN) origin isolate of aphids was substantially lower than the KY isolate, but again did not differ between *Arsenophonus*-infected and experimentally cured aphids in either the cage assay ($t = 0.86$, $df = 22$, $P = 0.40$), or the observation assay ($t = 0.12$, $df = 22$, $P = 0.90$).

There were no differences in *A. certus* parasitism of the KY isolate in the cage assay ($t = 0.38$, $df = 22$, $P = 0.71$) or the observation assay ($t = 0.52$, $df = 20$, $P = 0.61$), nor of the MN isolate in the cage assay ($t = 0.02$, $df = 19$, $P = 0.98$) or the observation assay ($t = 0.99$, $df = 18$, $P = 0.33$; Figure 1B). *A. certus* had the greatest disparity in performance between the two assays, with very low rates of parasitism for cage assays compared to the observation assays.

There were also no differences in proportion parasitism by *A. colemani* between infected and experimentally cured soybean aphid for either isolate or parasitism assay (KY cage assay: $t = 0.33$, $df = 20$, $P = 0.75$; KY observation assay: $t = 0.29$, $df = 24$, $P = 0.77$; MN cage assay: $t = 0.97$, $df = 20$, $P = 0.34$; MN observation assay: $t = 1.87$, $df = 18$, $P = 0.07$; Figure 1C).

Fungal assays

In a challenge using the entomopathogenic fungus *P. neoaphidis*, observed proportions of infection were highly variable, ranging from 0 to 0.76 per replicate. Mean (\pm SE) proportion *P. neoaphidis* infection in the *Arsenophonus* infected and uninfected aphids in the KY isolate were 0.15 ± 0.05 and 0.12 ± 0.06 respectively, and arcsine squareroot transformed values did not differ significantly from one another ($t = 0.58$, $df = 18$, $P = 0.57$). Likewise, *Arsenophonus* infected and uninfected aphids in the MN isolate had 0.22 ± 0.07 and 0.13 ± 0.05 proportion infected, and again did not differ significantly from one another ($t = 1.46$, $df = 18$, $P = 0.16$).

Discussion

Our primary goal was to assess whether *Arsenophonus* defends soybean aphid against natural enemies. Using three parasitoid wasp species, we found no evidence that *Arsenophonus* provides this defense in either of two genotypes of soybean aphid. All three species of parasitoids were able to successfully attack soybean aphid, and there were no significant differences in successful parasitism of *Arsenophonus*-infected versus cured aphids in either cage or observation assays. Likewise, we found no difference in aphid mortality from the fungus *P. neoaphidis* based on *Arsenophonus* infection.

Our aggregated results indicate that *Arsenophonus* is likely not a defensive symbiont in soybean aphid, but some caveats should be considered. First, we used only two genotypes of aphids, which were infected with the same strain type of *Arsenophonus*, based on identical *Arsenophonus* ribosomal and MLST sequences. It is possible that other *Arsenophonus* strains may provide protection to other genotypes of soybean aphid host. For example, different strains of the bacterial endosymbiont *Hamiltonella defensa* provide differential protection against parasitism to pea aphid based on the presence or absence and type of APSE phage [49]. Additionally, a strain of *Regiella insecticola* was recently shown to protect its aphid

Table 1. Soybean aphid, *Aphis glycines*, collection locations, year collected, collector, and *Arsenophonus* prevalence.

Locality	Year	Collector	<i>Arsenophonus</i> positive/Aphids screened
Native			
Hebei Province, China	2008	Wu Kongming	8/8
Shangdong Province, China	2008	Wu Kongming	9/10
Guangxi Province, China	2008	Wu Kongming	10/10
Hangzou District, China	2008	Wu Kongming	7/10
Yangling District, China	2008	Wu Kongming	9/10
Harbin Province, China	2008	Wu Kongming	5/8
Introduced			
Whitley Co., Indiana, USA	2008	Marc Rhainds	23/25
Tippecanoe Co., Indiana, USA	2008	Marc Rhainds	10/10
Wabash Co., Indiana, USA	2008	Marc Rhainds	5/5
Huntington Co., Indiana, USA	2008	Marc Rhainds	5/5
Olmsted Co., Minnesota, USA	2008	Fritz Breitenbach	5/5
Waseca Co., Minnesota, USA	2008	George Heimpel	5/5
Fayette Co., Kentucky, USA	2011	Jason Wulff	27/28

doi:10.1371/journal.pone.0062145.t001

host against parasitism, a trait not previously associated with the symbiont [46], indicating that bacterial strains can vary in their defensive properties. However, in soybean aphid our broad MLST survey of *Arsenophonus* did not identify any additional bacterial strains in either the native or introduced range, indicating that hypothetical alternate strain types are rare, if they exist at all. Furthermore, soybean aphid is a recent introduction to North America, and is notably lacking in genetic diversity [50]; consequently, it seems unlikely that additional sampling of aphid/symbiont genotypes in the invaded range would yield different results.

We limited our parasitism assays to wasp species relevant to the North American introduced range of soybean aphid. *B. communis* and *A. certus* are both of interest for biological control and represent two different families of parasitoids (Braconidae and Aphelinidae, respectively), the latter being a more generalized parasitoid species [45,51]. However, there is growing evidence that defensive symbiont-mediated selection can favor parasitoid genotypes that are insensitive to the symbiont [52]. The high prevalence of *Arsenophonus* infection in the field makes it likely that field-collected parasitoids of soybean aphid have encountered and potentially adapted to the symbiont. *A. colemani*, the third wasp we assayed, was commercially cultured on other aphid species and presumably naïve to soybean aphid, yet it was also unaffected by *Arsenophonus*.

Although our results indicate that *Arsenophonus* does not defend its host against these natural enemies, it does have a very high infection rate in both the introduced and native populations. Several possible explanations could underlie this widespread infection. First, *Arsenophonus* could manipulate host reproduction. Reproductive manipulation is a common means by which endosymbionts promote their own infection, and has recently been documented in the sexual generation of pea aphid by the endosymbiont *Spiroplasma* [53,54]. Second, *Arsenophonus* could be providing other context-specific benefits to soybean aphid, e.g. heat tolerance, defense against other pathogens [8,14], or general fecundity or longevity effects [55]. Third, *Arsenophonus* may be transmitted horizontally, either directly between aphids or

indirectly through the plant [56,57]. Finally, high fidelity vertical transmission, coupled with a very low metabolic cost to the host, could permit *Arsenophonus* to persist in a population without any benefit to the host [58]. However, other endosymbionts that had been considered previously to be neutral passengers were subsequently found to be extremely beneficial to their hosts under certain circumstances [13,59]. Given the very high prevalence of *Arsenophonus* in soybean aphid, it is therefore reasonable to presume that *Arsenophonus*, too, provides soybean aphid with a context-specific benefit that remains to be elucidated.

Materials and Methods

Geographic survey

To evaluate the prevalence of *Arsenophonus*, soybean aphids were collected from the Asian native range and North American invasive range. Collections were made either at university agricultural stations or on private lands with landowner permission (Table 1). For each population, 30 adult aphids were collected from plants at least 1 meter apart to minimize sampling of siblings, and immediately placed in 95% ethanol. Five aphids were selected at random from each introduced range population and ten aphids were selected from each native range population for molecular analysis. We extracted DNA by homogenizing individual aphids in 100 μ l of 10% w/v Chelex (Sigma-Aldrich, St Louis, MO, USA) in PCR-grade purified water. We added 6 μ l of proteinase K to each sample, vortexed, incubated overnight at 56°C, and then incubated samples at 96°C for ten minutes. We screened for the presence of *Arsenophonus* using a diagnostic PCR protocol modified from Thao and Baumann [26], which uses *Arsenophonus* specific primers to amplify the intervening region between 16S and 23S rDNA: Ars23S-1 (5'-CGTTTGATG ATTCATAGTCAAA-3') and Ars23S-2 (5'-GGTCCTCCAGTTAGTGTACCCAAC -3'). Reactions totaled 10 μ l, containing: 2.0 μ l of DNA template, 1.0 μ l of 25 mM MgCl₂, 1.0 μ l of 10 mM dNTP mixture, 1.0 μ l of Invitrogen 10 \times buffer (MgCl₂ free), 0.8 μ l of 5.0 pmole μ l⁻¹ of each primer, 0.1 μ l of 5 U/ μ l Invitrogen Taq polymerase, and ddH₂O to 10 μ l. PCR conditions were: initial denature at 95°C




Figure 1. Mean (\pm SE) proportion of soybean aphids parasitized by *Binodoxys communis* (A), *Aphelinus certus* (B), and *Aphis colemani* (C). Black bars represent naturally *Arsenophonus*-infected soybean aphids and white bars represent experimentally cured isolines with the same genetic background. Two isoline pairs (KY and MN) were each evaluated in two experiments (cage and observation assays) for each parasitoid species. No significant differences were detected in any assay.

doi:10.1371/journal.pone.0062145.g001

for 5 min; followed by 30 cycles of (95°C, 30 s; 55°C, 30 s; 72°C, 45 s); and final elongation at 70°C for 10 min. All PCRs included negative and positive controls. Product from multiple samples was sequenced to confirm *Arsenophonus*. All sequences were identical and the shared sequence was submitted to Genbank (Accession number KC019882). As a further control of extraction quality, we ran samples with the primers CAIF (5'-GCCTGATGCAGC-CATGCCGCGTGTATG-3') and CAIR (5'-GTCATCCC-CACCTTCC-3') with the same PCR conditions as previously listed. These primers were developed by Dale et al. [60] to target *Arsenophonus* 16S sequence in the hippoboscid fly, *Pseudolynchia canariensis*. However, they reliably detected 16S sequence from the obligate symbiont *Buchnera aphidicola* in soybean aphid, as confirmed by sequencing results (Accession number KC019881). Because this obligate symbiont should be present in all extractions, any samples that failed to amplify *B. aphidicola* were considered to be of poor quality and discarded. To compare *Arsenophonus* infection prevalence between the native and introduced ranges, we used logistic regression (Arc v. 1.06). To avoid overrepresentation of heavily sampled geographic regions, aphids collected from within the same county were considered to come from a single population, and pooled prior to statistical analysis.

MLST

We investigated potential genetic diversity in *Arsenophonus* using an MLST approach. We randomly selected a single extraction from each native and introduced population (Table 1), as well as from our two experimental colonies (KY and MN). We amplified DNA from each sample with the following primer sets: *fbaAf* (5'-GCYGCYAAAGTTTCRTTCCC-3') and *fbaAr2* (5'-GGCAAAT-TAAATTTCTGCGCAACG-3'), *ftsKf* (5'-GTTGTYATGGTY-GATGAATTTG-3') and *ftsKr* (5'-GCTCTTCATCACYTCA-WAACC-3'), *yaeTf* (5'-GCATACGGTTCAGACGGGTTTG-3') and *yaeTr* (5'-GCCGAAACGCCTTCAGA AAAG-3'). The PCR reaction recipe followed the protocol above and PCR conditions were: initial denature at 93°C for 3 min; 30 cycles of (93°C, 30 s; 52°C, 30 s; 72°C, 1 min); and final elongation at 72°C for 5 min [30,44]. Because sequences generated from each population were identical for each of the genes, *fbaA*, *ftsK*, and *yaeT*, a single sequence per gene was submitted to Genbank (KC701199, KC701198, KC701197).

Arsenophonus curing and colony maintenance

We used two soybean aphid clones for experimental manipulations. These clones were collected independently of the geographic survey specimens. One aphid clone, "KY", was initially collected in Fayette County, KY in 2009. The second clone, "MN", was originally collected in Ramsey County, MN and was maintained in culture at the University of Minnesota prior to transfer to Kentucky in 2010 (USDA Permit # P526P-10-00818). In addition to *Arsenophonus*, each aphid clone was screened diagnostically for other known bacterial symbionts of aphids [29], and examined for total bacterial diversity using denaturing gradient gel electrophoresis (DGGE) of bacterial 16S sequences [61]. The only bacterial endosymbionts detected were *Arsenophonus* and *Buchnera* (J. Wulff, unpublished data).

We cured these aphid clones of *Arsenophonus* infection using antibiotic microinjection, following a protocol modified from Oliver et al. [11]. Individual aphids from each clone were immobilized on a screen-covered pipette tip attached to vacuum, under a stereo microscope. Antibiotic was fed into a borosilicate microinjection needle attached to a syringe via tubing. Fourth-instar aphids were injected with 1.0 mg ml⁻¹ ampicillin solution [62]. *Arsenophonus* is susceptible to ampicillin, but the aphid's

primary symbiont, *Buchnera aphidicola*, is not [63]. After the initial injection, aphids were individually placed on excised soybean leaves maintained on 1% w/v agar, monitored for survivors, and a subset of offspring were checked for *Arsenophonus* via diagnostic PCR. This procedure was repeated for two subsequent generations using offspring of survivors from the previous bout of injections. Cured and infected isoline colonies were kept at $25 \pm 1^\circ\text{C}$ and 16L:8D on Asgrow® G4303 variety commercial soybeans in 10 cm pots. Plants were individually caged in 3.78 liter plastic jars that had panels of mesh to allow ventilation while preventing aphid escape. Aphids were transferred to new plants as needed, approximately twice per month, to avoid overcrowding and prevent alate production. All aphid isolines were maintained in this manner for at least 3 months prior to experiments. Five individuals from each soybean aphid isoline were screened with diagnostic PCR at least every 2 months to assure that the isoline retained the expected infection status. The cured aphid isolines never tested positive for *Arsenophonus*.

Parasitism assays

We evaluated the influence of *Arsenophonus* in soybean aphid on parasitism success by three parasitoid wasp species. The classical biological control agent *Binodoxys communis* was initially collected in August 2002 near Harbin, in the Chinese province of Heilongjiang, and was maintained in quarantine in St. Paul, Minnesota prior to initiation of our colony in Kentucky (USDA-APHIS permit P526P-10-01532) [64]. *Aphelinus certus* was collected locally in Lexington, KY in August 2010 from parasitized soybean aphids. *Aphidius colemani* is a commercially available biological control agent of aphids (APHIPAR, Koppert Biological Systems, The Netherlands). Each species of parasitoid was maintained in culture with *Arsenophonus*-cured soybean aphids and soybean plants at $25 \pm 1^\circ\text{C}$ and 16L:8D in the previously described culture jars with supplemental honey and water for at least two generations prior to use in parasitism assays.

Cage parasitism assays

We conducted cage parasitism assays using methodology adapted from Oliver et al. [11]. For each *Arsenophonus* infected/cured isoline pair, we assayed parasitism success by each of the three parasitoid species in separate experiments (6 assays total). For each assay, 12 vegetative stage 2 (V2) soybean plants were infested with *Arsenophonus*-infected aphids and 12 V2 soybean plants were infested with *Arsenophonus*-cured aphids. We transferred a leaf with >100 juvenile aphids to each experimental plant. Experimental plants were covered with cup cages, constructed from 947 ml translucent plastic containers, organza screening material, and weather stripping to provide a tight seal between cage and pot. After allowing 24 h for aphid establishment, we culled the aphids to either 30 aphids (*A. certus* assays), or 50 aphids (*B. communis* and *A. colemani* assays). *B. communis* and *A. certus* assays were conducted primarily with 2nd and 3rd instar aphids, whereas *A. colemani* assays were conducted primarily with 3rd and 4th instar aphids [64,65]. A single mated female wasp was introduced to each cup cage and removed after 24 h. If the wasp was dead or missing after this interval, the replicate was discarded. After 10 d, parasitized aphids (mummies) were counted, and proportion parasitism was calculated by dividing the number of mummies observed by the initial aphid number for that replicate. For each assay, the effect of aphid infection status on proportion parasitism was assessed using a t-test (IBM SPSS v20). Proportion data required an arcsine square-root transformation to satisfy the assumptions of the model.

Observation assays

Six observation assays were conducted in parallel to the cage assays, using the same three parasitoid species and two aphid genotypes. For each experiment, soybean leaves infested with either *Arsenophonus*-infected or cured aphids of all instars were embedded, adaxial side, in 1% agar in 100×15 mm petri dishes. Five to ten wasps of the same species were aspirated onto the embedded leaf. Wasps were allowed to settle and then culled to four actively parasitizing wasps. Wasps were observed continuously under a dissecting microscope. When oviposition was observed, each stung aphid was moved to a 35 mm leaf disk embedded in 1% agar, until a total of 10–15 aphids were parasitized, constituting a replicate. This procedure was repeated with fresh wasps until 10 replicates were generated per treatment per assay.

We regularly removed aphid progeny from leaf disks to avoid confusing progeny with the original stung aphids. Wasp mummies typically formed within 5–7 days, regardless of the parasitoid species. On day 10, we calculated proportion parasitism by dividing the number of mummies by the number of aphids that had survived until just prior to mummy formation. Aphids that died prior to day 5 were excluded from the data. Proportions were arcsine square-root transformed and analyzed using a t-test for each assay.

Fungal assays

To assess the effect of *Arsenophonus* infection status on soybean aphid susceptibility to the entomophthorean fungus *P. neoaphidis*, we conducted bioassays of *Arsenophonus*-infected versus cured aphids using the same two aphid genotypes as the parasitism assays. For each replicate, we transferred 25, 3rd–4th instar alate nymphs to a 100×15 mm, sterile, polystyrene petri dish containing moistened filter paper and an excised soybean leaflet (variety S19R5; NK, Golden Valley, MN). The petiole of each leaflet was placed in moist florist foam to prevent leaflet desiccation. To measure aphid exposure to fungal conidia, a glass cover slip was attached to each leaflet to allow for enumeration of conidia after aphid exposure to cultures.

We initiated a total of 20 replicates for each aphid isoline pair, 10 each from the infected and cured isolines. We used actively sporulating *P. neoaphidis* cultures to inoculate aphids. Subcultures used in the assays had been established 30–40 days prior to use and were only used after sporulation became evident (i.e., when conidia became visible on culture lids). All fungal cultures originated from the same *P. neoaphidis* isolate, which had been initially isolated from an infected, field-collected pea aphid (*Acyrtosiphon pisum*). The field collected isolate was used to infect soybean aphids in the laboratory, after which, the fungus was recovered from a single infected soybean aphid. The resulting isolate was periodically passed through and recovered from single soybean aphid individuals prior to use in the assays. Such periodic infection and recovery was necessary to maintain culture pathogenicity. Cultures used to infect soybean aphids in these assays originated from a single culture recovered from an infected soybean aphid immediately prior to assay initiation. The *P. neoaphidis* isolate has been deposited in the USDA, Agricultural Research Service's Collection of Entomopathogenic Fungal Cultures (ARSEF 11663).

Fungal cultures were inverted over each soybean aphid replicate. After 2 h, the fungal cultures and coverslip were removed from each replicate, and the dishes were sealed with parafilm to maintain the humidity required for fungal disease initiation. Each cover slip was stained with aceto-orcein stain, and examined at 200× magnification. Spores had been deposited on

all, indicating that all replicates were exposed to fungal conidia. We then counted spores in 10 randomly chosen fields of view per replicate, and calculated mean spore number per field as an estimate of fungal exposure.

We examined the aphids once per day over the next 5 days. Dead or apparently infected aphids were removed from the experimental dish and transferred to a 50 mm tissue culture dish containing 1% water agar to induce sporulation. If sporulation occurred, the aphid was considered to be infected. We confirmed fungal species identity for two aphids exhibiting successful sporulation on each of the 5 days that assays were monitored. Conidia were stained with aceto-orcein stain and species identity was confirmed via spore morphology at 200× magnification [66].

We calculated the proportion of aphids infected per replicate, and used Pearson's correlation coefficient to determine whether this value was significantly associated with fungal exposure per replicate. We observed substantial variation in both variables, but they were not strongly correlated ($R=0.067$, $P=0.72$), so we proceeded to compare fungal infection between treatments without including fungal exposure as a covariate. We arcsine

square-root transformed the proportion of aphids infected by *P. neophidis*, and performed t-tests (IBM SPSS v20) to determine whether this proportion differed as a function of *Arsenophonus* presence/absence in either aphid isolate.

Acknowledgments

We thank J. McCord for laboratory assistance, J. Obrycki and R. Perry for comments on earlier drafts of this manuscript, and R. Bessin for providing soybean seed. This is publication 13-08-029 of the Kentucky Agricultural Experiment Station. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

Author Contributions

Conceived and designed the experiments: JAW KAB JAW. Performed the experiments: JAW KAB. Analyzed the data: JAW KAB JAW. Contributed reagents/materials/analysis tools: JAW KAB KW GEH JAW. Wrote the paper: JAW JAW.

References

1. Hner P (1965) Endosymbiosis of animals with plant microorganisms. Interscience, New York.
2. Glas AE (1998) Nutritional interactions in insect-microbial symbioses: Aphids and their symbiotic bacteria *Buchnera*. *Ann Rev Entomol* 43: 17–37.
3. Dstrom JP, Russell JA, White JP, Moran NA (2001) Independent origins and horizontal transfer of bacterial symbionts of aphids. *Mol Ecol* 10: 217–228.
4. Duron O, Bouchon D, Boutin S, Bellamy L, Zhou LQ, et al. (2008) The diversity of reproductive parasites among arthropods: Wolbachia do not walk alone. *BMC Biol* 6: 12.
5. Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH (2008) How many species are infected with *Wolbachia*? - a statistical analysis of current data. *FEMS Microbiol Lett* 281: 215–220.
6. Akman L, Yamashita A, Watanabe H, Oshima K, Shiba T, et al. (2002) Genome sequence of the endocellular obligate symbiont of tsetse flies, *Wigglesworthia glossinidia*. *Nat Genet* 32: 402–407.
7. Oliver KM, Degnan PH, Burke GR, Moran NA (2010) Facultative symbionts in aphids and the horizontal transfer of ecologically important traits. *Annu Rev Entomol* 55: 247–266.
8. Russell JA, Moran NA (2006) Costs and benefits of symbiont infection in aphids: variation among symbionts and across temperatures. *Proc Biol Sci* 273: 603–610.
9. Tsuchida T, Koga R, Horikawa M, Tsunoda T, Maoka T, et al. (2010) Symbiotic bacterium modifies aphid body color. *Science* 330: 1102–1104.
10. Ferrari J, Scarborough CL, Godfray HCJ (2007) Genetic variation in the effect of a facultative symbiont on host-plant use by pea aphids. *Oecologia* 153: 323–329.
11. Oliver KM, Russell JA, Moran NA, Hunter MS (2003) Facultative bacterial symbionts in aphids confer resistance to parasitic wasps. *Proc Natl Acad Sci U S A* 100: 1803–1807.
12. Scarborough CL, Ferrari J, Godfray HCJ (2005) Aphid protected from pathogen by endosymbiont. *Science* 310: 1781–1781.
13. Hedges LM, Brownlie JC, O'Neill SL, Johnson KN (2008) *Wolbachia* and virus protection in insects. *Science* 322: 702–702.
14. Jaenike J, Unckless R, Cockburn SN, Boelio LM, Perlman SJ (2010) Adaptation via symbiosis: recent spread of a *Drosophila* defensive symbiont. *Science* 329: 212–215.
15. Novakova E, Hypsa V, Moran NA (2009) *Arsenophonus*, an emerging clade of intracellular symbionts with a broad host distribution. *BMC Microbiol* 9: 143.
16. Huger AM, Skinner SW, Werren JH (1985) Bacterial infections associated with the son-killer trait in the parasitoid wasp, *Nasonia* (= *Mormoniella*) *vitripennis* (Hymenoptera, Pteromalidae). *J Invertebr Pathol* 46:272–280.
17. Werren JH, Skinner SW, Huger AM (1986) Male-killing bacteria in a parasitoid wasp. *Science* 231: 990–992.
18. Gherna RL, Werren JH, Weisburg W, Cote R, Woese CR, et al. (1991) *Arsenophonus nasoniae* gen-nov, sp-nov, the causative agent of the son-killer trait in the parasitoid wasp *Nasonia vitripennis*. *Int J Syst Bacteriol* 41: 563–565.
19. Duron O, Wilkes TE, Hurst GD (2010) Interspecific transmission of a male-killing bacterium on an ecological timescale. *Ecol Lett* 13: 1139–1148.
20. Hypsa V (1993) Endocytobionts of *Triatoma infestans*- distribution and transmission. *J Invertebr Pathol* 61: 32–38.
21. Trowbridge RE, Dittmar K, Whiting MF (2006) Identification and phylogenetic analysis of *Arsenophonus*- and *Photorhabdus*-type bacteria from adult Hippoboscidae and Streblidae (Hippoboscoidea). *J Invertebr Pathol* 91: 64–68.
22. Perotti MA, Allen JM, Reed DL, Braig HR (2007) Host-symbiont interactions of the primary endosymbiont of human head and body lice. *FASEB J* 21: 1058–1066.
23. Zreik L, Bove JM, Garnier M (1998) Phylogenetic characterization of the bacterium-like organism associated with marginal chlorosis of strawberry and proposition of a Candidatus taxon for the organism, 'Candidatus *Phlomobacter fragariae*'. *Int J Syst Bacteriol* 48: 257–261.
24. Bressan A, Semetey O, Arceido J, Lherminier J, Boudon-Padieu E (2009) Vector transmission of a plant-pathogenic bacterium in the *Arsenophonus* clade sharing ecological traits with facultative insect endosymbionts. *Phytopathology* 99: 1289–1296.
25. Bressan A, Terlizzi F, Credi R (2012) Independent origins of vectored plant pathogenic bacteria from arthropod-associated *Arsenophonus* endosymbionts. *Microb Ecol* 63: 628–638.
26. Thao MLL, Baumann P (2004) Evidence for multiple acquisition of *Arsenophonus* by whitefly species (Sternorrhyncha: Aleyrodidae). *Curr Microbiol* 48: 140–144.
27. Subandiyah S, Nikoh N, Tsuyumu S, Somowiyarjo S, Fukatsu T (2000) Complex endosymbiotic microbiota of the citrus psyllid *Diuraphis citri* (Homoptera: Psylloidea). *Zoolog Sci* 17: 983–989.
28. Thao ML, Clark MA, Baumann L, Brennan EB, Moran NA, et al. (2000) Secondary endosymbionts of psyllids have been acquired multiple times. *Curr Microbiol* 41: 300–304.
29. Russell JA, Latorre A, Sabater-Munoz B, Moya A, Moran NA (2003) Side-stepping secondary symbionts: widespread horizontal transfer across and beyond the Aphidoidea. *Mol Ecol* 12: 1061–1075.
30. Jouselin E, Coeur d'acier A, Vanlerberghe-Masutti F, Duron O (2013). Evolution and diversity of *Arsenophonus* endosymbionts in aphids. *Mol Ecol* 22: 260–270.
31. Hansen AK, Jeong G, Paine TD, Stouthamer R (2007) Frequency of secondary symbiont infection in an invasive psyllid relates to parasitism pressure on a geographic scale in California. *Appl Environ Microbiol* 73: 7531–7535.
32. Clay K, Holah J, Rudgers JA (2005) Herbivores cause a rapid increase in hereditary symbiosis and alter plant community composition. *Proc Natl Acad Sci U S A* 102: 12465–12470.
33. Oliver KM, Campos J, Moran NA, Hunter MS (2008) Population dynamics of defensive symbionts in aphids. *Proc Biol Sci* 275: 293–299.
34. Heimpel GE and Lundgren JG (2000) Sex ratios of commercially reared biological control agents. *Biol. Control* 19:77–93.
35. Carletto J, Gueguen G, Fleury F, Vanlerberghe-Masutti F (2008) Screening the bacterial endosymbiotic community of sap-feeding insects by terminal-restriction fragment length polymorphism analysis. *Entomol Exp Appl* 129: 228–234.
36. Wille BD, Hartman GL (2009) Two species of symbiotic bacteria present in the soybean aphid (Hemiptera: Aphididae). *Environ Entomol* 38: 110–115.
37. Ragsdale DW, Landis DA, Brodeur J, Heimpel GE, and Desneux N (2011) Ecology and management of soybean aphid in North America. *Annu Rev Entomol* 56:375–399.
38. Costamagna AC, Landis DA, Brewer MJ (2008) The role of natural enemy guilds in *Aphis glycines* suppression. *Biol Control* 45: 368–379.
39. Kaiser ME, Noma T, Brewer MJ, Pike KS, Vockeroth JR, et al. (2007) Hymenopteran parasitoids and dipteran predators found using soybean aphid after its midwestern United States invasion. *Ann Entomol Soc Am* 100: 196–205.
40. Noma T, Brewer MJ (2008) Seasonal abundance of resident parasitoids and predatory flies and corresponding soybean aphid densities, with comments on

- classical biological control of soybean aphid in the Midwest. *J Econ Entomol* 101: 278–287.
41. Heimpel GE, Ragsdale DW, Venette R, Hopper KR, O'Neil RJ, et al. (2004) Prospects for importation biological control of the soybean aphid: Anticipating potential costs and benefits. *Ann Entomol Soc Am* 97: 249–258.
 42. Wychuys KAG, Koch RL, Kula RR, Heimpel GE (2009) Potential exposure of a classical biological control agent of the soybean aphid, *Aphis glycines*, on non-target aphids in North America. *Biol Invasions* 11: 857–871.
 43. Nielsen C, Hajek AE (2005) Control of invasive soybean aphid, *Aphis glycines* (Hemiptera: Aphididae), populations by existing natural enemies in New York State, with emphasis on entomopathogenic fungi. *Environ Entomol* 34: 1036–1047.
 44. Wilkes T, Duron O, Darby AC, Hypsa V, Novakova E, Hurst GDD (2011) The genus *Arsenophonus*. In: Bourtzis K, Zchori-Fein E, editors. *Manipulative Tenants*. Boca Raton, Florida: CRC Press. 225–244.
 45. Heimpel GE, Frelich LE, Landis DA, Hopper KR, Hoelmer KA, et al. (2010) European buckthorn and Asian soybean aphid as components of an extensive invasional meltdown in North America. *Biol Invasions* 12: 2913–2931.
 46. Vorburger C, Gehrler L, Rodriguez PA (2009) A strain of the bacterial symbiont *Regiella insecticola* protects aphids against parasitoids. *Biol Lett* 6: 109–111.
 47. Lukasik P, van Asch M, Guo HF, Ferrari J, Godfray HCJ (2013) Unrelated facultative endosymbionts protect aphids against a fungal pathogen. *Ecol Lett* 16: 214–218.
 48. Koch KA, Ragsdale DW (2011) Impacts of thiamethoxam seed treatment and host plant resistance on the soybean aphid fungal pathogen, *Pandora neoaphidis*. *J Econ Entomol* 104: 1824–1832.
 49. Oliver KM, Degnan PH, Hunter MS, Moran NA (2009) Bacteriophages encode factors required for protection in a symbiotic mutualism. *Science* 325: 992–994.
 50. Michel AP, Zhang W, Jung JK, Kang ST, Mian MAR (2009) Population genetic structure of *Aphis glycines*. *Environ Entomol* 38: 1301–1311.
 51. Desneux N, Barta RJ, Hoelmer KA, Hopper KR, and Heimpel GE (2009) Multifaceted determinants of host specificity in an aphid parasitoid. *Oecologia* 160:387–398.
 52. Vorburger C, Sandrock C, Gousskov A, Castaneda LE, Ferrari J (2009) Genotypic variation and the role of defensive endosymbionts in an all-parthenogenetic host-parasitoid interaction. *Evolution* 63: 1439–1450.
 53. Engelstadter J, Hurst GDD (2009) The ecology and evolution of microbes that manipulate host reproduction. *Annu Rev Ecol Evol Syst* 40: 127–149.
 54. Simon JC, Boutin S, Tsuchida T, Koga R, Le Gallie JF, et al. (2011) Facultative symbiont infections affect aphid reproduction. *PLoS One* 6: e21831.
 55. Himler AG, Adachi-Hagimori T, Bergen JE, Kozuch A, Kelly SE, et al. (2011) Rapid spread of a bacterial symbiont in an invasive whitefly is driven by fitness benefits and female bias. *Science* 332: 254–256.
 56. Moran NA, Dunbar HE (2006) Sexual acquisition of beneficial symbionts in aphids. *Proc Natl Acad Sci U S A* 103: 12803–12806.
 57. Caspi-Fluger A, Inbar M, Mozes-Daube N, Katzir N, Portnoy V, et al. (2012) Horizontal transmission of the insect symbiont *Rickettsia* is plant-mediated. *Proc Biol Sci* 279: 1791–1796.
 58. Hoffmann AA, Hercus M, Dagher H (1998) Population dynamics of the *Wolbachia* infection causing cytoplasmic incompatibility in *Drosophila melanogaster*. *Genetics* 148: 221–231.
 59. Brownlie JC, Cass BN, Riegler M, Witsenburg JJ, Iturbe-Ormaetxe I, et al. (2009) Evidence for metabolic provisioning by a common invertebrate endosymbiont, *Wolbachia pipiensis*, during periods of nutritional stress. *PLoS Pathog* 5: e1000368.
 60. Dale C, Becton M, Harbison C, Jones T, Pontes M, (2006) Isolation, pure culture, and characterization of "Candidatus *Arsenophonus arthropodicus*," an intracellular secondary endosymbiont from the hippoboscids louse fly *Pseudolynchia canariensis*. *Appl Environ Microbiol* 72: 2997–3004.
 61. Russell JA, Weldon S, Smith AH, Kim KL, Hu Y, et al. (2013) Uncovering symbiont-driven genetic diversity across North American pea aphids. *Mol Ecol* 22: 2045–59.
 62. Ruan YM, Xu J, Liu SS (2006) Effects of antibiotics on fitness of the B biotype and a non-B biotype of the whitefly *Bemisia tabaci*. *Entomol Exp Appl* 121: 159–166.
 63. Griffiths GW, Beck SD (1974) Effects of antibiotics on intracellular symbionts in the pea aphid, *Acyrtosiphon pisum*. *Cell Tissue Res* 148: 287–300.
 64. Wychuys KAG, Stone L, Desneux N, Hoelmer KA, Hopper KR, and Heimpel GE (2008) Parasitism of the soybean aphid, *Aphis glycines*, by *Binodoxys communis*: the role of aphid defensive behaviour and parasitoid reproductive performance. *Bull Entomol Res* 98:361–370.
 65. Lin LA, Ives AR (2003) The effect of parasitoid host-size preference on host population growth rates: an example of *Aphidius colemani* and *Aphis glycines*. *Ecol Entomol* 28: 542–550.
 66. Samson RA, Evans HC, Latgé JP (1988) Atlas of entomopathogenic fungi. The Netherlands: Springer-Verlag.