



University of Kentucky
UKnowledge

Internal Medicine Presentations

Internal Medicine

2-2-2011

CMR of LV non-compaction cardiomyopathy: association of clinical presentation and prognosis with cardiac phenotype

Mushabbar A. Syed
Loyola University

Steve W. Leung
University of Kentucky, steve.leung@uky.edu

Samy Claude Elayi
University of Kentucky

Richard J. Charnigo
University of Kentucky

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/internalmedicine_present

 Part of the [Medicine and Health Sciences Commons](#)

Repository Citation

Syed, Mushabbar A.; Leung, Steve W.; Elayi, Samy Claude; and Charnigo, Richard J., "CMR of LV non-compaction cardiomyopathy: association of clinical presentation and prognosis with cardiac phenotype" (2011). *Internal Medicine Presentations*. 4.
https://uknowledge.uky.edu/internalmedicine_present/4

This Presentation is brought to you for free and open access by the Internal Medicine at UKnowledge. It has been accepted for inclusion in Internal Medicine Presentations by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

POSTER PRESENTATION

Open Access

CMR of LV non-compaction cardiomyopathy: association of clinical presentation and prognosis with cardiac phenotype

Mushabbar A Syed^{1*}, Steve W Leung², Samy Claude Elayi³, Richard J Charnigo Jr.³

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Background

Left ventricular non-compaction (LVNC) is a rare congenital disorder characterized by two layered myocardium; trabeculated (non-compacted) and a non-trabeculated (compacted). LVNC is increasingly being recognized due to better imaging technology as a cause for heart failure and sudden cardiac death; however, data on clinical and imaging characteristics remains limited.

Objective

To investigate the association of clinical presentation and outcomes in LVNC with cardiac phenotype by CMR.

Methods

Fourteen patients (mean age 33.1 ± 17.6 years, 9 male) were retrospectively identified from CMR database between December 2007 and May 2010. CMR imaging included SSFP cine in standard views and late gadolinium enhancement. Quantitative analysis included left and right ventricular function, volumes, mass, LV wall motion score and non-compacted to compacted myocardium (NC/C) ratios in different segments. Number of involved LV segments and regions of maximum NC/C ratio were also recorded.

Patient's medical records were reviewed for clinical history including NYHA functional class, ECG, telemetry, Holter/event monitoring and electrophysiology studies.

Non-parametric U test, logistic regression analysis and parametric T-test were used to determine statistical significance as appropriate.

Results

Seven patients presented with acute heart failure including one in cardiogenic shock. Three patients presented with syncope, one with documented ventricular tachycardia (VT).

Mean LVEF was $36.2 \pm 22.8\%$ and mean RVEF $31.5 \pm 16.7\%$. LVEF $<50\%$ was present in 8 patients (57.1%), RVEF $<40\%$ in 7 (50%) and both in 6 (42.8%) patients. Mean NC/C myocardium ratio was 3.7 ± 0.8 with mean of 5.5 ± 3.1 LV segments involved. Patients with LV dysfunction were older, more symptomatic with higher NYHA class, had more myocardial segments involvement with non-compaction, and higher NC/C ratios (Table 1). No myocardial infarction or mid-wall fibrosis was seen on late gadolinium enhancement. One patient had thrombus in the right ventricle associated with severe RV dysfunction.

Four patients had non-sustained monomorphic VT. Two patients had premature ventricular complexes on

Table 1

	LVEF $>50\%$ (N=6)	LVEF $\leq 50\%$ (N=8)	p-value
Age	21.6 ± 11.0	42.5 ± 16.6	0.008
NYHA class	1 (IQR 1-1)	3 (IQR 2-4)	0.007
LV end-diastolic volume index (ml/m ²)	96.1 ± 15.3	155.4 ± 36.6	0.001
LV end-systolic volume index (ml/m ²)	38.6 ± 7.5	129.6 ± 38.2	<0.001
LV mass index (g/m ²)	55.4 ± 8.5	80.2 ± 17.3	0.004
Wall motion score index	1.0 ± 0.02	2.3 ± 0.3	<0.001
Non-compacted segments	3.8 ± 2.1	6.8 ± 3.2	<0.001
NC/C ratio (maximum)	3.2 ± 0.6	4.1 ± 0.8	<0.001
RV EF (%)	44.5 ± 6.8	21.8 ± 15.3	0.002
Heart failure	0 (0%)	7 (88%)	0.005

¹Loyola University, Stritch School of Medicine, Maywood, IL, USA
Full list of author information is available at the end of the article

telemetry. One patient had paroxysmal atrial fibrillation, and one had atrioventricular nodal reentry tachycardia (AVNRT).

There were no deaths over a mean follow-up of 7.0 ± 4.6 months. One patient received a heart transplant for severe refractory heart failure. Five patients received an ICD; 4 with non-sustained VT and 1 with severe LV dysfunction. Patient with AVNRT underwent successful ablation.

Conclusion

Patients with LVNC have a spectrum of cardiac phenotypes ranging from normal LV and RV to severe biventricular dysfunction. Clinical presentation and symptoms are associated with degree of non-compaction and ventricular dysfunction.

Author details

¹Loyola University, Stritch School of Medicine, Maywood, IL, USA. ²National Institutes of Health, Bethesda, MD, USA. ³University of Kentucky, Lexington, KY, USA.

Published: 2 February 2011

doi:10.1186/1532-429X-13-S1-P291

Cite this article as: Syed *et al.*: CMR of LV non-compaction cardiomyopathy: association of clinical presentation and prognosis with cardiac phenotype. *Journal of Cardiovascular Magnetic Resonance* 2011 **13** (Suppl 1):P291.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

