



University of Kentucky
UKnowledge

Chemistry Faculty Publications

Chemistry

10-19-2014


Interactive Comment on “Iodine Oxide in the Global Marine Boundary Layer” by C. Prados-Roman et al.

Marcelo I. Guzman

University of Kentucky, marcelo.guzman@uky.edu

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/chemistry_facpub

 Part of the [Atmospheric Sciences Commons](#), [Environmental Chemistry Commons](#), [Environmental Engineering Commons](#), and the [Environmental Sciences Commons](#)

Repository Citation

Guzman, Marcelo I., "Interactive Comment on “Iodine Oxide in the Global Marine Boundary Layer” by C. Prados-Roman et al." (2014). *Chemistry Faculty Publications*. 44.

https://uknowledge.uky.edu/chemistry_facpub/44

This Response or Comment is brought to you for free and open access by the Chemistry at UKnowledge. It has been accepted for inclusion in Chemistry Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Interactive Comment on “Iodine Oxide in the Global Marine Boundary Layer” by C. Prados-Roman et al.

Notes/Citation Information

Published in *Atmospheric Chemistry and Physics*, v. 14, issue 15, p. C8250-C8251.

© Author(s) 2014.

This work is distributed under the [Creative Commons Attribute 3.0 License](#).

Interactive comment on “Iodine oxide in the global marine boundary layer” by C. Prados-Roman et al.

M.I. Guzman

marcelo.guzman@uky.edu

Received and published: 19 October 2014

The manuscript “Iodine oxide in the global marine boundary layer” by Prados-Roman et al. presents multi-axis DOAS measurements of IO radical mixing ratios (< 1 pptv, altitude ≤ 600 m) performed over the marine boundary layer (MBL). A combined analysis with other field data suggests that iodine driven chemistry is of global importance over the oceans. A 3D CAM-Chem model discerning the contribution of organic and inorganic emissions (specifically hypiodous acid and molecular iodine), and their associated geographical dependence, estimates that 75% of the total iodine oxide budget is of abiotic origin in the global MBL. This manuscript is an important contribution to understand the oxidizing capacity of the atmosphere and presents new data to support an abiotic mechanism is operative over open ocean waters. However, it would be important to consider in this manuscript a recent laboratory study by Pillar et al. (En-

C8250

viron. Sci. Technol., 2013, 47, 10971–10979, <http://dx.doi.org/10.1021/es401700h>) that indicates how sea spray aerosol production and in-situ oxidation produces hypiodous acid and molecular iodine. Guzman et al. (J. Phys. Chem. A, 2012, 116, 5428–5435, <http://dx.doi.org/10.1021/jp3011316>) studied the enrichment of halides during aerosolization of seawater mimic samples providing new insights about how concentration effects could be included in a model. More importantly, it would be interesting to discuss in the final version of the manuscript to be published in ACP how reactions at the air–water interface of sea spray, followed by transfer of reactive products to the gas-phase (Environ. Sci. Technol., 2013, 47, 10971–10979, <http://dx.doi.org/10.1021/es401700h>) contributes to the model presented. In addition, it would be important to connect the manuscript with Pillar et al. previously proposal indicating that 1) the actual source of reactive iodine species will vary geographically, 2) the production of sea spray will be sensitive to local conditions, particularly surface winds, 3) the production of iodine will depend on factors such as temperature, humidity, and the concentration of halogen species, and 4) 3D models should be chosen over 1D models to approach this problem.

References Guzman, M. I.; Athalye, R. R.; and Rodriguez, J. M.: Concentration effects and ion properties controlling the fractionation of halides during aerosol formation, J. Phys. Chem. A, 116, 5428–5435, doi:10.1021/jp3011316. Pillar, E. A.; Guzman, M. I.; and Rodriguez, J. M.: Conversion of Iodide to Hypiodous Acid and Iodine in Aqueous Microdroplets Exposed to Ozone, Environ. Sci. Technol., 47, 10971–10979, doi:10.1021/es401700h, 2013. Prados-Roman, C.; Cuevas, C. A.; Hay, T.; Fernandez, R. P.; Mahajan, A. S.; Royer, S.-J.; Galí, M.; Simó, R.; Dachs, J.; Großmann, K.; Kinnison, D. E.; Lamarque, J.-F.; and Saiz-Lopez, A.: Iodine oxide in the global marine boundary layer, Atmos. Chem. Phys. Discuss., 14, 22217–22243, doi:10.5194/acpd-14-22217-2014, 2014.

Interactive comment on Atmos. Chem. Phys. Discuss., 14, 22217, 2014.

C8251