



University of Kentucky
UKnowledge

Saha Cardiovascular Research Center Faculty
Publications

Cardiovascular Research

4-2014

Bisphenol A Increases Atherosclerosis in Pregnane X Receptor-Humanized ApoE Deficient Mice

Yipeng Sui

University of Kentucky, yipeng.sui@uky.edu

Se-Hyung Park

University of Kentucky, sh.park@uky.edu

Robert N. Helsley

University of Kentucky, robert.helsley@uky.edu

Manjula Sunkara

University of Kentucky, manjula.sunkara@uky.edu

Frank J. Gonzalez

National Cancer Institute

See next page for additional authors

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/cvrc_facpub

 Part of the [Cardiology Commons](#), and the [Circulatory and Respiratory Physiology Commons](#)

Repository Citation

Sui, Yipeng; Park, Se-Hyung; Helsley, Robert N.; Sunkara, Manjula; Gonzalez, Frank J.; Morris, Andrew J.; and Zhou, Changcheng, "Bisphenol A Increases Atherosclerosis in Pregnane X Receptor-Humanized ApoE Deficient Mice" (2014). *Saha Cardiovascular Research Center Faculty Publications*. 22.

https://uknowledge.uky.edu/cvrc_facpub/22

This Article is brought to you for free and open access by the Cardiovascular Research at UKnowledge. It has been accepted for inclusion in Saha Cardiovascular Research Center Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Authors

Yipeng Sui, Se-Hyung Park, Robert N. Helsley, Manjula Sunkara, Frank J. Gonzalez, Andrew J. Morris, and Changcheng Zhou

Bisphenol A Increases Atherosclerosis in Pregnane X Receptor-Humanized ApoE Deficient Mice**Notes/Citation Information**

Published in *Journal of the American Heart Association*, v. 3, no. 2, article e000492, p. 1-11.

© 2014 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley Blackwell.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial License](#), which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Digital Object Identifier (DOI)

<http://dx.doi.org/10.1161/JAHA.113.000492>

Bisphenol A Increases Atherosclerosis in Pregnane X Receptor-Humanized ApoE Deficient Mice

Yipeng Sui, PhD; Se-Hyung Park, PhD; Robert N. Helsley, BS; Manjula Sunkara, MS; Frank J. Gonzalez, PhD; Andrew J. Morris, PhD; Changcheng Zhou, PhD

Background—Bisphenol A (BPA) is a base chemical used extensively in many consumer products. BPA has recently been associated with increased risk of cardiovascular disease (CVD) in multiple large-scale human population studies, but the underlying mechanisms remain elusive. We previously reported that BPA activates the pregnane X receptor (PXR), which acts as a xenobiotic sensor to regulate xenobiotic metabolism and has pro-atherogenic effects in animal models upon activation. Interestingly, BPA is a potent agonist of human PXR but does not activate mouse or rat PXR signaling, which confounds the use of rodent models to evaluate mechanisms of BPA-mediated CVD risk. This study aimed to investigate the atherogenic mechanism of BPA using a PXR-humanized mouse model.

Methods and Results—A PXR-humanized ApoE deficient (huPXR•ApoE^{-/-}) mouse line was generated that respond to human PXR ligands and feeding studies were performed to determine the effects of BPA exposure on atherosclerosis development. Exposure to BPA significantly increased atherosclerotic lesion area in the aortic root and brachiocephalic artery of huPXR•ApoE^{-/-} mice by 104% ($P<0.001$) and 120% ($P<0.05$), respectively. By contrast, BPA did not affect atherosclerosis development in the control littermates without human PXR. BPA exposure did not affect plasma lipid levels but increased CD36 expression and lipid accumulation in macrophages of huPXR•ApoE^{-/-} mice.

Conclusion—These findings identify a molecular mechanism that could link BPA exposure to increased risk of CVD in exposed individuals. PXR is therefore a relevant target for future risk assessment of BPA and related environmental chemicals in humans. (*J Am Heart Assoc.* 2014;3:e000492 doi: 10.1161/JAHA.113.000492)

Key Words: atherosclerosis • cells • receptors • risk factors

Bisphenol A (BPA), a base chemical used extensively in polycarbonate plastics in many consumer products, is among the world's highest production-volume chemicals, with more than 8 billion pounds produced each year.¹ More than 80 biomonitoring studies indicate that human exposure to BPA is ubiquitous, and over 95% of the U.S. population is exposed to BPA.^{1,2} BPA has been detected in human blood,

urine, tissues and other fluids,¹ and numerous animal studies show that exposure to BPA causes diverse adverse effects.^{3,4} Despite strong evidence for BPA's adverse effects in animals and, by extrapolation, in humans, recent evaluations of BPA safety by multiple panels have arrived at disparate conclusions and thus controversy remains about the specificity and mechanisms of the potential adverse effects of BPA.^{5,6}

Recent large and well-conducted cross-sectional and longitudinal studies have found that higher BPA exposure is consistently associated with increased risk of cardiovascular disease (CVD).⁷⁻⁹ Lang et al⁷ first reported positive associations between urinary BPA concentrations and the CVD, type 2 diabetes, and liver enzyme abnormalities using data from the National Health and Nutrition Examination Survey (NHANES) 2003-2004. Higher urinary BPA levels were significantly associated with increased diagnosis of CVD including coronary heart disease, myocardial infarction, and angina.⁷ Melzer et al⁸ replicated the early association between urinary BPA concentrations and coronary heart disease using a separate NHANES 2005-2006 database. A separate large-scale longitudinal study has confirmed associations between

From the Graduate Center for Nutritional Sciences (Y.S., S.-H.P., R.N.H., C.Z.), Saha Cardiovascular Research Center (A.J.M., C.Z.), Division of Cardiovascular Medicine (M.S., A.J.M.), University of Kentucky, Lexington, KY; Laboratory of Metabolism, Center for Cancer Research, National Cancer Institute, Bethesda, MD (F.J.G.).

Correspondence to: Changcheng Zhou, PhD, Graduate Center for Nutritional Sciences, Saha Cardiovascular Research Center, University of Kentucky, 900 S. Limestone Street, 517 Wethington Building, Lexington, KY 40536. E-mail: c.zhou@uky.edu

Received February 28, 2014; accepted March 13, 2014.

© 2014 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley Blackwell. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

higher BPA exposure levels and incident coronary artery disease during >10 years of follow-up of a group of healthy people who took part in the European Prospective Investigation of Cancer-Norfolk UK in the 1990s.⁹ Interestingly, exclusion of subjects with obesity and adjustment for blood lipid concentrations or levels of physical activity had little effect on the associations,^{7,9} suggesting that the associations are independent of traditional CVD risk factors. More recently, independent studies have associated BPA exposure with coronary atherosclerosis,¹⁰ carotid atherosclerosis,¹¹ and peripheral arterial disease,¹² indicating potential effects of BPA exposure on atherosclerosis, the most common cause of CVD. However, the underlying mechanisms responsible for these associations remain unclear, which continues to hamper rational assessment of the health risks of BPA exposure.

BPA, regarded as a xenoestrogen, is a weak agonist of the estrogen receptor (ER), and most health studies of BPA have focused on its estrogenic effects.^{4,13} However, the estrogenic effects of BPA probably do not explain the link between BPA exposure and CVD, as animal and human studies identify protective effects of estrogen against atherosclerosis or CVD.^{14–18} To date, BPA has not been reported to have atherogenic effects in any animal models. Further, despite compelling evidence about BPA's estrogenic activity, doubts remain whether BPA exerts adverse estrogenic effects in animals and humans.^{19–21} Thus, the endocrine-disrupting effects of BPA cannot be entirely attributed to its estrogenic activity and more mechanistic studies are urgently needed to explore the effect of BPA on other signaling pathways.

We previously reported that BPA and its analogs activate another nuclear receptor, the pregnane X receptor (PXR; also known as steroid and xenobiotic receptor, or SXR).²² PXR functions as a xenobiotic sensor that regulates genes involved in drug and xenobiotic transport and metabolism, including cytochromes P450 (CYP), conjugating enzymes (eg, glutathione transferase (GST)), and ABC family transporters (eg, multidrug resistance 1 (MDR1)).^{23,24} PXR is activated by endogenous hormones, dietary steroids, pharmaceutical agents, and other xenobiotic chemicals.^{24–26} In mammals, PXR also exhibits considerable differences in its pharmacology and its ligand-binding domain (LBD) is remarkably divergent across species.^{24,25} Interestingly, we found that BPA is a potent agonist for human PXR (hPXR) but not for mouse or rat PXR (mPXR and rPXR, respectively),²² underscoring the importance of species choice in predicting the human risk assessment of BPA.

We recently revealed the pro-atherogenic effects of PXR in animal models and found that chronic activation of PXR increases atherosclerosis in ApoE-deficient (ApoE^{-/-}) mice.²⁷ These observations suggest that BPA-mediated PXR activation could potentially accelerate atherosclerosis development and increase CVD risk in humans. Because BPA is a potent agonist

of human but not mouse or rat PXR,²² the choice of animal model is a paramount issue in conducting preclinical studies to evaluate the contribution of BPA exposure to CVD risk. To investigate the effects of BPA exposure on atherosclerosis development, a PXR-humanized ApoE-deficient (huPXR•ApoE^{-/-}) mouse line was generated. Here we report that BPA increases atherosclerosis in ApoE^{-/-} mice in a human PXR-dependent manner. BPA exposure does not affect plasma lipid levels but increases lipid accumulation and foam cell formation in macrophages of huPXR•ApoE^{-/-} mice.

Methods

Animals and Diets

ApoE^{-/-} mice on the C57BL/6 background were purchased from The Jackson Laboratory. PXR-humanized mice (huPXR, mouse PXR knockout/human PXR transgenic) were generated by transgenesis on a *Pxr*-null mice using a BAC clone containing the complete human PXR gene and including 5'- and 3'-flanking sequences as previously described.²⁸ huPXR mice have similar tissue distribution of PXR expression in liver and intestine as native PXR in humans and mice.²⁸ huPXR mice were backcrossed with C57BL/6 wild-type (WT) mice for at least 4 generations at NCI,²⁸ and backcrossed for 4 additional generations onto the C57BL/6 background and then bred with ApoE^{-/-} mice to generate huPXR•ApoE^{-/-} (hPXRtg-PXR^{-/-}•ApoE^{-/-}) and PXR^{-/-}•ApoE^{-/-} mice. All the mice used in this study have the same background (PXR and ApoE null alleles) except for one allele of huPXR•ApoE^{-/-} mice carrying the human PXR gene. All the animals were housed in a specific pathogen-free room with a 12-hour light-dark cycle in the University of Kentucky Division of Laboratory Animal Resources under a protocol approved by the Institutional Animal Care and Use Committee. BPA, rifampicin (RIF), and pregnenolone 16 α -carbonitrile (PCN) were purchased from Sigma-Aldrich. BPA was incorporated into a modified semisynthetic diet containing 4.2% fat and 0.02% cholesterol^{29,30} at a dose of 50 mg/kg by Harland Laboratories, Inc.^{31–33} Four-week-old experimental male huPXR•ApoE^{-/-} and PXR^{-/-}•ApoE^{-/-} littermates were weaned and fed with a control diet or a diet supplemented with BPA for 12 weeks until euthanization at 16 weeks of age (15 to 20 mice per group). Five to 10 mice were used for primary cell isolation and tissue analysis, and the rest were used for atherosclerosis analysis. The number of mice used in each study is listed in figure legends or shown by scatter points.

Blood Analysis and Atherosclerotic Lesion Quantification

Plasma total cholesterol and triglyceride concentrations were determined enzymatically by a colorimetric method.²⁹ Plasma

from multiple mice (n=6) was pooled and plasma lipoprotein cholesterol distribution was determined by fast-performance liquid chromatography (FPLC).³⁴ OCT-embedded hearts or brachiocephalic arteries were sectioned and stained with Oil-red-O, and atherosclerotic lesions were quantified as previously described.^{29,34} Immunohistochemistry were performed on sections of aortic roots with specific antibodies against PXR, monocyte/macrophage marker MOMA-2, or CD36 as previously described.^{29,34}

Analysis of Urinary BPA by LC-ESI-MS/MS

Analysis of urinary BPA was performed using a modified chemical derivatization liquid chromatography-electrospray ionization-tandem mass spectrometry (LC-ESI-MS/MS) method.³⁵ Labeled BPA-d₁₆ internal standard (Sigma-Aldrich) was added to each urine sample (50 μ L) prior to extraction. An amount of 500 μ L of cold acetonitrile was added to the samples and protein precipitation was collected by centrifugations. Supernatants were dried in 4 mL vials under N₂. An amount of 100 μ L of sodium bicarbonate buffer (0.1 mol/L, pH 10) was added to the vials, followed by 100 μ L of 1 mg/mL solution of pyridine-3-sulfonyl chloride (PSC) in acetone. Vials were vortexed and placed in a heater block at 70°C for 5 to 10 minutes for preparation of PSC derivatives. Reaction mixtures were then cooled on ice for 10 minutes and dried under N₂. The samples were then reconstituted with 100 μ L of methanol and transferred to autosampler vials for LC-ESI-MS/MS analysis. PSC derivatives of BPA and BPA-glucuronide samples were detected and quantitated by reverse phase HPLC using a Waters XTerra MS C8 column. The mobile phase consisted of 20% methanol with 1 mmol/L ammonium formate as solvent A and 100% methanol as solvent B. Analysis of BPA was achieved from 0% to 70% solvent B for 1 minute, which was gradually increased to 80% over 3 minutes and then to 90% over the next 4 minutes and maintained at 90% for the last 2 minutes. The column was equilibrated to initial conditions in 3 minutes. The flow rate was 0.5 mL/min with a column temperature of 30°C. The sample injection volume was 10 μ L. The mass spectrometer was operated in positive electrospray ionization mode with a declustering potential of 51 V, entrance potential of 10 V, collision energy of 37 V, collision cell exit potential of 12 V, curtain gas of 10 psi, ion spray voltage of 5500 V, ion source gas1/gas2 of 40 psi and temperature of 550°C. The instrument was operated in selected ion monitoring mode with the following precursor product ion pairs monitored for the indicated analytes: m/z 511.1/354, m/z 511.1/212, m/z 511.1/79 for PSC-BPA and m/z 527.2/223.2, m/z 527.2/367.1, m/z 527.2/79 for PSC-BPA-d₁₆ and m/z 546.2/213.2, m/z 546.2/276.1, m/z 546.2/79.2 for PSC-BPA-glucuronide.

Peritoneal Macrophage Isolation and Staining

Mice were injected intraperitoneally with 1 mL of 3% thioglycollate. Peritoneal macrophages were collected 4 days later and Oil red O/hematoxylin staining was performed as described before.^{27,34} Cells containing lipid droplets (>10) were counted as foam cells and at least 10 fields per condition were counted.³⁶

RNA Isolation and Quantitative Real-Time PCR Analysis

Total RNA was isolated from mouse tissues or cells using TRIzol Reagent (Life Technologies) and quantitative real-time PCR (QPCR) was performed using gene-specific primers and the SYBR green PCR kit (Life Technologies) as described previously.²² The primer sets used in this study are listed in Table 1.

Statistical Analysis

Statistical analysis was performed using a 2-sample, 2-tailed Student's *t* test for comparisons between 2 groups, in which *P*<0.05 was regarded as significant. One-way ANOVA analysis of variance was used when multiple comparisons were made followed by post hoc Bonferroni *t* test. All data were presented by mean \pm SD.

Results

Generation and Characterization of PXR-Humanized ApoE^{-/-} Mice for BPA Risk Assessment

Since BPA is a human PXR-selective ligand, one of the key challenges to study the effects of BPA-mediated PXR activation on atherosclerosis is development of a mouse model that recapitulates the human response to PXR ligands. To address this issue, PXR-humanized ApoE knockout mice (huPXR•ApoE^{-/-}) were generated. The huPXR mice, expressing the human *PXR* gene in place of mouse *Pxr* gene,^{28,37} were crossed with atherosclerosis-prone ApoE^{-/-} mice to generate huPXR•ApoE^{-/-} mice and PXR^{-/-}•ApoE^{-/-} mice (Figure 1A). The huPXR•ApoE^{-/-} and PXR^{-/-}•ApoE^{-/-} mice generated by this strategy have the same genetic background (mPXR and ApoE null alleles) except for one allele of huPXR•ApoE^{-/-} mice carrying the human PXR transgene (Figure 1A). The huPXR•ApoE^{-/-} mice were then treated with the hPXR-specific ligand rifampicin (RIF) or the mPXR-specific ligand pregnenolone 16 α -carbonitrile (PCN).²² As expected, huPXR•ApoE^{-/-} mice can respond to the human-specific PXR agonist RIF but not to the mouse-specific activator PCN

Table 1. Primer Sequences for Genomic PCR and QPCR

| Gene | Primer Sequence | Gene | Primer Sequence |
|---------------------------------|---------------------------------|------------------------------|--------------------------------|
| ApoE | 5'-GCCTAGCCGAGGGAGAGCCG-3' | CD36 | 5'-CAGTCGGAGACATGCT-3' |
| | 5'-TGTGACTTGGGAGCTCTGCAGC-3' | | 5'-CTCGGGGTCTGAGTT-3' |
| | 5'-GCCGCCCGACTGCATCT-3' | | SR-A |
| 5'-CTGGTCATCACTGTTGCTGTACCA-3' | 5'-GTCAATGGAGGCCCA-3' | | |
| 5'-GCAGCATAGACAAGTTATTCTAGAG-3' | SR-BI | 5'-CTCATCAAGCAGCAGGTGCTCA-3' | |
| 5'-CTAAAGCGCATGCTCCAGACTGC-3' | | 5'-GAGGATTCGGGTGTCATGAA-3' | |
| hPXR | 5'-GCACCTGCTGCTAGGAATA-3' | ABCA1 | 5'-CCGAGGAAGACGTGGACACCTTC-3' |
| | 5'-CTCCATTGCCCTCCTAAGT-3' | | 5'-CCTCAGCCATGACCTGCCTTGTAG-3' |
| CYP3A11 | 5'-CAGCTTGGTCTCCTCTACC-3' | ABCG1 | 5'-AGGTCTCAGCCTCTAAAGTTCCTC-3' |
| | 5'-TCAAACAACCCCATGTTTT-3' | | 5'-TCTCTCGAATGAAATTTATCG-3' |
| MDR1a | 5'-CCCCGAGATTGACAGCTAC-3' | GAPDH | 5'-AACTTTGGCATTGTGGAAGG-3' |
| | 5'-ACTCCACTAAATTGCACATTTCTTC-3' | | 5'-GGATGCAGGGATGATGTTCT-3' |

PCR indicates polymerase chain reaction; QPCR, quantitative real-time polymerase chain reaction.

(Figure 1B). RIF-mediated PXR target gene *Cyp3a11* upregulation in the liver was abolished in $PXR^{-/-}ApoE^{-/-}$ mice (Figure 1B). These results confirm the presence of the functional hPXR in $huPXR \cdot ApoE^{-/-}$ mice. Thus, $huPXR \cdot ApoE^{-/-}$ mice provides an in vivo system to assess atherogenic responses to relevant environment chemicals such as BPA, while allowing the use of a murine model to evaluate mechanisms of deleterious effects of BPA arising from human exposure.

BPA Activates hPXR and Stimulates PXR Target Gene Expression in $huPXR \cdot ApoE^{-/-}$ Mice

The main route of human exposure to BPA is oral and pharmacokinetic studies have demonstrated that exposure via diet is a more natural continuous exposure route than other methods commonly used in chronically exposed animals.³⁸ To determine the impact of chronic exposure to BPA on atherosclerosis development, 4-week-old male $huPXR \cdot ApoE^{-/-}$ and $PXR^{-/-}ApoE^{-/-}$ littermates were fed a control diet or a diet supplemented with BPA at a dose of 50 mg/kg. The choice of 50 mg BPA/kg feed weight was based on previous studies demonstrating that 50 mg BPA/kg feed weight represents a moderate or low dose exposure to BPA in experimental animal models.^{31–33,38}

To determine whether 50 mg/kg feed weight provides urinary BPA concentrations similar to that observed in humans, a chemical derivatization LC-ESI-MS/MS method was developed to measure urinary BPA concentrations (Figure 2). Consistent with previous reports that BPA undergoes metabolism (conjugation) and clearance from the body, we were able to detect both conjugated BPA-glucuronide and




Figure 1. Generation of $huPXR \cdot ApoE^{-/-}$ mice. A, Genotype analysis of $huPXR \cdot ApoE^{-/-}$ and $PXR^{-/-}ApoE^{-/-}$ mice by 3 different PCR assays. The presence of the human (h) PXR transgene was determined by hPXR primers (576 bp). Mouse (m) PXR primers were used to identify WT allele (348 bp) and PXR null allele (265 bp). Mouse ApoE primers were used to identify WT allele (155 bp) and ApoE null allele (245 bp). Mouse no. 1 is $huPXR \cdot ApoE^{-/-}$, no. 2 is $PXR^{-/-}ApoE^{-/-}$, and no. 3 is WT control. B, Six-week-old male $huPXR \cdot ApoE^{-/-}$ and $PXR^{-/-}ApoE^{-/-}$ mice were treated with DMSO vehicle control, mPXR-specific ligand pregnenolone 16 α -carbonitrile (PCN), or hPXR-specific ligand rifampicin (RIF) by intraperitoneal injection at the dose of 10 mg/kg per day for 3 days. Total RNA was extracted from the liver, and the mRNA levels of prototypic PXR activated gene *CYP3A11* were measured by QPCR (n=5 per group, ** $P < 0.01$). PCR indicates polymerase chain reaction; PXR, pregnane X receptor; QPCR, quantitative real-time polymerase chain reaction; WT, wild-type.




Figure 2. A novel LC-ESI-MS/MS method to quantitate BPA. A, BPA was derivatized by reaction with pyridine-3-sulfonyl chloride (PSC) in acetone and 1 pmol of the derivatized material was analyzed by reverse phase HPLC using a Waters XTerra MS C8 column. PSC-BPA was detected by positive mode electrospray ionization (ESI) selective reaction monitoring mode tandem MS using an ABSciex 4000 Q-Trap instrument as described in the methods monitoring the precursor product ion pairs shown in the calibration (B). The structure of the PSC derivative of BPA is shown (C) with a fragmentation scheme generating the product ions monitored in (A) and (B). BPA indicates bisphenol A; LC-ESI-MS/MS liquid chromatography-electrospray ionization-tandem mass spectrometry.

unconjugated BPA in the urine of BPA-exposed mice. While the concentrations of BPA-glucuronide were higher than unconjugated BPA, the urinary concentrations of unconjugated BPA were 1.19 and 2.33 ng/mL in $PXR^{-/-}$ ApoE $^{-/-}$ and huPXR•ApoE $^{-/-}$ mice, respectively (Table 2). These values are similar to those detected in human urine samples with the unconjugated BPA concentrations that ranged from undetectable

to 2.5 ng/mL.^{1,39,40} BPA feeding stimulated expression of the prototypic PXR target genes, *Cyp3a11*, *Mdr1a*, and *Cd36* in the liver of huPXR•ApoE $^{-/-}$ mice, but not in $PXR^{-/-}$ ApoE $^{-/-}$ mice (Figure 3), indicating that feeding huPXR•ApoE $^{-/-}$ mice 50 mg BPA/kg feed weight can efficiently activate human PXR in vivo.

Exposure to BPA Does Not Affect Plasma Lipid Levels but Increases Atherosclerosis in huPXR•ApoE $^{-/-}$ Mice

Exposure to BPA for 12 weeks did not affect the body weight of huPXR•ApoE $^{-/-}$ and $PXR^{-/-}$ ApoE $^{-/-}$ mice (Figure 4). The effect of BPA exposure on plasma lipid and lipoprotein levels revealed no changes in plasma triglyceride and cholesterol levels (Figure 5). In addition, FPLC analysis showed that huPXR•ApoE $^{-/-}$ and $PXR^{-/-}$ ApoE $^{-/-}$ mice had similar plasma cholesterol distribution patterns, which were not affected by BPA treatment (Figures 5C and 5F).

Atherosclerotic lesion areas were determined in the aortic root and brachiocephalic artery (BCA) as shown in Figure 6. Feeding huPXR•ApoE $^{-/-}$ mice BPA for 12 weeks significantly increased lesion areas in the aortic root by 104% ($P<0.001$; Student's *t* test) (Figure 6A). BPA feeding also accelerated atherosclerosis development in the BCA, an artery prone to developing advanced lesions (Figure 6B). Compared to mice fed control diet, BCA cross-section lesion areas were increased by 120% ($P<0.05$; Student's *t* test) in BPA-fed huPXR•ApoE $^{-/-}$ mice (Figure 6B). By contrast, exposure to BPA did not affect atherosclerotic lesion development in either the aortic root or BCA of $PXR^{-/-}$ ApoE $^{-/-}$ mice (Figures 6C and 6D). Thus, BPA increases atherosclerosis in these models in a human PXR-dependent manner.

BPA Increases Lipid Accumulation and Foam Cell Formation in Macrophages of huPXR•ApoE $^{-/-}$ Mice

Macrophages play a critical role in atherogenesis and accumulation of lipid-loaded macrophages is a hallmark of atherosclerosis.^{41,42} We previously reported that activation of

Table 2. Urinary BPA and BPA-Glucuronide Concentrations in Mice Fed the Control or BPA Diet for 12 Weeks

| Genotype | Diet | BPA (ng/mL) | BPA-Glucuronide (ng/mL) |
|---------------------------|---------|-------------|-------------------------|
| $PXR^{-/-}$ ApoE $^{-/-}$ | Control | N.D. | N.D. |
| huPXR•ApoE $^{-/-}$ | Control | N.D. | N.D. |
| $PXR^{-/-}$ ApoE $^{-/-}$ | BPA | 1.19±0.98 | 19.97±15.62 |
| huPXR•ApoE $^{-/-}$ | BPA | 2.33±1.93 | 11.60±6.48 |

All values shown are mean±SD (n=9). BPA indicates bisphenol A; PXR, pregnane X receptor; N.D., not detectable.




Figure 3. BPA increases PXR target gene expression in huPXR•ApoE^{-/-} mice. Four-week-old male PXR^{-/-}ApoE^{-/-} and huPXR•ApoE^{-/-} littermates were fed a control diet or supplemented with 50 mg/kg BPA (BPA) for 12 weeks. The expression of hepatic PXR target gene mRNAs was measured by QPCR (n=4 per group, *P<0.05). BPA indicates bisphenol A; PXR, pregnane X receptor.

PXR increases lipid accumulation in macrophages of ApoE^{-/-} mice, which contributes to PXR's pro-atherogenic effects.²⁷ To elucidate possible molecular mechanisms through which BPA increases atherosclerosis, exposure to BPA-affected macrophage functions was investigated. Peritoneal macrophages were isolated from huPXR•ApoE^{-/-} and PXR^{-/-}ApoE^{-/-} mice fed control diet or supplemented with BPA for 12 weeks and neutral lipid levels and foam cell formation in peritoneal macrophages were determined by oil-red-O staining. BPA feeding promoted lipid accumulation and foam cell formation in peritoneal macrophages of huPXR•ApoE^{-/-} mice but not in that of PXR^{-/-}ApoE^{-/-} mice (Figures 7A and 7B). Gene expression analysis showed that BPA exposure stimulated mRNA levels of the prototypic PXR activated

genes, *Mdr1a* (P<0.05; Student's *t* test) and *Cd36* (P<0.01; Student's *t* test), in the macrophages of huPXR•ApoE^{-/-} but not PXR^{-/-}ApoE^{-/-} mice (Figure 7C). CD36 is a member of the scavenger receptor class B family and plays an important role in mediating macrophage lipid uptake and foam cell formation.⁴³ In contrast, the expression levels of mRNA encoding scavenger receptors, SR-A and SR-BI, and ABC transporters, ABCA1, and ABCG1, were not affected by BPA exposure in the macrophages derived from either huPXR•ApoE^{-/-} or PXR^{-/-}ApoE^{-/-} mice (Figure 7C).

The significantly increased *Cd36* mRNA levels and elevated lipid accumulation in the macrophages of BPA-treated huPXR•ApoE^{-/-} mice promoted us to investigate the protein content of CD36 in the atherosclerotic lesions. Immunofluorescence staining showed that PXR is present in the lesions of huPXR•ApoE^{-/-} mice and expressed by lesional macrophages (Figure 7D). Consistent with macrophage gene expression analysis, BPA exposure substantially increased CD36 protein levels in atherosclerotic lesions of huPXR•ApoE^{-/-} mice (Figure 7D). Analysis of atherosclerotic lesions further confirmed that BPA significantly increased macrophage (P<0.05; Student's *t* test) and CD36 content (P<0.05; Student's *t* test) in plaques of huPXR•ApoE^{-/-} mice but not PXR^{-/-}ApoE^{-/-} littermates (Figure 7E). Thus, the increase in atherosclerotic lesions in BPA-fed huPXR•ApoE^{-/-} mice is associated with increased CD36 expression and foam cell formation in macrophages.

Discussion

Risk assessment of BPA is still hampered by large scientific uncertainties and the impact of BPA exposure on human health is not clearly understood. While >95% of the US population is exposed to BPA, there is an urgent need to understand the molecular mechanisms underlying the associations between BPA exposure and CVD. We recently




Figure 4. BPA exposure does not affect body weight. Growth curves of 4-week-old male PXR^{-/-}ApoE^{-/-} (A) and huPXR•ApoE^{-/-} (B) littermates fed a control diet or supplemented with 50 mg/kg BPA (BPA) for 12 weeks (n=11 to 17 per group). BPA indicates bisphenol A; PXR, pregnane X receptor.




Figure 5. BPA exposure does not affect plasma lipid levels and cholesterol distribution. Four-week-old male huPXR•ApoE^{-/-} and PXR^{-/-}•ApoE^{-/-} littermates were fed a control diet or supplemented with 50 mg/kg BPA (BPA) for 12 weeks. The plasma levels of triglyceride (A and D) and cholesterol (B and E) were measured by standard methods (n=11 to 12 per group) and plasma cholesterol distribution (C and F) was analyzed by FPLC. BPA indicates bisphenol A; FPLC, fast-performance liquid chromatography; HDL, high-density lipoprotein; LDL, low-density lipoprotein; PXR, pregnane X receptor.

reported that BPA is a potent agonist of PXR that has proatherogenic effects in animal models upon activation.^{22,24,27} BPA is an hPXR-selective agonist but does not affect rodent PXR activity; consequently, the choice of an appropriate animal model is paramount in predicting the human risk assessment of BPA. Since the ligand-binding specificity of PXR differs between humans and rodents, PXR-humanized mouse models have been generated to more faithfully predict xenobiotic effects and responses in humans.^{24,28,44} Compared with other humanized mice, one of the advantages of the huPXR mouse model is expression in a similar tissue distribution pattern as native PXR gene in humans and mice,²⁸ which enables the study of hPXR function in multiple tissues/cell types. This model has been successfully used in many studies to investigate human PXR ligand-mediated xenobiotic response in mice, and has been established as a useful tool for the prediction of human drug metabolism and toxicological risk assessment.^{37,45–47} Therefore, the huPXR mice were used to generate the huPXR•ApoE^{-/-} mice for studying the effects of BPA on atherogenic effects *in vivo*.

Since the main source of BPA exposure in humans is through the diet, mice were exposed to 50 mg BPA/kg feed weight in the current study. It was previously assumed that BPA undergoes rapid metabolism and clearance from the

body. However, unconjugated BPA has been detected in human urine and tissues, and recent human biomonitoring data demonstrated that the unconjugated BPA concentrations is higher than previously predicted given assumptions about the amount of BPA ingested by humans and its expected rate of clearance.^{1,48,49} Using our newly developed LC-ESI-MS/MS method, urinary unconjugated BPA in mice exposed to BPA were readily detected. While most human studies only report the total BPA concentration, several studies have detected unconjugated BPA in human urine with the concentrations ranging from undetectable to 2.5 ng/mL.^{1,39,40} Our results confirmed that the dose of 50 mg BPA/kg feed weight is appropriated for long-term exposure studies in mice, which can result in urinary BPA concentrations similar to that observed in human samples. Further, the previously described standard LC-MS/MS methods, although reportedly sensitive, were unable to detect unconjugated BPA in many human samples.^{50–52} Therefore, our LC-ESI-MS/MS method, which has significantly improved sensitivity compared to previously described methods may be applied to future biomonitoring studies for the evaluation of BPA exposure levels and safety in humans.

Interestingly, we found that chronic exposure to BPA increased atherosclerosis in huPXR•ApoE^{-/-} mice but not




Figure 6. BPA increases atherosclerosis in a human PXR-dependent manner. Four-week-old male huPXR•ApoE^{-/-} and PXR^{-/-}•ApoE^{-/-} littermates were fed a control diet or a diet supplemented with 50 mg/kg BPA for 12 weeks. Quantitative analysis of atherosclerotic lesion size in the aortic root and brachiocephalic artery (BCA) of huPXR•ApoE^{-/-} (A and B) and PXR^{-/-}•ApoE^{-/-} (C and D) mice (n=10 to 11 per group, **P*<0.05 and ****P*<0.001). Representative Oil red O-stained sections are shown as indicated. BPA indicates bisphenol A; PXR, pregnane X receptor.

their PXR^{-/-}•ApoE^{-/-} littermates without altering plasma lipid levels and cholesterol distribution patterns. Therefore, the increased atherosclerosis in huPXR•ApoE^{-/-} mice exposed to BPA could not be explained by the unchanged plasma lipid levels. PXR can directly regulate fatty acid transporter CD36 transcription, and activation of PXR promotes CD36-mediated hepatic lipid accumulation.⁵³ CD36 plays an important role in atherosclerosis-related processes such as macrophage lipid uptake and foam cell formation.^{43,54} We previously demonstrated that activation of PXR increases CD36 levels and lipid accumulation in peritoneal macrophages of ApoE^{-/-} mice.²⁷ In the current study, the expression levels of CD36 and lipid accumulation were significantly increased in the peritoneal macrophages of huPXR•ApoE^{-/-} mice exposed to BPA. We also observed that PXR is expressed by atherosclerotic lesional macrophages and that BPA exposure increased CD36 and macrophage content in plaques of

huPXR•ApoE^{-/-} mice but not that of PXR^{-/-}•ApoE^{-/-} mice. In addition, the expression levels of several other key receptors and transporters (eg, SR-A, ABCA1, ABCG1) involved in macrophage lipid uptake or efflux were not affected by BPA treatment in macrophages of either huPXR•ApoE^{-/-} or PXR^{-/-}•ApoE^{-/-} mice. Therefore, a plausible explanation for the increased atherosclerosis observed in huPXR•ApoE^{-/-} mice is the increased CD36 expression and CD36-mediated macrophage lipid uptake and foam cell formation.

BPA is a well-characterized xenoestrogen and the estrogenic effects of BPA have been extensively studied in animals. Many effects of BPA have been found to be similar to effects seen in response to estrogen in laboratory rodent models.⁴ Recent studies have also found some adverse effects of exposure to estrogen or BPA on rodent cardiac functions.^{55,56} It was reported that both estrogen and BPA can increase




Figure 7. BPA increases foam cell formation and CD36 expression in macrophages and atherosclerotic lesions of huPXR•ApoE^{-/-} mice. A, Freshly isolated peritoneal macrophages from huPXR•ApoE^{-/-} and PXR^{-/-}•ApoE^{-/-} mice fed a control diet or BPA diet were stained with Oil-red-O and haematoxylin. B, Foam cell quantification from peritoneal macrophages in studies described in panel A (n=4 per group, *P<0.05). C, The expression levels of macrophage genes were measured by QPCR (n=3 per group, *P<0.05 and **P<0.01). D, Sections of atherosclerotic lesion area in the aortic root of huPXR•ApoE^{-/-} mice were stained with anti-monocytes/macrophages (MOMA-2), anti-PXR or anti-CD36 primary antibodies, followed by fluorescein-labeled secondary antibodies. Nuclei were stained with DAPI. E, Quantification of macrophage (MOMA-2) and CD36 staining area in the aortic root of huPXR•ApoE^{-/-} or PXR^{-/-}•ApoE^{-/-} mice (n=5 to 6 per group, *P<0.05). BPA indicates bisphenol A; PXR, pregnane X receptor; QPCR, quantitative real-time polymerase chain reaction.

cardiac arrhythmias in isolated hearts from female but not male rodents⁵⁶ and alter cardiac calcium homeostasis via ER stimulation in isolated female rodent hearts.⁵⁵ However, BPA exposure has been associated with atherosclerosis in humans^{10–12} and numerous studies have confirmed that estrogen has atheroprotective effects in animals and humans.^{14–17} Therefore, it is unlikely that estrogenic activity of BPA can alone increase atherosclerosis and incidence of CVD in humans. It is still plausible that activation of both PXR and ER by BPA coordinately contribute to increased CVD risk in humans. It would be interesting to study the effects of BPA exposure on atherosclerosis development or cardiac functions in ER α - or ER β -deficient mice in the presence or absence of hPXR in the future.

In addition to BPA, we have previously identified several environmentally significant BPA analogs including BPB and BPAF as human PXR ligands.²² Further, we demonstrated that BPA and analogs can synergistically activate human PXR.²² The synergism between BPA and other environmental chemicals support the need to include mixtures for future in vivo studies, which may have important implications for environmental chemical risk assessment. Combinations of BPA and other environmental chemicals may produce significant effects on PXR activity and atherosclerosis development in humans, even when each chemical is present at low doses that individually do not induce observable effects. In addition, BPA has been implicated to have carcinogenic potential.^{18,57} Activation of PXR has been shown to induce tumor aggressiveness in humans and mice.⁵⁸ Future studies are needed to investigate whether BPA-mediated hPXR activation can induce tumorigenesis in PXR-humanized animal models.

In summary, we found that BPA increased atherosclerosis in ApoE^{-/-} mice in a human PXR-dependent manner. BPA exposure did not affect plasma lipid levels but increased CD36 expression and lipid accumulation in macrophages of huPXR \cdot ApoE^{-/-} mice. These findings demonstrate, for the first time, that BPA exposure increases atherosclerosis development in a laboratory animal model, and provide a potential molecular mechanism by which exposure to BPA increases atherosclerosis and CVD risk in humans. Activation of human PXR should be taken into consideration for future risk assessment of BPA and related environmental chemicals.

Sources of Funding

This work was supported in part by NIH grants (P30HL101300, P20GM103527, R01ES023470), and American Heart Association grant 09SDG2150176. Morris and Sunkara were supported by the University of Kentucky Superfund Research Center grant P42ES007380; Helsley

was supported by NIH training grant T32HL072743; and Sui was supported by American Heart Association Postdoctoral Fellowship 14POST18740064.

Disclosures

None.

References

- Vandenbergh LN, Chahoud I, Heindel JJ, Padmanabhan V, Paumgarten FJ, Schoenfelder G. Urinary, circulating, and tissue biomonitoring studies indicate widespread exposure to bisphenol A. *Environ Health Perspect*. 2010;118:1055–1070.
- Calafat AM, Ye X, Wong LY, Reidy JA, Needham LL. Exposure of the U.S. population to bisphenol A and 4-tertiary-octylphenol: 2003–2004. *Environ Health Perspect*. 2008;116:39–44.
- vom Saal FS, Akingbemi BT, Belcher SM, Birnbaum LS, Crain DA, Eriksen M, Farabolini F, Guillette LJ Jr, Hauser R, Heindel JJ, Ho SM, Hunt PA, Iguchi T, Jobling S, Kanno J, Keri RA, Knudsen KE, Lauffer H, LeBlanc GA, Marcus M, McLachlan JA, Myers JP, Nadal A, Newbold RR, Olea N, Prins GS, Richter CA, Rubin BS, Sonnenschein C, Soto AM, Talsness CE, Vandenbergh JG, Vandenbergh LN, Walser-Kuntz DR, Watson CS, Welshons WV, Wetherill Y, Zoeller RT. Chapel Hill bisphenol A expert panel consensus statement: integration of mechanisms, effects in animals and potential to impact human health at current levels of exposure. *Reprod Toxicol*. 2007;24:131–138.
- Richter CA, Birnbaum LS, Farabolini F, Newbold RR, Rubin BS, Talsness CE, Vandenbergh JG, Walser-Kuntz DR, vom Saal FS. In vivo effects of bisphenol A in laboratory rodent studies. *Reprod Toxicol*. 2007;24:199–224.
- Vandenbergh LN, Chahoud I, Padmanabhan V, Paumgarten FJ, Schoenfelder G. Biomonitoring studies should be used by regulatory agencies to assess human exposure levels and safety of bisphenol A. *Environ Health Perspect*. 2010;118:1051–1054.
- Vandenbergh LN, Maffini MV, Sonnenschein C, Rubin BS, Soto AM. Bisphenol-A and the great divide: a review of controversies in the field of endocrine disruption. *Endocr Rev*. 2009;30:75–95.
- Lang IA, Galloway TS, Scarlett A, Henley WE, Depledge M, Wallace RB, Melzer D. Association of urinary bisphenol A concentration with medical disorders and laboratory abnormalities in adults. *JAMA*. 2008;300:1303–1310.
- Melzer D, Rice NE, Lewis C, Henley WE, Galloway TS. Association of urinary bisphenol A concentration with heart disease: evidence from NHANES 2003/06. *PLoS One*. 2010;5:e8673.
- Melzer D, Osborne NJ, Henley WE, Cipelli R, Young A, Money C, McCormack P, Luben R, Khaw KT, Wareham NJ, Galloway TS. Urinary bisphenol A concentration and risk of future coronary artery disease in apparently healthy men and women. *Circulation*. 2012;125:1482–1490.
- Melzer D, Gates P, Osborn NJ, Henley WE, Cipelli R, Young A, Money C, McCormack P, Schofield P, Mosedale D, Grainger D, Galloway TS. Urinary bisphenol A concentration and angiography-defined coronary artery stenosis. *PLoS One*. 2012;7:e43378.
- Lind PM, Lind L. Circulating levels of bisphenol A and phthalates are related to carotid atherosclerosis in the elderly. *Atherosclerosis*. 2011;218:207–213.
- Shankar A, Teppala S, Sabanayagam C. Bisphenol A and peripheral arterial disease: results from the nhanes. *Environ Health Perspect*. 2012;120:1297–1300.
- Wetherill YB, Akingbemi BT, Kanno J, McLachlan JA, Nadal A, Sonnenschein C, Watson CS, Zoeller RT, Belcher SM. In vitro molecular mechanisms of bisphenol A action. *Reprod Toxicol*. 2007;24:178–198.
- Nathan L, Chaudhuri G. Estrogens and atherosclerosis. *Annu Rev Pharmacol Toxicol*. 1997;37:477–515.
- Hodgin JB, Kregel JH, Reddick RL, Korach KS, Smithies O, Maeda N. Estrogen receptor alpha is a major mediator of 17beta-estradiol's atheroprotective effects on lesion size in apoe^{-/-} mice. *J Clin Invest*. 2001;107:333–340.
- Billon-Gales A, Krust A, Fontaine C, Abot A, Flouriot G, Toutain C, Berges H, Gadeau AP, Lenfant F, Gourdy P, Chambon P, Arnal JF. Activation function 2 (AF2) of estrogen receptor-alpha is required for the atheroprotective action of estradiol but not to accelerate endothelial healing. *Proc Natl Acad Sci USA*. 2011;108:13311–13316.
- Arnal JF, Fontaine C, Billon-Gales A, Favre J, Laurell H, Lenfant F, Gourdy P. Estrogen receptors and endothelium. *Arterioscler Thromb Vasc Biol*. 2010;30:1506–1512.

18. Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM, Zoeller RT, Gore AC. Endocrine-disrupting chemicals: an endocrine society scientific statement. *Endocr Rev*. 2009;30:293–342.
19. Sharpe RM. Is it time to end concerns over the estrogenic effects of bisphenol A? *Toxicol Sci*. 2010;114:1–4.
20. Ryan BC, Hotchkiss AK, Crofton KM, Gray LE Jr. In utero and lactational exposure to bisphenol A, in contrast to ethinyl estradiol, does not alter sexually dimorphic behavior, puberty, fertility, and anatomy of female rats. *Toxicol Sci*. 2010;114:133–148.
21. Howdeshell KL, Furr J, Lambright CR, Wilson VS, Ryan BC, Gray LE Jr. Gestational and lactational exposure to ethinyl estradiol, but not bisphenol A, decreases androgen-dependent reproductive organ weights and epididymal sperm abundance in the male long evans hooded rat. *Toxicol Sci*. 2008;102:371–382.
22. Sui Y, Ai N, Park SH, Rios-Pilier J, Perkins JT, Welsh WJ, Zhou C. Bisphenol A and its analogues activate human pregnane X receptor. *Environ Health Perspect*. 2012;120:399–405.
23. Kliewer SA, Goodwin B, Willson TM. The nuclear pregnane X receptor: a key regulator of xenobiotic metabolism. *Endocr Rev*. 2002;23:687–702.
24. Zhou C, Verma S, Blumberg B. The steroid and xenobiotic receptor (SXR), beyond xenobiotic metabolism. *Nucl Recept Signal*. 2009;7:e001.
25. Blumberg B, Sabbagh W Jr, Juguilon H, Bolado J Jr, van Meter CM, Ong ES, Evans RM. SXR, a novel steroid and xenobiotic-sensing nuclear receptor. *Genes Dev*. 1998;12:3195–3205.
26. Kliewer SA, Moore JT, Wade L, Staudinger JL, Watson MA, Jones SA, McKee DD, Oliver BB, Willson TM, Zetterstrom RH, Perlmann T, Lehmann JM. An orphan nuclear receptor activated by pregnanes defines a novel steroid signaling pathway. *Cell*. 1998;92:73–82.
27. Zhou C, King N, Chen KY, Breslow JL. Activation of PXR induces hypercholesterolemia in wild-type and accelerates atherosclerosis in apoE deficient mice. *J Lipid Res*. 2009;50:2004–2013.
28. Ma X, Shah Y, Cheung C, Guo GL, Feigenbaum L, Krausz KW, Idle JR, Gonzalez FJ. The PREGnane X receptor gene-humanized mouse: a model for investigating drug-drug interactions mediated by cytochromes P450 3A. *Drug Metab Dispos*. 2007;35:194–200.
29. Zhou C, Pridgen B, King N, Xu J, Breslow JL. Hyperglycemic Ins2Akita^{ldlr^{-/-}} mice show severely elevated lipid levels and increased atherosclerosis: a model of type 1 diabetic macrovascular disease. *J Lipid Res*. 2011;52:1483–1493.
30. Teupser D, Persky AD, Breslow JL. Induction of atherosclerosis by low-fat, semisynthetic diets in LDL receptor-deficient C57BL/6J and FVB/NJ mice: comparison of lesions of the aortic root, brachiocephalic artery, and whole aorta (en face measurement). *Arterioscler Thromb Vasc Biol*. 2003;23:1907–1913.
31. Dolinoy DC, Huang D, Jirtle RL. Maternal nutrient supplementation counteracts bisphenol A-induced DNA hypomethylation in early development. *Proc Natl Acad Sci USA*. 2007;104:13056–13061.
32. Jasarevic E, Sieli PT, Twellman EE, Welsh TH Jr, Schachtman TR, Roberts RM, Geary DC, Rosenfeld CS. Disruption of adult expression of sexually selected traits by developmental exposure to bisphenol A. *Proc Natl Acad Sci USA*. 2011;108:11715–11720.
33. Rosenfeld CS, Sieli PT, Warzak DA, Ellersieck MR, Pennington KA, Roberts RM. Maternal exposure to bisphenol A and genistein has minimal effect on A(vy)/a offspring coat color but favors birth of agouti over nonagouti mice. *Proc Natl Acad Sci USA*. 2013;110:537–542.
34. Park SH, Sui Y, Gizard F, Xu J, Rios-Pilier J, Hellsley RN, Han SS, Zhou C. Myeloid-specific ikappaB kinase beta deficiency decreases atherosclerosis in low-density lipoprotein receptor-deficient mice. *Arterioscler Thromb Vasc Biol*. 2012;32:2869–2876.
35. Xu L, Spink DC. Analysis of steroidal estrogens as pyridine-3-sulfonyl derivatives by liquid chromatography electrospray tandem mass spectrometry. *Anal Biochem*. 2008;375:105–114.
36. Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, Feldstein AE, Britt EB, Fu X, Chung YM, Wu Y, Schauer P, Smith JD, Allayee H, Tang WH, DiDonato JA, Lusis AJ, Hazen SL. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature*. 2011;472:57–63.
37. Li F, Lu J, Cheng J, Wang L, Matsubara T, Csanaky IL, Klaassen CD, Gonzalez FJ, Ma X. Human PXR modulates hepatotoxicity associated with rifampicin and isoniazid co-therapy. *Nat Med*. 2013;19:418–420.
38. Sieli PT, Jasarevic E, Warzak DA, Mao J, Ellersieck MR, Liao C, Kannan K, Collet SH, Toutain PL, Vom SAAL FS, Rosenfeld CS. Comparison of serum bisphenol A concentrations in mice exposed to bisphenol A through the diet versus oral bolus exposure. *Environ Health Perspect*. 2011;119:1260–1265.
39. Volkel W, Kiranoglu M, Fromme H. Determination of free and total bisphenol A in human urine to assess daily uptake as a basis for a valid risk assessment. *Toxicol Lett*. 2008;179:155–162.
40. Schorringhumer K, Cichna-Markl M. Sample clean-up with sol-gel enzyme and immunoaffinity columns for the determination of bisphenol A in human urine. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2007;850:361–369.
41. Lusis AJ. Atherosclerosis. *Nature*. 2000;407:233–241.
42. Glass CK, Witztum JL. Atherosclerosis. The road ahead. *Cell*. 2001;104:503–516.
43. Rahaman SO, Lennon DJ, Febbraio M, Podrez EA, Hazen SL, Silverstein RL. A cd36-dependent signaling cascade is necessary for macrophage foam cell formation. *Cell Metab*. 2006;4:211–221.
44. Xie W, Barwick JL, Downes M, Blumberg B, Simon CM, Nelson MC, Neuschwander-Tetri BA, Brunt EM, Guzelian PS, Evans RM. Humanized xenobiotic response in mice expressing nuclear receptor SXR. *Nature*. 2000;406:435–439.
45. Holmstock N, Gonzalez FJ, Baes M, Annaert P, Augustijns P. PXR/CYP3A4-humanized mice for studying drug-drug interactions involving intestinal P-glycoprotein. *Mol Pharm*. 2013;10:1056–1062.
46. Johnson CH, Bonzo JA, Cheng J, Krausz KW, Kang DW, Luecke H, Idle JR, Gonzalez FJ. Cytochrome P450 regulation by alpha-tocopherol in Pxr-null and PXR-humanized mice. *Drug Metab Dispos*. 2013;41:406–413.
47. Cheung C, Gonzalez FJ. Humanized mouse lines and their application for prediction of human drug metabolism and toxicological risk assessment. *J Pharmacol Exp Ther*. 2008;327:288–299.
48. Vandenberg LN, Hauser R, Marcus M, Olea N, Welshons WV. Human exposure to bisphenol A (BPA). *Reprod Toxicol*. 2007;24:139–177.
49. Betts KS. Body of proof: biomonitoring data reveal widespread bisphenol A exposures. *Environ Health Perspect*. 2010;118:a353.
50. Volkel W, Colnot T, Csanady GA, Filsler JG, Dekant W. Metabolism and kinetics of bisphenol A in humans at low doses following oral administration. *Chem Res Toxicol*. 2002;15:1281–1287.
51. Teeguarden JG, Calafat AM, Ye X, Doerge DR, Churchwell MI, Gunawan R, Graham MK. Twenty-four hour human urine and serum profiles of bisphenol A during high-dietary exposure. *Toxicol Sci*. 2011;123:48–57.
52. Vom SAAL FS, Prins GS, Welshons WV. Report of very low real-world exposure to bisphenol A is unwarranted based on a lack of data and flawed assumptions. *Toxicol Sci*. 2012;125:318–320; author reply 321–315.
53. Zhou J, Zhai Y, Mu Y, Gong H, Uppal H, Toma D, Ren S, Evans RM, Xie W. A novel pregnane X receptor-mediated and sterol regulatory element-binding protein-independent lipogenic pathway. *J Biol Chem*. 2006;281:15013–15020.
54. Kennedy DJ, Kuchibhotla SD, Guy E, Park YM, Nimako G, Vanegas D, Morton RE, Febbraio M. Dietary cholesterol plays a role in CD36-mediated atherogenesis in LDLR-knockout mice. *Arterioscler Thromb Vasc Biol*. 2009;29:1481–1487.
55. Belcher SM, Chen Y, Yan S, Wang HS. Rapid estrogen receptor-mediated mechanisms determine the sexually dimorphic sensitivity of ventricular myocytes to 17beta-estradiol and the environmental endocrine disruptor bisphenol A. *Endocrinology*. 2012;153:712–720.
56. Yan S, Chen Y, Dong M, Song W, Belcher SM, Wang HS. Bisphenol A and 17beta-estradiol promote arrhythmia in the female heart via alteration of calcium handling. *PLoS One*. 2011;6:e25455.
57. Soto AM, Sonnenschein C. Environmental causes of cancer: endocrine disruptors as carcinogens. *Nat Rev Endocrinol*. 2010;6:363–370.
58. Wang H, Venkatesh M, Li H, Goetz R, Mukherjee S, Biswas A, Zhu L, Kaubisch A, Wang L, Pullman J, Whitney K, Kuro-o M, Roig AI, Shay JW, Mohammadi M, Mani S. Pregnane X receptor activation induces FGF19-dependent tumor aggressiveness in humans and mice. *J Clin Invest*. 2011;121:3220–3232.



Bisphenol A Increases Atherosclerosis in Pregnane X Receptor-Humanized ApoE Deficient Mice

Yipeng Sui, Se-Hyung Park, Robert N. Helsley, Manjula Sunkara, Frank J. Gonzalez, Andrew J. Morris and Changcheng Zhou

J Am Heart Assoc. 2014;3:e000492; originally published April 22, 2014;

doi: 10.1161/JAHA.113.000492

The *Journal of the American Heart Association* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231
Online ISSN: 2047-9980

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://jaha.ahajournals.org/content/3/2/e000492>

Subscriptions, Permissions, and Reprints: The *Journal of the American Heart Association* is an online only Open Access publication. Visit the Journal at <http://jaha.ahajournals.org/> for more information.