



4-5-2011

High-Activity Mutants of Butyrylcholinesterase for Cocaine Hydrolysis and Method of Generating the Same

Chang-Guo Zhan

University of Kentucky, chang-guo.zhan@uky.edu

Hoon Cho

University of Kentucky

Hsin-Hsiung Tai

University of Kentucky, htai1@uky.edu

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/ps_patents

 Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Zhan, Chang-Guo; Cho, Hoon; and Tai, Hsin-Hsiung, "High-Activity Mutants of Butyrylcholinesterase for Cocaine Hydrolysis and Method of Generating the Same" (2011). *Pharmaceutical Sciences Faculty Patents*. 2.

https://uknowledge.uky.edu/ps_patents/2

This Patent is brought to you for free and open access by the Pharmaceutical Sciences at UKnowledge. It has been accepted for inclusion in Pharmaceutical Sciences Faculty Patents by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.



US007919082B1

(12) **United States Patent**
Zhan et al.

(10) **Patent No.:** **US 7,919,082 B1**
(45) **Date of Patent:** **Apr. 5, 2011**

(54) **HIGH-ACTIVITY MUTANTS OF BUTYRYLCHOLINESTERASE FOR COCAINE HYDROLYSIS AND METHOD OF GENERATING THE SAME**

(75) Inventors: **Chang-Guo Zhan**, Lexington, KY (US);
Hoon Cho, Lexington, KY (US);
Hsin-Hsiung Tai, Lexington, KY (US)

(73) Assignee: **University of Kentucky Research Foundation**, Lexington, KY (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/752,920**

(22) Filed: **Apr. 1, 2010**

Related U.S. Application Data

(62) Division of application No. 12/192,394, filed on Aug. 15, 2008, now Pat. No. 7,731,957, which is a division of application No. 11/243,111, filed on Oct. 4, 2005, now Pat. No. 7,438,904.

(51) **Int. Cl.**
A61K 38/46 (2006.01)
C12N 9/16 (2006.01)

(52) **U.S. Cl.** **424/94.6**; 435/196; 435/197

(58) **Field of Classification Search** 424/94.6;
435/196, 197

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,049,121 B2 5/2006 Watkins et al.
7,731,957 B1 * 6/2010 Zhan et al. 424/94.6
2003/0153062 A1 8/2003 Watkins et al.
2004/0120939 A1 6/2004 Watkins et al.

OTHER PUBLICATIONS

Gao, et al.; Modeling effects of oxyanion hole on the ester hydrolyses catalyzed by human cholinesterases; Phys. Chem. B; 2005; 109; pp. 23070-23076.

Gao, et al; Computational design of a human butyrylcholinesterase mutant for accelerating cocaine hydrolysis based on the transition-state simulation; Angew. Chem. Int. Ed.; 2006; 45; pp. 653-657.

Gao, et al; Modeling evolution of hydrogen bonding and stabilization of transition states in the process of cocaine hydrolysis catalyzed by human butyrylcholinesterase; Proteins; 2006; 62; pp. 99-110.

Hamza, et al.; Molecular dynamics simulation of cocaine binding with human butyrylcholinesterase and its mutants; J. Phys. Chem. B.; 2005; 109; pp. 4776-4782.

Pan, et al.; Computational redesign of human butyrylcholinesterase for anti-cocaine medication; G. Proc. Natl. Acad. Sci. USA; 2005; 102; pp. 16656-16661.

Zhan, et al.; Fundamental reaction mechanism for cocaine metabolism in human butyrylcholinesterase; J. Am. Chem. Soc.; 2003; 125; pp. 2462-2474.

Zhan, et al.; Catalytic Mechanism and energy barriers for butyrylcholinesterase-catalyzed hydrolysis of cocaine; Biophysical Journal; 2005; 89; 3863-3872.

Xie, et al.; An improved cocaine hydrolase: the A328Y mutant of human butyrylcholinesterase is 4-fold more efficient; Moleculare Pharm; 1999; 55; pp. 83-91.

* cited by examiner

Primary Examiner — Tekchand Saidha

(74) *Attorney, Agent, or Firm* — Stites & Harbison PLLC; Mandy Wilson Decker

(57) **ABSTRACT**

A novel computational method and generation of mutant butyrylcholinesterase for cocaine hydrolysis is provided. The method includes molecular modeling a possible BChE mutant and conducting molecular dynamics simulations and hybrid quantum mechanical/molecular mechanical calculations thereby providing a screening method of possible BChE mutants by predicting which mutant will lead to a more stable transition state for a rate determining step. Site-directed mutagenesis, protein expression, and protein activity is conducted for mutants determined computationally as being good candidates for possible BChE mutants, i.e., ones predicted to have higher catalytic efficiency as compared with wild-type BChE. In addition, mutants A199S/A328W/Y332G, A199S/F227A/A328W/Y332G, A199S/S287G/A328W/Y332G, A199S/F227A/S287G/A328W/Y332G, and A199S/F227A/S287G/A328W/E441D all have enhanced catalytic efficiency for (-)-cocaine compared with wild-type BChE.

5 Claims, 6 Drawing Sheets







FIGURE 1




(A) BChE-(-)-cocaine
non-prereactive complex



(B) BChE-(-)-cocaine
prereactive complex




(C) BChE-(+)-cocaine
non-prereactive complex



(D) BChE-(+)-cocaine
prereactive complex

FIGURE 2



(A)

(B)

FIGURE 3




FIGURE 4




FIGURE 5

(A)




FIGURE 6

1
HIGH-ACTIVITY MUTANTS OF
BUTYRYLCHOLINESTERASE FOR
COCAINE HYDROLYSIS AND METHOD OF
GENERATING THE SAME

RELATED APPLICATIONS

This application is a division of and claims benefit to U.S. patent application Ser. No. 12/192,394, now allowed, filed Aug. 15, 2008 now U.S. Pat. No. 7,731,957, which is a division of and claims benefit to U.S. patent application Ser. No. 11/243,111, now issued as U.S. Pat. No. 7,438,904, filed Oct. 4, 2005. The contents of which are incorporated herein by reference in their entirety.

GOVERNMENT INTEREST

Subject matter described herein was made with government support under Grant Number R01DA013930 awarded by the National Institute on Drug Abuse (NIDA) of the National Institutes of Health (NIH). The government has certain rights in the described subject matter.



FIELD OF THE INVENTION

The present invention relates to butyrylcholinesterase variant polypeptides, and in particular, butyrylcholinesterase mutants having amino acid substitutions.

BACKGROUND OF THE INVENTION

Cocaine abuse is a major medical and public health problem that continues to defy treatment. The disastrous medical and social consequences of cocaine addiction, such as violent crime, loss in individual productivity, illness and death, have made the development of an effective pharmacological treatment a high priority. However, cocaine mediates its reinforcing and toxic effects by blocking neurotransmitter reuptake and the classical pharmacodynamic approach has failed to yield small-molecule receptor antagonists due to the difficulties inherent in blocking a blocker. An alternative to receptor-based approaches is to interfere with the delivery of cocaine to its receptors and accelerate its metabolism in the body.

The dominant pathway for cocaine metabolism in primates is butyrylcholinesterase (BChE)-catalyzed hydrolysis at the benzoyl ester group (Scheme 1).



Scheme 1. Schematic representation of BChE-catalyzed hydrolysis at the benzoyl ester group.

Only 5% of the cocaine is deactivated through oxidation by the liver microsomal cytochrome P450 system. Cocaine hydrolysis at benzoyl ester group yields ecgonine methyl ester, whereas the oxidation produces norcocaine. The metabolite ecgonine methyl ester is a biologically inactive metabolite, whereas the metabolite norcocaine is hepatotoxic and a local anesthetic. BChE is synthesized in the liver and widely distributed in the body, including plasma, brain, and lung. Extensive experimental studies in animals and humans demonstrate that enhancement of BChE activity by administration of exogenous enzyme substantially decreases cocaine half-life.

Enhancement of cocaine metabolism by administration of BChE has been recognized to be a promising pharmacokinetic approach for treatment of cocaine abuse and dependence. However, the catalytic activity of this plasma enzyme is three orders-of-magnitude lower against the naturally occurring (-)-cocaine than that against the biologically inactive (+)-cocaine enantiomer. (+)-cocaine can be cleared from plasma in seconds and prior to partitioning into the central nervous system (CNS), whereas (-)-cocaine has a plasma half-life of approximately 45-90 minutes, long enough for manifestation of the CNS effects which peak in minutes. Hence, BChE mutants with high activity against (-)-cocaine are highly desired for use in humans. Although some BChE mutants with increased catalytic activity over wild-type BChE have previously been generated, there exists a need for mutant BChE with even higher catalytic activity. Thus, prior mutants provide limited enhancement in catalytic activity over wild-type BChE.

Previous studies such as (a) Masson, P.; Legrand, P.; Bartels, C. F.; Froment, M.-T.; Schopfer, L. M.; Lockridge, O. *Biochemistry* 1997, 36, 2266 (b) Masson, P.; Xie, W., Froment, M.-T.; Levitsky, V.; Fortier, P.-L.; Albaret, C.; Lockridge, O. *Biochim. Biophys. Acta* 1999, 1433, 281, (c) Xie, W.; Altamirano, C. V.; Bartels, C. F.; Speirs, R. J.; Cashman, J. R.; Lockridge, O. *Mol. Pharmacol.* 1999, 55, 83, (d) Duyssen, E. G.; Bartels, C. F.; Lockridge, O. *J. Pharmacol. Exp. Ther.* 2002, 302, 751, (e) Nachon, F.; Nicolet, Y.; Viguie, N.; Masson, P.; Fontecilla-Camps, J. C.; Lockridge, O. *Eur. J. Biochem.* 2002, 269, 630, (f) Zhan, C.-G.; Landry, D. W. *J. Phys. Chem. A* 2001, 105, 1296; Berkman, C. E.; Underiner, G. E.; Cashman, J. R. *Biochem. Pharmacol.* 1997, 54, 1261; (g) Sun, H.; Yazal, J. E.; Lockridge, O.; Schopfer, L. M.; Brimijoin, S.; Pang, Y.-P. *J. Biol. Chem.* 2001, 276, 9330, (h) Sun, H.; Shen, M. L.; Pang, Y. P.; Lockridge, O.; Brimijoin, S. *J. Pharmacol. Exp. Ther.* 2002, 302, 710, (i) Sun, H.; Pang, Y. P.; Lockridge, O.; Brimijoin, S. *Mol. Pharmacol.* 2002, 62, 220 (hereinafter "Sun et al"); and (j) Zhan, C.-G.; Zheng, F.; Landry, D. W. *J. Am. Chem. Soc.* 2003, 125, 2462 (hereinafter "Zhan et al"), herein all incorporated by reference, suggested that, for both (-)-cocaine and (+)-cocaine, the BChE-substrate binding involves two different types of complexes: non-prereactive and prereactive BChE-substrate complexes. Whereas the non-prereactive BChE-cocaine complexes were first reported by Sun et al, Zhan et al were the first reporting the prereactive BChE-cocaine complexes and reaction coordinate calculations, disclosed in Zhan et al.

It was demonstrated that (-)/(+)-cocaine first slides down the substrate-binding gorge to bind to W82 and stands vertically in the gorge between D70 and W82 (non-prereactive complex) and then rotates to a position in the catalytic site within a favorable distance for nucleophilic attack and hydrolysis by S198 (prereactive complex). In the prereactive complex, cocaine lies horizontally at the bottom of the gorge. The main structural difference between the BChE-(-)-cocaine complexes and the corresponding BChE-(+)-cocaine complexes exists in the relative position of the cocaine methyl ester group. Reaction coordinate calculations revealed that the rate-determining step of BChE-catalyzed hydrolysis of (+)-cocaine is the chemical reaction process, whereas for (-)-cocaine the change from the non-prereactive complex to the prereactive complex is rate determining. A further analysis of the structural changes from the non-prereactive complex to the prereactive complex reveals specific amino acid residues hindering the structural changes, providing initial clues for the rational design of BChE mutants with improved catalytic activity for (-)-cocaine.

Previous molecular dynamics (MD) simulations of prereactive BChE-cocaine binding were limited to wild-type BChE. Even for the non-prereactive BChE-cocaine complex, only one mutant (A328W/Y332A) BChE binding with (-)-cocaine was simulated and its catalytic activity for (-)-cocaine was reported by Sun et al. No MD simulation was performed on any prereactive enzyme-substrate complex for (-) or (+)-cocaine binding with a mutant BChE. In addition, all previous computational studies of Sun et al and Zhan et al of BChE interacting with cocaine were performed based on a homology model of BChE when three-dimensional (3D) X-ray crystal structure was not available for BChE, as taught by Nicolet, Y.; Lockridge, O.; Masson, P.; Fontecilla-Camps, J. C.; Nachon, F. *J. Biol. Chem.* 2003, 278, 41141 (hereinafter "Nicolet et al"), recently reported 3D X-ray crystal structures of BChE. As expected, the structure of BChE is similar to a previously published theoretical model of this enzyme and to the structure of acetylcholinesterase.

The main difference between the experimentally determined BChE structure and its model was found at the acyl binding pocket (acyl loop) that is significantly bigger than expected. It is unclear whether the structural difference at the acyl binding pocket significantly affect BChE binding with (-)-cocaine and (+)-cocaine. Although previous MD simulations of cocaine binding with wild-type BChE and the reaction coordinate calculations point to some amino acid residues that might need to be mutated for the purpose of improving the catalytic activity for (-)-cocaine hydrolysis, it remained unknown which exact amino acid mutations will result in a BChE with a higher catalytic activity for (-)-cocaine.

Computational studies of wild-type BChE and cocaine from Sun, et al, based on a "homology model," suggest that the rate-determining step for BChE-catalyzed hydrolysis of cocaine is the rotation of the cocaine in the active site of BChE. By decreasing the hindrance of the rotation, the rate of the hydrolysis may be enhanced. Sun, et al describes creating an A328W/Y332A BChE mutant by: (1) replacing Tyr332 with Ala, "to reduce the steric hindrance and the π - π interaction that impede rotation," and (2) replacing Ala328 with Trp "to provide a cation- π interaction to restore substrate affinity lost in disabling the π - π interaction."

Sun et al studied the A328W/Y332A BChE mutant using enzyme assays and kinetics. In vitro studies were conducted using human plasma and in vivo studies were conducted using male Sprague-Dawley rats. The mutant was found to have enhanced catalytic properties. The mutant was further studied using molecular modeling. The three dimensional (3D) structure of A328W/Y332A was generated from the computationally generated 3D model of wild-type BChE and changing the relevant residues using commercially available software. Cocaine was docked to the catalytic gorge of the mutant BChE using other commercially available software. The cocaine-enzyme complex was refined by molecular dynamic simulation. The data generated by the molecular modeling studies were consistent with enzyme assays and kinetic data.

It should be noted that all prior computational techniques (molecular docking and molecular dynamics simulation) used by other researchers are based on an empirical force field which cannot be used to perform any necessary reaction coordinate calculation for the detailed understanding of the complicated catalytic reaction process. As it is well-known, it is particularly challenging to model and simulate the detailed reaction pathway and predict the kinetics of such an enzymatic reaction.

U.S. Patent Application Publication Nos. 2004/0121970; 2004/0120939; and 2003/0153062, describe 20+BChE mutants, or "variants," from human and other animals, each having from one to six amino acid alterations and increased cocaine hydrolysis activity. For example, mutants include F227A/A328W; F227A/S287G/A328W; A119S/S287G/A328W; A328W/Y332M/S287G/F227A, A199S/F227A/S/287G/A328W and A119S/F227A/S287G/A328W/Y332M. The mutants have varying increases in catalytic activity, up to 100-fold increase relative to wild-type BChE.

There exists a need in the art for determining which proposed mutant BChEs should have ever increasing catalytic activity and for generating those mutants which should have enhanced catalytic activity.

SUMMARY OF THE INVENTION

The present invention includes five novel human BChE mutants that have unexpected increased catalytic efficiency

for cocaine hydrolysis. The mutants have various unique amino acid residue substitutions which provide the surprising enhanced catalytic activity. These mutants are (1) A199S/A328W/Y332G mutant (SEQ ID NO: 2), which has an approximately 65-fold improved catalytic efficiency against (-)-cocaine; (2) A199S/F227A/A328W/Y332G mutant (SEQ ID NO: 8), which has an approximately 148-fold improved catalytic efficiency against (-)-cocaine; (3) A199S/S287G/A328W/Y332G mutant (SEQ ID NO: 14), which has an approximately 456-fold improved catalytic efficiency against (-)-cocaine; (4) A199S/F227A/S287G/A328W/Y332G mutant (SEQ ID NO: 20), which has an approximately 1.003-fold improved catalytic efficiency against (-)-cocaine; and (5) A199S/F227A/S287G/A328W/E441D mutant (SEQ ID NO: 26), which has an approximately 445-fold improved catalytic efficiency against (-)-cocaine.

In addition, the aforementioned mutant amino acid sequences can be truncated without substantially affecting the catalytic activity so that amino acid residues 1-67 and 443-574 can be removed without substantially affecting the catalytic activity of the enzyme. SEQ ID NOS: 4, 10, 16, 22 and 28 are the amino acid sequences for residue 68-142 corresponding to mutants 1-5, respectively. In addition, with regard to mutants 1-4, it was found that amino acid residues before 117 and after 438 could be removed without substantially changing the activity of the mutant enzymes, resulting in truncated amino acid sequences having SEQ ID NOS: 6, 12, 18 and 24, respectively. Finally with regard to mutant 5, amino acid residues before 117 and after 441 could be removed without substantially changing its activity resulting in SEQ ID NO: 30.

These aforementioned truncated sequences all of with similar catalytic activity is based on protein structures.

Further, the present invention is directed to a novel and unique pharmaceutical composition which comprises a butyrylcholinesterase variant, namely mutants 1-5, along with a suitable pharmaceutical carrier. The pharmaceutical composition can be administered to an individual in an effective amount to lower the patient's cocaine blood concentration and in particular (-)-cocaine blood concentration.

In addition, the present invention is directed to a novel and unique method for developing mutants which have enhanced catalytic efficiency. The generation method includes both a computational portion and an experimental portion. With regard to the computational portion, a variety of state of the art computational techniques including molecular modeling, molecular dynamics (MD) simulations and hybrid quantum mechanical/molecular mechanical (QM/MM) calculations, provide a virtual screening of possible BChE mutants. This virtual screening predicts which mutation will lead to a more stable transition state for a rate-determining step compared to the corresponding separated reactants, i.e., free cocaine and free enzyme. The more stable the transition state, the lower the energy barrier, and the higher the catalytic efficiency. Following the computational portion, an experimental test is then conducted on the possible mutants of the computation portion. The experimental test includes site-directed mutagenesis, protein expression, and enzyme activity assay. The experimental tests are conducted on mutants which are predicted to have a high catalytic efficiency against (-)-cocaine than the wild-type BChE and/or other known BChE mutants against (-)-cocaine. Thus, the present method identifies or predicts mutants having high catalytic activity for cocaine hydrolysis by performing molecular modeling and MD simulations on the transition state structures of possible mutants of BChE. This method is an improvement over traditional random-search approaches, which, given the com-

plex catalytic mechanism of cocaine hydrolysis, makes it difficult to improve the catalytic activity of BChE for cocaine hydrolysis.

The present invention in one form, concerns a butyrylcholinesterase variant peptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, and 30.

The present invention in another form thereof concerns a nucleic acid molecule comprising a nucleic acid sequence which encodes a butyrylcholinesterase variant peptide, the nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, and 29.

The present invention in another form thereof concerns a pharmaceutical composition comprising a butyrylcholinesterase variant polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS. 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, and 30; and a suitable pharmaceutical carrier.

The present invention in another form thereof concerns a method for treating a cocaine-induced condition comprising administering to an individual an effective amount of butyrylcholinesterase variant peptide having an amino acid sequence selected from the group consisting of SEQ ID NOS. 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, and 30, to lower blood cocaine concentration.

The present invention is another form thereof concerns a method for treating a cocaine induced condition comprising administering to an individual an effective amount of a pharmaceutical composition comprising a butyrylcholinesterase variant having an amino acid sequence selected from the group consisting of SEQ ID NOS. 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, and 30, and suitable pharmaceutical carrier of claim 3 to an individual in an effective amount to lower blood cocaine concentration.

The present invention in yet another form thereof concerns a method for generating butyrylcholinesterase mutants. The method includes generating an initial structure of the transition state structure for the rate-determining step of the cocaine hydrolysis catalyzed by a possible butyrylcholinesterase mutant. A sufficiently long time molecular dynamics simulation is performed on the transition state structure in water to have a stable molecular dynamics trajectory. The molecular dynamics trajectory is analyzed and the hydrogen bonding energies are estimated between the carboxyl oxygen of the (-)-cocaine benzyl ester and the oxyanion hole of the possible butyrylcholinesterase mutant. If the overall hydrogen binding energy between the carboxyl oxygen of the (-)-cocaine benzyl ester and the possible butyrylcholinesterase mutant, in the transition state, is stronger than the overall hydrogen binding energy between the carboxyl oxygen of the (-)-cocaine benzyl ester and the wild-type butyrylcholinesterase, optionally, hybrid quantum mechanical/molecular mechanical (QM/MM) geometry optimization is performed to refine the molecular dynamics-simulated structure, the hydrogen binding energies are calculated and the energy barrier is evaluated. Finally, the butyrylcholinesterase mutant is generated.

In various alternative embodiments, the generating an initial structure of the transition state structure is based on reaction coordinate calculations for the wild-type butyrylcholinesterase. The generating butyrylcholinesterase mutant includes performing site-directed mutagenesis on a nucleic acid sequence which includes wild-type butyrylcholinesterase to generate the mutant butyrylcholinesterase nucleic acid sequence. Using the mutant butyrylcholinesterase nucleic acid sequence, the protein encoded by the mutant

nucleic acid sequences is expressed to produce mutant butyrylcholinesterase and catalytic activity assay is performed on the mutant butyrylcholinesterase.

The hybrid quantum mechanical/molecular mechanical geometry optimization may include calculating the hydrogen binding energies and evaluating the energy barriers only if the overall hydrogen binding energy between the carboxyl oxygen of the (-)-cocaine benzyl ester and the possible butyrylcholinesterase mutant, in the transition state, is stronger than known butyrylcholinesterase mutants against (-)-cocaine.

In yet another alternative further embodiment, the method for generating butyrylcholinesterase mutants further includes determining the rate-limiting step in the hydrolysis of (-)-cocaine by the possible butyrylcholinesterase mutant by conducting molecular dynamics simulations and quantum mechanical/molecular mechanical calculations relating to the transition states for other reaction steps between (-)-cocaine by the possible butyrylcholinesterase mutant and calculating respective energy barriers, thereby establishing which of the reaction steps is the rate-determining one.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 are plots of distances in the MD simulation (-)-cocaine binding with A328W/Y332G BChE versus the simulation time, along with root-mean-square deviation (RMSD) in the enzyme-substrate complexes where FIG. 1A represents the non-prereactive enzyme-substrate complexes; and FIG. 1B represents the prereactive enzyme-substrate complexes in accordance with the present invention.

FIG. 2 shows the binding of structures of the simulated non-prereactive and prereactive complexes of wild-type BChE binding with two entomers of cocaine in which FIG. 2A depicts BChE (-)-cocaine non-prereactive complex; FIG. 2B depicts BChE (-)-cocaine prereactive complex, FIG. 2C depicts BChE (+)-cocaine non-prereactive complex; and FIG. 2D depicts BChE (+)-cocaine prereactive complex.

FIG. 3A shows the (-)-cocaine rotation in the BChE active site for the non-prereactive complex to the prereactive complex hindered by some residues at positions Y332, A328, and F329 residues in the non-prereactive complexes which are significantly different from the corresponding positions in the prereactive complex; and FIG. 3B shows the (+)-cocaine rotation in the BChE active site where none of the aforementioned residues hinders the (+)-cocaine rotation in the BChE

active site from the non-prereactive complex to the prereactive complex in accordance with the present invention.

FIG. 4A depicts the (-)-cocaine rotation in the active site of A328W/Y332A; FIG. 4B depicts the (-)-cocaine rotation in the active site of A328W/Y332G BChE from the non-prereactive complex to the prereactive complex; and FIG. 4C depicts the (-)-cocaine rotation in the active site of wild-type BChE.

FIG. 5A is a plot of the key internuclear distances (in Å) versus the time in the simulated TS1 structure for (-)-cocaine hydrolysis catalyzed by A328W/Y332A; and FIG. 5B for A199S/A328W/Y332G BChE.

FIG. 6 is a plot of key internuclear distances (in Å) versus the time in the simulated TS1 structure for (-)-cocaine hydrolysis catalyzed by A199S/S287G/A328W/Y332G BChE.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention has two major improvements over the prior art. The first is the presently discovered BChE mutants, mutant 1, A199S/A328W/Y332G; mutant 2, A199S/F227A/A328W/Y332G; mutant 3, A199S/S287G/A328W/Y332G; mutant 4, A199S/F227A/S287G/A328W/Y332G; and mutant 5, A199S/F227A/S287G/A328W/E441D each have a significantly higher catalytic efficiency. The second improvement is concerning the mutant designing or discovering process.

The BChE mutants 1-5 have full length amino acid sequences, SEQ ID NOS: 2, 8, 14, 20, and 26, respectively, which are encoded by nucleic acid sequences having SEQ ID NOS: 1, 7, 13, 19, and 25, respectively. Table 1 summarizes the catalytic efficiency against (-)-cocaine for the five mutants.

In addition to the full length BChE mutants, the respective amino acid sequence can be truncated without substantially affecting the respective catalytic activity. With all mutants, residues 1-67 and 443-574 can be removed without substantially affecting the catalytic activity of the respective mutant BChE. Further, with regard to mutant 1-4, amino acids 1-116 and 439-574 can be omitted without substantially affecting its respective catalytic activity. With regard to mutant 5, amino acid residues 1-116 and 442-574 can be omitted without substantially affecting its catalytic activity. Table 1 also provides a summary of amino acid SEQ ID NOS and corresponding nucleic acid SEQ ID NOS for the aforementioned truncated mutant BChE sequences.

TABLE 1

Mutant Number	Amino Acid Substituting	Nucleic Acid SEQ ID NO.	Amino Acid SEQ ID NO.	Partially Truncated Nucleic Acid Sequence		SEQ ID NO		Catalytic Efficiency Against (-)-cocaine
				Corresponding To Amino Acid Residues 68-442	SEQ ID NO. for Amino Acid Residues 68-442	Corresponding To Amino Acid Residues 117-438/441*	SEQ ID NO. for Amino Acid Residues 117-438/441*	
1	A199S/A328W/Y332G	1	2	3	4	5	6	65-fold
2	A199S/F227A/A328W/Y332G	7	8	9	10	11	12	148-fold
3	A199S/S287G/A328W/Y332G	13	14	15	16	17	18	456-fold

TABLE 1-continued

Mutant Number	Amino Acid Substituting	Nucleic Acid SEQ ID NO.	Amino Acid SEQ ID NO.	Partially Truncated	SEQ ID NO. for Amino Acid Residues 68-442	SEQ ID NO	SEQ ID NO. for Amino Acid Residues 117-438/441*	Catalytic Efficiency Against (-)-cocaine
				Sequence Corresponding To Amino Acid Residues 68-442		Corresponding To Amino Acid Residues 117-438/441*		
4	A199S/F 227A/S2 87G/A32 8W/Y33 2G	19	20	21	22	22	24	1,003-fold
5	A199S/F 227A/S2 87G/A32 8W/E44 1D	25	26	27	28	29	30	445-fold

(*Amino acid residues 117-438 for mutants 1-4 and residues 117-441 for mutant 5.)

The BChE variant polypeptide, e.g., SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, and 30 can be formulated in a pharmaceutical composition along with a suitable pharmaceutical carrier known to one skilled in the art.

The present BChE variant polypeptides can be used in treating a cocaine-induced condition by administering to an individual, an effective amount of one of the BChE variant polypeptides, i.e., SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, and 30, to lower blood cocaine concentration. The BChE variant polypeptide may be administered in the form of a pharmaceutical composition in which the BChE variant is included with a suitable pharmaceutical carrier. Treatment of a cocaine induced condition using one of the aforementioned BChE variant polypeptides can be done in accordance with Zhan et al., page 2463.

The preferred dose for administration of a butyrylcholinesterase or peptide composition in accordance with the present invention is that amount which will be effective in lowering (-)-cocaine concentration in a patient's bloodstream, and one would readily recognize that this amount will vary greatly depending on the nature of cocaine consumed, e.g., injected or inhaled, and the condition of a patient. An "effective amount" of butyrylcholinesterase mutant or pharmaceutical agent to be used in accordance with the invention is intended to mean a nontoxic but sufficient amount of the agent, such that the desired prophylactic or therapeutic effect is produced. Thus, the exact amount of the enzyme or a particular agent that is required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the condition being treated, the particular carrier or adjuvant being used and its mode of administration, and the like. Similarly, the dosing regimen should also be adjusted to suit the individual to whom the composition is administered and will once again vary with age, weight, metabolism, etc. of the individual. Accordingly, the "effective amount" of any particular butyrylcholinesterase composition will vary based on the particular circumstances, and an appropriate effective amount may be determined in each case of application by one of ordinary skill in the art using only routine experimentation.

A unique method was used to determine potential BChE mutants with projected increased catalytic activity for the hydrolysis of cocaine. The method provides a unique approach which first models the potential BChE mutant interaction with cocaine followed by generating the BChE mutant if the predicted model indicates that the BChE variant should have enhanced catalytic activity. The method includes gener-

ating an initial structure of the transition state structure for the rate-determining step for the cocaine hydrolysis catalyzed by a possible BChE mutant. A sufficiently long time molecular dynamics simulation is performed on the transition state structure in water to have a stable molecular dynamics trajectory. The molecular dynamics trajectory is analyzed and the hydrogen binding energies are estimated between the carboxyl oxygen of the (-)-cocaine benzoyl ester and the oxyanion hole of the possible BChE mutant. If the overall hydrogen binding energy between the carboxyl oxygen of the (-)-cocaine benzoyl ester and the possible BChE mutant, in the transition state, is stronger than the overall hydrogen binding energy between the carboxyl oxygen of the (-)-cocaine benzoyl ester and the wild-type BChE, hybrid quantum mechanical/molecular mechanical (QM/MM) geometry optimization is performed to refine the molecular dynamics-simulated structure, the hydrogen binding energies are calculated and the energy barrier is evaluated. The QM/MM calculations make the computational predictions more reliable. Finally, the BChE mutant is generated.

With regard to the molecular dynamics (MD) simulations and quantum mechanical/molecular mechanical (QM/MM) calculations, the first chemical reaction step of (-)-cocaine hydrolysis catalyzed by butyrylcholinesterase (BChE) mutants and, when needed, other reaction steps are modeled and calculated using molecular dynamics simulations and QM/MM calculations. Following this modeling, mutant BChE's are created using site-directed mutagenesis followed by protein expression. The aforementioned five mutants were identified by computational analysis and generated by site-directed mutagenesis which have significantly enhanced (-)-cocaine hydrolysis catalytic efficiency compared with wild-type BChE.

In the present method computational analysis in the form of molecular modeling of a potential BChE mutant and MD simulations and QM/MM calculations provide virtual screening of possible BChE mutants which have predicted enhanced catalytic activity for (-)-cocaine. For example, the MD simulations and QM/MM calculations predict which mutation will lead to a more stable transition state for the rate determining step compared to the corresponding separated reagents, i.e., free cocaine and free possible mutant BChE, where the more stable transition state leads to a lower energy barrier and higher predicted catalytic efficiency. Only after the computational analysis predicts enhanced catalytic efficiency, is site-directed mutagenesis conducted on wild-type BChE nucleic acid sequence to generate a mutant nucleic acid sequence which is then used to express a mutant BChE protein. The

mutant BChE protein is then used in catalytic assays to determine the catalytic efficiency against (-)-cocaine.

The use of predictive, computational modeling of the present method for identifying mutant BChE candidates and the resulting mutant BChE are novel and unexpected over prior conventional methods which will now be readily apparent to one of ordinary skill in the art.

Using the present method, most discovered new mutants include a specific mutation (Y332G) on residue #332. No prior BChE mutant having the Y332G mutation had ever been reported previously; only mutations Y332A and Y332M on residue #332 had been tested previously by other researchers. Prior to the present invention, there was no reason to expect that a mutant including Y332G mutation should be better than the corresponding mutant including Y332A mutation or Y332M. Thus, the present mutants with a Y332G mutation which have enhanced catalytic activity represent a surprising and unexpected result over prior BChE mutants.

A Y332G mutant (single mutation) was first tested and found that the Y332G mutant had a slightly low (or approximately equal) catalytic efficiency than the wild-type. So, only an appropriate combination of different mutations on different residues could make the enzyme more active. As seen below, the prior art did not reveal that any of the particular combinations tested was expected to have an improved catalytic efficiency. The present method is based on the present unique, extensive computational modeling and simulations of the detailed catalytic mechanism for both the wild-type BChE and the mutants.

The primary improvement of the present method over the prior art is that high-performance computational modeling and simulations of the detailed catalytic mechanism are performed, which includes modeling how cocaine binds with BChE and the subsequent structural transformation and chemical reaction process. The prior art only considered the cocaine binding with the enzyme (BChE) and was unable to examine the detailed catalytic reaction process after the BChE-cocaine binding. When molecular modeling was limited to studying the BChE-cocaine binding, one could only design a mutation to improve the BChE-cocaine binding without knowing whether the mutation will also speedup the subsequent chemical reaction process or not.

To overcome the obstacles of prior challenges to using computational techniques such as molecular docking and molecular dynamics simulation previously used by others which were based on an empirical force field which cannot be used to perform necessary reaction coordinate calculations for the catalytic reaction process, a variety of state-of-the-art computational techniques of homology modeling, molecular docking, molecular dynamics, quantum mechanics (QM), and hybrid quantum mechanics/molecular mechanics (QM/MM) were used for the rational design of the BChE mutants. The combined use of these computational techniques, including QM and QM/MM, led to the study of the detailed reaction coordinate of the BChE-catalyzed hydrolysis of cocaine which, for the first time, provided the detailed structures of all transition states and intermediates existing in the reaction process and the corresponding energetics. These extensive computational modeling and simulation studies provided for the rational design of possible BChE mutants that not only can improve the BChE-cocaine binding, but also can speedup the subsequent chemical reaction process. As a result, one can now quickly discover the BChE mutants with the significantly improved catalytic efficiency.

In addition, to the differences mentioned above, the present molecular modeling of the BChE-cocaine binding also differs from the prior modeling. The molecular modeling in the prior

art considered only one binding mode for each BChE-cocaine system, without modeling the possible cocaine rotation in the BChE active site after the binding. The present method considers two different binding modes for each BChE-cocaine system and the structural transformation between them: non-prereactive and prereactive BChE-cocaine complexes. The present modeling provides more detailed information about the BChE-cocaine binding and the subsequent structural transformation.

The present method includes molecular dynamics simulations performed on the cocaine binding with both the wild-type BChE and the mutants whereas prior molecular dynamics simulations were only performed on the cocaine binding with the wild-type BChE. As is shown below in the following experiments, the computational prediction could be completely wrong without directly modeling and simulating cocaine binding with the proposed mutants.

The present invention will now be discussed with regard to the following non-limiting examples in the form of experiments which are provided to enhance understanding of the present invention but in no way limit its scope or applicability.

Experiment 1

Computational Study of Cocaine Binding with WILD-Type and Mutant BChE's for A328W/Y332G, A328W/Y332A, and A328W/Y332A/Y419S.

A detailed computational study of cocaine binding with wild-type and mutant BChE's starting from the available X-ray crystal structure of wild-type BChE was performed. The simulated mutants include A328W/Y332G, A328W/Y332A, and A328W/Y332A/Y419S, as simple geometric consideration of the binding site suggests that these mutations could be important for changing the (-)-cocaine rotation from the non-prereactive complex to the prereactive complex. Wet experimental tests were conducted on the catalytic activity of these mutants for (-)-cocaine in order to verify the computational predictions. All of the obtained results clearly demonstrate that molecular modeling and MD simulations of cocaine binding with BChE mutants provide a reliable computational approach to the rational design of high-activity mutants of BChE for the (-)-cocaine hydrolysis.

3D model of BChE. The initial coordinates of human BChE used in the computational studies came from the X-ray crystal structure deposited in the Protein Data Bank (pdb code: 1POP). The missing residues (D2, D3, E255, D378, D379, N455, L530, E531, and M532) in the X-ray crystal structure were built using the automated homology modeling tool Modeler disclosed by Sali, A.; Blundell, T. L. *J. Mol. Biol.* 1990, 212 403, and Sali, A.; Blundell, T. L. *J. Mol. Biol.* 1993, 234, 779, herein incorporated by reference, and InsightII software (Accelrys, Inc.) with the default parameters.

Molecular docking. Molecular docking was performed for each non-prereactive protein-ligand binding complex. The binding site was defined as a sphere with an approximately 15 Å radius around the active site residue S198. The amino acid residues included in the binding site model are not contiguous in the protein. Cocaine, considered as a ligand, was initially positioned at 17 Å in front of S198 of the binding site. Each BChE-cocaine binding complex was energy-minimized by using the steepest descent algorithm first until the maximum energy derivative is smaller than 4 kcal/mol/Å and then the conjugated gradient algorithm until the maximum energy derivative is smaller than 0.001 kcal/mol/Å. The energy mini-

mization was followed by a 300 ps molecular dynamics (MD) simulation at T=298 K with a time step of 1 fs. During the energy minimization and MD simulation, only cocaine and the residues of BChE included in the binding site were allowed to move, while the remaining part of the protein was fixed. The energy-minimization and MD simulation for these processes were performed by using the Amber force field implemented in the Discover_3/InsightII calculation engine, disclosed by Cornell, W. D.; Cieplak, P.; Bayly, C. I.; Gould, I. R.; Merz, Jr., K. M.; Ferguson, D. M.; Spellmeyer, D. C.; Fox, T.; Caldwell, J. W.; Kollman, P. A. *J. Am. Chem. Soc.* 1995, 117, 5179. The non-bonded cut-off method and the dielectric constant were set up to group based (12 Å cut-off distance) and distance dependent, respectively ($\epsilon=4r$) in accordance with Harvey, S. C. *Proteins* 1989, 5, 78-92, herein incorporated by reference.

Molecular dynamic simulation in water. The initial coordinates used in the MD simulation of the non-prereactive complexes were determined by using the molecular docking procedure described above, whereas the initial coordinates used in the MD simulation of the prereactive complexes were obtained from superimposing backbone of the X-ray crystal structure to that of the previously disclosed simulated prereactive complex of Zhan et al between cocaine and a homology model of wild-type BChE. Each BChE-cocaine binding complex was neutralized by adding two chloride counterions and was solvated in a rectangular box of TIP3P water molecules with a minimum solute-wall distance of 10 Å. The general procedure for carrying out the MD simulations in water is similar to that used in our previously reported other computational studies such as those in Zhan et al and (a) Zhan, C.-G.; Norberto de Souza, O.; Rittenhouse, R.; Ornstein, R. L. *J. Am. Chem. Soc.* 1999, 121, 7279, (b) Koca, J.; Zhan, C.-G.; Rittenhouse, R.; Ornstein, R. L. *J. Am. Chem. Soc.* 2001, 123, 817, (c) Koca, J.; Zhan, C.-G.; Rittenhouse, R. C.; Ornstein, R. L. *J. Comput. Chem.* 2003, 24, 368, herein all incorporated by reference. These simulations were performed by using the Sander module of Amber7 program as taught by Case, D. A.; Pearlman, D. A.; Caldwell, J. W.; Cheatham III, T. E.; Wang, J.; Ross, W. S.; Simmerling, C. L.; Darden, T. A.; Merz, K. M.; Stanton, R. V.; Cheng, A. L.; Vincent, J. J.; Crowley, M.; Tsui, V.; Gohlke, H.; Radmer, R. J.; Duan, Y.; Pitner, J.; Massova, I.; Seibel, G. L.; Singh, U. C.; Weiner, P. K.; Kollman, P. A. (2002), AMBER 7, University of California, San Francisco, herein incorporated by reference. The solvated system was optimized prior to the MD simulation. First, the protein-ligand was frozen and the solvent molecules with counterions were allowed to move during a 5000-step minimization with the conjugate gradient algorithm and a 5 ps MD run at T=300 K. After full relaxation and the entire solvated system was energy-minimized, the system was slowly heated from T=10 K to T=300 K in 30 ps before the production MD simulation for 500 ps. The full minimization and equilibration procedure was repeated for each mutant. The MD simulations were performed with a periodic boundary condition in the NPT ensemble at T=300 K with Berendsen temperature coupling and constant pressure (P=1 atm) with isotropic molecule-based scaling disclosed in Berendsen, H. C.; Postma, J. P. M.; van Gunsteren, W. F.; DiNola, A.; Haak, J. R. *J. Comp. Phys.* 1984, 81, 3684, herein incorporated by reference. The SHAKE algorithm of Ryckaert, J. P.; Ciccotti, G.; Berendsen, H. C. *J. Comp. Phys.* 1977, 23, 327 (herein incorporated by reference) was applied to fix all covalent bonds containing a hydrogen atom, a time step of 2 fs was used, and the non-bond pair list was updated every 10 steps. The pressure was adjusted by isotropic position scaling. The particle mesh Ewald (PME) method of Essmann, U.; Perera, L.; Berkowitz,

M. L.; Darden, T. A.; Lee, H.; Pedersen; L. G. *J. Chem. Phys.* 1995, 98, 10089, herein incorporated by reference, was used to treat long-range electrostatic interactions. A residue-based cutoff of 10 Å was applied to the noncovalent interactions. During the 500 ps production MD simulation, the coordinates of the simulated complex were saved every 1 ps.

Molecular docking and MD simulation procedures described above were performed to study cocaine binding with wild-type BChE and three mutants, i.e., A328W/Y332A, A328W/Y332A/Y419S, and A328W/Y332G. For each protein system (wild-type or mutant BChE), the protein binding with cocaine was considered in both the non-prereactive and prereactive enzyme-substrate complexes.

Most of the MD simulations in water were performed on a supercomputer, Superdome (shared-memory, with 4 nodes and 256 processors), at the Center for Computational Sciences, University of Kentucky. The other computations were carried out on SGI Fuel workstations and a 34-processors IBM x335 Linux cluster.

Experimental procedure. Site-directed mutagenesis of human BChE cDNA was performed by the QuikChange method of Braman, J.; Papworth, C.; Greener, A. *Methods Mol. Biol.* 1996, 57, 5731, herein incorporated by reference. Mutations were generated from wild-type human BChE in a pRc/CMV expression plasmid in accordance with Xie, W.; Altamirano, C. V.; Bartels, C. F.; Speirs, R. J.; Cashman, J. R.; Lockridge, O. *Mol. Pharmacol.* 1999, 55, 83, all herein incorporated by reference, kindly provided by Dr. Lockridge at University of Nebraska Medical Center. Using plasmid DNA as template and primers with specific base-pair alterations, mutations were made by polymerase chain reaction with Pfu DNA polymerase, for replication fidelity. The PCR product was treated with Dpn I endonuclease to digest the parental DNA template. Modified plasmid DNA was transformed into *Escherichia coli*, amplified, and purified. The DNA sequences of the mutants were confirmed by DNA sequencing. BChE mutants were expressed in human embryonic kidney cell line 293T/17. Cells were grown to 80-90% confluence in 6-well dishes and then transfected by Lipofectamine 2000 complexes of 4 µg plasmid DNA per each well. Cells were incubated at 37° C. in a CO₂ incubator for 24 hours and cells were moved to 60-mm culture vessel and cultured for four more days. The culture medium [10% fetal bovine serum in Dulbecco's modified Eagle's medium (DMEM)] was harvested for a BChE activity assay. To measure cocaine and benzoic acid, the product of cocaine hydrolysis by BChE, we used sensitive radiometric assays based on toluene extraction of [³H]cocaine labeled on its benzene ring were used in accordance with Masson, P.; Xie, W.; Froment, M.-T.; Levitsky, V.; Fortier, P.-L.; Albaret, C.; Lockridge, O. *Biochim. Biophys. Acta* 1999, 1433, 281, herein incorporated by reference. In brief, to initiate reactions, 100 nCi of [³H]cocaine was mixed with 100 µl of culture medium. Reactions proceeded at 37° C. for varying times. Reactions were stopped by adding 300 µl of 0.02 M HCl, which neutralized the liberated benzoic acid while ensuring a positive charge on the residual cocaine. [³H]benzoic acid was extracted by 1 ml of toluene and measured by scintillation counting. Finally, the measured time-dependent radiometric data were fitted to the kinetic equation so that the catalytic efficiency (k_{cat}/K_M) was determined.

Depicted in FIG. 1 are plots of some important distances in the MD-simulated (–)–cocaine binding with A328W/Y332G BChE versus the simulation time, along with root-mean-square deviation (RMSD) of the coordinates of backbone atoms in the simulated structure from those in the X-ray crystal structure. MD trajectories for other complexes were similar to these two in FIG. 1, although the simulated average

15

distances are different. Summarized in Table 2 are the average values of some important geometric parameters in the simulated complexes.

TABLE 2

BChE-cocaine binding ^a	Average values of the geometric parameters ^c						RMSD ^d	
	<D1> _{non}	D1>	D2>	D3>	D4>	Θ>	r onpre	re
wild-type	5						1	
	.60	.27	.77	.71	.37	67	.14	.27
wild-type with (+)-cocaine ^b	7						1	
A328W/Y332A	.64	.69	.88	.30	.83	61	.15	.13
	.11	.87	.30	.14	.01	51	.58	1.65
A328W/Y332G	7						1	
	.06	.96	.28	.52	.42	60	.20	.35
A328W/Y332A/	5						2	
Y419S	.18	.84	.64	.56	.97	64	.66	.62


^aRefers to (-)-cocaine binding with wild-type human BChE or (-)-cocaine binding with a mutant BChE, unless indicated otherwise.

^bRefers to (+)-cocaine binding with wild-type human BChE.

^c<D1>_{non} and <D1> represent the average distances between the S198 O^γ atom and the carbonyl carbon of the cocaine benzoyl ester in the simulated non-prereactive and prereactive BChE-cocaine complexes, respectively. <D2>, <D3>, <D4> refer to the average values of the simulated distances from the carbonyl oxygen of the cocaine benzoyl ester to the NH hydrogen atoms of G116, G117, and A199 residues, respectively. <Θ> is the average value of the dihedral angle formed by the S198 O^γ atom and the plane of the carboxylate group of the cocaine benzoyl ester. See Scheme 2.

^dThe root-mean-square deviation (RMSD) of the coordinates of backbone atoms in the simulated structure from those in the X-ray crystal structure of BChE. "nonpre" and "pre" refer to the non-prereactive and prereactive BChE-cocaine complexes, respectively.

(-)- and (+)-cocaine binding with wild-type BChE. FIG. 2 shows the binding structures of the simulated non-prereactive and prereactive complexes of wild-type BChE binding with the two enantiomers of cocaine. In the non-prereactive complexes with (-)- and (+)-cocaine, the methyl ester group of cocaine is positioned at the top of the H438 backbone, while the cocaine benzoyl ester moiety is quasi-parallel to the C—O^γ side chain of S198 with a dihedral angle Θ of -8° and 140°, respectively.



Scheme 2. Hydrolysis of (-)-cocaine and (+)-cocaine.

Here, Θ refers to the dihedral angle formed by S198 O^γ and the plane of carboxylate group of the cocaine benzoyl ester as shown in the structure diagram below.

The simulated internuclear distances between the carbonyl oxygen of cocaine benzoyl ester group and the NH hydrogen of G116, G117, and A199 are comparable for the two enantiomers. The simulated average distances between the carbonyl carbon of the benzoyl ester and S198 O^γ are 5.60 Å and

16

5.18 Å for (-)- and (+)-cocaine, respectively. Comparing the simulated protein backbone structures to the X-ray crystal structure of Nicolet et al, one can see from FIG. 1 that the RMSD values are all smaller than 1.3 Å for the whole protein structures.

The MD simulations of the prereactive complexes reveal that wild-type BChE binding with (-)-cocaine is essentially the same as the binding with (+)-cocaine in the binding site, except for the different positions of methyl ester group of the substrates. The simulated average distances between the carbonyl carbon of the benzoyl ester and S198 O^γ are 3.27 and 3.69 Å for (-)-cocaine and (+)-cocaine, respectively. Moreover, the (+)-cocaine is stabilized more effectively by the formation of strong hydrogen bonds with the backbone NH of residues G116, G117, and A199 as summarized above in Table 2. The cocaine benzoyl ester moiety is positioned quasi-perpendicular to S198 C—O^γ with a dihedral angle Θ of ~67° and ~61° for (-)- and (+)-cocaine, respectively.

A comparison was made between the currently simulated structures of the BChE-cocaine binding with those simulated previously by using a homology model of BChE and it was noted that two major differences between the two sets of structures. By using the X-ray crystal structure in accordance with Nicolet et al, the acyl loop is positioned on the top of the cocaine benzoyl ester moiety of the cocaine, whereas the acyl loop is far from the cocaine benzoyl ester moiety in the structure simulated starting from the homology model of Zhan et al. The RMSD of the coordinates of backbone atoms in the previously simulated prereactive BChE(-)-cocaine complex from those in the X-ray crystal structure of BChE is ~2.0 Å for the entire protein and ~3.0 Å for the acyl loop. The RMSD value became ~2.4 Å for the entire protein and ~3.3 Å for the acyl loop, when the X-ray crystal structure was replaced by the MD-simulated prereactive BChE(-)-cocaine complex starting from the X-ray crystal structure. Despite these structural differences, the benzoyl ester group of the ligand is still close to the key residues (S197, G116, and G117) in the BChE binding site. Some significant differences are associated with the distances between the S198 O^γ atom and the carbonyl carbon of the cocaine benzoyl ester in non-prereactive complexes. The average values of this distance in the non-prereactive complexes were ~9.5 and ~8.5 Å for (-)- and (+)-cocaine, respectively, when a homology model was used. Using the X-ray crystal structure to conduct the analysis, corresponding average values became ~5.6 and ~5.2 Å, respectively. Therefore, both (-)- and (+)-cocaine became closer to the binding site when the homology model was replaced by the X-ray crystal structure. However, no significant changes of the binding in the prereactive complexes were observed when the used homology model was replaced by the X-ray crystal structure. The average values of the simulated distance between the S198 O^γ atom and the carbonyl carbon of the cocaine benzoyl ester in the prereactive complexes are always close to ~3.5 Å for both (-)- and (+)-cocaine no matter whether the X-ray crystal structure or homology model of BChE was used as the starting structure. The similar computational results obtained from the use of the X-ray crystal structure and homology model of BChE provides evidence that the fundamental structural and mechanistic insights obtained from the previous computational studies of Zhan et al are reliable, despite the previous simulations were performed by using the homology model when the X-ray crystal structure was not available.

Further, the simulated structures of the non-prereactive BChE-cocaine complexes were superimposed with the corresponding prereactive complexes. As shown in FIG. 3, the (-)-cocaine rotation in the BChE active site from the non-

prereactive complex to the prereactive complex is hindered by some residues as the positions of Y332, A328, and F329 residues in the non-prereactive complex are significantly different from the corresponding positions in the prereactive complex, whereas none of these residues hinders the (+)-cocaine rotation in the BChE active site from the non-prereactive complex to the prereactive complex because these residues stay in nearly the same positions in the two BChE-(+)-cocaine complexes.

(-)-cocaine binding with BChE mutants. Now that the (-)-cocaine rotation from the non-prereactive complex to the prereactive complex has been known to be the rate-determining step of the BChE-catalyzed hydrolysis of (-)-cocaine as shown by Zhan et al, useful BChE mutants should be designed to specifically accelerate the change from the non-prereactive BChE(-)-cocaine complex to the prereactive complex. The question is whether MD simulation can be performed to help design BChE mutants that have higher catalytic activity for (-)-cocaine hydrolysis.

In the simulated non-prereactive complex, the average distance between the carbonyl carbon of cocaine benzoyl ester and S198 O^γ is 7.6 Å for A328W/Y332A BChE and 7.1 Å for A328W/Y332G BChE, as seen in Table 2 above. In the simulated prereactive complex, the average values of this important internuclear distance become 3.87 and 3.96 Å for A328W/Y332A and A328W/Y332G BChE's, respectively. Compared to the simulated wild-type BChE(-)-cocaine prereactive complex, the average distances between the carbonyl carbon of the cocaine benzoyl ester and S198 O^γ in the prereactive complex of (-)-cocaine with A328W/Y332A and A328W/Y332G BChE's are all slightly longer, whereas the average distances between the carbonyl oxygen of the cocaine benzoyl ester and the NH of G116, G117, and A199 residues are all shorter. This provides evidence that (-)-cocaine more strongly bind with A328W/Y332A and A328W/Y332G BChE's in the prereactive complexes. More importantly, the (-)-cocaine rotation in the active site of A328W/Y332A and A328W/Y332G BChE's from the non-prereactive complex to the prereactive complex did not cause considerable changes of the positions of A332 (or G332), W328, and F329 residues as seen in FIG. 4, compared to the (-)-cocaine rotation in the active site of wild-type BChE. These results provide evidence that A328W/Y332A and A328W/Y332G BChE's should be associated with lower energy barriers than the wild-type for the (-)-cocaine rotation from the non-prereactive complex to the prereactive complex. Further, (-)-cocaine binding with A328W/Y332G BChE is very similar to the binding with A328W/Y332A BChE, but the position change of F329 residue caused by the (-)-cocaine rotation was significant only in A328W/Y332A BChE, thus suggesting that the energy barrier for the (-)-cocaine rotation in A328W/Y332G BChE should be slightly lower than that in A328W/Y332A BChE.

Concerning (-)-cocaine binding with A328W/Y332A/Y419S BChE, Y419 stays deep inside the protein and does not directly contact with the cocaine molecule. The Y419S mutation was tested because it was initially expected that this mutation would further increase the free space of the active site pocket so that the (-)-cocaine rotation could be easier. However, as seen in Table 2 above, the average distance between the carbonyl carbon of cocaine benzoyl ester and S198 O^γ atom in the simulated prereactive complex was as long as 5.84 Å. The average distances between the carbonyl oxygen of the cocaine benzoyl ester and the NH hydrolysis atoms of G116, G117, and A199 residues are between 4.56 and 6.97 Å; no any hydrogen bond between them. In addition to the internuclear distances, another interesting geometric

parameter is the dihedral angle, Θ , formed by S198 O^γ and the plane of the carboxylate group of the cocaine benzoyl ester. As seen in Table 2, the Θ values in the prereactive complexes of cocaine with wild-type BChE and all of the BChE mutants other than A328W/Y332A/Y419S BChE all slightly deviate from the ideal value of 90° for the nucleophilic attack of S198 O^γ at the carbonyl carbon of cocaine. The Θ value in the prereactive complex of (-)-cocaine with A328W/Y332A/Y419S BChE is 164°, which is considerably different from the ideal value of 90°.

Catalytic activity. The aforementioned discussion provides evidence that the energy barriers for the (-)-cocaine rotation in A328W/Y332A and A328W/Y332G BChE's from the non-prereactive complex to the prereactive complex, the rate-determining step for the BChE-catalyzed hydrolysis of (-)-cocaine, should be lower than that in the wild-type BChE. Thus, the MD simulations predict that both A328W/Y332A and A328W/Y332G BChE's should have a higher catalytic activity than the wild-type BChE for (-)-cocaine hydrolysis. Further, the MD simulations also suggest that the energy barrier for the (-)-cocaine rotation in A328W/Y332G BChE should be slightly lower than that in A328W/Y332A BChE and, therefore, the catalytic activity of A328W/Y332G BChE for the (-)-cocaine hydrolysis should be slightly higher than the activity of A328W/Y332A BChE. In addition, the MD simulations predict that A328W/Y332A/Y419S BChE should have no catalytic activity, or have a considerably lower catalytic activity than the wild-type, for (-)-cocaine hydrolysis because (-)-cocaine binds with the mutant BChE in a way that is not suitable for the catalysis.

The catalytic efficiency (k_{cat}/K_M) of A328W/Y332A BChE for (-)-cocaine hydrolysis was reported to be 8.56×10^6 M min⁻¹ which is 9.39 times of the k_{cat}/K_M value (9.11×10^5 M min⁻¹) of the wild-type BChE. To examine these theoretical predictions of the relative activity for A328W/Y332G and A328W/Y332A/Y419S BChE's, a A328W/Y332A, A328W/Y332G, and A328W/Y332A/Y419S BChE was produced through site-directed mutagenesis. To minimize the possible systematic experimental errors of the kinetic data, kinetic studies were performed with all of three mutants under the same condition and compared the catalytic efficiency of the A328W/Y332G and A328W/Y332A/Y419S to that of the A328W/Y332A for (-)-cocaine hydrolysis at benzoyl ester group. Based on the kinetic analysis of the measured time-dependent radiometric data, the ratio of the k_{cat}/K_M value of A328W/Y332G BChE to the k_{cat}/K_M value of A328W/Y332A BChE for the (-)-cocaine hydrolysis was determined to be ~2.08, or A328W/Y332G BChE has a k_{cat}/K_M value of $\sim 1.78 \times 10^7$ M min⁻¹ for the (-)-cocaine hydrolysis. The radiometric data show no significant catalytic activity for A328W/Y332A/Y419S BChE. These experimental data are consistent with the theoretical predictions based on the MD simulations.

Conclusion. Molecular modeling, molecular docking, and molecular dynamics (MD) simulations were performed to study cocaine binding with human butyrylcholinesterase (BChE) and its mutants, based on a recently reported X-ray crystal structure of human BChE. The MD simulations of cocaine binding with wild-type BChE led to average BChE-cocaine binding structures similar to those obtained recently from the MD simulations based on a homology model of BChE, despite the significant difference found at the acyl binding pocket. This confirms the fundamental structural and mechanistic insights obtained from the prior computational studies of Zhan et al based on a homology model of BChE, e.g., the rate-determining step for BChE-catalyzed hydrolysis of biologically active (-)-cocaine is the (-)-cocaine rotation

19

in the BChE active site from the non-prereactive BChE-(-)-cocaine complex to the prereactive complex.

The MD simulations further reveal that the (-)-cocaine rotation in the active site of wild-type BChE from the non-prereactive complex to the prereactive complex is hindered by some residues such that the positions of Y332, A328, and F329 residues in the non-prereactive complex are significantly different from those in the prereactive complex. Compared to (-)-cocaine binding with wild-type BChE, (-)-cocaine more strongly bind with A328W/Y332A and A328W/Y332G BChE's in the prereactive complexes. More importantly, the (-)-cocaine rotation in the active site of A328W/Y332A and A328W/Y332G BChE's from the non-prereactive complex to the prereactive complex did not cause considerable changes of the positions of A332 or G332, W328, and F329 residues. These results provide evidence that A328W/Y332A and A328W/Y332G BChE's are associated with lower energy barriers than wild-type BChE for the (-)-cocaine rotation from the non-prereactive complex to the prereactive complex. Further, (-)-cocaine binding with A328W/Y332G BChE is very similar to the binding with A328W/Y332A BChE, but the position change of F329 residue caused by the (-)-cocaine rotation was significant only in A328W/Y332A BChE, thus suggesting that the energy barrier for (-)-cocaine rotation in A328W/Y332G BChE should be slightly lower than that in A328W/Y332A BChE. It has also been demonstrated that (-)-cocaine binds with A328W/Y332A/Y419S BChE in a way that is not suitable for the catalysis.

Based on the computational results, both A328W/Y332A and A328W/Y332G BChE's have catalytic activity for (-)-cocaine hydrolysis higher than that of wild-type BChE and


20

the activity of A328W/Y332G BChE should be slightly higher than that of A328W/Y332A BChE, whereas A328W/Y332A/Y419S BChE is expected to lose the catalytic activity. The computational predictions are completely consistent with the experimental kinetic data, providing evidence that the used computational protocol, including molecular modeling, molecular docking, and MD simulations, is reliable in prediction of the catalytic activity of BChE mutants for (-)-cocaine hydrolysis.




Experiment 2

MD Simulations and Quantum Mechanical/Molecular Mechanical (QM/MM) Calculations Relating to a 199S/A328W/Y332G Mutant (Mutant 1) (SEQ ID NO: 1).


Generally speaking, for rational design of a mutant enzyme with a higher catalytic activity for a given substrate, one needs to design a mutation that can accelerate the rate-determining step of the entire catalytic reaction process while the other steps are not slowed down by the mutation. Reported computational modeling and experimental data indicated that the formation of the prereactive BChE-(-)-cocaine complex (ES) is the rate-determining step of (-)-cocaine hydrolysis catalyzed by wild-type BChE as disclosed by Sun et al, Zhan et al and Hamza, A.; Cho, H.; Tai, H.-H.; Zhan, C.-G. *J. Phys. Chem. B* 2005, 109, 4776, herein incorporated by reference, whereas the rate-determining step of the corresponding (+)-cocaine hydrolysis is the chemical reaction process consisting of four individual reaction steps disclosed by Zhan et al and shown in Scheme 3 and Scheme 4 below.



-continued



-continued



Scheme 3. Schematic representation of BChE-catalyzed hydrolysis of (-)-cocaine. Only QM-treated high-layer part of the reaction system in the QM/MM calculations are drawn. Notation [H] refers to a non-hydrogen atom in the MM-treated low-layer part of the protein and the cut covalent bond with this atom is saturated by a hydrogen atom. The dash lines in the transition state structures represent the transition bonds.

This mechanistic understanding is consistent with the experimental observation of Sun et al, that the catalytic rate constant of wild-type BChE against (+)-cocaine is pH-dependent, whereas that of the same enzyme against (-)-cocaine is independent of the pH. The pH-dependence of the rate constant for (+)-cocaine hydrolysis is clearly associated with the protonation of H438 residue in the catalytic triad (S198, H438, and E325). For the first and third steps of the reaction process, when H438 is protonated, the catalytic triad cannot function and, therefore, the enzyme becomes inactive. The lower the pH of the reaction solution is, the higher the concentration of the protonated H438 is, and the lower the concentration of the active enzyme is. Hence, the rate constant was found to decrease with decreasing the pH of the reaction solution for the enzymatic hydrolysis of (+)-cocaine.

Based on the above mechanistic understanding, the efforts for rational design of the BChE mutants reported in literature have been focused on how to improve the ES formation process. Indeed, several BChE mutants, including A328W, A328W/Y332A, A328W/Y332G, and F227A/S287G/A328W/Y332M, have been found to have a significantly higher catalytic efficiency (k_{cat}/K_M) against (-)-cocaine; these mutants of BChE have an approximate 9 to 34-fold improved catalytic efficiency against (-)-cocaine. Experimental observation also indicated that the catalytic rate constant of A328W/Y332A BChE is pH-dependent for both (-)- and (+)-cocaine. The pH-dependence reveals that for both (-)- and (+)-cocaine, the rate-determining step of the hydrolysis catalyzed by A328W/Y332A BChE should be either the first or the third step of the reaction process. Further, if the third step were rate determining, then the catalytic efficiency of the A328W/Y332A mutant against (-)-cocaine should be as high as that of the same mutant against (+)-cocaine because the (-)- and (+)-cocaine hydrolyses share the same third and fourth steps (see Scheme 3). However, it has also been observed that the A328W/Y332A mutant only has a ~9-fold improved catalytic efficiency against (-)-cocaine, whereas the A328W/Y332A mutation does not change the high catalytic activity against (+)-cocaine. This analysis of the experimental and computational data available in literature clearly shows that the rate-determining step of (-)-cocaine hydrolysis catalyzed by the A328W/Y332A mutant should be the first step of the chemical reaction process. Further, recently reported computational modeling also suggests that the formation of the prereactive BChE-(-)-cocaine complex (ES) is hindered mainly by the bulky side chain of Y332 residue in wild-type BChE, but the hindering can be removed by the Y332A or Y332G mutation. Therefore, starting from the A328W/Y332A or A328W/Y332G mutant, the truly rational design of further mutation(s) to improve the catalytic efficiency of BChE against (-)-cocaine should aim to decrease the energy barrier for the first reaction step without significantly affecting the ES formation and other chemical reaction steps.

The following rational design of a high-activity mutant of BChE against (-)-cocaine is based on detailed computational modeling of the transition state for the rate-determining step (i.e., the first step of the chemical reaction process). Molecular dynamics (MD) simulations and hybrid quantum mechanical/molecular mechanical (QM/MM) calculations

were performed to model the protein environmental effects on the stabilization of the transition-state structure for BChE-catalyzed hydrolysis of (-)-cocaine. The simulated and calculated results indicate that the transition-state structure can be stabilized better by the protein environment in A199S/A328W/Y332G mutant of BChE than that in the wild-type. The computational modeling led to a prediction of the higher catalytic efficiency for the A199S/A328W/Y332G mutant against (-)-cocaine. The prediction has been confirmed by wet experimental tests showing that the A199S/A328W/Y332G mutant has a significantly improved catalytic efficiency against (-)-cocaine. All of the obtained results clearly demonstrate that directly modeling the transition-state structure provides a reliable computational approach to the rational design of a high-activity mutant of BChE against (-)-cocaine.

MD simulations. It should be stressed that a critical issue exists with regard to any MD simulation on a transition state. In principle, MD simulation using a classical force field (molecular mechanics) can only simulate a stable structure corresponding to a local minimum on the potential energy surface, whereas a transition state during a reaction process is always associated with a first-order saddle point on the potential energy surface. Hence, MD simulation using a classical force field cannot directly simulate a transition state without any restraint on the geometry of the transition state. Nevertheless, if one can technically remove the freedom of imaginary vibration in the transition state structure, then the number of vibrational freedoms (normal vibration modes) for a nonlinear molecule will decrease from $3N-6$. The transition state structure is associated with a local minimum on the potential energy surface within a subspace of the reduced vibrational freedoms, although it is associated with a first-order saddle point on the potential energy surface with all of the $3N-6$ vibrational freedoms. Theoretically, the vibrational freedom associated with the imaginary vibrational frequency in the transition state structure can be removed by appropriately freezing the reaction coordinate. The reaction coordinate corresponding to the imaginary vibration of the transition state is generally characterized by a combination of some key geometric parameters. These key geometric parameters are bond lengths of the forming and breaking covalent bonds for BChE-catalyzed hydrolysis of cocaine, as seen in Scheme 3. Thus, one just needs to maintain the bond lengths of the forming and breaking covalent bonds during the MD simulation on a transition state. Technically, one can maintain the bond lengths of the forming and breaking covalent bonds by simply fixing all atoms within the reaction center, by using some constraints on the forming and breaking covalent bonds, or by redefining the forming and breaking covalent bonds. It should be pointed out that the only purpose of performing such type of MD simulation on a transition state is to examine the dynamic change of the protein environment surrounding the reaction center and the interaction between the reaction center and the protein environment. For this study, of interest is the simulated structures, as the total energies calculated in this way are meaningless.

The initial BChE structures used in the MD simulations were prepared based on the previous MD simulation in accordance with Hamza et al on the prereactive ES complex for wild-type BChE with (-)-cocaine in water by using Amber7 program package. The previous MD simulations on the prereactive BChE-(-)-cocaine complex (ES) started from the X-ray crystal structure of Nicolet et al deposited in the Protein Data Bank (pdb code: 1P0P). The present MD simulation on the transition state for the first step (TS1) was performed in such a way that bond lengths of the partially formed and


partially broken covalent bonds in the transition state were all constrained to be the same as those obtained from our previous ab initio reaction coordinate calculations on the model reaction system of wild-type BChE in accordance with Zhan et al. The partially formed and partially broken covalent bonds in the transition state will be called "transition" bonds below, for convenience. A sufficiently long MD simulation with the transition bonds constrained should lead to a reasonable protein environment stabilizing the reaction center in the transition-state structure simulated. Further, the simulated TS1 structure for wild-type BChE with (-)-cocaine was used to build the initial structures of TS1 for the examined BChE mutants with (-)-cocaine; only the side chains of mutated residues needed to be changed.

The partial atomic charges for the non-standard residue atoms, including cocaine atoms, in the TS1 structures were calculated by using the RESP protocol implemented in the Antechamber module of the Amber7 package following electrostatic potential (ESP) calculations at ab initio HF/6-31G* level using Gaussian03 program known in the art. The geometries used in the ESP calculations came from those obtained from the previous ab initio reaction coordinate calculations of Zhan et al, but the functional groups representing the oxyanion hole were removed. Thus, residues G116, G117, and A199 were the standard residues as supplied by Amber7 in the MD simulations. The general procedure for carrying out the MD simulations in water is essentially the same as that used in our previously reported other computational studies including Zhan et al, Hamza et al and (a) Zhan, C.-G.; Norberto de Souza, O.; Rittenhouse, R.; Ornstein, R. L. *J. Am. Chem. Soc.* 1999, 121, 7279, (b) Koca, J.; Zhan, C.-G.; Rittenhouse, R.; Ornstein, R. L. *J. Am. Chem. Soc.* 2001, 123, 817, (c) Koca, J.; Zhan, C.-G.; Rittenhouse, R. C.; Ornstein, R. L. *J. Comput. Chem.* 2003, 24, 368, (d) Hamza, A.; Cho, H.; Tai, H.-H.; Zhan, C.-G. *Bioorg. Med. Chem.* 2005, 13, 4544, herein all incorporated by reference. Each aforementioned starting TS1 structure was neutralized by adding chloride counterions and was solvated in a rectangular box of TIP3P water molecules with a minimum solute-wall distance of 10 Å as described by Jorgensen, W. L.; Chandrasekhar, J.; Madura, J.; Klein, M. L. *J. Chem. Phys.* 1983, 79, 926, herein incorporated by reference. The total numbers of atoms in the solvated protein structures for the MD simulations are nearly 70,000, although the total number of atoms of BChE and (-)-cocaine is only 8417 (for the wild-type BChE). All of the MD simulations were performed by using the Sander module of Amber7 package. The solvated systems were carefully equilibrated and fully energy minimized. These systems were gradually heated from T=10 K to T=298.15 K in 30 ps before running the MD simulation at T=298.15 K for 1 ns or longer, making sure that a stable MD trajectory was obtained for each of the simulated TS1 structures. The time step used for the MD simulations was 2 fs. Periodic boundary conditions in the NPT ensemble at T=298.15 K with Berendsen temperature coupling and P=1 atm with isotropic molecule-based scaling were applied. The SHAKE algorithm was used to fix all covalent bonds containing hydrogen atoms. The non-bonded pair list was updated every 10 steps. The particle mesh Ewald (PME) method in accordance with Essmann, U.; Perera, L.; Berkowitz, M. L.; Darden, T. A.; Lee, H.; Pedersen, L. G. *J. Chem. Phys.* 1995, 98, 10089, herein incorporated by reference, was used to treat long-range electrostatic interactions. A

residue-based cutoff of 10 Å was utilized to the non-covalent interactions. The coordinates of the simulated systems were collected every 1 ps during the production MD stages.

QM/MM calculations. For each TS1 structure examined, after the MD simulation was completed and a stable MD trajectory was obtained, all of the collected snapshots of the simulated structure, excluding those before the trajectory was stabilized, were averaged and further energy-minimized. The energy-minimized average structure (with the transition bonds constrained) was used as an initial geometry to carry out a further geometry optimization by using the ONIOM approach of Dapprich, S.; Komaromi, I.; Byun, K. S.; Morokuma, K.; Frisch, M. J. *J. Mol. Struct. (Theochem)* 1999, 461, 1-21, herein incorporated by reference, implemented in the Gaussian03 program of Frisch, M. J. et al *Gaussian 03*, Revision A.1, Gaussian, Inc., Pittsburgh, Pa., 2003, herein incorporated by reference. Two layers were defined in the present ONIOM calculation: the high layer, as depicted in Scheme 3 above, was calculated quantum mechanically at the ab initio HF/3-21G level, whereas the low layer was calculated molecular mechanically by using the Amber force field as used in our MD simulation and energy minimization with the Amber7 program. The ONIOM calculations at the HF/3-21G: Amber level in this study are a type of QM/MM calculations of Vreven, T.; Morokuma, K. *J. Chem. Phys.* 2000, 113, 2969-2975 and Frisch, M.; Vreven, T.; Schlegel, H. B.; Morokuma, K. *J. Comput. Chem.* 2003, 24, 760-769, herein both incorporated by reference. Previous reaction coordinate calculations with an active site model of wild-type BChE demonstrate that the HF/3-21G level is adequate for the geometry optimization of this enzymatic reaction system shown by Zhan et al, although the final energy calculations for calculating the energy barriers must be carried out at a higher level. As depicted in Scheme 3, for all of these QM/MM calculations, the same part of the enzyme was included in the QM-treated high layer. So, the QM-treated high layer included (-)-cocaine, key functional groups from the catalytic triad (S198, H438, and E325), and the three residues, i.e., G116, G117, and A199 (or S199 for a particular mutant, see Scheme 4), of the possible oxyanion hole, whereas the entire enzyme structure of BChE was included in the MM-treated low layer. A language computer program was developed to automatically generate the input files for the ONIOM calculations following the MD simulations and subsequent energy minimizations in order to make sure that the atom types used for all low-layer atoms are the same as what were used in the Amber7. During the TS1 geometry optimization using the two-layer ONIOM, the length of a key transition C—O bond was fixed which dominates the reaction coordinate; all of the other transition bond lengths were relaxed. The C and O atoms in the key transition C—O bond are the carbonyl carbon of (-)-cocaine benzoyl ester and the O' atom of S198, respectively, according to the previous reaction coordinate calculations with an active site model of wild-type BChE of Zhan et al.

Most of the MD simulations and QM/MM calculations were performed in parallel on an HP supercomputer (Superdome, with 256 shared-memory processors) at the Center for Computational Sciences, University of Kentucky. Some of the computations were carried out on a 34-processors IBM x335 Linux cluster and SGI Fuel workstations.



Scheme 4. Schematic representation of the first reaction step for (-)-cocaine hydrolysis catalyzed by a BChE mutant with an A199S mutation.

Experimental materials. Cloned pfu DNA polymerase and Dpn I endonuclease were obtained from Stratagene (La Jolla, Calif.). ³H-(-)-cocaine (50 Ci/mmol) was purchased from PerkinElmer Life Sciences (Boston, Mass.). The expression plasmid pRc/CMV was a gift from Dr. O. Lockridge, Univer-

sity of Nebraska Medical Center (Omaha, Nebr.). All oligonucleotides were synthesized by the Integrated DNA Technologies, Inc. The QIAprep Spin Plasmid Miniprep Kit and Qiagen plasmid purification kit and QIAquick PCR purification kit were obtained from Qiagen (Santa Clarita, Calif.). Human embryonic kidney 293T/17 cells were from ATCC (Manassas, Va.). Dulbecco's modified Eagle's medium (DMEM) was purchased from Fisher Scientific (Fairlawn,

N.J.). Oligonucleotide primers were synthesized by the Integrated DNA Technologies and Analysis Facility of the University of Kentucky. 3,3',5,5'-Tetramethylbenzidine (TMB) was obtained from Sigma (Saint Louis, Mo.). Anti-butyrylcholinesterase (mouse monoclonal antibody, Product # HAH002-01) was purchased from AntibodyShop (Gentofte, Denmark) and Goat anti-mouse IgG HRP conjugate from Zymed (San Francisco, Calif.).

Site-directed mutagenesis, protein expression, and BChE activity assay. Site-directed mutagenesis of human BChE cDNA was performed by using the QuikChange method of Braman et al. Mutations were generated from wild-type human BChE in a pRc/CMV expression plasmid in accordance with (a) Masson, P.; Legrand, P.; Bartels, C. F.; Froment, M.-T.; Schopfer, L. M.; Lockridge, O. *Biochemistry* 1997, 36, 2266, (b) Masson, P.; Xie, W.; Froment, M.-T.; Levitsky, V.; Fortier, P.-L.; Albaret, C.; Lockridge, O. *Biochim. Biophys. Acta* 1999, 1433, 281, (c) Xie, W.; Altamirano, C. V.; Bartels, C. F.; Speirs, R. J.; Cashman, J. R.; Lockridge, O. *Mol. Pharmacol.* 1999, 55, 83, (d) Duysen, E. G.; Bartels, C. F.; Lockridge, O. *J. Pharmacol. Exp. Ther.* 2002, 302, 751, (e) Nachon, F.; Nicolet, Y.; Viguie, N.; Masson, P.; Fontecilla-Camps, J. C.; Lockridge, O. *Eur. J. Biochem.* 2002, 269, 630, herein all incorporated by reference. Using plasmid DNA as template and primers with specific base-pair alterations, mutations were made by polymerase chain reaction with Pfu DNA polymerase, for replication fidelity. The PCR product was treated with Dpn I endonuclease to digest the parental DNA template. Modified plasmid DNA was transformed into *Escherichia coli*, amplified, and purified. The DNA sequences of the mutants were confirmed by DNA sequencing. BChE mutants were expressed in human embryonic kidney cell line 293T/17. Cells were grown to 80-90% confluence in 6-well dishes and then transfected by Lipofectamine 2000 complexes of 4 µg plasmid DNA per each well. Cells were incubated at 37° C. in a CO₂ incubator for 24 hours and cells were moved to 60-mm culture vessel and cultured for four more days. The culture medium [10% fetal bovine serum in Dulbecco's modified Eagle's medium (DMEM)] was harvested for a BChE activity assay. To measure (-)-cocaine and benzoic acid, the product of (-)-cocaine hydrolysis catalyzed by BChE, sensitive radiometric assays were used based on toluene extraction of [³H]-(-)-cocaine labeled on its benzene ring in accordance with Sun et al. In brief, to initiate the enzymatic reaction, 100 nCi of [³H]-(-)-cocaine was mixed with 100 µl of culture medium. The enzymatic reactions proceeded at room temperature (25° C.) for varying time. The reactions were stopped by adding 300 µl of 0.02 M HCl, which neutralized the liberated benzoic acid while ensuring a positive charge on the residual (-)-cocaine. [³H]benzoic acid was extracted by 1 ml of toluene and measured by scintillation counting. Finally, the measured time-dependent radiometric data were fitted to the kinetic equation so that the catalytic efficiency (k_{cat}/K_M) was determined along with the use of an enzyme-linked immunosorbent assay (ELISA) described below.

Enzyme-linked immunosorbent assay (ELISA). The ELISA buffers used in the present study are the same as those described in the literature such as (a) Brock, A.; Mortensen, V.; Loft, A. G. R.; Nergaard-Pedersen, B. *J. Clin. Chem. Clin. Biochem.* 1990, 28, 221-224, (b) Khattab, A. D.; Walker, C. H.; Johnston, G.; Siddiqui, M. K. Saphier, P. W. *Environmental Toxicology and Chemistry* 1994, 13, 1661-1667, herein both incorporated by reference. The coating buffer was 0.1 M sodium carbonate/bicarbonate buffer, pH 9.5. The diluent buffer (EIA buffer) was potassium phosphate monobasic/potassium phosphate monohydrate buffer, pH 7.5, containing

0.9% sodium chloride and 0.1% bovine serum albumin. The washing buffer (PBS-T) was 0.01 M potassium phosphate monobasic/potassium phosphate monohydrate buffer, pH 7.5, containing 0.05% (v/v) Tween-20. All the assays were performed in triplicate. Each well of an ELISA microtiter plate was filled with 100 µl of the mixture buffer consisting of 20 µl culture medium and 80 µl coating buffer. The plate was covered and incubated overnight at 4° C. to allow the antigen to bind to the plate. The solutions were then removed and the wells were washed four times with PBS-T. The washed wells were filled with 200 µl diluent buffer and kept shaking for 1.5 h at room temperature (25° C.). After washing with PBS-T for four times, the wells were filled with 100 µl antibody (1:8000) and were incubated for 1.5 h, followed by washing for four times. Then, the wells were filled with 100 µl goat anti-mouse IgG HRP conjugate complex diluted to a final 1:3000 dilution, and were incubated at room temperature for 1.5 h, followed by washing for four times. The enzyme reactions were started by addition of 100 µl substrate (TMB) solution. The reactions were stopped after 15 min by the addition of 100 µl of 2 M sulfuric acid, and the absorbance was read at 460 nm using a Bio-Rad ELISA plate reader.

Hydrogen bonding revealed by the MD simulations. In accordance with one aspect of the present invention, namely the generation of high-activity mutants of BChE against (-)-cocaine, the present invention includes predicting some possible mutations that can lower the energy of the transition state for the first chemical reaction step (TS1) and, therefore, lower the energy barrier for this critical reaction step. Apparently, a mutant associated with the stronger hydrogen bonding between the carbonyl oxygen of (-)-cocaine benzoyl ester and the oxyanion hole of the BChE mutant in the TS1 structure may potentially have a more stable TS1 structure and, therefore, a higher catalytic activity for (-)-cocaine hydrolysis. Hence, the hydrogen bonding with the oxyanion hole in the TS1 structure is a crucial factor affecting the transition state stabilization and the catalytic activity. The possible effects of some mutations on the hydrogen bonding were examined by performing MD simulations and QM/MM calculations on the TS1 structures for (-)-cocaine hydrolysis catalyzed by the wild-type and various mutants BChE's.

The MD simulation in water was performed for 1 ns or longer to make sure a stable MD trajectory was obtained for each simulated TS1 structure with the wild-type or mutant BChE. The MD trajectories actually became stable quickly, so were the H . . . O distances involved in the potential hydrogen bonds between the carbonyl oxygen of (-)-cocaine and the oxyanion hole of BChE.

Depicted in FIG. 5 are plots of four important H . . . O distances in the MD-simulated TS1 structure versus the simulation time for (-)-cocaine hydrolysis catalyzed by A328W/Y332A or A199S/A328W/Y332G BChE, along with the root-mean-square deviation (RMSD) of the simulated positions of backbone atoms from those in the corresponding initial structure. Traces D1, D2, and D3 refer to the distances between the carbonyl oxygen of (-)-cocaine and the NH hydrogen of G116, G117, and S199, respectively. Trace D4 is the internuclear distance between the carbonyl oxygen of (-)-cocaine and the hydroxyl hydrogen of the S199 side chain which exists only in A199S/A328W/Y332G BChE. RMSD represents the root-mean-square deviation (in Å) of the simulated positions of the protein backbone atoms from those in the initial structure. The H . . . O distances in the simulated TS1 structures corresponding to the wild-type BChE and the two mutants are summarized in Table 3. The H . . . O distances between the carbonyl oxygen of (-)-cocaine and the peptidic NH hydrogen atoms of G116, G117, and A199 (or S199) of

BChE are denoted by D1, D2, and D3, respectively, in Table 2 and FIG. 1. D4 in Table 3 and FIG. 5 refers to the H . . . O distance between the carbonyl oxygen of (-)-cocaine and the hydroxyl hydrogen of S199 side chain in the simulated TS1 structure corresponding to the A199S/A328W/Y332G mutant, mutant (1) (SEQ ID NO: 2).

Table 3. Summary of the MD-simulated and QM/MM-optimized key distances (in Å) and the calculated total hydrogen-bonding energies (HBE, in kcal/mol) between the oxyanion hole and the carbonyl oxygen of (-)-cocaine benzoyl ester in the first transition state (TS1).

Transition State	Method	Distance ^a				Total HBE ^b	
		D1	D2	D3	D4		
TS1 for (-)-cocaine hydrolysis catalyzed by wild-type BChE	MD	Average	4.59	2.91	1.92	-5.5(-4.6)	
		Maximum	5.73	4.14	2.35		
		Minimum	3.35	1.97	1.16		
		Fluctuation	0.35	0.35	0.12		
TS1 for (-)-cocaine hydrolysis catalyzed by A328W/Y332A mutant of BChE	QM/MM	Average	4.10	2.21	2.05	-4.2	
		Maximum	3.62	2.35	1.95		
		Minimum	4.35	3.37	3.02		
		Fluctuation	2.92	1.78	1.61		
TS1 for (-)-cocaine hydrolysis catalyzed by A199S/A328W/Y332G mutant of	MD	Average	0.23	0.27	0.17	-3.3	
		Maximum	3.39	2.05	2.47		
		Minimum	5.30	2.21	1.94		2.15
		Fluctuation	6.08	3.06	2.47		3.27
Y332G mutant of	QM/MM	Average	4.57	1.71	1.66	-9.7(-7.4)	
		Maximum	0.22	0.20	0.13		0.29
		Minimum	4.83	2.09	1.91		2.59
		Fluctuation					

^aD1, D2, and D3 represent the internuclear distances between the carbonyl oxygen of cocaine benzoyl ester and the NH hydrogen of residues #116 (i.e., G116), #117 (i.e., G117), and #199 (i.e., A199 or S199) of BChE, respectively. D4 is the internuclear distance between the carbonyl oxygen of cocaine benzoyl ester and the hydroxyl hydrogen of S199 side chain in the A199S/A328W/Y332G mutant.

^bThe Total HBE value under MD is the average of the HBE values calculated by using the instantaneous distances in all of the snapshots. The value in parenthesis is the total HBE value calculated by using the MD-simulated average distances. The total HBE value under QM/MM was evaluated by using the QM/MM-optimized distances.

35

As seen in Table 3, the simulated H . . . O distance D1 is always too long for the peptidic NH of G116 to form a N—H . . . O hydrogen bond with the carbonyl oxygen of (-)-cocaine. In the simulated TS1 structure corresponding to wild-type BChE, the carbonyl oxygen of (-)-cocaine formed a firm N—H . . . O hydrogen bond with the peptidic NH hydrogen atom of A199 residue; the simulated H . . . O distance was 1.61 to 2.35 Å, with an average value of 1.92 Å. Meanwhile, the carbonyl oxygen of (-)-cocaine also had a partial N—H . . . O hydrogen bond with the peptidic NH hydrogen atom of G117 residue; the simulated H . . . O distance was 1.97 to 4.14 Å (the average value: 2.91 Å). In the simulated TS1 structure corresponding to the A328W/Y332A mutant, the simulated average H . . . O distances with the peptidic NH hydrogen of G117 and A199 are 2.35 and 1.95 Å, respectively. These distances suggest a slightly weaker N—H . . . O hydrogen bond with A199, but a stronger N—H . . . O hydrogen bond with G117, in the simulated TS1 structure corresponding to the A328W/Y332A mutant. The overall strength of the hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole of the enzyme is not expected to change considerably when wild-type BChE is replaced by the A328W/Y332A mutant.

However, the story for the simulated TS1 structure associated with the A199S/A328W/Y332G mutant was remarkably different. As one can see from Scheme 4, FIG. 5, and Table 3, when residue #199 becomes a serine (i.e., S199), the hydroxyl group on the side chain of S199 can also hydrogen-bond to the carbonyl oxygen of (-)-cocaine to form an O—H . . . O hydrogen bond, in addition to the two N—H . . . O hydrogen bonds with the peptidic NH of G117 and S199.

The simulated average H . . . O distances with the peptidic NH hydrogen of G117, peptidic NH hydrogen of S199, and hydroxyl hydrogen of S199 are 2.21, 1.94, and 2.15 Å, respectively. Due to the additional O—H . . . O hydrogen bond, the overall strength of the hydrogen bonding with the modified oxyanion hole of A199S/A328W/Y332G BChE should be significantly stronger than that of the wild-type and A328W/Y332A BChE's.

To better represent the overall strength of hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole in a MD-simulated TS1 structure, the hydrogen

bonding energy (HBE) associated with each simulated H . . . O distance was estimated by using the empirical HBE equation implemented in AutoDock 3.0 program suite in accordance with (a) Morris, G. M.; Goodsell, D. S.; Halliday, R. S.; Huey, R.; Hart, W. E.; Belew, R. K.; Olson, A. J. *J. Comput. Chem.* 1998, 19, 1639-1662, and based on the general HBE equation, $HBE(r) \approx 5 \epsilon r_0^{12}/r^{12} - 6 \epsilon r_0^{10}/r^{10}$, in which r is the H . . . O distance in the considered hydrogen bond and r_0 is the minimum value of the H . . . O distance for which the HBE equation can be used. For the calculation, $r_0 = 1.52$ Å, because it is the shortest H . . . O distance found in all of our MD simulations. The ϵ value was determined by using the condition that $HBE(r) = -5.0$ kcal/mol when $r = 1.90$ Å, herein incorporated by reference. Specifically, for each hydrogen bond with the carbonyl oxygen of (-)-cocaine, a HBE value can be evaluated with each snapshot of the MD-simulated structure. The final HBE of the MD-simulated hydrogen bond is considered to be the average HBE value of all snapshots taken from the stable MD trajectory. The estimated total HBE value for the hydrogen bonds between the carbonyl oxygen of (-)-cocaine and the oxyanion hole in each simulated TS1 structure is given in Table 3.

The HBE for each hydrogen bond was estimated by using the MD-simulated average H . . . O distance. As seen in Table 3, the total hydrogen-bonding energies (i.e., -4.6, -4.9, and -7.4 kcal/mol for the wild-type, A328W/Y332A, and A199S/A328W/Y332G BChE's, respectively) estimated in this way are systematically higher (i.e., less negative) than the corresponding total hydrogen-bonding energies (i.e., -5.5, -6.2, and -9.7 kcal/mol) estimated in the aforementioned way. However, the two sets of total HBE values are qualitatively

consistent in terms of the relative hydrogen-bonding strengths in the three simulated TS1 structures. In particular, the two sets of total HBE values consistently reveal that the overall strength of the hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole in the simulated TS1 structure for A199S/A328W/Y332G BChE is significantly higher than that for the wild-type and A328W/Y332A BChE's.

Hydrogen bonding based on the QM/MM calculations. The above conclusion obtained from the MD simulations was further examined by carrying out QM/MM calculations. Although this enzymatic reaction system is too large to calculate the QM/MM force constant matrix required in the automated search for a first-order saddle point corresponding to TS1, a partial geometry optimization was performed by fixing the length of the transition C-O bond between the carbonyl carbon of (-)-cocaine and the O' atom of S198 in the QM/MM calculation. This transition C—O bond length dominates the reaction coordinate, according to the previous first-principle reaction coordinate calculations, in accordance with Zhan et al, with an active model of wild-type BChE. In the partial geometry optimization, the transition C—O bond length was fixed at that in the TS1 geometry optimized previously by performing the first-principle reaction coordinate calculation with an active site model of wild-type BChE. This partially optimized geometry should be close to the precisely defined TS1 geometry associated with a first-order saddle point on the potential energy surface, particularly for the hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole of BChE.

The QM/MM results summarized in Table 3 demonstrate two hydrogen bonds in each of the QM/MM-optimized TS1 structures. Specifically, D2=2.21 Å and D3=2.05 Å in the optimized TS1 structure for wild-type BChE; D2=2.05 Å and D3=2.47 Å in the optimized TS1 structure for the A328W/Y332A mutant; D2=2.09 Å, D3=1.91 Å, and D4=2.59 Å in the optimized TS1 structure for the A199S/A328W/Y332G mutant. Although the QM/MM-optimized individual H . . . O distances and the estimated HBE values are different from the corresponding values for individual hydrogen bonds, the relative total HBE values (i.e., -4.2, -3.3, and -7.46 kcal/mol for the wild-type, A328W/Y332A, and A199S/A328W/Y332G BChE's, respectively) estimated from these optimized distances are qualitatively consistent with the corresponding total HBE values (i.e., -4.6, -4.9, and -7.4 kcal/mol) estimated from the MD-simulated average H . . . O distances.

It should be pointed out that the absolute HBE values estimated in this study are not expected to be accurate, as different computational approaches led different HBE values. Nevertheless, for the purpose of our computational design of a high-activity mutant of BChE, one only needs to know the relative strength of the hydrogen bonding based on the estimated relative total HBE values of different mutants. The three sets of total HBE values (Table 3) estimated from the MD-simulated and QM/MM-optimized H . . . O distances all consistently demonstrate: (1) the overall strength of hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole in the TS1 structure for (-)-cocaine hydrolysis catalyzed by A328W/Y332A BChE should be close to that in the TS1 structure for (-)-cocaine hydrolysis catalyzed by wild-type BChE; (2) the overall hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole in the TS1 structure for (-)-cocaine hydrolysis catalyzed by A199S/A328W/Y332G BChE should be significantly stronger than that in the TS1 structure for (-)-cocaine hydrolysis catalyzed by A328W/Y332A BChE.

Catalytic activity. The computational results discussed above suggest that the transition state for the first chemical reaction step (TS1) of (-)-cocaine hydrolysis catalyzed by the A199S/A328W/Y332G mutant should be significantly more stable than that by the A328W/Y332A mutant, due to the significant increase of the overall hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole of the enzyme in the TS1 structure. The aforementioned analysis of the literature, namely Sun et al and Hamza et al, also indicate that the first chemical reaction step associated with TS1 should be the rate-determining step of (-)-cocaine hydrolysis catalyzed by a BChE mutant including Y332A or Y332G mutation, although the formation of the prereactive enzyme-substrate complex (ES) is the rate-determining step for (-)-cocaine hydrolysis catalyzed by wild-type BChE. This provides evidence of a clear correlation between the TS1 stabilization and the catalytic activity of A328W/Y332A and A199S/A328W/Y332G BChE's for (-)-cocaine hydrolysis: the more stable the TS1 structure, the lower the energy barrier, and the higher the catalytic activity. Thus, both the MD simulations and QM/MM calculations predict that A199S/A328W/Y332G BChE should have a higher catalytic activity than A328W/Y332A BChE for (-)-cocaine hydrolysis.

The catalytic efficiency (k_{cat}/K_M) of A328W/Y332A BChE for (-)-cocaine hydrolysis was reported by Sun et al to be $\sim 8.6 \times 10^6 \text{ M min}^{-1}$ which is ~ 9.4 times of the k_{cat}/K_M value ($\sim 9.1 \times 10^5 \text{ M min}^{-1}$) of wild-type BChE. To examine the theoretical prediction of the higher catalytic activity for A199S/A328W/Y332G BChE, A328W/Y332A and A199S/A328W/Y332G mutants of BChE were produced through site-directed mutagenesis. To minimize the possible systematic experimental errors of the kinetic data, kinetic studies were performed with the two mutants and wild-type BChE under the same condition and compared the catalytic efficiency of A328W/Y332A and A199S/A328W/Y332G BChE's to that of the wild-type for (-)-cocaine hydrolysis at benzoyl ester group. Based on the kinetic analysis of the measured time-dependent radiometric data and the ELISA data, the ratio of the k_{cat}/K_M value of A328W/Y332A BChE to the k_{cat}/K_M value of wild-type BChE for (-)-cocaine hydrolysis was determined to be ~ 8.6 . The determined catalytic efficiency ratio of ~ 8.6 is in good agreement with the ratio of ~ 9.4 determined by Sun et al. Further, by using the same experimental protocol, the ratio of the k_{cat}/K_M value of A199S/A328W/Y332G BChE to the k_{cat}/K_M value of A328W/Y332A BChE for (-)-cocaine hydrolysis was determined to be ~ 7.2 . These data indicate that the ratio of the k_{cat}/K_M value of A199S/A328W/Y332G BChE to the k_{cat}/K_M value of wild-type BChE for (-)-cocaine hydrolysis should be $\sim 7.2 \times 8.6 = 62$ or $\sim 7.2 \times 9.4 = 68$. Thus, A199S/A328W/Y332G BChE has a $\sim (65 \pm 6)$ -fold improved catalytic efficiency against (-)-cocaine compared to the wild-type, or A199S/A328W/Y332G BChE has a k_{cat}/K_M value of $\sim (5.9 \pm 0.5) \times 10^7 \text{ M min}^{-1}$ for (-)-cocaine hydrolysis.

Very recently reported F227A/S287G/A328W/Y332M BChE (i.e., AME-359, $k_{cat}/K_M = 3.1 \times 10^7 \text{ M min}^{-1}$) has a ~ 34 -fold improved catalytic efficiency for (-)-cocaine hydrolysis. AME-359 has the highest catalytic efficiency against (-)-cocaine within all of the BChE mutants reported in literature prior to the present study. The catalytic efficiency for our A199S/A328W/Y332G BChE is about two times of that for AME-359 against (-)-cocaine.

Conclusion. Molecular dynamics (MD) simulations and hybrid quantum mechanical/molecular mechanical (QM/MM) calculations on the transition state for the first chemical reaction step (TS1) of (-)-cocaine hydrolysis catalyzed by butyrylcholinesterase (BChE) mutants lead to a better under-

standing of the effects of protein environment on the transition state stabilization. All of the computational results consistently demonstrate that the overall hydrogen bonding between the carbonyl oxygen of (-)-cocaine benzoyl ester and the oxyanion hole of BChE in the TS1 structure for (-)-cocaine hydrolysis catalyzed by A199S/A328W/Y332G BChE should be significantly stronger than that in the TS1 structure for (-)-cocaine hydrolysis catalyzed by the wild-type and A328W/Y332A BChE's. Thus, both the MD simulations and QM/MM calculations predict that A199S/A328W/Y332G BChE should have a lower energy barrier for the chemical reaction process and, therefore, a higher catalytic efficiency (k_{cat}/K_M) for (-)-cocaine hydrolysis; A328W/Y332A BChE has been known to have a ~9-fold improved catalytic efficiency for (-)-cocaine hydrolysis. The theoretical prediction has been confirmed by wet experimental tests showing a ~65±6-fold improved catalytic efficiency for A199S/A328W/Y332G BChE against (-)-cocaine compared to the wild-type BChE. The k_{cat}/K_M value determined for A199S/A328W/Y332G BChE is about two times of the k_{cat}/K_M value for F227A/S287G/A328W/Y332M BChE (i.e., AME-359, which has the highest catalytic efficiency within all BChE mutants reported prior to the present study) against (-)-cocaine. The encouraging outcome of this study suggests that the transition-state modeling is a promising approach for rational design of high-activity mutants of BChE as a therapeutic treatment of cocaine abuse.

Experiment 3

MD Simulation and Generation of Mutant A199S/S287G/A328W/Y332G BChE (Mutant 3) (SEQ ID NO: 14)

The following simulation and mutant generation relate to A199S/S287G/A328W/Y332G (mutant 3) as a rational design of a high-activity mutant of BChE against (-)-cocaine based on detailed computational modeling of the transition state for the rate-determining step (i.e., the first step of the chemical reaction process). Molecular dynamics (MD) simulations were performed to model the protein environmental effects on the stabilization of the transition-state structure for BChE-catalyzed hydrolysis of (-)-cocaine as described above for mutant A199S/A328W/Y332G (mutant 1). The simulated results indicate that the transition-state structure can be stabilized much better by the protein environment in A199S/S287G/A328W/Y332G BChE than that in wild-type BChE and in other BChE mutants examined. The computational modeling led to a prediction of the higher catalytic efficiency for the A199S/S287G/A328W/Y332G mutant against (-)-cocaine. The prediction has been confirmed by wet experimental tests showing that the A199S/S287G/A328W/Y332G mutant has a remarkably improved catalytic efficiency against (-)-cocaine. All of the obtained results

clearly demonstrate that directly modeling the transition-state structure provides a reliable computational approach to the rational design of a high-activity mutant of BChE against (-)-cocaine.

MD simulations. As with the A199S/A328W/Y332G mutant, when performing any MD simulation on a transition state, in principle, MD simulation using a classical force field (molecular mechanics) can only simulate a stable structure corresponding to a local minimum on the potential energy surface, whereas a transition state during a reaction process is always associated with a first-order saddle point on the potential energy surface.

The initial BChE structures used in the MD simulations were prepared based on our previous MD simulation as above with the prior mutant and the prereactive BChE-(-)-cocaine complex (ES) started from the X-ray crystal structure deposited in the Protein Data Bank (pdb code: 1POP).

The general procedure for carrying out the MD simulations in water is essentially the same as that used in the computational studies for mutant A199S/A328W/Y332G (mutant 1). Site-directed mutagenesis of human BChE cDNA was performed as described before.

The MD simulation in water was performed as described above, on this mutant. Depicted in FIG. 6 are plots of four important H...O distances in the MD-simulated TS1 structure versus the simulation time for (-)-cocaine hydrolysis catalyzed by A199S/S287G/A328W/Y332G BChE, along with the root-mean-square deviation (RMSD) of the simulated positions of backbone atoms from those in the corresponding initial structure. Traces D1, D2, and D3 refer to the distances between the carbonyl oxygen of (-)-cocaine and the NH hydrogen of G116, G117, and S199, respectively. Trace D4 is the internuclear distance between the carbonyl oxygen of (-)-cocaine and the hydroxyl hydrogen of the S199 side chain in A199S/S287G/A328W/Y332G BChE (mutant 3). RMSD represents the root-mean-square deviation (in Å) of the simulated positions of the protein backbone atoms from those in the initial structure.

The H...O distances in the simulated TS1 structures for wild-type BChE and its three mutants are summarized in Table 4. The H...O distances between the carbonyl oxygen of (-)-cocaine and the peptidic NH hydrogen atoms of G116, G117, and A199 (or S199) of BChE are denoted by D1, D2, and D3, respectively, in Table 4 and FIG. 6. D4 in Table 4 and FIG. 6 refers to the H...O distance between the carbonyl oxygen of (-)-cocaine and the hydroxyl hydrogen of S199 side chain in the simulated TS1 structure corresponding to the A199S/S287G/A328W/Y332G mutant.

Table 4. Summary of the MD-simulated key distances (in Å) and the calculated total hydrogen-bonding energies (HBE, in kcal/mol) between the oxyanion hole and the carbonyl oxygen of (-)-cocaine benzoyl ester in the first transition state (TS1).

Transition State		Distance ^a				Total HBE ^b
		D1	D2	D3	D4	
TS1 structure for (-)-cocaine hydrolysis catalyzed by wild-type BChE	Average	4.59	2.91	1.92		-5.5(-4.6)
	Maximum	5.73	4.14	2.35		
	Minimum	3.35	1.97	1.16		
	Fluctuation	0.35	0.35	0.12		
TS1 structure for (-)-cocaine hydrolysis catalyzed by A328W/Y332A mutant of BChE	Average	3.62	2.35	1.95		-6.2(-4.9)
	Maximum	4.35	3.37	3.02		
	Minimum	2.92	1.78	1.61		
	Fluctuation	0.23	0.27	0.17		

-continued


Transition State		Distance ^a				Total HBE ^b
		D1	D2	D3	D4	
TS1 structure for (-)-cocaine hydrolysis catalyzed by A328W/Y332G mutant of BChE	Average	3.60	2.25	1.97		-6.4(-5.0)
	Maximum	4.24	3.17	2.76		
	Minimum	2.89	1.77	1.62		
	Fluctuation	0.23	0.24	0.17		
TS1 structure for (-)-cocaine hydrolysis catalyzed by A199S/S287G/A328W/Y332G mutant of BChE	Average	4.39	2.60	2.01	1.76	-14.0(-12.0)
	Maximum	5.72	4.42	2.68	2.50	
	Minimum	2.87	1.76	1.62	1.48	
	Fluctuation	0.48	0.36	0.17	0.12	

^aD1, D2, and D3 represent the internuclear distances between the carbonyl oxygen of cocaine benzoyl ester and the NH hydrogen of residues #116 (i.e., G116), #117 (i.e., G117), and #199 (i.e., A199 or S199) of BChE, respectively. D4 is the internuclear distance between the carbonyl oxygen of cocaine benzoyl ester and the hydroxyl hydrogen of S199 side chain in the A199S/S287G/A328W/Y332G mutant.

^bThe Total HBE value is the average of the HBE values calculated by using the instantaneous distances in all of the snapshots. The value in parenthesis is the total HBE value calculated by using the MD-simulated average distances.

As seen in Table 4, the simulated H . . . O distance D1 is always too long for the peptidic NH of G116 to form a N—H . . . O hydrogen bond with the carbonyl oxygen of (-)-cocaine in all of the simulated TS1 structures. In the simulated TS1 structure for wild-type BChE, the carbonyl oxygen of (-)-cocaine formed a firm N—H . . . O hydrogen bond with the peptidic NH hydrogen atom of A199 residue; the simulated H . . . O distance (D2) was 1.61 to 2.35 Å, with an average D2 value of 1.92 Å. Meanwhile, the carbonyl oxygen of (-)-cocaine also had a partial N—H . . . O hydrogen bond with the peptidic NH hydrogen atom of G117 residue; the simulated H . . . O distance (D3) was 1.97 to 4.14 Å (the average D3 value: 2.91 Å). The average D2 and D3 values became 2.35 and 1.95 Å, respectively, in the simulated TS1 structure for the A328W/Y332A mutant. These distances

However, the story for the simulated TS1 structure for the A199S/S287G/A328W/Y332G mutant was remarkably different. As one can see from Scheme 5, FIG. 6, and Table 4, when residue #199 becomes a serine (i.e., S199), the hydroxyl group on the side chain of S199 can also hydrogen-bond to the carbonyl oxygen of (-)-cocaine to form an O—H . . . O hydrogen bond, in addition to the two N—H . . . O hydrogen bonds with the peptidic NH of G117 and S199. The simulated average H . . . O distances with the peptidic NH hydrogen of G117, peptidic NH hydrogen of S199, and hydroxyl hydrogen of S199 are 2.60, 2.01, and 1.76 Å, respectively. Due to the additional O—H . . . O hydrogen bond, the overall strength of the hydrogen bonding with the modified oxyanion hole of A199S/S287G/A328W/Y332G BChE should be significantly stronger than that of wild-type, A328W/Y332A, and A328W/Y332G BChE's.



TS1

suggest a slightly weaker N—H . . . O hydrogen bond with A199, but a stronger N—H . . . O hydrogen bond with G117, in the simulated TS1 structure for the A328W/Y332A mutant that the corresponding N—H . . . O hydrogen bonds for the wild-type. The average D2 and D3 values (2.25 and 1.97 Å, respectively) in the simulated TS1 structure for the A328W/Y332G mutant are close to the corresponding distances for the A328W/Y332A mutant. The overall strength of the hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole of the enzyme is not expected to change considerably when wild-type BChE is replaced by the A328W/Y332A or A328W/Y332G mutant.

Scheme 5. Schematic representation of the transition-state structure for first reaction step for (-)-cocaine hydrolysis catalyzed by a BChE mutant with an A199S mutation.

To better represent the overall strength of hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole in a MD-simulated TS1 structure, the hydrogen bonding energy (HBE) associated with each simulated H . . . O distance was estimated by using the empirical HBE equation implemented in AutoDock 3.0 program suite of (a) Masson, P.; Legrand, P.; Bartels, C. F.; Froment, M.-T.; Schopfer, L. M.; Lockridge, O. *Biochemistry* 1997, 36, 2266, (b) Masson, P.; Xie, W.; Froment, M.-T.; Levitsky, V.; Fortier, P.-L.; Albaret, C.; Lockridge, O. *Biochim. Biophys. Acta* 1999,

1433, 281, (c) Xie, W.; Altamirano, C. V.; Bartels, C. F.; Speirs, R. J.; Cashman, J. R.; Lockridge, O. *Mol. Pharmacol.* 1999, 55, 83, (d) Duysen, E. G.; Bartels, C. F.; Lockridge, O. *J. Pharmacol. Exp. Ther.* 2002, 302, 751, (e) Nachon, F.; Nicolet, Y.; Viguie, N.; Masson, P.; Fontecilla-Camps, J. C.; Lockridge, O. *Eur. J. Biochem.* 2002, 269, 630, herein all incorporated by reference. Specifically, for each hydrogen bond with the carbonyl oxygen of (-)-cocaine, a HBE value can be evaluated with each snapshot of the MD-simulated structure. The final HBE of the MD-simulated hydrogen bond is considered to be the average HBE value of all snapshots taken from the stable MD trajectory. The estimated total HBE value for the hydrogen bonds between the carbonyl oxygen of (-)-cocaine and the oxyanion hole in each simulated TS1 structure is also listed in Table 4.

The HBE for each hydrogen bond was estimated by using the MD-simulated average H...O distance. As seen in Table 4, the total hydrogen-bonding energies (i.e., -4.6, -4.9, -5.0, and -12.0 kcal/mol for the wild-type, A328W/Y332A, A328W/Y332G, and A199S/S287G/A328W/Y332G BChE's, respectively) estimated in this way are systematically higher (i.e., less negative) than the corresponding total hydrogen-bonding energies (i.e., -5.5, -6.2, -6.4, and -14.0 kcal/mol) estimated in the aforementioned way. However, the two sets of total HBE values are qualitatively consistent with each other in terms of the relative hydrogen-bonding strengths in the three simulated TS1 structures. In particular, the two sets of total HBE values consistently reveal that the overall strength of the hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole in the simulated TS1 structure for A199S/S287G/A328W/Y332G BChE is significantly higher than that for wild-type, A328W/Y332A, and A328W/Y332G BChE's.

Catalytic activity. The computational results discussed above provides evidence that the transition state for the first chemical reaction step (TS1) of (-)-cocaine hydrolysis catalyzed by the A199S/S287G/A328W/Y332G mutant should be significantly more stable than that by the A328W/Y332A or A328W/Y332G mutant, due to the significant increase of the overall hydrogen bonding between the carbonyl oxygen of (-)-cocaine and the oxyanion hole of the enzyme in the TS1 structure. The aforementioned analysis of the literature of Sun et al and Hamza et al also indicates that the first chemical reaction step associated with TS1 should be the rate-determining step of (-)-cocaine hydrolysis catalyzed by a BChE mutant including Y332A or Y332G mutation, although the formation of the prereactive enzyme-substrate complex (ES) is the rate-determining step for (-)-cocaine hydrolysis catalyzed by wild-type BChE. This suggests a clear correlation between the TS1 stabilization and the catalytic activity of A328W/Y332A, A328W/Y332G, and A199S/S287G/A328W/Y332G BChE's for (-)-cocaine hydrolysis: the more stable the TS1 structure, the lower the energy barrier, and the higher the catalytic activity. Thus, the MD simulations predict that A199S/S287G/A328W/Y332G BChE should have a higher catalytic activity than A328W/Y332A or A328W/Y332G BChE for (-)-cocaine hydrolysis.

The catalytic efficiency (k_{cat}/K_M) of A328W/Y332A BChE for (-)-cocaine hydrolysis was reported to be $\sim 8.6 \times 10^6$ M min⁻¹ which is ~ 9.4 times of the k_{cat}/K_M value ($\sim 9.1 \times 10^5$ M min⁻¹) of wild-type BChE for (-)-cocaine hydrolysis. The catalytic efficiency of A328W/Y332G BChE was found to be slightly higher than that of A328W/Y332A BChE for (-)-cocaine hydrolysis. To examine the theoretical prediction of the higher catalytic activity for A199S/S287G/A328W/Y332G BChE, the A328W/Y332A and A199S/S287G/A328W/Y332G mutants of BChE were produced as previ-

ously discussed through site-directed mutagenesis. To minimize the possible systematic experimental errors of the kinetic data, kinetic studies were performed with the two mutants and wild-type BChE under the same condition and compared the catalytic efficiency of A328W/Y332A and A199S/S287G/A328W/Y332G BChE's to that of the wild-type for (-)-cocaine hydrolysis at benzoyl ester group. Based on the kinetic analysis of the measured time-dependent radiometric data and the ELISA data, the ratio of the k_{cat}/K_M value of A328W/Y332A BChE to the k_{cat}/K_M value of wild-type BChE for (-)-cocaine hydrolysis was determined to be ~ 8.6 . The determined catalytic efficiency ratio of ~ 8.6 is in good agreement with the ratio of ~ 9.4 determined by Sun et al. Further, by using the same experimental protocol, the ratio of the k_{cat}/K_M value of A199S/S287G/A328W/Y332G BChE to the k_{cat}/K_M value of A328W/Y332A BChE for (-)-cocaine hydrolysis was determined to be ~ 50.6 . These data indicate that the ratio of the k_{cat}/K_M value of A199S/S287G/A328W/Y332G BChE to the k_{cat}/K_M value of wild-type BChE for (-)-cocaine hydrolysis should be $\sim 50.6 \times 8.6 \sim 435$ or $\sim 50.6 \times 9.4 \sim 476$. Thus, A199S/S287G/A328W/Y332G BChE has a $\sim (456 \pm 41)$ -fold improved catalytic efficiency against (-)-cocaine compared to the wild-type, or A199S/S287G/A328W/Y332G BChE has a k_{cat}/K_M value of $\sim (4.15 \pm 0.37) \times 10^8$ M min⁻¹ for (-)-cocaine hydrolysis. The catalytic efficiency of A199S/S287G/A328W/Y332G BChE against (-)-cocaine is much higher than that of AME-359 (i.e., F227A/S287G/A328W/Y332M BChE, $k_{cat}/K_M = 3.1 \times 10^7$ M min⁻¹, whose catalytic efficiency against (-)-cocaine is the highest within all of the previously reported BChE mutants) which has a ~ 34 -fold improved catalytic efficiency against (-)-cocaine compared to wild-type BChE.

By using the designed A199S/S287G/A328W/Y332G BChE as an exogenous enzyme in human, when the concentration of this mutant is kept the same as that of the wild-type BChE in plasma, the half-life time of (-)-cocaine in plasma should be reduced from the ~ 45 -90 min to only ~ 6 -12 seconds, considerably shorter than the time required for cocaine crossing the blood-brain barrier to reach CNS. Hence, the outcome of this study could eventually result in a valuable, efficient anti-cocaine medication.

Conclusion. The transition-state simulations demonstrate that the overall hydrogen bonding between the carbonyl oxygen of (-)-cocaine benzoyl ester and the oxyanion hole of BChE in the TS1 structure for (-)-cocaine hydrolysis catalyzed by A199S/S287G/A328W/Y332G BChE should be significantly stronger than that in the TS1 structure for (-)-cocaine hydrolysis catalyzed by the wild-type BChE and other BChE mutants simulated. Thus, the MD simulations predict that A199S/S287G/A328W/Y332G BChE should have a significantly lower energy barrier for the chemical reaction process and, therefore, a significantly higher catalytic efficiency (k_{cat}/K_M) for (-)-cocaine hydrolysis. The theoretical prediction has been confirmed by wet experimental tests showing a $\sim (456 \pm 41)$ -fold improved catalytic efficiency for A199S/S287G/A328W/Y332G BChE against (-)-cocaine compared to the wild-type BChE. The k_{cat}/K_M value determined for A199S/S287G/A328W/Y332G BChE is much higher than the k_{cat}/K_M value for AME-359 (i.e., F227A/S287G/A328W/Y332M BChE, whose catalytic efficiency against (-)-cocaine is the highest within all of the BChE mutants reported previously in literature) which has a ~ 34 -fold improved catalytic efficiency against (-)-cocaine compared to the wild-type BChE. The outcome of this study provides evidence that the transition-state simulation is a novel and unique approach for rational enzyme redesign and drug discovery.

Although the invention has been described in detail with respect to preferred embodiments thereof, it will be apparent that the invention is capable of numerous modifications and variations, apparent to those skilled in the art, without departing from the spirit and scope of the invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 30

<210> SEQ ID NO 1

<211> LENGTH: 1722

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 14-1 mutant A199S/A328W/Y332G of Human butyrylcholinesterase (BChE)

<400> SEQUENCE: 1

```

gaagatgaca tcataattgc aacaagaat ggaaaagtca gagggatgaa cttgacagtt      60
tttgggtgga cggtaacagc ctttcttggga attccctatg cacagccacc tcttggtaga    120
cttcgattca aaaagccaca gtctctgacc aagtggctctg atatttggaa tgccacaaaa    180
tatgcaaatt cttgctgtca gaacatagat caaagttttc caggcttcca tggatcagag     240
atgtggaacc caaacactga cctcagttaa gactgtttat atctaaatgt atggattcca     300
gcacctaaac caaaaaatgc cactgtattg atatggattt atgggtgggg ttttcaaact     360
ggaacatcat ctttacctgt ttatgatggc aagtttctgg ctcggttga aagagtattt     420
gtagtgtcaa tgaactatag ggtgggtgcc ctaggattct tagctttgcc aggaaatcct     480
gaggctccag ggaacatggg tttatttgat caacagttgg ctcttcagtg ggttcaaaaa    540
aatatagcag cctttggtgg aaatcctaaa agtgtaactc tctttggaga aagttccgga    600
gcagcttcag ttagcctgca tttgctttct cctggaagcc attcattggt caccagagcc    660
attctgcaaa gtggttcctt taatgctcct tgggcggtaa catctcttta tgaagctagg    720
aacagaacgt tgaacttagc taaattgact ggttgctcta gagagaatga gactgaaata    780
atcaagtgtc ttagaaataa agatccocaa gaaattcttc tgaatgaagc atttgttgtc    840
ccctatggga ctcttttgtc agtaaaactt ggtccgaccg tggatggtga ttttctcact    900
gacatgccag acatattact tgaacttggc caatttaaaa aaaccagat tttggtgggt    960
gttaataaag atgaagggac atgggtttta gtcggtgggt ctctctggtt cagcaaagat   1020
aacaatagta tcataactag aaaagaattt caggaagggt taaaaatatt tttccagga   1080
gtgagtgagt ttgaaaagga atccatcctt tttcattaca cagactgggt agatgatcag   1140
agacctgaaa actaccgtga ggccttgggt gatgtgttg gggattataa tttcatatgc   1200
cctgccttgg agttcaccia gaagtctca gaatgggaa ataatgcctt tttctactat   1260
tttgaacacc gatcctccaa acttccgtgg ccagaatgga tgggagtgat gcatggctat   1320
gaaattgaat ttgtctttgg tttacctctg gaaagaagag ataattacac aaaagccgag   1380
gaaattttga gtagatccat agtgaaacgg tgggcaaatt ttgcaaaata tgggaatcca   1440
aatgagactc agaacaatag cacaagctgg cctgtcttca aaagcactga acaaaaatat   1500
ctaaccttga atacagagtc aacaagaata atgacgaaac tacgtgctca acaatgtcga   1560
ttctggacat cattttttcc aaaagtcttg gaaatgacag gaaatattga tgaagcagaa   1620
tgggagtgga aagcaggatt ccatcgctgg aacaattaca tgatggactg gaaaaatcaa   1680
tttaacgatt acactagcaa gaaagaaagt tgtgtgggtc tc                          1722

```

<210> SEQ ID NO 2

<211> LENGTH: 574

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 14-1 mutant (A199S/A328W/Y332G) amino acid

-continued

sequence modified BChE

<400> SEQUENCE: 2

Glu Asp Asp Ile Ile Ile Ala Thr Lys Asn Gly Lys Val Arg Gly Met
 1 5 10 15
 Asn Leu Thr Val Phe Gly Gly Thr Val Thr Ala Phe Leu Gly Ile Pro
 20 25 30
 Tyr Ala Gln Pro Pro Leu Gly Arg Leu Arg Phe Lys Lys Pro Gln Ser
 35 40 45
 Leu Thr Lys Trp Ser Asp Ile Trp Asn Ala Thr Lys Tyr Ala Asn Ser
 50 55 60
 Cys Cys Gln Asn Ile Asp Gln Ser Phe Pro Gly Phe His Gly Ser Glu
 65 70 75 80
 Met Trp Asn Pro Asn Thr Asp Leu Ser Glu Asp Cys Leu Tyr Leu Asn
 85 90 95
 Val Trp Ile Pro Ala Pro Lys Pro Lys Asn Ala Thr Val Leu Ile Trp
 100 105 110
 Ile Tyr Gly Gly Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr
 115 120 125
 Asp Gly Lys Phe Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met
 130 135 140
 Asn Tyr Arg Val Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro
 145 150 155 160
 Glu Ala Pro Gly Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln
 165 170 175
 Trp Val Gln Lys Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val
 180 185 190
 Thr Leu Phe Gly Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu
 195 200 205
 Leu Ser Pro Gly Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser
 210 215 220
 Gly Ser Phe Asn Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg
 225 230 235 240
 Asn Arg Thr Leu Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn
 245 250 255
 Glu Thr Glu Ile Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile
 260 265 270
 Leu Leu Asn Glu Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Ser Val
 275 280 285
 Asn Phe Gly Pro Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp
 290 295 300
 Ile Leu Leu Glu Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly
 305 310 315 320
 Val Asn Lys Asp Glu Gly Thr Trp Phe Leu Val Gly Gly Ala Pro Gly
 325 330 335
 Phe Ser Lys Asp Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu
 340 345 350
 Gly Leu Lys Ile Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser
 355 360 365
 Ile Leu Phe His Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn
 370 375 380
 Tyr Arg Glu Ala Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys
 385 390 395 400
 Pro Ala Leu Glu Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala

-continued

	405		410		415
Phe Phe Tyr Tyr Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu	420		425		430
Trp Met Gly Val Met His Gly Tyr Glu Ile Glu Phe Val Phe Gly Leu	435		440		445
Pro Leu Glu Arg Arg Asp Asn Tyr Thr Lys Ala Glu Glu Ile Leu Ser	450		455		460
Arg Ser Ile Val Lys Arg Trp Ala Asn Phe Ala Lys Tyr Gly Asn Pro	465		470		480
Asn Glu Thr Gln Asn Asn Ser Thr Ser Trp Pro Val Phe Lys Ser Thr	485		490		495
Glu Gln Lys Tyr Leu Thr Leu Asn Thr Glu Ser Thr Arg Ile Met Thr	500		505		510
Lys Leu Arg Ala Gln Gln Cys Arg Phe Trp Thr Ser Phe Phe Pro Lys	515		520		525
Val Leu Glu Met Thr Gly Asn Ile Asp Glu Ala Glu Trp Glu Trp Lys	530		535		540
Ala Gly Phe His Arg Trp Asn Asn Tyr Met Met Asp Trp Lys Asn Gln	545		550		555
Phe Asn Asp Tyr Thr Ser Lys Lys Glu Ser Cys Val Gly Leu	565		570		

<210> SEQ ID NO 3

<211> LENGTH: 1125

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: truncated 14-1 (A199S/A328/Y332G) nucleic Acid
Sequence for modified BChE corresponding to a.a. residues 68-442

<400> SEQUENCE: 3

```

aacatagatc aaagttttcc aggccttccat ggatcagaga tgtggaaccc aaacactgac      60
ctcagtgaag actgtttata tctaaatgta tggattccag cacctaaacc aaaaaatgcc      120
actgtattga tatggattta tgggtgggtg tttcaaactg gaacatcadc tttacatggt      180
tatgatggca agtttctggc tcgggttgaa agagtattg tagtgtcaat gaactatagg      240
gtgggtgccc taggattcct agccttgcca ggaaatcctg aggcctccagg gaacatgggt      300
ttatttgatc aacagttggc tcttcagtgg gttcaaaaaa atatagcagc ctttggtgga      360
aatcctaaaa gtgtaactct ctttgagaaa agttccggag cagcttcagt tagcctgcat      420
ttgctttctc ctggaagcca ttcattgttc accagagcca ttctgcaaag tggttccttt      480
aatgctcctt gggcggtaac atctctttat gaagctagga acagaacggt gaacttagct      540
aaattgactg gttgctctag agagaatgag actgaaataa tcaagtgtct tagaaataaa      600
gatccccaag aaattcttct gaatgaagca tttgtgtgcc cctatgggac tcctttgtca      660
gtaaactttg gtccgaccgt ggatgggtgat tttctcactg acatgccaga catattactt      720
gaacttgga c aatttaaaaa aaccagatt ttggtgggtg ttaataaaga tgaagggaca      780
tggtttttag tcggtgggtc tcctggcttc agcaaagata acaatagtat cataactaga      840
aaagaatttc aggaaggttt aaaaatattt tttccaggag tgagtgagtt tggaaaggaa      900
tccatccttt ttcattacac agactgggta gatgatcaga gacctgaaaa ctaccgtgag      960
gccttgggtg atgtttgttg ggattataat ttcatatgcc ctgccttgga gttcaccaag     1020
aagttctcag aatggggaaa taatgccttt ttctactatt ttgaacaccg atcctccaaa     1080
cttcogtggc cagaatggat gggagtgatg catggctatg aaatt                               1125

```

-continued

<210> SEQ ID NO 4
 <211> LENGTH: 375
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: truncated 14-1 mutant (A199S/A328/Y332G) amino
 acid sequence of modified BChE

<400> SEQUENCE: 4

```

Asn Ile Asp Gln Ser Phe Pro Gly Phe His Gly Ser Glu Met Trp Asn
1           5           10           15
Pro Asn Thr Asp Leu Ser Glu Asp Cys Leu Tyr Leu Asn Val Trp Ile
20           25           30
Pro Ala Pro Lys Pro Lys Asn Ala Thr Val Leu Ile Trp Ile Tyr Gly
35           40           45
Gly Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys
50           55           60
Phe Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg
65           70           75           80
Val Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro
85           90           95
Gly Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln
100          105          110
Lys Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe
115          120          125
Gly Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro
130          135          140
Gly Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Phe
145          150          155          160
Asn Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg Asn Arg Thr
165          170          175
Leu Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn Glu Thr Glu
180          185          190
Ile Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile Leu Leu Asn
195          200          205
Glu Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Ser Val Asn Phe Gly
210          215          220
Pro Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp Ile Leu Leu
225          230          235          240
Glu Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly Val Asn Lys
245          250          255
Asp Glu Gly Thr Trp Phe Leu Val Gly Gly Ala Pro Gly Phe Ser Lys
260          265          270
Asp Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu Gly Leu Lys
275          280          285
Ile Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser Ile Leu Phe
290          295          300
His Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn Tyr Arg Glu
305          310          315          320
Ala Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys Pro Ala Leu
325          330          335
Glu Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala Phe Phe Tyr
340          345          350
Tyr Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu Trp Met Gly
355          360          365
  
```

-continued

Val Met His Gly Tyr Glu Ile
370 375

<210> SEQ ID NO 5
<211> LENGTH: 966
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 14-1 mutant (A199S/A328W/Y332G) BChE nucleic acid sequence corresponding to a.a. residues 117-438

<400> SEQUENCE: 5

```

ggttttcaaa ctggaacatc atctttacat gtttatgatg gcaagtttct ggctcggggt    60
gaaagagtta ttgtagtgtc aatgaactat aggggtgggtg ccctaggatt cttagctttg    120
ccaggaaatc ctgaggctcc agggaacatg ggtttatttg atcaacagtt ggctcttcag    180
tgggttcaaa aaaatatagc agcctttggt ggaaatccta aaagtgtaac tctctttgga    240
gaaagttccg gagcagcttc agttagcctg catttgcttt ctctggaag ccattcattg    300
ttcaccagag ccattctgca aagtggttcc tttaatgctc cttgggcggg aacatctctt    360
tatgaagcta ggaacagaac gttgaactta gctaaattga ctggttgctc tagagagaat    420
gagactgaaa taatcaagtg tcttagaagt aaagatcccc aagaaattct tctgaatgaa    480
gcatttggtg tcccctatgg gactcctttg tcagtaaaact ttggtccgac cgtggatggt    540
gattttctca ctgacatgcc agacatatta cttgaacttg gacaatttaa aaaaaccag    600
atthttggtg gtgtaataa agatgaaggg acatggtttt tagtcgggtg tgctcctggc    660
ttcagcaaa ataacaatag taccataact agaaaagaat ttcaggaagg tttaaaata    720
tttttccag gagtgagtga gtttgaaaag gaatccatcc tttttcatta cacagactgg    780
gtagatgate agagacctga aaactaccgt gaggccttgg gtgatgttgt tggggattat    840
aatttcatat gccttcctt ggagttcacc aagaagttct cagaatgggg aaataatgcc    900
ttttctact atthttgaaca ccgatcctcc aaacttccgt ggccagaatg gatgggagtg    960
atgcat    966

```

<210> SEQ ID NO 6
<211> LENGTH: 322
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 14-1 mutant (A199S/A328W/Y332G) BChE a.a. residues 117-438

<400> SEQUENCE: 6

```

Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys Phe
1           5           10           15
Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg Val
20           25           30
Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro Gly
35           40           45
Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln Lys
50           55           60
Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe Gly
65           70           75           80
Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro Gly
85           90           95
Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Phe Asn
100          105          110

```

-continued

Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg Asn Arg Thr Leu
 115 120 125
 Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn Glu Thr Glu Ile
 130 135 140
 Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile Leu Leu Asn Glu
 145 150 155 160
 Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Ser Val Asn Phe Gly Pro
 165 170 175
 Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp Ile Leu Leu Glu
 180 185 190
 Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly Val Asn Lys Asp
 195 200 205
 Glu Gly Thr Trp Phe Leu Val Gly Gly Ala Pro Gly Phe Ser Lys Asp
 210 215 220
 Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu Gly Leu Lys Ile
 225 230 235 240
 Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser Ile Leu Phe His
 245 250 255
 Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn Tyr Arg Glu Ala
 260 265 270
 Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys Pro Ala Leu Glu
 275 280 285
 Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala Phe Phe Tyr Tyr
 290 295 300
 Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu Trp Met Gly Val
 305 310 315 320
 Met His

<210> SEQ ID NO 7
 <211> LENGTH: 1722
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 14-2 mutant (A199S/F227A/A328W/Y332G) BChE,
 full nucleic acid sequence

<400> SEQUENCE: 7

gaagatgaca tcataattgc aacaagaat ggaaaagtca gagggatgaa cttgacagtt 60
 tttggtggca cggtaacagc ctttcttgga attcctatg cacagccacc tcttggtaga 120
 cttcgattca aaaagccaca gtctctgacc aagtggctcg atatttggaa tgccacaaaa 180
 tatgcaaatt cttgctgtca gaacatagat caaagtttcc caggcttcca tggatcagag 240
 atgtggaacc caaacactga cctcagttaa gactgtttat atctaaatgt atggattcca 300
 gcacctaaac caaaaaatgc cactgtattg atatggattt atggtggtgg ttttcaaact 360
 ggaacatcat ctttcatagt ttatgatggc aagtttctgg ctcggttga aagagttatt 420
 gtagtgtaa tgaactatag ggtgggtgcc ctaggattct tagctttgcc aggaaatcct 480
 gaggtccag ggaacatggg tttatttgat caacagttgg ctcttcagtg ggttcaaaaa 540
 aatatagcag cctttggtgg aaatcctaaa agtgtaactc tctttggaga aagttccgga 600
 gcagctcag ttagcctgca tttgctttct cctggaagcc attcattgtt caccagagcc 660
 attctgcaaa gtggttccgc taatgctcct tggcggttaa catctcttta tgaagctagg 720
 aacagaacgt tgaacttagc taaattgact ggttgctcta gagagaatga gactgaaata 780
 atcaagtgtc ttagaataaa agatccocaa gaaattcttc tgaatgaagc atttgtgtgc 840

-continued

```

ccctatggga ctctttgtc agtaaacttt ggtccgaccg tggatggtga tttctcact   900
gacatgccag acatattact tgaacttgga caatttaaaa aaaccagat tttggtgggt   960
gttaataaag atgaaggac atggttttta gtcggtggtg ctcttggtt cagcaaagat  1020
aacaatagta tcataactag aaaagaatth caggaaggtt taaaaatatt tttccagga  1080
gtgagtgagt ttggaagga atccatcctt tttcattaca cagactgggt agatgatcag  1140
agacctgaaa actaccgtga ggccttgggt gatgttgttg gggattataa tttcatatgc  1200
cctgccttgg agttcaccaa gaagtcttca gaatggggaa ataatgcctt tttctactat  1260
ttgaaacacc gatcctccaa acttccgttg ccagaatgga tgggagtgat gcatggctat  1320
gaaattgaat ttgtctttgg tttacctctg gaaagaagag ataattacac aaaagccgag  1380
gaaattttga gtagatccat agtgaacagg tgggcaaatt ttgcaaaata tgggaatcca  1440
aatgagactc agaacaatag cacaagctgg cctgtcttca aaagcactga acaaaaatat  1500
ctaaccttga atacagagtc aacaagaata atgacgaaac tacgtgctca acaatgtcga  1560
ttctggacat cattttttcc aaaagtcttg gaaatgacag gaaatattga tgaagcagaa  1620
tgggagtgga aagcaggatt ccatcgctgg aacaattaca tgatggactg gaaaaatcaa  1680
tttaacgatt aactagcaa gaaagaaagt tgtgtgggtc tc                       1722

```

<210> SEQ ID NO 8

<211> LENGTH: 574

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 14-2 mutant (A199S/F227A/A328W/Y332G), full amino acid sequence

<400> SEQUENCE: 8

```

Glu Asp Asp Ile Ile Ile Ala Thr Lys Asn Gly Lys Val Arg Gly Met
1           5           10           15
Asn Leu Thr Val Phe Gly Gly Thr Val Thr Ala Phe Leu Gly Ile Pro
20          25          30
Tyr Ala Gln Pro Pro Leu Gly Arg Leu Arg Phe Lys Lys Pro Gln Ser
35          40          45
Leu Thr Lys Trp Ser Asp Ile Trp Asn Ala Thr Lys Tyr Ala Asn Ser
50          55          60
Cys Cys Gln Asn Ile Asp Gln Ser Phe Pro Gly Phe His Gly Ser Glu
65          70          75          80
Met Trp Asn Pro Asn Thr Asp Leu Ser Glu Asp Cys Leu Tyr Leu Asn
85          90          95
Val Trp Ile Pro Ala Pro Lys Pro Lys Asn Ala Thr Val Leu Ile Trp
100         105         110
Ile Tyr Gly Gly Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr
115         120         125
Asp Gly Lys Phe Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met
130         135         140
Asn Tyr Arg Val Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro
145         150         155         160
Glu Ala Pro Gly Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln
165         170         175
Trp Val Gln Lys Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val
180         185         190
Thr Leu Phe Gly Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu
195         200         205

```

-continued

Leu Ser Pro Gly Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser
 210 215 220
 Gly Ser Ala Asn Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg
 225 230 235 240
 Asn Arg Thr Leu Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn
 245 250 255
 Glu Thr Glu Ile Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile
 260 265 270
 Leu Leu Asn Glu Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Ser Val
 275 280 285
 Asn Phe Gly Pro Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp
 290 295 300
 Ile Leu Leu Glu Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly
 305 310 315 320
 Val Asn Lys Asp Glu Gly Thr Trp Phe Leu Val Gly Gly Ala Pro Gly
 325 330 335
 Phe Ser Lys Asp Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu
 340 345 350
 Gly Leu Lys Ile Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser
 355 360 365
 Ile Leu Phe His Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn
 370 375 380
 Tyr Arg Glu Ala Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys
 385 390 395 400
 Pro Ala Leu Glu Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala
 405 410 415
 Phe Phe Tyr Tyr Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu
 420 425 430
 Trp Met Gly Val Met His Gly Tyr Glu Ile Glu Phe Val Phe Gly Leu
 435 440 445
 Pro Leu Glu Arg Arg Asp Asn Tyr Thr Lys Ala Glu Glu Ile Leu Ser
 450 455 460
 Arg Ser Ile Val Lys Arg Trp Ala Asn Phe Ala Lys Tyr Gly Asn Pro
 465 470 475 480
 Asn Glu Thr Gln Asn Asn Ser Thr Ser Trp Pro Val Phe Lys Ser Thr
 485 490 495
 Glu Gln Lys Tyr Leu Thr Leu Asn Thr Glu Ser Thr Arg Ile Met Thr
 500 505 510
 Lys Leu Arg Ala Gln Gln Cys Arg Phe Trp Thr Ser Phe Phe Pro Lys
 515 520 525
 Val Leu Glu Met Thr Gly Asn Ile Asp Glu Ala Glu Trp Glu Trp Lys
 530 535 540
 Ala Gly Phe His Arg Trp Asn Asn Tyr Met Met Asp Trp Lys Asn Gln
 545 550 555 560
 Phe Asn Asp Tyr Thr Ser Lys Lys Glu Ser Cys Val Gly Leu
 565 570

<210> SEQ ID NO 9

<211> LENGTH: 1125

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

 <223> OTHER INFORMATION: 14-2 mutant (A199/F227A/A328W/Y322G) BChE
 nucleic acid sequence for a. a. residues 68-442

<400> SEQUENCE: 9

-continued

```

aacatagatc aaagttttcc aggcttccat ggatcagaga tgtggaaccc aaacctgac    60
ctcagtgaag actgtttata tctaaatgta tggattccag cacctaaacc aaaaaatgcc  120
actgtattga tatggattta tgggtggtgt tttcaaactg gaacatcatc tttacatggt  180
tatgatggca agtttctggc tcgggttgaa agagtatttg tagtgtcaat gaactatagg  240
gtgggtgccc taggattcct agccttgcca ggaatcctg aggctccagg gaacatgggt  300
ttatttgatc aacagttggc tcttcagtgg gttcaaaaaa atatagcagc ctttggtgga  360
aatcctaaaa gtgtaactct ctttgaggaa agttccggag cagcttcagt tagcctgcat  420
ttgctttctc ctggaagcca ttcattgttc accagagcca ttctgcaaag tggttccgct  480
aatgctcctt gggcggtaac atctctttat gaagctagga acagaacggt gaacttagct  540
aaattgactg gttgctctag agagaatgag actgaaataa tcaagtgtct tagaaataaa  600
gatccccaag aaattcttct gaatgaagca tttgttgtcc cctatgggac tcctttgtca  660
gtaaactttg gtccgaccgt ggatgggtgat tttctcactg acatgccaga catattactt  720
gaacttggac aatttaaaaa aaccagatt ttggtgggtg ttaataaaga tgaagggaca  780
tggtttttag tcggtggtgc tcctggcttc agcaaagata acaatagtat cataactaga  840
aaagaatttc aggaagggtt aaaaatattt tttccaggag tgagtgagtt tggaaaggaa  900
tccatccttt ttcattacac agactgggta gatgatcaga gacctgaaaa ctaccgtgag  960
gccttgggtg atgttgttgg ggattataat ttcatatgcc ctgccttggg gttcaccaag 1020
aagttctcag aatggggaaa taatgccttt ttctactatt ttgaacaccg atcctccaaa 1080
cttcctgtgc cagaatggat gggagtgatg catggctatg aaatt                    1125

```

<210> SEQ ID NO 10

<211> LENGTH: 375

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 14-2 mutant (A199S/F227A/A328W/Y332G) BChE,
amino acid sequence for residues 68-442

<400> SEQUENCE: 10

```

Asn Ile Asp Gln Ser Phe Pro Gly Phe His Gly Ser Glu Met Trp Asn
1           5           10           15
Pro Asn Thr Asp Leu Ser Glu Asp Cys Leu Tyr Leu Asn Val Trp Ile
          20           25           30
Pro Ala Pro Lys Pro Lys Asn Ala Thr Val Leu Ile Trp Ile Tyr Gly
          35           40           45
Gly Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys
          50           55           60
Phe Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg
65           70           75           80
Val Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro
          85           90           95
Gly Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln
          100          105          110
Lys Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe
          115          120          125
Gly Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro
          130          135          140
Gly Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Ala
          145          150          155          160

```

-continued

Asn	Ala	Pro	Trp	Ala	Val	Thr	Ser	Leu	Tyr	Glu	Ala	Arg	Asn	Arg	Thr
				165					170					175	
Leu	Asn	Leu	Ala	Lys	Leu	Thr	Gly	Cys	Ser	Arg	Glu	Asn	Glu	Thr	Glu
			180					185					190		
Ile	Ile	Lys	Cys	Leu	Arg	Asn	Lys	Asp	Pro	Gln	Glu	Ile	Leu	Leu	Asn
		195					200					205			
Glu	Ala	Phe	Val	Val	Pro	Tyr	Gly	Thr	Pro	Leu	Ser	Val	Asn	Phe	Gly
	210					215					220				
Pro	Thr	Val	Asp	Gly	Asp	Phe	Leu	Thr	Asp	Met	Pro	Asp	Ile	Leu	Leu
225					230					235					240
Glu	Leu	Gly	Gln	Phe	Lys	Lys	Thr	Gln	Ile	Leu	Val	Gly	Val	Asn	Lys
				245					250					255	
Asp	Glu	Gly	Thr	Trp	Phe	Leu	Val	Gly	Gly	Ala	Pro	Gly	Phe	Ser	Lys
			260					265					270		
Asp	Asn	Asn	Ser	Ile	Ile	Thr	Arg	Lys	Glu	Phe	Gln	Glu	Gly	Leu	Lys
		275					280					285			
Ile	Phe	Phe	Pro	Gly	Val	Ser	Glu	Phe	Gly	Lys	Glu	Ser	Ile	Leu	Phe
	290					295					300				
His	Tyr	Thr	Asp	Trp	Val	Asp	Asp	Gln	Arg	Pro	Glu	Asn	Tyr	Arg	Glu
305					310					315					320
Ala	Leu	Gly	Asp	Val	Val	Gly	Asp	Tyr	Asn	Phe	Ile	Cys	Pro	Ala	Leu
				325					330					335	
Glu	Phe	Thr	Lys	Lys	Phe	Ser	Glu	Trp	Gly	Asn	Asn	Ala	Phe	Phe	Tyr
			340					345					350		
Tyr	Phe	Glu	His	Arg	Ser	Ser	Lys	Leu	Pro	Trp	Pro	Glu	Trp	Met	Gly
	355						360					365			
Val	Met	His	Gly	Tyr	Glu	Ile									
	370					375									

<210> SEQ ID NO 11

<211> LENGTH: 966

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 14-2 mutant (A199S/F227A/A328W/Y332G) BChE nucleic acid sequence for residues 117-438

<400> SEQUENCE: 11

```

ggttttcaaa ctggaacatc atctttacat gtttatgatg gcaagtttct ggetcggggt    60
gaaagagtta ttgtagtgtc aatgaactat aggggtgggtg ccctaggatt ctagctttg    120
ccaggaaatc ctgaggctcc agggaaacatg ggtttatttg atcaacagtt ggetcttcag    180
tggggtcaaa aaaatatagc agcctttggt ggaaatccta aaagtgtaac tctctttgga    240
gaaagtcccg gaggcagctc agtttagcctg catttgcttt ctctggaag ccattcattg    300
ttcaccagag ccattctgca aagtgggtcc gctaagtctc cttgggcggt aacatctctt    360
tatgaagcta ggaacagaac gttgaactta gctaaattga ctggttgctc tagagagaat    420
gagactgaaa taatcaagtg tcttagaagt aaagatcccc aagaaattct tctgaatgaa    480
gcatttggtg tcccctatgg gactcctttg tcagtaaact ttggtccgac cgtggatggt    540
gattttctca ctgacatgcc agacatatta cttgaacttg gacaatttaa aaaaaccag    600
atthttggtg gtgtaataa agatgaaggg acatggtttt tagtcggtgg tgctcctggc    660
ttcagcaaaag ataacaatag taccataact agaaaagaat ttcaggaagg tttaaaaata    720
ttttttccag gagtgagtga gtttggaag gaatccatcc tttttcatta cacagactgg    780
gtagatgatc agagacctga aaactaccgt gaggccttgg gtgatgttgt tggggattat    840

```


-continued

```

aatttcatat gcctgcctt ggagttcacc aagaagttct cagaatgggg aaataatgcc 900
ttttctact attttgaaca cggatcctcc aaacttccgt ggccagaatg gatgggagtg 960
atgcat 966

```

```

<210> SEQ ID NO 12
<211> LENGTH: 322
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 14-2 (A199S/F227A/A328W/Y332G) a.a residues
117-438

```

```

<400> SEQUENCE: 12

```

```

Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys Phe
1 5 10 15
Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg Val
20 25 30
Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro Gly
35 40 45
Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln Lys
50 55 60
Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe Gly
65 70 75 80
Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro Gly
85 90 95
Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Ala Asn
100 105 110
Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg Asn Arg Thr Leu
115 120 125
Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn Glu Thr Glu Ile
130 135 140
Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile Leu Leu Asn Glu
145 150 155 160
Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Ser Val Asn Phe Gly Pro
165 170 175
Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp Ile Leu Leu Glu
180 185 190
Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly Val Asn Lys Asp
195 200 205
Glu Gly Thr Trp Phe Leu Val Gly Gly Ala Pro Gly Phe Ser Lys Asp
210 215 220
Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu Gly Leu Lys Ile
225 230 235 240
Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser Ile Leu Phe His
245 250 255
Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn Tyr Arg Glu Ala
260 265 270
Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys Pro Ala Leu Glu
275 280 285
Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala Phe Phe Tyr Tyr
290 295 300
Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu Trp Met Gly Val
305 310 315 320
Met His

```

-continued

```

<210> SEQ ID NO 13
<211> LENGTH: 1722
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 14-3 mutant (A199S/S287G/A328W/Y332G) BChE
      nucleic acid sequence

<400> SEQUENCE: 13

gaagatgaca tcataattgc aacaagaat ggaaaagtca gagggatgaa cttgacagtt      60
tttgggtggca cggtaacagc ctttcttggga attccctatg cacagccacc tcttggtaga    120
cttcgattca aaaagccaca gtctctgacc aagtggctctg atatttggaa tgccacaaaa    180
tatgcaaatt cttgctgtca gaacatagat caaagttttc caggcttcca tggatcagag     240
atgtggaacc caaacactga cctcagtgaa gactgtttat atctaagtgt atggattcca    300
gcacctaaac caaaaaatgc cactgtattg atatggattt atggtggtgg ttttcaaact    360
ggaacatcat ctttacaatgt ttatgatggc aagtttctgg ctcggttga aagagttatt    420
gtagtgtcaa tgaactatag ggtgggtgcc ctaggattct tagctttgcc aggaaatcct    480
gaggctccag ggaacatggg tttatttgat caacagttgg ctcttcagtg ggttcaaaaa    540
aatatagcag cctttggtgg aaatcctaaa agtghtaactc tctttggaga aagttccgga    600
gcagcttcag ttagcctgca tttgctttct cctggaagcc attcattgtt caccagagcc    660
attctgcaaa gtggttcctt taatgctcct tgggcggtaa catctcttta tgaagctagg    720
aacagaacgt tgaacttagc taaattgact ggttgctcta gagagaatga gactgaaata    780
atcaagtgtc ttagaataaa agatccccaa gaaattcttc tgaatgaagc atttgtgtgc    840
ccctatggga ctcccttggg tgtaaacttt ggtccgaccg tggatggtga ttttctcact    900
gacatgccag acatattact tgaacttggga caatttaaaa aaaccagat tttggtgggt    960
gttaataaa atgaagggac atgggtttta gtcggtggtg ctctggtctt cagcaaagat   1020
aacaatagta tcataactag aaaagaattt caggaaggtt taaaaatatt tttccagga   1080
gtgagtgagt ttggaaggga atccatcctt tttcattaca cagactgggt agatgatcag   1140
agacctgaaa actaccgtga ggccttgggt gatgtgttg gggattataa tttcatatgc   1200
cctgccttgg agttcaccaa gaagtcttca gaatggggaa ataatgcctt tttctactat   1260
tttgaacacc gatcctcaa acttccgtgg ccagaatgga tgggagtgat gcatggctat   1320
gaaattgaat ttgtctttgg tttacctctg gaaagaagag ataattacac aaaagccgag   1380
gaaattttga gtagatccat agtgaaacgg tgggcaaatt ttgcaaaata tgggaatcca   1440
aatgagactc agaacaatag cacaagctgg cctgtcttca aaagcactga acaaaaaatat   1500
ctaaccttga atacagagtc aacaagaata atgacgaaac tacgtgctca acaatgtcga   1560
ttctggacat cattttttcc aaaagtcttg gaaatgacag gaaatattga tgaagcagaa   1620
tgggagtgga aagcaggatt ccatcgctgg aacaattaca tgatggactg gaaaaatcaa   1680
tttaacgatt acactagcaa gaaagaaagt tgtgtgggtc tc                          1722

```

```

<210> SEQ ID NO 14
<211> LENGTH: 574
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 14-3 mutant (A199S/S287G/A328W/Y332G) BChE
      amino acid sequence

<400> SEQUENCE: 14

```

-continued

Glu	Asp	Asp	Ile	Ile	Ile	Ala	Thr	Lys	Asn	Gly	Lys	Val	Arg	Gly	Met	1	5	10	15
Asn	Leu	Thr	Val	Phe	Gly	Gly	Thr	Val	Thr	Ala	Phe	Leu	Gly	Ile	Pro	20	25	30	
Tyr	Ala	Gln	Pro	Pro	Leu	Gly	Arg	Leu	Arg	Phe	Lys	Lys	Pro	Gln	Ser	35	40	45	
Leu	Thr	Lys	Trp	Ser	Asp	Ile	Trp	Asn	Ala	Thr	Lys	Tyr	Ala	Asn	Ser	50	55	60	
Cys	Cys	Gln	Asn	Ile	Asp	Gln	Ser	Phe	Pro	Gly	Phe	His	Gly	Ser	Glu	65	70	75	80
Met	Trp	Asn	Pro	Asn	Thr	Asp	Leu	Ser	Glu	Asp	Cys	Leu	Tyr	Leu	Asn	85	90	95	
Val	Trp	Ile	Pro	Ala	Pro	Lys	Pro	Lys	Asn	Ala	Thr	Val	Leu	Ile	Trp	100	105	110	
Ile	Tyr	Gly	Gly	Gly	Phe	Gln	Thr	Gly	Thr	Ser	Ser	Leu	His	Val	Tyr	115	120	125	
Asp	Gly	Lys	Phe	Leu	Ala	Arg	Val	Glu	Arg	Val	Ile	Val	Val	Ser	Met	130	135	140	
Asn	Tyr	Arg	Val	Gly	Ala	Leu	Gly	Phe	Leu	Ala	Leu	Pro	Gly	Asn	Pro	145	150	155	160
Glu	Ala	Pro	Gly	Asn	Met	Gly	Leu	Phe	Asp	Gln	Gln	Leu	Ala	Leu	Gln	165	170	175	
Trp	Val	Gln	Lys	Asn	Ile	Ala	Ala	Phe	Gly	Gly	Asn	Pro	Lys	Ser	Val	180	185	190	
Thr	Leu	Phe	Gly	Glu	Ser	Ser	Gly	Ala	Ala	Ser	Val	Ser	Leu	His	Leu	195	200	205	
Leu	Ser	Pro	Gly	Ser	His	Ser	Leu	Phe	Thr	Arg	Ala	Ile	Leu	Gln	Ser	210	215	220	
Gly	Ser	Phe	Asn	Ala	Pro	Trp	Ala	Val	Thr	Ser	Leu	Tyr	Glu	Ala	Arg	225	230	235	240
Asn	Arg	Thr	Leu	Asn	Leu	Ala	Lys	Leu	Thr	Gly	Cys	Ser	Arg	Glu	Asn	245	250	255	
Glu	Thr	Glu	Ile	Ile	Lys	Cys	Leu	Arg	Asn	Lys	Asp	Pro	Gln	Glu	Ile	260	265	270	
Leu	Leu	Asn	Glu	Ala	Phe	Val	Val	Pro	Tyr	Gly	Thr	Pro	Leu	Gly	Val	275	280	285	
Asn	Phe	Gly	Pro	Thr	Val	Asp	Gly	Asp	Phe	Leu	Thr	Asp	Met	Pro	Asp	290	295	300	
Ile	Leu	Leu	Glu	Leu	Gly	Gln	Phe	Lys	Lys	Thr	Gln	Ile	Leu	Val	Gly	305	310	315	320
Val	Asn	Lys	Asp	Glu	Gly	Thr	Trp	Phe	Leu	Val	Gly	Gly	Ala	Pro	Gly	325	330	335	
Phe	Ser	Lys	Asp	Asn	Asn	Ser	Ile	Ile	Thr	Arg	Lys	Glu	Phe	Gln	Glu	340	345	350	
Gly	Leu	Lys	Ile	Phe	Phe	Pro	Gly	Val	Ser	Glu	Phe	Gly	Lys	Glu	Ser	355	360	365	
Ile	Leu	Phe	His	Tyr	Thr	Asp	Trp	Val	Asp	Asp	Gln	Arg	Pro	Glu	Asn	370	375	380	
Tyr	Arg	Glu	Ala	Leu	Gly	Asp	Val	Val	Gly	Asp	Tyr	Asn	Phe	Ile	Cys	385	390	395	400
Pro	Ala	Leu	Glu	Phe	Thr	Lys	Lys	Phe	Ser	Glu	Trp	Gly	Asn	Asn	Ala	405	410	415	
Phe	Phe	Tyr	Tyr	Phe	Glu	His	Arg	Ser	Ser	Lys	Leu	Pro	Trp	Pro	Glu	420	425	430	

-continued

Trp Met Gly Val Met His Gly Tyr Glu Ile Glu Phe Val Phe Gly Leu
 435 440 445

Pro Leu Glu Arg Arg Asp Asn Tyr Thr Lys Ala Glu Glu Ile Leu Ser
 450 455 460

Arg Ser Ile Val Lys Arg Trp Ala Asn Phe Ala Lys Tyr Gly Asn Pro
 465 470 475 480

Asn Glu Thr Gln Asn Asn Ser Thr Ser Trp Pro Val Phe Lys Ser Thr
 485 490 495

Glu Gln Lys Tyr Leu Thr Leu Asn Thr Glu Ser Thr Arg Ile Met Thr
 500 505 510

Lys Leu Arg Ala Gln Gln Cys Arg Phe Trp Thr Ser Phe Phe Pro Lys
 515 520 525

Val Leu Glu Met Thr Gly Asn Ile Asp Glu Ala Glu Trp Glu Trp Lys
 530 535 540

Ala Gly Phe His Arg Trp Asn Asn Tyr Met Met Asp Trp Lys Asn Gln
 545 550 555 560

Phe Asn Asp Tyr Thr Ser Lys Lys Glu Ser Cys Val Gly Leu
 565 570

<210> SEQ ID NO 15
 <211> LENGTH: 1125
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 14-3 mutant (A199S/S287G/A328W/Y332G) BChE
 nucleic acid sequence for amino acid residues 68-442

<400> SEQUENCE: 15

aacatagatc aaagttttcc aggcctccat ggatcagaga tgtggaaccc aaacctgac 60
 ctcaagtgaag actgtttata tctaaatgta tggattccag cacctaaacc aaaaaatgcc 120
 actgtattga tatggattta tgggtgggtg tttcaaactg gaacatcatc tttacatggt 180
 tatgatggca agtttctggc tggggttgaa agagttattg tagtgtcaat gaactatagg 240
 gtgggtgccc taggattcct agccttgcca ggaaatcctg aggcctccagg gaacatgggt 300
 ttatttgatc aacagttggc tcttcagtgg gttcaaaaaa atatagcagc ctttggtgga 360
 aatcctaaaa gtgtaactct ctttgagaaa agttccggag cagcttcagt tagcctgcat 420
 ttgctttctc ctggaagcca ttcattgttc accagagcca ttctgcaaag tggttccttt 480
 aatgctcctt gggcggtaac atctctttat gaagctagga acagaacgtt gaacttagct 540
 aaattgactg gttgctctag agagaatgag actgaaataa tcaagtgtct tagaaataaa 600
 gatccccaag aaattcttct gaatgaagca tttgtgtgcc cctatgggac tcctttgggt 660
 gtaaactttg gtccgaccgt ggatgggtgat tttctcactg acatgccaga catattactt 720
 gaacttggac aatttaaaaa aaccagatt ttggtgggtg ttaataaaga tgaagggaca 780
 tggtttttag tgggtgggtc tcctggcttc agcaaagata acaatagtat cataactaga 840
 aaagaatttc aggaaggttt aaaaatattt tttccaggag tgagttagtt tggaaaggaa 900
 tccatccttt ttcattacac agactgggta gatgatcaga gacctgaaaa ctaccgtgag 960
 gccttgggtg atgttgttgg ggattataat ttcatatgcc ctgccttggg gttcaccaag 1020
 aagttctcag aatggggaaa taatgccttt ttctactatt tgaacaccg atcctccaaa 1080
 ctccctgggc cagaatggat gggagtgatg catggctatg aaatt 1125

<210> SEQ ID NO 16
 <211> LENGTH: 375

-continued

<212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 14-3 mutant (A199S/S287G/A328W/Y332G) BChE
 amino acid sequence for residues 68-442

<400> SEQUENCE: 16

Asn Ile Asp Gln Ser Phe Pro Gly Phe His Gly Ser Glu Met Trp Asn
 1 5 10 15
 Pro Asn Thr Asp Leu Ser Glu Asp Cys Leu Tyr Leu Asn Val Trp Ile
 20 25 30
 Pro Ala Pro Lys Pro Lys Asn Ala Thr Val Leu Ile Trp Ile Tyr Gly
 35 40 45
 Gly Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys
 50 55 60
 Phe Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg
 65 70 75 80
 Val Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro
 85 90 95
 Gly Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln
 100 105 110
 Lys Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe
 115 120 125
 Gly Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro
 130 135 140
 Gly Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Phe
 145 150 155 160
 Asn Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg Asn Arg Thr
 165 170 175
 Leu Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn Glu Thr Glu
 180 185 190
 Ile Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile Leu Leu Asn
 195 200 205
 Glu Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Gly Val Asn Phe Gly
 210 215 220
 Pro Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp Ile Leu Leu
 225 230 235 240
 Glu Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly Val Asn Lys
 245 250 255
 Asp Glu Gly Thr Trp Phe Leu Val Gly Gly Ala Pro Gly Phe Ser Lys
 260 265 270
 Asp Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu Gly Leu Lys
 275 280 285
 Ile Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser Ile Leu Phe
 290 295 300
 His Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn Tyr Arg Glu
 305 310 315 320
 Ala Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys Pro Ala Leu
 325 330 335
 Glu Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala Phe Phe Tyr
 340 345 350
 Tyr Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu Trp Met Gly
 355 360 365
 Val Met His Gly Tyr Glu Ile
 370 375

-continued

<210> SEQ ID NO 17
 <211> LENGTH: 966
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 14-3 mutant (A199S/S287G/A328W/Y332G) BChE
 nucleic acid sequence for amino acid residues 117-438

<400> SEQUENCE: 17

```

ggttttcaaa ctggaacatc atcctttacat gtttatgatg gcaagtttct ggctcggggt    60
gaaagagtta ttgtagtgtc aatgaactat aggggtgggtg ccctaggatt cttagctttg    120
ccaggaaatc ctgaggctcc agggaacatg ggtttatttg atcaacagtt ggctcttcag    180
tgggttcaaa aaaatatagc agcctttggt ggaaatccta aaagtgtaac tctctttgga    240
gaaagtcccg gagcagcttc agttagcctg catttgcttt ctctggaag ccattcattg    300
ttcaccagag ccattctgca aagtggttcc tttaatgctc cttgggcggg aacatctctt    360
tatgaagcta ggaacagaac gttgaactta gctaaattga ctggttgctc tagagagaat    420
gagactgaaa taatcaagtg tcttagaagt aaagatcccc aagaaattct tctgaatgaa    480
gcatttggtt tcccctatgg gactcctttg ggtgtaaact ttggtccgac cgtggatggt    540
gattttctca ctgacatgcc agacatatta cttgaacttg gacaatttaa aaaaaccag    600
attttggtgg gtgtaataa agatgaaggg acatggtttt tagtcggtgg tgctcctggc    660
ttcagcaaag ataacaatag taccataact agaaaagaat ttcaggaagg tttaaaaata    720
ttttttccag gagtgagtga gtttgaaaag gaatccatcc tttttcatta cacagactgg    780
gtagatgatc agagacctga aaactaccgt gaggccttgg gtgatgttgt tggggattat    840
aatttcatat gccttcctt ggagttcacc aagaagttct cagaatgggg aaataatgcc    900
tttttctact attttgaaca ccgatctccc aaacttccgt ggccagaatg gatgggagtg    960
atgcat    966
  
```

<210> SEQ ID NO 18
 <211> LENGTH: 322
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 14-3 mutant (A199S/S287G/A328W/Y332G) BChE
 amino acid sequence for residues 117-438

<400> SEQUENCE: 18

```

Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys Phe
 1          5          10          15
Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg Val
 20          25          30
Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro Gly
 35          40          45
Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln Lys
 50          55          60
Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe Gly
 65          70          75          80
Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro Gly
 85          90          95
Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Phe Asn
 100         105         110
Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg Asn Arg Thr Leu
 115         120         125
  
```

-continued

Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn Glu Thr Glu Ile
 130 135 140
 Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile Leu Leu Asn Glu
 145 150 155 160
 Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Gly Val Asn Phe Gly Pro
 165 170 175
 Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp Ile Leu Leu Glu
 180 185 190
 Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly Val Asn Lys Asp
 195 200 205
 Glu Gly Thr Trp Phe Leu Val Gly Gly Ala Pro Gly Phe Ser Lys Asp
 210 215 220
 Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu Gly Leu Lys Ile
 225 230 235 240
 Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser Ile Leu Phe His
 245 250 255
 Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn Tyr Arg Glu Ala
 260 265 270
 Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys Pro Ala Leu Glu
 275 280 285
 Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala Phe Phe Tyr Tyr
 290 295 300
 Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu Trp Met Gly Val
 305 310 315 320
 Met His

<210> SEQ ID NO 19
 <211> LENGTH: 1722
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 12-7 mutant (A199S/F227A/S287G/A328W/Y332G)
 BChE nucleic acid sequence

<400> SEQUENCE: 19

gaagatgaca tcataattgc aacaagaat ggaaaagtca gagggatgaa cttgacagtt 60
 tttggtggca cggtaacagc ctttcttggga attccctatg cacagccacc tcttggtaga 120
 cttcgattca aaaagccaca gtctctgacc aagtggctctg atatttggaa tgcacaaaa 180
 tatgcaaatt cttgctgtca gaacatagat caaagttttc caggcttcca tggatcagag 240
 atgtggaacc caaacactga cctcagttaa gactgtttat atctaaatgt atggattcca 300
 gcacctaac caaaaaatgc cactgtattg atatggattt atggtggtgg ttttcaaact 360
 ggaacatcat ctttcatatg ttatgatggc aagtttctgg ctcggttga aagagttatt 420
 gtagtgtaaa tgaactatag ggtgggtgcc ctaggattct tagctttgcc aggaaatcct 480
 gaggtccag ggaacatggg tttatttggat caacagttgg ctcttcagtg ggttcaaaaa 540
 aatatagcag cctttggtgg aaatcctaaa agtgtaactc tctttggaga aagttccgga 600
 gcagcttcag ttagectgca tttgctttct cctggaagcc attcattggt caccagagcc 660
 attctgcaaa gtggttccgc taatgctcct tgggcggtaa catctcttta tgaagctagg 720
 aacagaacgt tgaacttagc taaattgact ggttgctcta gagagaatga gactgaaata 780
 atcaagtgtc ttagaaataa agatcccaaa gaaattcttc tgaatgaagc atttgtgtc 840
 ccctatggga ctcccttggg tgtaaacctt ggtccgaccg tggatggtga ttttctcact 900
 gacatgccag acatattact tgaacttggg caatttaaaa aaaccagat tttggtgggt 960

-continued

```

gttaataaag atgaaggac atggtttita gtcggtggtg ctctggctt cagcaaagat 1020
aacaatagta tcataactag aaaagaattt caggaagggt taaaaatatt tttccagga 1080
gtgagtgagt ttggaagga atccatcctt tttcattaca cagactgggt agatgatcag 1140
agacctgaaa actaccgtga ggccttgggt gatgttgttg gggattataa tttcatatgc 1200
cctgccttgg agttcaccaa gaagtcttca gaatggggaa ataatgcctt tttctactat 1260
tttgaacacc gatcctccaa acttccgtgg ccagaatgga tgggagtgat gcatggctat 1320
gaaattgaat ttgtcttttg tttacctctg gaaagaagag ataattacac aaaagccgag 1380
gaaattttga gtagatccat agtgaacagg tgggcaaatt ttgcaaaata tgggaatcca 1440
aatgagactc agaacaatag cacaagctgg cctgtcttca aaagcactga acaaaaatat 1500
ctaaccttga atacagagtc aacaagaata atgacgaaac tacgtgctca acaatgtcga 1560
ttctggacat cattttttcc aaaagtcttg gaaatgacag gaaatattga tgaagcagaa 1620
tgggagtgga aagcaggatt ccacgctgg aacaattaca tgatggactg gaaaaatcaa 1680
ttaaactgatt aactagcaa gaaagaaagt tgtgtgggtc tc 1722

```

<210> SEQ ID NO 20

<211> LENGTH: 574

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 12-7 mutant (A199S/F227A/S287G/A328W/Y332G)
BChE amino acid sequence

<400> SEQUENCE: 20

```

Glu Asp Asp Ile Ile Ile Ala Thr Lys Asn Gly Lys Val Arg Gly Met
1           5           10           15
Asn Leu Thr Val Phe Gly Gly Thr Val Thr Ala Phe Leu Gly Ile Pro
20          25          30
Tyr Ala Gln Pro Pro Leu Gly Arg Leu Arg Phe Lys Lys Pro Gln Ser
35          40          45
Leu Thr Lys Trp Ser Asp Ile Trp Asn Ala Thr Lys Tyr Ala Asn Ser
50          55          60
Cys Cys Gln Asn Ile Asp Gln Ser Phe Pro Gly Phe His Gly Ser Glu
65          70          75          80
Met Trp Asn Pro Asn Thr Asp Leu Ser Glu Asp Cys Leu Tyr Leu Asn
85          90          95
Val Trp Ile Pro Ala Pro Lys Pro Lys Asn Ala Thr Val Leu Ile Trp
100         105         110
Ile Tyr Gly Gly Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr
115         120         125
Asp Gly Lys Phe Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met
130         135         140
Asn Tyr Arg Val Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro
145         150         155         160
Glu Ala Pro Gly Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln
165         170         175
Trp Val Gln Lys Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val
180         185         190
Thr Leu Phe Gly Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu
195         200         205
Leu Ser Pro Gly Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser
210         215         220

```


-continued

Gly Ser Ala Asn Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg
 225 230 235 240
 Asn Arg Thr Leu Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn
 245 250 255
 Glu Thr Glu Ile Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile
 260 265 270
 Leu Leu Asn Glu Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Gly Val
 275 280 285
 Asn Phe Gly Pro Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp
 290 295 300
 Ile Leu Leu Glu Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly
 305 310 315 320
 Val Asn Lys Asp Glu Gly Thr Trp Phe Leu Val Gly Gly Ala Pro Gly
 325 330 335
 Phe Ser Lys Asp Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu
 340 345 350
 Gly Leu Lys Ile Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser
 355 360 365
 Ile Leu Phe His Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn
 370 375 380
 Tyr Arg Glu Ala Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys
 385 390 395 400
 Pro Ala Leu Glu Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala
 405 410 415
 Phe Phe Tyr Tyr Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu
 420 425 430
 Trp Met Gly Val Met His Gly Tyr Glu Ile Glu Phe Val Phe Gly Leu
 435 440 445
 Pro Leu Glu Arg Arg Asp Asn Tyr Thr Lys Ala Glu Glu Ile Leu Ser
 450 455 460
 Arg Ser Ile Val Lys Arg Trp Ala Asn Phe Ala Lys Tyr Gly Asn Pro
 465 470 475 480
 Asn Glu Thr Gln Asn Asn Ser Thr Ser Trp Pro Val Phe Lys Ser Thr
 485 490 495
 Glu Gln Lys Tyr Leu Thr Leu Asn Thr Glu Ser Thr Arg Ile Met Thr
 500 505 510
 Lys Leu Arg Ala Gln Gln Cys Arg Phe Trp Thr Ser Phe Phe Pro Lys
 515 520 525
 Val Leu Glu Met Thr Gly Asn Ile Asp Glu Ala Glu Trp Glu Trp Lys
 530 535 540
 Ala Gly Phe His Arg Trp Asn Asn Tyr Met Met Asp Trp Lys Asn Gln
 545 550 555 560
 Phe Asn Asp Tyr Thr Ser Lys Lys Glu Ser Cys Val Gly Leu
 565 570

<210> SEQ ID NO 21

<211> LENGTH: 1125

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 12-7 mutant (A199S/F227A/S287G/A328W/Y332G)
 EChE nucleic acid sequence for amino acid residues 68-442

<400> SEQUENCE: 21

aacatagatc aaagttttcc aggcctccat ggatcagaga tgtggaaccc aaacactgac 60

ctcagtgaag actgtttata tctaaatgta tggattccag cacctaaacc aaaaaatgcc 120

-continued

```

actgtattga tatggattta tgggtgggtggt tttcaaactg gaacatcadc tttacatggt 180
tatgatggca agtttctggc tcgggttgaa agagtatttg tagtgtcaat gaactatagg 240
gtgggtgccc taggattctt agccttgcca ggaaatcctg aggctccagg gaacatgggt 300
ttatttgatc aacagttggc tcttcagtgg gttcaaaaaa atatagcagc ctttggtgga 360
aatcctaaaa gtgtaactct ctttgagaaa agttccggag cagcttcagt tagcctgcat 420
ttgctttctc ctggaagcca ttcattgttc accagagcca ttctgcaaag tggttccgct 480
aatgctcctt gggcggtaac atctctttat gaagctagga acagaacgtt gaacttagct 540
aaattgactg gttgctctag agagaatgag actgaaataa tcaagtgtct tagaaataaa 600
gatccccaag aaattctctt gaatgaagca tttgtgttcc cctatgggac tcctttgggt 660
gtaaactttg gtccgaccgt ggatgggtgat tttctcactg acatgccaga catattactt 720
gaacttggac aatttaaaaa aaccagatt ttggtgggtg ttaataaaga tgaagggaca 780
tggtttttag tcggtgggtc tcttggttc agcaaagata acaatagtat cataactaga 840
aaagaatttc aggaaggttt aaaaatattt tttccaggag tgagttagtt tggaaaggaa 900
tccatccttt ttcattacac agactgggta gatgatcaga gacctgaaaa ctaccgtgag 960
gccttgggtg atgttgggtg ggattataat ttcatatgcc ctgccttggg gttcaccaag 1020
aagtctcag aatggggaaa taatgctttt ttctactatt ttgaacaccg atcctccaaa 1080
cttccgtggc cagaatggat gggagtgatg catggctatg aaatt 1125

```

<210> SEQ ID NO 22

<211> LENGTH: 375

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: 12-7 mutant (A199S/F227A/S287G/A328W/Y332G)
 BChE amino acid sequence for residues 68-442

<400> SEQUENCE: 22

```

Asn Ile Asp Gln Ser Phe Pro Gly Phe His Gly Ser Glu Met Trp Asn
1           5           10           15
Pro Asn Thr Asp Leu Ser Glu Asp Cys Leu Tyr Leu Asn Val Trp Ile
20          25          30
Pro Ala Pro Lys Pro Lys Asn Ala Thr Val Leu Ile Trp Ile Tyr Gly
35          40          45
Gly Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys
50          55          60
Phe Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg
65          70          75          80
Val Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro
85          90          95
Gly Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln
100         105         110
Lys Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe
115         120         125
Gly Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro
130         135         140
Gly Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Ala
145         150         155         160
Asn Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg Asn Arg Thr
165         170         175
Leu Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn Glu Thr Glu

```

-continued

180					185					190					
Ile	Ile	Lys	Cys	Leu	Arg	Asn	Lys	Asp	Pro	Gln	Glu	Ile	Leu	Leu	Asn
		195					200					205			
Glu	Ala	Phe	Val	Val	Pro	Tyr	Gly	Thr	Pro	Leu	Gly	Val	Asn	Phe	Gly
	210					215					220				
Pro	Thr	Val	Asp	Gly	Asp	Phe	Leu	Thr	Asp	Met	Pro	Asp	Ile	Leu	Leu
	225					230					235				240
Glu	Leu	Gly	Gln	Phe	Lys	Lys	Thr	Gln	Ile	Leu	Val	Gly	Val	Asn	Lys
			245						250					255	
Asp	Glu	Gly	Thr	Trp	Phe	Leu	Val	Gly	Gly	Ala	Pro	Gly	Phe	Ser	Lys
			260					265					270		
Asp	Asn	Asn	Ser	Ile	Ile	Thr	Arg	Lys	Glu	Phe	Gln	Glu	Gly	Leu	Lys
		275					280					285			
Ile	Phe	Phe	Pro	Gly	Val	Ser	Glu	Phe	Gly	Lys	Glu	Ser	Ile	Leu	Phe
	290					295					300				
His	Tyr	Thr	Asp	Trp	Val	Asp	Asp	Gln	Arg	Pro	Glu	Asn	Tyr	Arg	Glu
	305					310					315				320
Ala	Leu	Gly	Asp	Val	Val	Gly	Asp	Tyr	Asn	Phe	Ile	Cys	Pro	Ala	Leu
				325					330					335	
Glu	Phe	Thr	Lys	Lys	Phe	Ser	Glu	Trp	Gly	Asn	Asn	Ala	Phe	Phe	Tyr
			340					345					350		
Tyr	Phe	Glu	His	Arg	Ser	Ser	Lys	Leu	Pro	Trp	Pro	Glu	Trp	Met	Gly
		355					360					365			
Val	Met	His	Gly	Tyr	Glu	Ile									
	370					375									

<210> SEQ ID NO 23
 <211> LENGTH: 966
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 12-7 mutant (A199S/F227A/S287G/A328W/Y332G)
 BChE nucleic acid sequence for amino acid residues 117-438

<400> SEQUENCE: 23

```

ggttttcaaa ctggaacatc atctttacat gtttatgatg gcaagtttct ggctcggggt    60
gaaagagtta ttgtagtgtc aatgaactat aggggtgggtg ccctaggatt cttagctttg    120
ccaggaaatc ctgaggtccc agggaacatg gggttatttg atcaacagtt ggetcttcag    180
tgggttcaaa aaaatatagc agcctttggt ggaaatccta aaagtgtaac tctctttgga    240
gaaagtcccg gagcagcttc agttagcctg catttgcttt ctctggaag ccattcattg    300
ttcaccagag ccattctgca aagtggttcc gctaatgctc cttgggcggt aacatctctt    360
tatgaagcta ggaacagaac gttgaactta gctaaattga ctggttgctc tagagagaat    420
gagactgaaa taatcaagtg tcttagaagt aaagatcccc aagaaattct tctgaatgaa    480
gcatttggtg tcccctatgg gactcctttg ggtgtaaact ttggtccgac cgtggatggt    540
gattttctca ctgacatgcc agacatatta cttgaacttg gacaatttaa aaaaccag    600
atthttggtg gtgtaataaa agatgaaggg acatggtttt tagtcggtgg tgctcctggc    660
ttcagcaaag ataacaatag taccataact agaaaagaat ttcaggaagg tttaaaaata    720
tttttccag gagtgagtga gtttgaaaag gaatccatcc tttttcatta cacagactgg    780
gtagatgata agagacctga aaactaccgt gaggccttgg gtgatgttgt tggggattat    840
aatttcatat gccctgcctt ggagttcacc aagaagttct cagaatgggg aaataatgcc    900
tttttctact atthttgaaca ccgatcctcc aaacttccgt ggccagaatg gatgggagtg    960
  
```

-continued

atgcat

966

<210> SEQ ID NO 24
 <211> LENGTH: 322
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 12-7 mutant (A199S/F227A/S287G/A328W/Y332G)
 BChE amino acid sequence for residues 117-438

<400> SEQUENCE: 24

Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys Phe
 1 5 10 15
 Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg Val
 20 25 30
 Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro Gly
 35 40 45
 Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln Lys
 50 55 60
 Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe Gly
 65 70 75 80
 Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro Gly
 85 90 95
 Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Ala Asn
 100 105 110
 Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg Asn Arg Thr Leu
 115 120 125
 Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn Glu Thr Glu Ile
 130 135 140
 Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile Leu Leu Asn Glu
 145 150 155 160
 Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Gly Val Asn Phe Gly Pro
 165 170 175
 Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp Ile Leu Leu Glu
 180 185 190
 Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly Val Asn Lys Asp
 195 200 205
 Glu Gly Thr Trp Phe Leu Val Gly Gly Ala Pro Gly Phe Ser Lys Asp
 210 215 220
 Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu Gly Leu Lys Ile
 225 230 235 240
 Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser Ile Leu Phe His
 245 250 255
 Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn Tyr Arg Glu Ala
 260 265 270
 Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys Pro Ala Leu Glu
 275 280 285
 Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala Phe Phe Tyr Tyr
 290 295 300
 Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu Trp Met Gly Val
 305 310 315 320
 Met His

<210> SEQ ID NO 25
 <211> LENGTH: 1722
 <212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 12-4 mutant (A199S/F227A/S287G/A328W/E441D)
      Bche nucleic acid sequence

<400> SEQUENCE: 25
gaagatgaca tcataattgc aacaaagaat ggaaaagtca gagggatgaa cttgacagtt      60
tttgggtggca cggtaacagc ctttcttgga attccctatg cacagccacc tcttggtaga      120
cttcgattca aaaagccaca gtctctgacc aagtggctctg atatttggaa tgccacaaaa      180
tatgcaaatt cttgctgtca gaacatagat caaagttttc caggcttcca tggatcagag      240
atgtggaacc caaacactga cctcagtga gactgtttat atctaaatgt atggattcca      300
gcacctaaac caaaaaatgc cactgtattg atatggattt atgggtgggg ttttcaaact      360
ggaacatcat ctttacatgt ttatgatggc aagtttctgg ctcgggttga aagagttatt      420
gtagtgtaaa tgaactatag ggtgggtgcc ctaggattct tagctttgcc aggaaatcct      480
gaggtccag ggaacatggg tttatttgat caacagttgg ctcttcagtg ggttcaaaaa      540
aatatagcag cctttgggtg aaatcctaaa agtgtaactc tctttggaga aagttccgga      600
gcagcttcag ttagcctgca tttgctttct cctggaagcc attcattgtt caccagagcc      660
attctgcaaa gtggttccgc taatgctcct tggggcggtaa catctcttta tgaagctagg      720
aacagaacgt tgaacttagc taaatgact ggttgctcta gagagaatga gactgaata      780
atcaagtgtc ttagaaaata agatcccca gaaattcttc tgaatgaagc atttgtgtc      840
ccctatggga ctctttggg tgtaaacctt ggtccgaccg tggatgggta ttttctcaact      900
gacatgccag acatattact tgaacttggc caatttaaaa aaaccagat tttggtgggt      960
gttaataaag atgaagggac atggttttta gtctatggtg ctctggcctt cagcaaagat      1020
aacaatagta tcataactag aaaagaattt caggaaggtt taaaaatatt tttccagga      1080
gtgagtgagt ttggaagga atccatcctt tttcattaca cagactgggt agatgatcag      1140
agacctgaaa actaccgtga ggccttgggt gatgttgttg gggattataa tttcatatgc      1200
cctgccttgg agttcaccaa gaagtctca gaatggggaa ataatgcctt tttctactat      1260
tttgaacacc gatcctccaa acttccgtgg ccagaatgga tgggagtgat gcatggctat      1320
gacattgaat ttgtcttttg tttacctctg gaaagaagag ataattacac aaaagccgag      1380
gaaattttga gtagatccat agtgaacggt tgggcaaatt ttgcaaaata tgggaatcca      1440
aatgagactc agaacaatag cacaagctgg cctgtcttca aaagcactga acaaaaatat      1500
ctaaccctga atacagatgc aacaagaata atgacgaaac tacgtgctca acaatgtcga      1560
ttctggacat cttttttcc aaaagtcttg gaaatgacag gaaatattga tgaagcagaa      1620
tgggagtgga aagcaggatt ccatcgctgg aacaattaca tgatggactg gaaaaatcaa      1680
tttaacgatt aactagcaa gaaagaaagt tgtgtgggtc tc      1722

```

```

<210> SEQ ID NO 26
<211> LENGTH: 574
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 12-4 mutant (A199S/F227A/S287G/A328W/E441D)
      Bche amino acid sequence

```

```

<400> SEQUENCE: 26

```

```

Glu Asp Asp Ile Ile Ala Thr Lys Asn Gly Lys Val Arg Gly Met
1           5           10          15

```

```

Asn Leu Thr Val Phe Gly Gly Thr Val Thr Ala Phe Leu Gly Ile Pro

```

-continued

20					25					30					
Tyr	Ala	Gln	Pro	Pro	Leu	Gly	Arg	Leu	Arg	Phe	Lys	Lys	Pro	Gln	Ser
		35					40					45			
Leu	Thr	Lys	Trp	Ser	Asp	Ile	Trp	Asn	Ala	Thr	Lys	Tyr	Ala	Asn	Ser
	50					55					60				
Cys	Cys	Gln	Asn	Ile	Asp	Gln	Ser	Phe	Pro	Gly	Phe	His	Gly	Ser	Glu
65					70					75					80
Met	Trp	Asn	Pro	Asn	Thr	Asp	Leu	Ser	Glu	Asp	Cys	Leu	Tyr	Leu	Asn
				85					90					95	
Val	Trp	Ile	Pro	Ala	Pro	Lys	Pro	Lys	Asn	Ala	Thr	Val	Leu	Ile	Trp
			100					105					110		
Ile	Tyr	Gly	Gly	Gly	Phe	Gln	Thr	Gly	Thr	Ser	Ser	Leu	His	Val	Tyr
		115					120					125			
Asp	Gly	Lys	Phe	Leu	Ala	Arg	Val	Glu	Arg	Val	Ile	Val	Val	Ser	Met
	130					135					140				
Asn	Tyr	Arg	Val	Gly	Ala	Leu	Gly	Phe	Leu	Ala	Leu	Pro	Gly	Asn	Pro
145					150					155					160
Glu	Ala	Pro	Gly	Asn	Met	Gly	Leu	Phe	Asp	Gln	Gln	Leu	Ala	Leu	Gln
				165					170						175
Trp	Val	Gln	Lys	Asn	Ile	Ala	Ala	Phe	Gly	Gly	Asn	Pro	Lys	Ser	Val
			180					185						190	
Thr	Leu	Phe	Gly	Glu	Ser	Ser	Gly	Ala	Ala	Ser	Val	Ser	Leu	His	Leu
		195					200					205			
Leu	Ser	Pro	Gly	Ser	His	Ser	Leu	Phe	Thr	Arg	Ala	Ile	Leu	Gln	Ser
	210					215					220				
Gly	Ser	Ala	Asn	Ala	Pro	Trp	Ala	Val	Thr	Ser	Leu	Tyr	Glu	Ala	Arg
225					230					235					240
Asn	Arg	Thr	Leu	Asn	Leu	Ala	Lys	Leu	Thr	Gly	Cys	Ser	Arg	Glu	Asn
				245					250					255	
Glu	Thr	Glu	Ile	Ile	Lys	Cys	Leu	Arg	Asn	Lys	Asp	Pro	Gln	Glu	Ile
			260					265					270		
Leu	Leu	Asn	Glu	Ala	Phe	Val	Val	Pro	Tyr	Gly	Thr	Pro	Leu	Gly	Val
		275					280					285			
Asn	Phe	Gly	Pro	Thr	Val	Asp	Gly	Asp	Phe	Leu	Thr	Asp	Met	Pro	Asp
		290				295					300				
Ile	Leu	Leu	Glu	Leu	Gly	Gln	Phe	Lys	Lys	Thr	Gln	Ile	Leu	Val	Gly
305					310					315					320
Val	Asn	Lys	Asp	Glu	Gly	Thr	Trp	Phe	Leu	Val	Tyr	Gly	Ala	Pro	Gly
				325					330					335	
Phe	Ser	Lys	Asp	Asn	Asn	Ser	Ile	Ile	Thr	Arg	Lys	Glu	Phe	Gln	Glu
			340					345					350		
Gly	Leu	Lys	Ile	Phe	Phe	Pro	Gly	Val	Ser	Glu	Phe	Gly	Lys	Glu	Ser
		355					360					365			
Ile	Leu	Phe	His	Tyr	Thr	Asp	Trp	Val	Asp	Asp	Gln	Arg	Pro	Glu	Asn
	370					375					380				
Tyr	Arg	Glu	Ala	Leu	Gly	Asp	Val	Val	Gly	Asp	Tyr	Asn	Phe	Ile	Cys
385					390					395					400
Pro	Ala	Leu	Glu	Phe	Thr	Lys	Lys	Phe	Ser	Glu	Trp	Gly	Asn	Asn	Ala
				405					410					415	
Phe	Phe	Tyr	Tyr	Phe	Glu	His	Arg	Ser	Ser	Lys	Leu	Pro	Trp	Pro	Glu
			420					425					430		
Trp	Met	Gly	Val	Met	His	Gly	Tyr	Asp	Ile	Glu	Phe	Val	Phe	Gly	Leu
		435					440					445			

-continued

Pro Leu Glu Arg Arg Asp Asn Tyr Thr Lys Ala Glu Glu Ile Leu Ser
 450 455 460

Arg Ser Ile Val Lys Arg Trp Ala Asn Phe Ala Lys Tyr Gly Asn Pro
 465 470 475 480

Asn Glu Thr Gln Asn Asn Ser Thr Ser Trp Pro Val Phe Lys Ser Thr
 485 490 495

Glu Gln Lys Tyr Leu Thr Leu Asn Thr Glu Ser Thr Arg Ile Met Thr
 500 505 510

Lys Leu Arg Ala Gln Gln Cys Arg Phe Trp Thr Ser Phe Phe Pro Lys
 515 520 525

Val Leu Glu Met Thr Gly Asn Ile Asp Glu Ala Glu Trp Glu Trp Lys
 530 535 540

Ala Gly Phe His Arg Trp Asn Asn Tyr Met Met Asp Trp Lys Asn Gln
 545 550 555 560

Phe Asn Asp Tyr Thr Ser Lys Lys Glu Ser Cys Val Gly Leu
 565 570

<210> SEQ ID NO 27
 <211> LENGTH: 1125
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 12-4 mutant (A199S/F227A/S287G/A328W/E441D)
 BChE nucleic acid sequence for amino acid residues 68-442

<400> SEQUENCE: 27

```

aacatagatc aaagttttcc aggcttccat ggatcagaga tgtggaaccc aaacactgac      60
ctcagtgaag actgtttata tctaaatgta tggattccag cacctaaacc aaaaaatgcc      120
actgtattga tatggattta tgggtgggtt tttcaaacctg gaacatcatc tttacatggt      180
tatgatggca agtttctggc tcgggttgaa agagtattg tagtgtcaat gaactatagg      240
gtgggtgccc taggattctt agctttgccca ggaaatcctg aggctccagg gaacatgggt      300
ttatttgatc aacagttggc tcttcagtgg gttcaaaaaa atatagcagc ctttggtgga      360
aatcctaaaa gtgtaactct ctttggagaa agttccggag cagcttcagt tagcctgcat      420
ttgctttctc ctggaagcca ttcattgttc accagagcca tctgcaaag tggttccgct      480
aatgctcctt gggcggtaac atctctttat gaagctagga acagaacggt gaacttagct      540
aaattgactg gttgctctag agagaatgag actgaaataa tcaagtgtct tagaaataaa      600
gatccccaag aaattcttct gaatgaagca tttgtgttcc cctatgggac tcctttgggt      660
gtaaactttg gtccgaccgt ggatgggtgat tttctcactg acatgccaga catattactt      720
gaacttgga c aatttaaaaa aaccagatt ttggtgggtg ttaataaaga tgaagggaca      780
tggtttttag tctatggtgc tcctggcttc agcaaagata acaatagtat cataactaga      840
aaagaatttc aggaagggtt aaaaatattt tttccaggag tgagtgagtt tggaaaggaa      900
tccatccttt ttcattacac agactgggta gatgatcaga gacctgaaaa ctaccgtgag      960
gccttgggtg atgttgttgg ggattataat ttcatatgcc ctgccttggga gttaccaag     1020
aagtctcag aatggggaaa taatgccttt ttctactatt ttgaacaccg atcctccaaa     1080
cttccgtggc cagaatggat gggagtgatg catggctatg acatt                       1125

```

<210> SEQ ID NO 28
 <211> LENGTH: 375
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: 12-4 mutant (A199S/F227A/S287G/A328W/E441D)

-continued

EChE amino acid sequence for residues 68-442

<400> SEQUENCE: 28

Asn Ile Asp Gln Ser Phe Pro Gly Phe His Gly Ser Glu Met Trp Asn
 1 5 10 15
 Pro Asn Thr Asp Leu Ser Glu Asp Cys Leu Tyr Leu Asn Val Trp Ile
 20 25 30
 Pro Ala Pro Lys Pro Lys Asn Ala Thr Val Leu Ile Trp Ile Tyr Gly
 35 40 45
 Gly Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys
 50 55 60
 Phe Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg
 65 70 75 80
 Val Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro
 85 90 95
 Gly Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln
 100 105 110
 Lys Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe
 115 120 125
 Gly Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro
 130 135 140
 Gly Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Ala
 145 150 155 160
 Asn Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg Asn Arg Thr
 165 170 175
 Leu Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn Glu Thr Glu
 180 185 190
 Ile Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile Leu Leu Asn
 195 200 205
 Glu Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Gly Val Asn Phe Gly
 210 215 220
 Pro Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp Ile Leu Leu
 225 230 235 240
 Glu Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly Val Asn Lys
 245 250 255
 Asp Glu Gly Thr Trp Phe Leu Val Tyr Gly Ala Pro Gly Phe Ser Lys
 260 265 270
 Asp Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu Gly Leu Lys
 275 280 285
 Ile Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser Ile Leu Phe
 290 295 300
 His Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn Tyr Arg Glu
 305 310 315 320
 Ala Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys Pro Ala Leu
 325 330 335
 Glu Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala Phe Phe Tyr
 340 345 350
 Tyr Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu Trp Met Gly
 355 360 365
 Val Met His Gly Tyr Asp Ile
 370 375

<210> SEQ ID NO 29

<211> LENGTH: 975

<212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 12-4 mutant (A199S/F227A/S287G/A328W/E441D)
      BChE nucleic acid sequence for amino acid residues 117-441

<400> SEQUENCE: 29
ggttttcaaa ctggaacatc atctttacat gtttatgatg gcaagtttct ggctcggggt    60
gaaagagtta ttgtagtgtc aatgaactat aggggtgggtg ccctaggatt cttagctttg    120
ccaggaaatc ctgaggtccc agggaacatg ggtttatttg atcaacagtt ggctcttcag    180
tgggttcaaa aaaatatagc agcctttggt ggaaatccta aaagtgtaac tctctttgga    240
gaaagtccg gagcagcttc agttagcctg catttgcttt ctctggaag ccattcattg    300
ttcaccagag ccattctgca aagtgggtcc gctaatgctc cttgggcggt aacatctctt    360
tatgaagcta ggaacagaac gttgaactta gctaaattga ctggttgctc tagagagaat    420
gagactgaaa taatcaagtg tcttagaagt aaagatcccc aagaaattct tctgaatgaa    480
gcatttggtg tccccatagg gactcctttg ggtgtaaact ttggtccgac cgtggatggt    540
gattttctca ctgacatgcc agacatatta cttgaacttg gacaatttaa aaaaacccag    600
attttggtgg gtgtaataaa agatgaaggg acatggtttt tagtctatgg tgctcctggc    660
ttcagcaaag ataacaatag tatcataact agaaaagaat ttcaggaagg tttaaaaata    720
ttttttccag gagtgagtga gtttgaaaag gaatccatcc tttttcatta cacagactgg    780
gtagatgatc agagacctga aaactaccgt gaggccttgg gtgatgttgt tggggattat    840
aatttcatat gccttcgctt ggagttcacc aagaagttct cagaatgggg aaataatgcc    900
tttttctact attttgaaca ccgatcctcc aaacttccgt ggccagaatg gatgggagtg    960
atgcatggct atgac                                                    975

```

```

<210> SEQ ID NO 30
<211> LENGTH: 325
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: 12-4 mutant (A199S/F227A/S287G/A328W/E441D)
      BChE amino acid sequence for residues 117-441

```

```

<400> SEQUENCE: 30
Gly Phe Gln Thr Gly Thr Ser Ser Leu His Val Tyr Asp Gly Lys Phe
1           5           10          15
Leu Ala Arg Val Glu Arg Val Ile Val Val Ser Met Asn Tyr Arg Val
20          25          30
Gly Ala Leu Gly Phe Leu Ala Leu Pro Gly Asn Pro Glu Ala Pro Gly
35          40          45
Asn Met Gly Leu Phe Asp Gln Gln Leu Ala Leu Gln Trp Val Gln Lys
50          55          60
Asn Ile Ala Ala Phe Gly Gly Asn Pro Lys Ser Val Thr Leu Phe Gly
65          70          75          80
Glu Ser Ser Gly Ala Ala Ser Val Ser Leu His Leu Leu Ser Pro Gly
85          90          95
Ser His Ser Leu Phe Thr Arg Ala Ile Leu Gln Ser Gly Ser Ala Asn
100         105         110
Ala Pro Trp Ala Val Thr Ser Leu Tyr Glu Ala Arg Asn Arg Thr Leu
115         120         125
Asn Leu Ala Lys Leu Thr Gly Cys Ser Arg Glu Asn Glu Thr Glu Ile
130         135         140
Ile Lys Cys Leu Arg Asn Lys Asp Pro Gln Glu Ile Leu Leu Asn Glu

```

-continued

145	150	155	160
Ala Phe Val Val Pro Tyr Gly Thr Pro Leu Gly Val Asn Phe Gly Pro	165	170	175
Thr Val Asp Gly Asp Phe Leu Thr Asp Met Pro Asp Ile Leu Leu Glu	180	185	190
Leu Gly Gln Phe Lys Lys Thr Gln Ile Leu Val Gly Val Asn Lys Asp	195	200	205
Glu Gly Thr Trp Phe Leu Val Tyr Gly Ala Pro Gly Phe Ser Lys Asp	210	215	220
Asn Asn Ser Ile Ile Thr Arg Lys Glu Phe Gln Glu Gly Leu Lys Ile	225	230	235
Phe Phe Pro Gly Val Ser Glu Phe Gly Lys Glu Ser Ile Leu Phe His	245	250	255
Tyr Thr Asp Trp Val Asp Asp Gln Arg Pro Glu Asn Tyr Arg Glu Ala	260	265	270
Leu Gly Asp Val Val Gly Asp Tyr Asn Phe Ile Cys Pro Ala Leu Glu	275	280	285
Phe Thr Lys Lys Phe Ser Glu Trp Gly Asn Asn Ala Phe Phe Tyr Tyr	290	295	300
Phe Glu His Arg Ser Ser Lys Leu Pro Trp Pro Glu Trp Met Gly Val	305	310	315
Met His Gly Tyr Asp	325		

What is claimed is:

1. A butyrylcholinesterase variant peptide comprising the amino acid sequence of SEQ ID NO: 26.

2. A pharmaceutical composition comprising:
 a butyrylcholinesterase variant peptide comprising the amino acid sequence of SEQ ID NO: 26, and
 a suitable pharmaceutical carrier.

3. A method of treating a cocaine-induced condition comprising administering to an individual an effective amount of the butyrylcholinesterase variant polypeptide of claim 1 to lower blood cocaine concentration.

4. The method of claim 3, wherein said butyrylcholinesterase variant polypeptide exhibits a one-hundred-fold or more increase in cocaine hydrolysis catalytic efficiency compared to butyrylcholinesterase.

5. A method of treating a cocaine-induced condition comprising administering to an individual an effective amount of the BChE butyrylcholinesterase pharmaceutical composition of claim 2 to lower blood cocaine concentration.

* * * * *