



[The VAD Journal: The journal of mechanical assisted circulation and heart failure](#)

Review

Inotropes are linked to Increased Mortality in Heart Failure

M Chadi Alraies¹, Bill Tran¹ and Sirtaz Adatya^{2*}

¹ Department of Medicine, Division of Cardiovascular Medicine, University of Minnesota, Minneapolis, Minnesota, USA

² Department of Medicine, Division of Cardiovascular Medicine, University of Chicago, Chicago Illinois.

* Corresponding author: sadatya@gmail.com

Keywords

Inotropes; Heart failure; ventricular assist device

Introduction

With the advancement of technology and availability of mechanical circulatory support (MCS) devices for the treatment of acute decompensated systolic HF in the 21st century, the role of inotropes is becoming obsolete. Medical therapy for acute decompensated heart failure (HF) has not changed since 1960's, we still use supplemental oxygen, diuretics, vasodilators and inotropes to improve congestion, cardiac output, end organ perfusion and symptoms related to elevate filling pressures¹. Despite demonstrated improvements in hemodynamics, the uses of inotropes have not demonstrated any improvements in mortality. Mechanical circulatory support (MCS) use is growing rapidly in the USA, in patients with stage D HF both as destination therapy and as a bridge to cardiac transplantation. In both, transplant-eligible and non-transplant-eligible patients, improvement in end-organ perfusion, functional capacity, quality of life and most importantly mortality have been demonstrated. In this perspective paper we are going to discuss the current lack of evidence for inotropic therapy and the evolving benefit of MCS in end-stage systolic HF patients.

Citation: Alrais, M. et al. (2015). Inotropes are linked to Increased Mortality. *VAD Journal*, 1. doi: <http://dx.doi.org/10.13023/VAD.2015.08>

Editor-in-Chief: Maya Guglin, University of Kentucky

Received: June 17, 2015

Accepted: June 25, 2015

Published: June 29, 2015, 2015

© 2015 The Author(s). This is an open access article published under the terms of the [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/) (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided that the original author(s) and the publication source are credited.

Funding: Not applicable

Competing interests: Not applicable



Inotropes in End-Stage HF

Advanced HF with cardiogenic shock, are syndromes characterized in many patients by a reduction in myocardial contractile force. Medical therapy is still considered the mainstay for acute decompensated HF. The guidelines recommend diuretic therapy, which decreases intracardiac filling pressures and relieve congestion²⁻⁴. This has to be accompanied with medications favoring left ventricular reverse remodeling such as angiotensin-converting enzyme inhibitors (ACEI)⁵, angiotensin receptors blockers⁶, beta blockers⁷, aldosterone blockers⁸, and nitrate/hydralazine⁹. As the disease progresses, compensatory mechanisms fail, and cardiac output decreases (stage D HF).

Inotropic therapies are utilized to stabilize acute decompensated HF patients with low cardiac output. The use of inotropes can be divided into three distinct patterns of use: hospital admission for patients presenting with acute decompensated HF and end organ involvement, intermittent home infusions (several times per week at the infusion center), and continuous home infusion as a bridge to cardiac transplantation or for palliation¹⁰⁻¹².

Inotropes have been used for more than fifty years and continues to be the used in patients presenting with low perfusion state due to cardiogenic shock.¹ The Acute HF Global Survey of Standard Treatment (ALARM-HF) global survey of 666 hospitals in nine countries showed that inotropes were used in 39% of all admissions for acute HF¹³. Furthermore, in the mechanical circulatory support (MCS) studies, 72% of patients in the medical arm and 65% of patients in the ventricular assist device arm were on inotropes¹⁴. Inotropes successfully improve cardiac hemodynamics and patients' symptoms by increasing myocardial contractility mainly by increasing intracellular calcium concentrations. Inotropes are crucial in bridging selected patients to cardiac transplantation¹⁵.

Despite improving hemodynamic compromise, inotropic agents are accompanied by increased morbidity and mortality due to increased tachycardia and myocardial oxygen consumption leading to arrhythmia, and myocardial ischemia¹⁶. This negative effect has been demonstrated in both the hospital and outpatient setting.² Beside digoxin, the current American College of Cardiology/American Heart Association, European Society of Cardiology and Heart Failure Society of America guidelines endorse that inotropic agents should be reserved for patients presenting with acute decompensated HF and low-output states and reduced end-organ perfusion, who typically are admitted to an intensive care unit²⁻⁴. Furthermore, the ACCF/AHA guidelines state specifically that use of parenteral inotropic agents in hospitalized patients without documented severe systolic dysfunction, low blood pressure, or impaired perfusion, and evidence of significantly depressed cardiac output, with or without congestion, is potentially harmful. This is based on multiple studies showing evidence for harm in patients with advanced HF¹⁶.

Chronic HF patients who are not eligible for advanced therapies have been shown to have poor survival when treated with inotropes^{17,18}. These finding



are based on retrospective studies and post hoc analysis of prospective studies¹⁶. However, despite the reduced survival, many patients prefer improved quality of life, even at the potential expense of a shorter survival.^{18,19} The decision to use inotropic therapy and the selection of inotropic agent should reflect the realistic goals of therapy for the individual with HF. These goals should be discussed in great detail with the patient and family.

Evidence is sufficient to link inotropes to increased mortality

Based on the current studies that showed increased mortality in patients with advanced HF treated with inotropes, the use of inotropes should be limited to patients with advanced HF who are not candidates to MCS or heart transplantation. In the ADHERE (Acute Decompensated Heart Failure National Registry) study¹⁷, Abraham et al. performed a retrospective analysis of observational patient data from the ADHERE registry where they reviewed over 65,000 cases admitted for acute decompensated HF from 263 US hospitals during October 2001 to July 2003 where they included patients whom had received nitroglycerin, nesiritide, milrinone or dobutamine (n= 15,230) therapy and assessed risk factor and propensity-adjusted odds ratios for in-hospital mortality. What they demonstrated was that use of inotropes doubled the in-hospital mortality compared to use of vasodilators; 12.3% and 13.9% for patients receiving milrinone and dobutamine, respectively, compared to 4.7% and 7.1% for patients receiving nitroglycerin and nesiritide, respectively. Moreover, they also found that length of stay was longer compared to those on vasodilator therapy, 10.9 and 10 days for patients receiving milrinone and dobutamine, respectively, compared to 7.1 and 7.9 days for patients receiving nitroglycerin and nesiritide, respectively.

Inotropic dependence is defined as the failure to wean from inotropes because of imminent (minutes to hours) worsening of the patient's clinical status such that death appeared impending, and the patient is deemed highly unlikely to survive inotrope withdrawal to permit hospital discharge²⁰. In other words, withdrawal of inotropic support in this cohort of patients can be acutely life-threatening^{2,20}.

Furthermore, in the OPTIME-CHF (Outcomes of a Prospective Trial of Intravenous Milrinone for Exacerbations of Chronic HF) trial²¹, Cuff et al. demonstrated in this prospective study that intravenous (IV) milrinone used in chronic HF patients, admitted for acute decompensated HF, increased morbidity associated with hypotension and new atrial arrhythmias but no significant difference in in-hospital mortality or 60-day mortality. As an extension of the OPTIME-CHF trial, Felker et al.²² analyzed the interaction of the etiology of HF (ischemic vs. non-ischemic) with IV milrinone and found that patients with ischemic HF were not only hospitalized longer (13 days vs. 11.7 days) but also had increased 60-day mortality (11.6% vs. 7.5%) compared to patients with non-ischemic cardiomyopathy. In this study, 22% of patients were started on beta-blockers on admission. Thus, the "neutralizing" effect of beta-blockers for milrinone infusion was suspected. A post-hoc analysis was done looking into the



neutralizing effect of milrinone and showed decrease incidence of death or re-hospitalization in non-ischemic group suggesting neutralizing effect. However, the study had limitation since it was retrospective post-hoc analysis and authors did not clarify the definition of ischemic cardiomyopathy. Furthermore, double the patients in the non-ischemic cardiomyopathy group were lost to follow up²³.

In spite of increased mortality seen in the inpatient setting as seen in the ADHERE study, the consideration of intermittent inotropic therapy has also been studied in the outpatient setting. With a varying array of treatment regimens as well as administration from daily to weekly regimens, some with concomitant use of amiodarone as an antiarrhythmic, there has been some evidence of improved symptoms, decreased hospitalization and short term mortality, but the studies were small and observational in nature and failed to demonstrate “true” benefit of intermittent inotropic therapy^{18,24,25}.

Inotropes Use in Palliative and Hospice Care

Due to lack of evidence for mortality benefit association with inotropes in ADHF patients, inotropes were studied in hospice care and palliative care settings.²⁰ In one prospective study, the author suggested that intermittent infusion of intravenous inotropes is can modify end-stage HF symptoms, improve functional class and may also facilitate better utilization of hospice and palliative care resources among patients with end-stage heart.²⁶ The current HF guidelines suggest that long-term, continuous intravenous inotropic support may be considered as palliative therapy for symptom control in select patients with stage D HF despite optimal medical therapy and device therapy who are not eligible for either MCS or cardiac transplantation.²

Mechanical Circulatory Assist Devices – A Game Changer in End-Stage HF management

Prior to the advent of left ventricular assist devices (LVAD), patients with end-stage HF who were ineligible for heart transplant did not have many options except for palliative care or palliative inotropic therapy which was not a viable long term treatment strategy. To evaluate the efficacy of LVADs, the REMATCH (Randomized Evaluation of Mechanical Assistance for the Treatment of Congestive HF) trial from May 1998 to July 2001 randomized 128 patients to either LVAD or optimal medical therapy¹⁴. Their findings were remarkable. Survival at one year was 52% in the device group and 25% in the medical group. The observed mortality of 75% at 1 year in the medical group approaches mortality rates seen in pancreatic cancer. Of note, however, patients in the device group were twice as likely to have a serious adverse effect, namely, infection, bleeding or malfunction of device, but overall had improved quality of life compared to medical group. Moreover, the INTrEPID (Investigation of Non-transplant Eligible Patients who are Inotropic Dependent)²⁷ trial which was a prospective nonrandomized clinical trial comparing LVAD to optimal medical therapy enrolled 55 patients between March 2000 and May 2003 who were inotropic dependent, demonstrated that survival at 6 months was 46% in the



device group and 22% in the medical group and survival at 1 year was 27% in the device group and 11% in the medical group. Further, 85% of the LVAD patients demonstrated either no symptoms or minimal HF symptoms compared to the medical group, whom experienced no improvement in functional NYHA class. Results of the REMATCH and INTrEPID trial demonstrated that patients with end-stage HF have dismal survival rates on optimal medical therapy, but do far much better with mechanical support as well as have improved quality of life.

With evidence of the benefits of mechanical support compared to inotropic therapy for end-stage HF, it is essential to consider mechanical support as the first line treatment in patients with end-stage HF, with inotropes being reserved to acutely stabilize the patient as a bridge to MCS or for palliation. Further, future research should not be directed to development of novel inotropic agents as by definition any drug that increases myocardial contractility will result in increase myocardial oxygen demand, which is associate with worse outcomes. Medical therapy in end stage HF is obsolete and this is the era of mechanical circulatory support. Although inotropes may be useful as a short-term strategy to help temporize acute hemodynamic instability in cardiogenic shock, the clear increase in mortality and morbidity with its use should be weighed against the benefit of a more durable, long-term treatment strategy with mechanical assist devices.

References

1. Ramirez A, Abelmann WH. Cardiac decompensation. *N Engl J Med.* 1974;290(9):499-501.
2. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: A report of the american college of cardiology foundation/american heart association task force on practice guidelines. *J Am Coll Cardiol.* 2013;62(16):e147-239.
3. McMurray JJ, Adamopoulos S, Anker SD, et al. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The task force for the diagnosis and treatment of acute and chronic heart failure 2012 of the european society of cardiology. developed in collaboration with the heart failure association (HFA) of the ESC. *Eur Heart J.* 2012;33(14):1787-1847.
4. Heart Failure Society of America, Lindenfeld J, Albert NM, et al. HFSA 2010 comprehensive heart failure practice guideline. *J Card Fail.* 2010;16(6):e1-194.
5. Cohn JN, Johnson GR, Shabetai R, et al. Ejection fraction, peak exercise oxygen consumption, cardiothoracic ratio, ventricular arrhythmias, and plasma norepinephrine as determinants of prognosis in heart failure. the V-HeFT VA cooperative studies group. *Circulation.* 1993;87(6 Suppl):VI5-16.
6. McMurray JJ, Östergren J, Swedberg K, et al. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function



taking angiotensin-converting-enzyme inhibitors: The CHARM-added trial. *The Lancet*. 2003;362(9386):767-771.

7. Poole-Wilson PA, Swedberg K, Cleland JG, et al. Comparison of carvedilol and metoprolol on clinical outcomes in patients with chronic heart failure in the carvedilol or metoprolol european trial (COMET): Randomised controlled trial. *The Lancet*. 2003;362(9377):7-13.

8. Pitt B, Bakris G, Ruilope LM, DiCarlo L, Mukherjee R, EPHEBUS Investigators. Serum potassium and clinical outcomes in the eplerenone post-acute myocardial infarction heart failure efficacy and survival study (EPHEBUS). *Circulation*. 2008;118(16):1643-1650.

9. Cohn JN, Johnson G, Ziesche S, et al. A comparison of enalapril with hydralazine–isosorbide dinitrate in the treatment of chronic congestive heart failure. *N Engl J Med*. 1991;325(5):303-310.

10. Packer M, Carver JR, Rodeheffer RJ, et al. Effect of oral milrinone on mortality in severe chronic heart failure. the PROMISE study research group. *N Engl J Med*. 1991;325(21):1468-1475.

11. Alraies MC, Eckman P. Adult heart transplant: Indications and outcomes. *J Thorac Dis*. 2014;6(8):1120-1128.

12. Metra M, Eichhorn E, Abraham WT, et al. Effects of low-dose oral enoximone administration on mortality, morbidity, and exercise capacity in patients with advanced heart failure: The randomized, double-blind, placebo-controlled, parallel group ESSENTIAL trials. *Eur Heart J*. 2009;30(24):3015-3026.

13. Follath F, Yilmaz MB, Delgado JF, et al. Clinical presentation, management and outcomes in the acute heart failure global survey of standard treatment (ALARM-HF). *Intensive Care Med*. 2011;37(4):619-626.

14. Rose EA, Gelijns AC, Moskowitz AJ, et al. Long-term use of a left ventricular assist device for end-stage heart failure. *N Engl J Med*. 2001;345(20):1435-1443.

15. Biddle TL, Benotti JR, Creager MA, et al. Comparison of intravenous milrinone and dobutamine for congestive heart failure secondary to either ischemic or dilated cardiomyopathy. *Am J Cardiol*. 1987;59(15):1345-1350.

16. Francis GS, Bartos JA, Adatya S. Inotropes. *J Am Coll Cardiol*. 2014;63(20):2069-2078.

17. Abraham WT, Adams KF, Fonarow GC, et al. In-hospital mortality in patients with acute decompensated heart failure requiring intravenous vasoactive medications: An analysis from the acute decompensated heart failure national registry (ADHERE). *J Am Coll Cardiol*. 2005;46(1):57-64.



18. Gorodeski EZ, Chu EC, Reese JR, Shishehbor MH, Hsich E, Starling RC. Prognosis on chronic dobutamine or milrinone infusions for stage D heart failure. *Circ Heart Fail.* 2009;2(4):320-324.
19. MacIver J, Rao V, Delgado DH, et al. Choices: A study of preferences for end-of-life treatments in patients with advanced heart failure. *J Heart Lung Transplant.* 2008;27(9):1002-1007.
20. Hershberger RE, Nauman D, Walker TL, Dutton D, Burgess D. Care processes and clinical outcomes of continuous outpatient support with inotropes (COSI) in patients with refractory endstage heart failure. *J Card Fail.* 2003;9(3):180-187.
21. Cuffe MS, Califf RM, Adams KF, Jr, et al. Short-term intravenous milrinone for acute exacerbation of chronic heart failure: A randomized controlled trial. *JAMA.* 2002;287(12):1541-1547.
22. Felker GM, Benza RL, Chandler AB, et al. Heart failure etiology and response to milrinone in decompensated heart failure: Results from the OPTIME-HF study. *J Am Coll Cardiol.* 2003;41(6):997-1003.
23. Gattis WA, O'Connor CM, Leimberger JD, Felker GM, Adams KF, Gheorghiade M. Clinical outcomes in patients on beta-blocker therapy admitted with worsening chronic heart failure. *Am J Cardiol.* 2003;91(2):169-174.
24. Drakos SG, Kanakakis JV, Nanas S, et al. Intermittent inotropic infusions combined with prophylactic oral amiodarone for patients with decompensated end-stage heart failure. *J Cardiovasc Pharmacol.* 2009;53(2):157-161.
25. Cesario D, Clark J, Maisel A. Beneficial effects of intermittent home administration of the inotrope/vasodilator milrinone in patients with end-stage congestive heart failure: A preliminary study. *Am Heart J.* 1998;135(1):121-129.
26. López - Candales A, Carron C, Schwartz J. Need for hospice and palliative care services in patients with end - stage heart failure treated with intermittent infusion of inotropes. *Clin Cardiol.* 2004;27(1):23-28.
27. Rogers JG, Butler J, Lansman SL, et al. Chronic mechanical circulatory support for inotrope-dependent heart failure patients who are not transplant candidates: Results of the INTrEPID trial. *J Am Coll Cardiol.* 2007;50(8):741-747.

