



University of Kentucky
UKnowledge

Molecular and Cellular Biochemistry Faculty
Patents

Molecular and Cellular Biochemistry

12-10-1985

Inhibitors of Bacterial Collagenase

Richard E. Galardy
University of Kentucky

Damian Grobelny
University of Kentucky

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/biochem_patents

 Part of the [Medical Biochemistry Commons](#)

Recommended Citation


Galardy, Richard E. and Grobelny, Damian, "Inhibitors of Bacterial Collagenase" (1985). *Molecular and Cellular Biochemistry Faculty Patents*. 20.
https://uknowledge.uky.edu/biochem_patents/20

This Patent is brought to you for free and open access by the Molecular and Cellular Biochemistry at UKnowledge. It has been accepted for inclusion in Molecular and Cellular Biochemistry Faculty Patents by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

3

wherein n is 0 or 1; R is hydrogen, lower alkyl, phenyl lower alkyl, hydroxy phenyl lower alkyl, hydroxy lower alkyl, aminolower alkyl, guanidino lower alkyl, imidazolyl lower alkyl, indolyl lower alkyl, mercapto lower alkyl, lower alkyl mercapto lower alkyl; R₃ is hydrogen; R₄ is hydrogen, lower alkyl, phenyl lower alkyl, hydroxy phenyl lower alkyl, hydroxy lower alkyl, amino lower alkyl, guanidino lower alkyl, imidazolyl lower alkyl, indolyl lower alkyl, mercapto lower alkyl, lower alkyl mercapto lower alkyl; R₃ and R₄ may be connected together to form an alkylene bridge of from 2 to 4 carbon atoms or an alkylene bridge of from 2 to 3 carbon atoms and one sulfur atom; X is O, NR⁵, S where R₅=H or lower alkyl; R₁ is hydrogen, lower alkyl, aralkyl or aryl; and R₂ is hydrogen, lower alkyl, aralkyl or aryl and pharmaceutically acceptable salts thereof.


U.S. Pat. No. 4,379,146 to Greenlee et al discloses a substituted phosphonamide and related compounds useful as converting enzyme inhibitors and as antihypertensives. Compounds are disclosed of the general formula:



wherein R₁ is alkyl or substituted alkyl of C₁-C₆ wherein the substituent is halo, amino, acylamino; aralkyl wherein the alkyl is C₁-C₄ optionally substituted by amino or acylamino and wherein the aryl function is phenyl or naphthyl optionally substituted by halo or hydroxyl; or, heteroaralkyl wherein the alkyl is C₁-C₄ optionally substituted by amino or acylamino and wherein the heteroaryl group can be indolyl or thienyl; R₂ is H, lower alkyl of C₁-C₄, aralkyl such as benzyl; R₃ is lower alkyl of C₁-C₆ optionally substituted by an amino group; R₄ is H, lower alkyl of C₁-C₆, aralkyl such as benzyl; and X is (CH₂)_n wherein n is 1 or 2, CH-OCH₃, CH-OH, or S. Other phosphonamides are taught by U.S. Pat. Nos. 4,100,275; 4,143,134; and 4,316,896.

SUMMARY OF THE INVENTION


In a composition aspect, the present invention refers to a bacterial collagenase inhibitor that is an amino acid derivative of the general formula



wherein R is alkyl, aralkyl, aryl, alkoxy, aryloxy, hydroxy or pharmaceutically acceptable salts thereof; R₁ is hydrogen, alkali metal, lower alkyl, phenyl lower alkyl, or phenyl, or substituted phenyl wherein the substituent is halo, lower alkoxy or lower alkyl or nitro or carboxy or cyano, and pharmaceutically acceptable salts thereof; R₂, R₃ and R₄ is hydrogen or alkyl; R₂ and R₄ may be the side groups found on the 20 common α-amino acids; R₃ and R₄ may be connected together to form an alkylene bridge of from 2 to 4 carbon atoms; Y

4

is alkoxy, aryloxy, primary amide, amino acid derivative or




where R₅ and R₆ is hydrogen or alkyl; R₅ and R₆ may be connected together to form an alkylene bridge of from 2 to 4 carbon atoms; and X is hydroxyl or pharmaceutically acceptable salts thereof, or an amino acid derivative.

In a further composition aspect, this invention relates to pharmaceutical compositions wherein the compounds disclosed above comprise the effective ingredient. In a method of use aspect, the present invention relates to the use of compounds as disclosed above for the in vivo inhibition of bacterially produced collagenase.


Further aspects of the present invention relate to the use of the above-disclosed compounds as a ligand for chromatographic packing materials, and as a collagenase inhibitor in the preparation of cell dispersions.

DETAILED DESCRIPTION OF THE INVENTION

Bacterial collagenase is inhibited by amino acid derivatives of the general formula I



wherein R is alkyl, aralkyl, aryl, alkoxy, aryloxy, hydroxy or pharmaceutically acceptable salts thereof; R₁ is hydrogen, alkali metal, lower alkyl, phenyl lower alkyl, or phenyl, or substituted phenyl wherein the substituent is halo, lower alkoxy or lower alkyl or nitro or carboxy or cyano, and pharmaceutically acceptable salts thereof; R₂, R₃ and R₄ is hydrogen or alkyl; R₂ and R₄ may be the side groups found on the 20 common α-amino acids; R₃ and R₄ may also be connected together to form an alkylene bridge of from 2 to 4 carbon atoms; Y is alkoxy, aryloxy, primary amide, amino acid derivative or



where R₅ and R₆ is hydrogen or alkyl; R₅ and R₆ may be connected together to form an alkylene bridge of from 2 to 4 carbon atoms; and X is hydroxyl or pharmaceutically acceptable salts thereof, or an amino acid derivative.

The term "lower alkyl" when used herein and in the appended claims includes straight and branched chain hydrocarbon radicals having from 1 to about 6 carbon atoms. Illustrative examples of which are methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, n-pentyl, isopentyl (isoamyl), n-hexyl, 3-methyl pentyl, 2,3-dimethyl butyl, etc. The term "aryl" includes both carbocyclic and heterocyclic (that is, containing one or more O, S, and N atoms as ring members) aromatic ring systems which may be substituted, for example, with one or more of halogen, fluorine, chlorine, bromine or iodine, trifluoro-

methyl, lower alkyl, lower alkoxy, lower alkanoyloxy, carbonyloxyalkyl, nitro, and cyano. Illustrative aromatic ring systems are phenyl, naphthyl, furyl, thienyl, pyrrolyl, imidazolyl, pyridyl, pyrimidinyl, indolyl, quinolyl, oxazolyl, and isooxazolyl. Lower alkoxy is a lower alkyl as defined above linked by an ether oxygen atom. Examples of lower alkoxy groups are methoxy, ethoxy, n-propoxy, n-butoxy, isopropoxy, tert-butoxy and so forth. The term "pharmacologically acceptable salt" contemplates the usual and customary forms of medicaments formulated in a solid or oral dosage form for pharmacological purposes. It also includes those employed to formulate either in unit dose or multidose form, oral and injectable suspensions and solutions, either directly or for reconstitution before administration.

The compounds of this invention can be utilized to block the action of collagenase of bacterial origin in mammals, including humans. Thus the compounds of this invention can be administered in the form of a physiologically acceptable salt in a physiologically acceptable vehicle or carrier by injection or in the form of tablets or capsules, for example.


The preferred dosage range of the compositions of the present invention is about 5 mg/kg to about 15 mg/kg. The standard dosage for oral administration preferably ranges from 100 to 500 mg, and most preferably 250-500 mg. Reduced dosages for children and infants are preferred. Administration may be repeated during a 24-hour period.

Dosages for administration according to this invention may be compounded into oral dosage forms such as tablets, capsules, and the like. This is done by combining the compounds with conventional carriers and other excipients, such as magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, low melting wax, cocoa butter, and the like. Diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, tabletdisintegrating agents and the like may also be compounded with the compositions of the present invention. Active ingredients may be encapsulated with or without other carriers. In all cases, the proportion of active ingredients in these compositions, whether solid or liquid, will be at least sufficient to impart collagenase inhibitory activity in vivo after oral or parenteral administration.

The dosage requirements vary with the particular compositions employed, the route of administration, the severity of symptoms presented, and the particular subject being treated. Treatment will generally be initiated with small dosages less than the optimum dose of the compound. Throughout, the dosage may be increased until the optimum effect under the circumstances is reached. In general, the tangible embodiments of the invention are most desirably administered at a concentration that will generally afford effective results without causing any harmful or deleterious side effects. The dose will vary depending on these and other such factors which a person skilled in the art will recognize.

The ability of the compounds of the present invention to inhibit bacterial collagenase may be demonstrated by in vitro testing in an enzyme assay using collagenase produced by *Clostridium histolyticum*, for example.

The following examples show the preparation of representative compounds of the present invention. The compounds of the general formula I described above can be prepared by the following equation:



Compound (III) in a salt form is condensed in an inert solvent such as methylene chloride with phosphochloridate II in the presence of a base, such as triethylamine, to obtain ester IV. Specific hydrolysis or hydrogenolysis of ester IV affords the compound of formula I. Amino acids such as alanine or their derivatives such as alanine benzyl ester may be attached at Y at their α -amino group.

In the general formula I, R₂ and R₄ may be the side groups found on the 20 α -amino acids commonly found in proteins (also known as the standard amino acids). These side groups are well known in the art. See for example, FIGS. 4-2 to 4-4 in A. Lehninger, *Biochemistry* 2nd Ed., Worth Publishers, Inc. (1975), which figures are incorporated herein by reference. These side groups include straight and branched chain alkyl, hydrodroxy alkyl, carboxy alkyl, aralkyl, amino alkyl, carboxamide alkyl, mercapto alkyl, phenyl alkyl, hydroxy phenyl alkyl, guanidino alkyl, imidazolyl alkyl, indolyl alkyl, and pyrrolidinyl. The compounds of the present invention exhibit enhanced binding characteristics when R₂ and R₄ bear these side groups, thereby increasing their resemblance to a collagen or other protein fragment.

Also, specific compounds within the generic grouping of the present invention are also effective as inhibitors of other bacterial enzymes that degrade collagen, such as the elastase produced by the genus *Pseudomonas*. One example of such an elastase inhibitor is the amino acid derivative of the generic formula wherein R=ethyl, R₁=K⁺ or H, R₂=isoamyl, R₃=H, R₄=benzyl, and Y=NH₂.

Products of general formula I have asymmetric carbon atoms to which R₂, R₄ and R₆ are attached, when R₂, R₄ and R₆ are other than hydrogen. The compounds exist in diastereoisomeric forms or racemic mixtures thereof. All of these are within the scope of this invention, the L-isomer with respect to the carbon atom of the amino acids is generally preferred. The compounds of this invention form basic salts with various inorganic and organic bases which are also within the scope of this invention. Such salts include ammonium salts, alkali metal salts, alkaline earth metal salts, salts with organic bases, e.g. dicyclohexylamine, triethylamine and the like. The pharmaceutically acceptable salts are preferred, although other salts are also useful, e.g., in isolating or purifying the product.

Contemplated equivalents of the compounds employed in the compositions of this invention are those phosphorous-containing amino acid derivatives which are effective to inhibit bacterial collagenase. Because the inhibiting effect of the compound, rather than its exact chemical structure, is critical to this invention, it will be apparent to those skilled in the art that other

R-groups can meet the efficacy requirements of the present invention.

The following examples illustrate representative examples of formulations of the compounds of the present invention. In these examples and throughout the specification, parts are by weight unless otherwise indicated.

EXAMPLE 1

N^α-(Dibenzylphosphoryl)-Glycyl-L-Proline Amide

To a fresh solution of dibenzylphosphochloridate prepared from 1.02 ml (4.6 mmoles) of dibenzyl phosphite by the method of Atherton et al., *J. Chem. Soc.*, 1948, 1106-1110, and 0.9 g (8.86 mmoles) of trifluoroacetate salt of glycyl-L-proline amide, 1.11 ml (0.8 mmoles) of triethylamine is added at 0° C. After 20 h at 0° C., the reaction mixture is diluted to 150 ml with methylene chloride and washed successively with 0.1N hydrochloric acid (2×25 ml), water (1×25 ml), 0.1 N sodium hydroxide (2×25 ml), water (2×30 ml) and saturated sodium chloride solution, dried over anhydrous sodium sulfate and evaporated yielding 1.4 g (84% yield) of N^α-(dibenzylphosphoryl-glycyl-L-proline amide, melting point 91°-95° C. H¹-NMR (CDCl₃, TMS): δ1.7-2.2 (m, 4H, Pro CH₂β,γ); 3.0-3.7 (m, 4H, Pro CH₂δ, Gly CH₂); 4.1-4.6 (m, 2H, Pro CH_α, P-NH-); 5.05 (d, 4H, PhCH₂); 6.3 (broad m, 1H, NH₂); 7.3 (m, 11H, Ph, NH₂). Mass spectrometry (methane chemical ionization (m/e): 432 (M+1), 450 (M+29), 472 (M+41).

EXAMPLE 2

N^α-(Phosphoryl)-Glycyl-L-Proline Amide Dipotassium Salt

Hydrogen gas is bubbled for 1 h through a mixture of 0.43 g (1 mmole) of N^α-(dibenzylphosphoryl)-glycyl-L-proline amide and 2 ml of 1.0N potassium hydroxide (2 mmoles) in 10 ml of 50% (v/v) aqueous tetrahydrofuran in the presence of 0.22 g of 10% palladium on carbon. After removal of catalyst by filtration, the filtrate is partially evaporated, frozen and lyophilized to give 0.3 g (91%) of the dipotassium salt of N^α-phosphoryl-glycyl-L-proline amide. H¹-NMR (D₂O): δ2.1 (m, 4H, Pro CH₂β,γ); 3.7 (m, 4H, Gly CH₂, Pro CH₂δ), 4.3 (m, 1H, Pro CH_α).

EXAMPLE 3

N^α-(Dibenzylphosphoryl)-Glycyl-L-Prolyl-L-Alanine Benzyl Ester

When benzyl-glycyl-L-prolyl-L-alaninate hydrochloride is substituted for glycyl-L-proline amide salt in Example 1, the identical process affords N^α-(dibenzylphosphoryl)-glycyl-L-prolyl-L-alanine benzyl ester with yield 82%, melting point 77°-80° C. H¹-NMR (CDCl₃) δ1.35 (d, 3H, Ala CH₃); 1.8-2.2 (m, 4H, Pro CH₂β,γ) 3.2-3.8 (m, 4H, Pro CH₂γ, Gly CH₂) 3.9-4.3 (m, 2H, Ala CH, P-NH); 4.6 (m, 1H, Pro CH_α), 5-5.3 (m, 2H, Ph CH₂); 7.35 (m, 16 H, Ph, Ala NH); MS (m/e): 594 (m+1).

EXAMPLE 4

N^α-(Phosphoryl)-Glycyl-L-Prolyl-L-Alanine Tripotassium Salt

When N^α-(dibenzylphosphoryl)-glycyl-L-prolyl-L-alanine benzyl ester is substituted for N^α-(dibenzylphosphoryl)-glycyl-L-proline amide with 3 ml of 1.0N potassium hydroxide (3 mmoles) in Example 2 the identical process affords N^α-phosphoryl-glycyl-L-prolyl-L-

alanine tripotassium salt. H¹-NMR (D₂O) δ1.61 d, 3H, Ala CH₃), 2.1 (m, 4H, Pro CH₂β,γ); 3.7 (m, 4H, Gly CH₂, Pro CH₂δ); 4.2 (m, 1H, Ala CH_α) 4.45 (m, 1H, Pro CH_α).

EXAMPLE 5

N^α-(O-Benzyl-P-Ethylphosphonyl)-Glycyl-L-Prolyl-L-Alanine Benzyl Ester

When O-benzyl-P-ethylphosphochloridate, prepared from ethylphosphonate dibenzylester, by the method of Petrov et al, 1959, *J. Gen. Chem. USSR* 29, 1465-1467, is substituted for dibenzyl phosphochloridate in Example 3, the identical process affords N^α-(O-benzyl-ethylphosphonyl)-glycyl-L-prolyl-L-alanine benzyl ester as an oil, with yield 65% after purification of the crude product by column chromatography [Silica gel, CHCl₃, (2) CH₃CN(2): i-PrOH (1)] H¹-NMR(CDCl₃ TMS) δ0.8-2.2 (m, 8H, P-CH₂CH₃, Ala CH₃, Pro CH₂β,γ) 2.9-4.0 (m, 5H, Pro CH₂δ, Gly CH₂ (d, 3.6, P-NH); 4.45 (m, 2H, Pro CH, Ala CH); 5.0 (m, 4H, PhCH₂); 7.3 (s, 10H, Ph) 7.45 (broad m, 1H, Ala, NH). MS (m/e: 516 (M+1), 544 (M+29).

EXAMPLE 6

N^α-(Ethylphosphonyl)-Glycyl-L-Prolyl-L-Alanine Dipotassium Salt

When N^α-(O-benzyl-P-ethylphosphonyl)-glycyl-L-prolyl-L-alanine benzyl ester is substituted for N^α-(dibenzylphosphoryl)-glycyl-L-proline amide in Example 2 the identical process affords the titled compound with 100% yield. H¹-NMR (D₂O) δ0.8-1.75 (m, 12H, P CH₂CH₃, Ala CH₃), 2.05 (m, 4H, Pro CH₂β,γ), 3.5 (m, 2H, Pro CH₂δ) 3.65 (d, 2H, Gly CH₂), 4.15 (m, 1H, Ala CH), 4.45 (m, 1H, Pro CH_α).

EXAMPLE 7

N^α-(O-Benzyl-P-iso-Amylphosphonyl)-Glycyl-L-Prolyl-L-Alanine Benzyl Ester

When iso-amylphosphonate dibenzyl ester is substituted for ethylphosphonate dibenzyl ester in Example 5 the identical process affords N^α-(O-benzyl-P-iso-amylphosphonyl)-glycyl-L-prolyl-L-alanine benzyl ester as an oil, with 40% yield after purification of crude product on column chromatography [silica gel, CHCl₃(3): CH₃CN(1): hexane (1): n-ProH(1)]. H¹-NMR(CDCl₃) δ0.8 (d, 6H, i-Am CH₃); 1.25-2.2 (m, 12H, Ala CH₃, i-Am CH₂α,β, CH_γ, Pro CH₂β,γ); 3.15-3.85 (m, 5H, Pro CH₂δ, Gly CH₂, P-NH); 4.5 (m, 2H, Pro CH_α, Ala CH_α); 5.0 (d, 2H, PhCH₂OP); 5.15 (s, 2H, PhCH₂OC); 7.35 (s, 11H, Ph, Ala NH). MS (m/e) 558 (M+1); 586 (M+29).

EXAMPLE 8

N^α-(iso-Amylphosphonyl)-Glycyl-L-Prolyl-L-Alanine Dipotassium Salt

When N^α-(O-benzyl-P-iso-amylphosphonyl)-glycyl-L-prolyl-L-alanine benzyl ester is substituted for N^α-(ethylphosphonyl)-glycyl-L-prolyl-L-alanine benzyl ester in Example 6 the identical process affords the titled compound with 80% yield. H¹-NMR (D₂O) δ0.75 (d, 6H, i-Am CH₃); 1.2-1.6 (m, 5H, i-Am CH₂α,β, CH_γ); 1.95 (m, 4H, Pro CH₂β,γ); 3.3-3.68 (m, 5H, 4H, Pro CH₂δ, Gly CH₂); 3.72-4.4 (m, 3H, Gly, CH₂, Pro CH_α).

EXAMPLE 9

N^α-(O-Benzyl-P-n-Decylphosphonyl)-Glycyl-L-Prolyl-L-Alanine Benzyl Ester

When n-decylphosphonate dibenzyl ester is substituted for iso-amylphosphonate dibenzyl ester in Example 7 the identical process affords the titled compound as an oil with 67% yield. H¹-NMR (CDCl₃) δ0.08-2.3 (m, 28H, N-dec, Ala CH₃, Pro CH₂β,γ); 3.0-4.0 (m, 5H, Pro CH₂δ, Gly CH₂, P-NH) 4.4-4.8 (m 2, H, Ala CH, Pro CH); 5.05 (d, 2H, PhCH₂OP) 5.2 (s, 2H, PhCH₂OC); 7.3 (s, 10H, Ph); 7.5 (broad s, 1H, Ala NH). MS (m/e) 6.28 (M+1), 656 (M+29).

EXAMPLE 10

N^α-(n-Decylphosphonyl)-Glycyl-L-Prolyl-L-Alanine Dipotassium Salt

When N^α-(O-Benzyl-P-n-decylphosphonyl)-glycyl-L-prolyl-L-alanine benzyl ester is substituted for N^α-(O-benzyl-P-iso-amylphosphonyl)-glycyl-L-prolyl-L-alanine benzyl ester in Example 8 the identical process affords the titled compound with 89% yield. H¹-NMR (D₂O) δ0.75-1.65 (m, 24H, n-decyl, Ala CH₃); 1.95 (m, 4H, Pro CH₂β,γ); 3.45 (m, 2H, Pro CH₂δ); 4.0 (m, 1H, Ala CHα); 4.25 (m, 1H, Pro CHα).

EXAMPLE 11

N^α-(O-Benzyl-P-iso-Amylphosphonyl)-Glycyl-Glycyl-L-Proline Benzyl Ester

When benzyl glycyl-glycyl-L-prolinate hydrochloride is substituted for benzyl glycyl-L-prolyl-L-alaninate hydrochloride in Example 7 the identical process affords the titled compound as an oil with 33% yield. H¹-NMR (CDCl₃,TMS): δ0.8 (d, 6H, i-Am CH₃); 1.3-2.2 (m, 9H, i-Am CH₂α,β, CHγ, Pro CH₂β,γ); 3.3-4.5 (m, 8H, pro CH₂δ, CHα, Gly CH₂, P-NH); 5.0-5.3 (m, 4H, Ph CH₂); 7.3 (s, 10H, Ph) 7.6 (broad m, 1H, Gly NH). MS (m/e): 544 (M+1); 572 (M+29); 584 (M+41).

EXAMPLE 12

N^α-(iso-Amylphosphonyl)-Glycyl-Glycyl-L-Proline Dipotassium Salt

When N^α-(o-benzyl-P-iso-amylphosphonyl)-glycyl-glycyl-L-proline benzyl ester is substituted for N^α-(O-benzyl-P-n-decylphosphonyl)-glycyl-L-prolyl-L-alanine benzyl ester in Example 10 the identical process affords the title compound with 80% yield. H¹-NMR (D₂O) δ0.75 (d, 6H, i-Am CH₃); 1.2-1.6 (m, 5H, i-Am CH₂α,β, CHγ); 1.95 (m, 4H, Pro CH₂β,γ); 3.3-3.68 (m, 4H, Pro CH₂δ, Gly CH₂); 3.72-4.4 (m, 3H, Gly CH₂, Pro CHα).

EXAMPLE 13

In a conventional manner, fill gelatin capsules with, or prepare compressed tablets containing 150, 350 or 500 mg of N^α-(Dibenzylphosphoryl)-Glycyl-L-Prolyl-L-Alanine benzyl ester per capsule or tablet. Administer one to two capsules or tablets three or four times daily to an adult in order to treat the effects of a Clostridium infection.

EXAMPLE 14

Prepare an alcoholic (8½%) mint-flavored and sweetened solution of 500, 750 or 1000 mg of N^α-(Ethylphosphonyl)-Glycyl-L-Prolyl-L-Alanine Dipotassium Salt per fluid ounce (30 ml). Administer ½ fluid ounce up to

six times daily, or one fluid ounce up to four times daily, every four to six hours for the treatment of bacterial tissue infection.

EXAMPLE 15

The collagenase inhibitors of the present invention may also be useful as in vitro reagents in the production of unicellular suspensions from tissue samples.

Under sterile conditions, excise the distal ⅓ of a raccoon tail. With a sterile razor blade scrape the surface of the tail in order to remove hairs present thereon. Next place the tail portion in a solution of clorox or other bleach for approximately 30 minutes in order to dissolve any remaining hair or surface contaminants. After removing the tail from this bath, rinse three times serially in sterile distilled water. Then dissect away the epidermal portion of the tissue sample, and with forceps and a sterile scalpel, tease away remaining tissue present. This should result in approximately 5 to 15 grams of raccoon tail tissue.

In a sterile petri dish, using scissors and forceps, gently mince the tissue into small fragments. Then with a magnetic stirring bar and stirrer apparatus place the minced tissue in a volume of approximately 20 ml of sterile Ringers lactate, and add 10 ml of 0.05% collagenase solution. Allow the stir bar to rotate for approximately 1 hour, or until the tissue sample is sufficiently disaggregated. Filter the suspension through sterile gauze and isolate further clumps by low speed centrifugation.

To the supernatant add 5.0 ml of 0.1% collagenase inhibitor as disclosed in the present invention, mix thoroughly, and then pellet the cells by higher speed centrifugation. Wash by resuspending the cell pellet three times with serial recentrifugations, resuspend in sterile RPMI 1640 culture solution and add aliquots of appropriate volume to cell culturing vessels and incubate.

EXAMPLE 16


The bacterial collagenase inhibitors of the present invention are also useful in the formation of affinity chromatography columns to bind collagenase. Affinity columns using collagenase inhibitors as a ligand may be prepared in the method of Komiyama et al., "Inhibitory Effects of Phosphoramidon on Neutral Metalloendopeptidases and its Application on Affinity Chromatography", 65(1)352: 357 (1975). One gram of AH-sepharose 4B is washed with 0.5M NaCl and with water is mixed with 77 mg of 1-ethyl-3(3-dimethylaminopropyl) carbodiimide hydrochloride and 50 mg of N^α-(P-iso-amylphosphonyl)-Glycyl-L-Prolyl-L-alanine in 5.0 ml of aqueous solution at pH 5.0. This preparation is left at room temperature with gentle stirring for one day. After reaction, the gel is packed into a small column (1×5 cm), and washed successively with 10 ml of each of the following solutions: 1M NaCl, 0.1M tris-HCl containing 1M NaCl, pH 8.0, 50 mM formic acid containing 1M NaCl, pH 3.0; 0.1M tris-HCl containing 1M NaCl, pH 8.0; 50 mM tris-HCl, pH 8.0; and then equilibrated with 10 mM tris-HCl containing 5 mM CaCl₂, pH 7.0. Under these conditions about 10 micromoles of the inhibitor is coupled with 1 ml of swelled gel. Elutions of bacterial collagenase from the column are performed in a conventional manner.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and without departing from the spirit and


scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. The foregoing preferred specific embodiments are, therefore, to be construed as merely illustrative and not limitative of the disclosure in any way whatsoever.

We claim:

1. A method for inhibiting bacterial collagenase in vivo wherein the bacteria are selected from the group consisting of Clostridium and Pseudomonas, which comprises administering to a human being a collagenase inhibiting effective amount of a pharmaceutical composition including a compound of the formula:



wherein R is alkyl, aralkyl, aryl, alkoxy, aryloxy, hydroxy or pharmaceutically acceptable salts thereof; R₁ is hydrogen, alkali metal, lower alkyl, phenyl lower alkyl, or phenyl, or substituted phenyl wherein the substituent is halo, lower alkoxy or lower alkyl or nitro or carboxy or cyano, and pharmaceutically acceptable salts thereof; R₂, R₃ and R₄ are hydrogen or alkyl; R₂ and R₄ are selected from the group consisting of straight and branched chain alkyl, hydroxy alkyl, carboxy alkyl, aralkyl, amino alkyl, carboxamide alkyl, mercapto alkyl, phenyl alkyl, hydroxy phenyl alkyl, guanidino alkyl, imidazolyl alkyl, indolyl alkyl, and pyrrolidinyl; R₃ and R₄ may be connected together to form an alkylene bridge of from 2 to 4 carbon atoms; Y is alkoxy, aryloxy, primary amide, alanine benzyl ester or



where R₅ and R₆ is hydrogen or alkyl; R₅ and R₆ may be connected together to form an alkylene bridge of from 2 to 4 carbon atoms; and X is hydroxyl or pharmaceutically acceptable salts thereof, in admixture with a pharmaceutically acceptable carrier.


2. A method according to claim 1, wherein the composition is adapted for oral ingestion and is administered orally.

3. A method according to claim 1, wherein the composition is adapted for parenteral administration and is administered parenterally.


4. A method according to claim 1, in the form of a tablet or capsule.

5. A method according to claim 1, in the form of an aqueous solution.

6. A method of inhibiting the action of a bacterial collagenase in vitro wherein the bacteria are selected from the group consisting of clostridium and Pseudomonas, by allowing the collagenase to react with an inhibition-effective amount of a compound of the formula:



wherein R is alkyl, aralkyl, aryl, alkoxy, aryloxy, hydroxy or pharmaceutically acceptable salts thereof; R₁ is hydrogen, alkali metal, lower alkyl, phenyl lower alkyl, or phenyl, or substituted phenyl wherein the substituent is halo, lower alkoxy or lower alkyl or nitro or carboxy or cyano, and pharmaceutically acceptable salts thereof; R₂, R₃ and R₄ are hydrogen or alkyl; R₂ and R₄ are selected from the group consisting of straight and branched chain alkyl, hydroxy alkyl, carboxy alkyl, aralkyl, amino alkyl, carboxamide alkyl, mercapto alkyl, phenyl alkyl, hydroxy phenyl alkyl, guanidino alkyl, imidazolyl alkyl, indolyl alkyl, and pyrrolidinyl; R₃ and R₄ may be connected together to form an alkylene bridge of from 2 to 4 carbon atoms; Y is alkoxy, aryloxy, primary amide, alanine benzyl ester or



where R₅ and R₆ is hydrogen or alkyl; R₅ and R₆ may be connected together to form an alkylene bridge of from 2 to 4 carbon atoms; and X is hydroxyl or pharmaceutically acceptable salts thereof, in admixture with a pharmaceutically acceptable carrier in an appropriate solution.

7. The method of claim 6, wherein R=ethyl, R₁=K⁺ or H, R₂=isoamyl, R₃=H, R₄=benzyl, and Y=NH₂.

8. The method of claim 1, wherein R=isoamyl, R₁=K⁺ or H, R₂=H, R₃+R₄=-CH₂CH₂CH₂-, and Y=Alanine.

9. The method of claim 1, wherein R=ethyl, R₁=K⁺ or H, R₂=H, R₃+R₄=-CH₂CH₂CH₂-, and Y=Alanine.

10. The method of claim 1, wherein R=isoamyl, R₁=benzyl, R₂=H, R₃+R₄=-CH₂CH₂CH₂-, and Y=Alanine benzyl ester.

11. The method of claim 1, wherein R=ethyl, R₁=benzyl, R₂=H, R₃+R₄=-CH₂CH₂CH₂-, and Y=Alanine benzyl ester.

* * * * *

60

65