



University of Kentucky
UKnowledge

Agronomy Notes

Plant and Soil Sciences

5-1968

Chemical Weed Control in Tobacco Fields

James W. Herron
University of Kentucky

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/pss_notes

 Part of the [Agronomy and Crop Sciences Commons](#)

Repository Citation

Herron, James W., "Chemical Weed Control in Tobacco Fields" (1968). *Agronomy Notes*. 181.
https://uknowledge.uky.edu/pss_notes/181

This Report is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in Agronomy Notes by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

*
630.717
Ag 86

CROPS
*SOILS

UNIVERSITY of KENTUCKY • COLLEGE of AGRICULTURE

AGRONOMY NOTES

DEPARTMENT of AGRONOMY Lexington 40506

MAY 10 1968

Vol. 1, No. 16

May 1968

CHEMICAL WEED CONTROL IN TOBACCO FIELDS

James W. Herron

Enide 50W and Vernam 6E are the two herbicides recommended in 1968 for weed control in tobacco fields.

Enide 50W

Enide 50W should be applied at the rate of 12 pounds in 30-50 gallons of water per acre of actual area treated. It can be applied as a broadcast treatment over the entire field or in 20-inch bands over the row. Small grains and grasses sown after tobacco harvest will be damaged by residues of this chemical in the treated soil. Band treatments are often applied to reduce cost and to permit growing a small grain or cover crop in the middles to reduce winter erosion on rolling land.

For small areas, apply 3 ounces of Enide 50W in 3 to 4 quarts of water for band-spraying approximately 400 feet of row.

Enide 50W can be applied over the top or directionally to the base of the plants after transplanting before weed seeds germinate.

Vernam 6E

Vernam 6E should be applied at the rate of 2 quarts in 20-50 gallons of water per acre. It should not be applied until at least 7 days after transplanting. If some weed seeds have germinated, the field should be cleanly cultivated. After clean cultivation, apply as a broadcast spray and incorporate with the soil immediately (within minutes) by shallow cultivation. Another shallow cultivation at lay-by may be needed to destroy escaped weeds.

Enide 50W and Vernam 6E control annual grasses and most annual broad-leaved weeds for a period of 4 to 6 weeks.

For best results, it is important to get uniform coverage of the area treated. Before using any pesticide, be sure to read the precautions on the label.

1,250-5-68