



4-21-2017

Discovery and Fine-Mapping of Adiposity Loci Using High Density Imputation of Genome-Wide Association Studies in Individuals of African Ancestry: African Ancestry Anthropometry Genetics Consortium

Maggie C. Y. Ng
Wake Forest University

Mariaelisa Graff
University of North Carolina at Chapel Hill

Yingchang Lu
Icahn School of Medicine at Mount Sinai

Anne E. Justice
University of North Carolina at Chapel Hill

Poorva Mudgal
Wake Forest University

Repository Citation

Ng, Maggie C. Y.; Graff, Mariaelisa; Lu, Yingchang; Justice, Anne E.; Mudgal, Poorva; Liu, Ching-Ti; Young, Kristin; Yanek, Lisa R.; Feitosa, Mary F.; Wojczynski, Mary K.; Rand, Kristin; Brody, Jennifer A.; Cade, Brian E.; Dimitrov, Latchezar; Duan, Qing; Guo, Xiuqing; Lange, Leslie A.; Nalls, Michael A.; Okut, Hayretin; Tajuddin, Salman M.; Tayo, Bamidele O.; Vedantam, Sailaja; Bradfield, Jonathan P.; Chen, Guanjie; Chen, Wei-Min; Chesi, Alessandra; Irvin, Marguerite R.; Padhukasahasram, Badri; Smith, Jennifer A.; Zheng, Wei; and Arnett, Donna K., "Discovery and Fine-Mapping of Adiposity Loci Using High Density Imputation of Genome-Wide Association Studies in Individuals of African Ancestry: African Ancestry Anthropometry Genetics Consortium" (2017). *Epidemiology Faculty Publications*. 20.
https://uknowledge.uky.edu/epidemiology_facpub/20

See next page for additional authors

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/epidemiology_facpub

 Part of the [African Studies Commons](#), [Eastern European Studies Commons](#), [Epidemiology Commons](#), and the [Genetics and Genomics Commons](#)

Authors

Maggie C. Y. Ng, Mariaelisa Graff, Yingchang Lu, Anne E. Justice, Poorva Mudgal, Ching-Ti Liu, Kristin Young, Lisa R. Yanek, Mary F. Feitosa, Mary K. Wojczynski, Kristin Rand, Jennifer A. Brody, Brian E. Cade, Latchezar Dimitrov, Qing Duan, Xiuqing Guo, Leslie A. Lange, Michael A. Nalls, Hayrettin Okut, Salman M. Tajuddin, Bamidele O. Tayo, Sailaja Vedantam, Jonathan P. Bradfield, Guanjie Chen, Wei-Min Chen, Alessandra Chesi, Marguerite R. Irvin, Badri Padhukasahasram, Jennifer A. Smith, Wei Zheng, and Donna K. Arnett

Discovery and Fine-Mapping of Adiposity Loci Using High Density Imputation of Genome-Wide Association Studies in Individuals of African Ancestry: African Ancestry Anthropometry Genetics Consortium**Notes/Citation Information**

Published in *PLOS Genetics*, v. 13, 4, e1006719, p. 1-25.

This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](#) public domain dedication.

Due to the large number of authors, only the first 30 and the authors affiliated with the University of Kentucky are listed in the author section above. For the complete list of authors, please download this article.


Digital Object Identifier (DOI)

<https://doi.org/10.1371/journal.pgen.1006719>

RESEARCH ARTICLE

Discovery and fine-mapping of adiposity loci using high density imputation of genome-wide association studies in individuals of African ancestry: African Ancestry Anthropometry Genetics Consortium

Maggie C. Y. Ng^{1,2}*, Mariaelisa Graff³*, Yingchang Lu⁴, Anne E. Justice³, Poorva Mudgal², Ching-Ti Liu⁵, Kristin Young³, Lisa R. Yanek⁶, Mary F. Feitosa⁷, Mary K. Wojczynski⁷, Kristin Rand⁸, Jennifer A. Brody⁹, Brian E. Cade^{10,11}, Latchezar Dimitrov¹, Qing Duan¹², Xiuqing Guo¹³, Leslie A. Lange¹², Michael A. Nalls^{14,15}, Hayrettin Okut², Salman M. Tajuddin¹⁶, Bamidele O. Tayo¹⁷, Sailaja Vedantam^{18,19}, Jonathan P. Bradfield²⁰, Guanjie Chen²¹, Wei-Min Chen²², Alessandra Chesi²³, Marguerite R. Irvin²⁴, Badri Padhukasahasram²⁵, Jennifer A. Smith²⁶, Wei Zheng²⁷, Matthew A. Allison²⁸, Christine B. Ambrosone²⁹, Elisa V. Bandera³⁰, Traci M. Bartz³¹, Sonja I. Berndt³², Leslie Bernstein³³, William J. Blot³⁴, Erwin P. Bottinger⁴, John Carpten³⁵, Stephen J. Chanock³², Yii-Der Ida Chen¹³, David V. Conti⁸, Richard S. Cooper¹⁷, Myriam Fornage³⁶, Barry I. Freedman³⁷, Melissa Garcia¹⁶, Phyllis J. Goodman³⁸, Yu-Han H. Hsu^{18,19,39}, Jennifer Hu^{40,41}, Chad D. Huff⁴², Sue A. Ingles^{8,43}, Esther M. John^{44,45}, Rick Kittles⁴⁶, Eric Klein⁴⁷, Jin Li⁴⁸, Barbara McKnight⁴⁹, Uma Nayak⁵⁰, Barbara Nemesure⁵¹, Adesola Ogunniyi⁵², Andrew Olshan^{3,53}, Michael F. Press⁵⁴, Rebecca Rohde³, Benjamin A. Rybicki⁵⁵, Babatunde Salako⁵², Maureen Sanderson⁵⁶, Yaming Shao³, David S. Siscovick⁵⁷, Janet L. Stanford^{58,59}, Victoria L. Stevens⁶⁰, Alex Stram⁸, Sara S. Strom⁴², Dhananjay Vaidya^{6,61}, John S. Witte^{62,63}, Jie Yao¹³, Xiaofeng Zhu⁶⁴, Regina G. Ziegler⁶⁵, Alan B. Zonderman¹⁶, Adebowale Adeyemo²¹, Stefan Ambs⁶⁶, Mary Cushman⁶⁷, Jessica D. Faul⁶⁸, Hakon Hakonarson^{20,69}, Albert M. Levin⁵⁵, Katherine L. Nathanson⁷⁰, Erin B. Ware^{26,68}, David R. Weir⁶⁸, Wei Zhao²⁶, Degui Zhi⁷¹, The Bone Mineral Density in Childhood Study (BMDCS) Group¹, Donna K. Arnett⁷², Struan F. A. Grant^{20,23,69,73}, Sharon L. R. Kardia²⁶, Olufunmilayo I. Oloapde⁷⁴, D. C. Rao⁷⁵, Charles N. Rotimi²¹, Michele M. Sale²², L. Keoki Williams^{25,76}, Babette S. Zemel^{69,77}, Diane M. Becker⁶, Ingrid B. Borecki^{7,78}, Michele K. Evans¹⁶, Tamara B. Harris¹⁶, Joel N. Hirschhorn^{18,19,79}, Yun Li^{12,80,81}, Sanjay R. Patel⁸², Bruce M. Psaty^{83,84}, Jerome I. Rotter^{13,85}, James G. Wilson⁸⁶, Donald W. Bowden^{1,2,87}, L. Adrienne Cupples^{5,88}, Christopher A. Haiman^{8,43†*}, Ruth J. F. Loos^{4,89†*}, Kari E. North^{3†*}



OPEN ACCESS

Citation: Ng MCY, Graff M, Lu Y, Justice AE, Mudgal P, Liu C-T, et al. (2017) Discovery and fine-mapping of adiposity loci using high density imputation of genome-wide association studies in individuals of African ancestry: African Ancestry Anthropometry Genetics Consortium. *PLoS Genet* 13(4): e1006719. <https://doi.org/10.1371/journal.pgen.1006719>

Editor: Gregory P. Copenhaver, The University of North Carolina at Chapel Hill, UNITED STATES

Received: December 8, 2016

Accepted: March 29, 2017

Published: April 21, 2017

Copyright: This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the [Creative Commons CC0](https://creativecommons.org/licenses/by/4.0/) public domain dedication.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files. Summary association statistics from meta-analyses are available from dbGAP at accession number phs000930.

Funding: AABC-MEC: The MEC and genotyping of samples for the GWAS of breast and prostate cancer was supported by National Institutes of Health grants CA63464, CA54281, CA164973,

1 Center for Genomics and Personalized Medicine Research, Wake Forest School of Medicine, Winston-Salem, NC, United States of America, **2** Center for Diabetes Research, Wake Forest School of Medicine, Winston-Salem, NC, United States of America, **3** Department of Epidemiology, University of North Carolina, Chapel Hill, NC, United States of America, **4** The Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, United States of America, **5** Department of Biostatistics, Boston University School of Public Health, Boston, MA, United States of America, **6** Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, United States of America, **7** Division of Statistical Genomics, Department of Genetics, Washington University School of Medicine, St. Louis MO, United States of America, **8** Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA, United States of America, **9** Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle, WA, United States of America, **10** Division of Sleep and Circadian Disorders, Brigham and Women's Hospital, Boston, MA, United States of America, **11** Harvard Medical School, Boston, MA, United States of America, **12** Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC, United States of America, **13** Institute for Translational Genomics and Population Sciences, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Torrance, CA, United States of America, **14** Laboratory of Neurogenetics, National Institute on Aging, National Institutes of Health, Bethesda, MD, United States of America, **15** Data Tecnica

CA1326792, CA148085, HG004726 and a Department of Defense Breast Cancer Research Program Era of Hope Scholar Award to CAH (W81XWH-08-1-0383) and the Norris Foundation. AABC-CARE: CARE was supported by National Institute for Child Health and Development contract NO1-HD-3-3175. AABC-WCHS: WCHS is supported by a U.S. Army Medical Research and Materiel Command (USAMRMC) grant DAMD-17-01-0-0334, National Institutes of Health grant CA100598, and the Breast Cancer Research Foundation. AABC-SFBCS: SFBCS was supported by National Institutes of Health grant CA77305 and United States Army Medical Research Program grant DAMD17-96-6071. AABC-NC-BCFR: The Breast Cancer Family Registry (BCFR) was supported by the National Cancer Institute, National Institutes of Health under RFA CA-06-503 and through cooperative agreements with members of the Breast Cancer Family Registry and Principal Investigators, including the Cancer Prevention Institute of California (U01 CA69417). The content of this manuscript does not necessarily reflect the views or policies of the National Cancer Institute or any of the collaborating centers in the BCFR, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government or the BCFR. AABC-CBCS: CBCS is supported by National Institutes of Health Specialized Program of Research Excellence in Breast Cancer grant CA58223 and Center for Environmental Health and Susceptibility, National Institute of Environmental Health Sciences, National Institutes of Health, grant ES10126. AABC-PLCO: Genotyping of the PLCO samples was funded by the Intramural Research Program of the Division of Cancer Epidemiology and Genetics, National Cancer Institute, the National Institutes of Health. The authors thank Drs. Christine Berg and Philip Prorok, Division of Cancer Prevention, National Cancer Institute, the screening center investigators and staff of the PLCO Cancer Screening Trial, Mr. Thomas Riley and staff at Information Management Services, Inc., and Ms. Barbara O'Brien and staff at Westat, Inc. for their contributions to the PLCO Cancer Screening Trial. AABC-NBHS: NBHS is support by National Institutes of Health grant CA100374. NBHS sample preparation was conducted at the Biospecimen Core Lab that is supported in part by the Vanderbilt-Ingram Cancer Center (CA68485). AABC-WFBC: WFBC is supported by National Institutes of Health grant CA73629. AAPC-CPS-II: CPSII is supported by the American Cancer Society. AAPC-KCPCS: KCPCS was supported by National Institutes of Health grants CA056678, CA082664 and CA092579, with additional support

International, Glen Echo, MD, United States of America, **16** National Institute on Aging, National Institutes of Health, Baltimore, MD, United States of America, **17** Department of Public Health Sciences, Stritch School of Medicine, Loyola University Chicago, Maywood, IL, United States of America, **18** Division of Endocrinology and Center for Basic and Translational Obesity Research, Boston Children's Hospital, Boston, MA, United States of America, **19** Broad Institute of MIT and Harvard, Cambridge, MA, United States of America, **20** Center for Applied Genomics, The Children's Hospital of Philadelphia, Philadelphia, PA, United States of America, **21** Center for Research on Genomics and Global Health, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD, United States of America, **22** Department of Public Health Sciences and Center for Public Health Genomics, University of Virginia School of Medicine, Charlottesville, VA, United States of America, **23** Division of Human Genetics, The Children's Hospital of Philadelphia, Philadelphia, PA, United States of America, **24** Department of Epidemiology, University of Alabama at Birmingham, Birmingham, AL, United States of America, **25** Center for Health Policy and Health Services Research, Henry Ford Health System, Detroit, MI, United States of America, **26** Department of Epidemiology, University of Michigan, Ann Arbor, MI, United States of America, **27** Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt University School of Medicine, Nashville, TN, United States of America, **28** Division of Preventive Medicine, Department of Family Medicine and Public Health, University of California San Diego, La Jolla, CA, United States of America, **29** Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, NY, United States of America, **30** Department of Population Science, Rutgers Cancer Institute of New Jersey, New Brunswick, NJ, United States of America, **31** Cardiovascular Health Research Unit, Departments of Medicine and Biostatistics, University of Washington, Seattle, WA, United States of America, **32** Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, MD, United States of America, **33** Beckman Research Institute of the City of Hope, Duarte, CA, United States of America, **34** International Epidemiology Institute, Rockville, MD, United States of America, **35** Department of Translational Genomics, Keck School of Medicine, University of Southern California, Los Angeles, CA, United States of America, **36** Center for Human Genetics, University of Texas Health Science Center at Houston, Houston, TX, United States of America, **37** Department of Internal Medicine, Wake Forest School of Medicine, Winston-Salem, NC, United States of America, **38** SWOG Statistical Center, Fred Hutchinson Cancer Research Center, Seattle, WA, United States of America, **39** Program in Bioinformatics and Integrative Genomics, Harvard Medical School, Boston, MA, United States of America, **40** Sylvester Comprehensive Cancer Center, University of Miami Leonard Miller School of Medicine, Miami, FL, United States of America, **41** Department of Public Health Sciences, University of Miami Leonard Miller School of Medicine, Miami, FL, United States of America, **42** Department of Epidemiology, University of Texas M.D. Anderson Cancer Center, Houston, TX, United States of America, **43** Norris Comprehensive Cancer Center, University of Southern California, Los Angeles, CA, United States of America, **44** Cancer Prevention Institute of California, Fremont, CA, United States of America, **45** Department of Health Research and Policy (Epidemiology) and Stanford Cancer Institute, Stanford University School of Medicine, Stanford, CA, United States of America, **46** Division of Urology, Department of Surgery, The University of Arizona, Tucson, AZ, United States of America, **47** Glickman Urological and Kidney Institute, Cleveland Clinic, Cleveland, OH, United States of America, **48** Division of Cardiovascular Medicine, Department of Medicine, Stanford University School of Medicine, Palo Alto, CA, United States of America, **49** Cardiovascular Health Research Unit, Department of Biostatistics, University of Washington, Seattle, WA, United States of America, **50** Center for Public Health Genomics, University of Virginia School of Medicine, Charlottesville, VA, United States of America, **51** Department of Preventive Medicine, Stony Brook University, Stony Brook, NY, United States of America, **52** Department of Medicine, University of Ibadan, Ibadan, Nigeria, **53** Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, Chapel Hill, NC, United States of America, **54** Department of Pathology and Norris Comprehensive Cancer Center, University of Southern California Keck School of Medicine, Los Angeles, CA, United States of America, **55** Department of Public Health Sciences, Henry Ford Health System, Detroit, MI, United States of America, **56** Department of Family and Community Medicine, Meharry Medical College, Nashville, TN, United States of America, **57** The New York Academy of Medicine, New York, NY, United States of America, **58** Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, United States of America, **59** Department of Epidemiology, School of Public Health, University of Washington, Seattle, WA, United States of America, **60** Epidemiology Research Program, American Cancer Society, Atlanta, GA, United States of America, **61** Department of Epidemiology, Bloomberg School of Public Health, Baltimore, MD, United States of America, **62** Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, CA, United States of America, **63** Institute for Human Genetics, University of California, San Francisco, San Francisco, CA, United States of America, **64** Department of Epidemiology and Biostatistics, Case Western Reserve University, Cleveland, OH, United States of America, **65** Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Bethesda, MD, United States of America, **66** Laboratory of Human Carcinogenesis, National Cancer Institute, Bethesda, MD, United States of America, **67** Department of Medicine, University of Vermont College of Medicine, Burlington, VT, United States of America, **68** Survey Research Center, Institute for Social Research, University of Michigan, Ann Arbor, MI, United States of America, **69** Department of Pediatrics, Perelman School of

from the Fred Hutchinson Cancer Research Center. AAPC-LAAPC: LAAPC was funded by grant 99-00524V-10258 from the Cancer Research Fund, under Interagency Agreement #97-12013 (University of California contract #98-00924V) with the Department of Health Services Cancer Research Program. AAPC-DCPC: DCPC was supported by National Institutes of Health grant S06GM08016 and Department of Defense grants DAMD W81XWH-07-1-0203 and DAMD W81XWH-06-1-0066. AAPC-MEC: The MEC and genotyping of samples for the GWAS of prostate cancer was supported by National Institutes of Health grants CA63464, CA54281, CA164973, CA1326792, CA148085 and HG004726. AAPC-SELECT trial: The SELECT trial is supported by National Cancer Institute grants U10CA37429 and 5UM1CA182883. AAPC-PLCO: See AABC-PLCO. AAPC-GECAP: GECAP was supported by National Institutes of Health grant ES011126. AAPC-SCCS: SCCS is funded by National Institutes of Health grant CA092447. SCCS sample preparation was conducted at the Biospecimen Core Lab that is supported in part by the Vanderbilt-Ingram Cancer Center (CA68485). AAPC-MDA: MDA was supported by grants CA68578, ES007784, DAMD W81XWH-07-1-0645, and CA140388. AAPC-CaP Genes: CaP Genes was supported by CA88164 and CA127298. ARIC: The Atherosclerosis Risk in Communities Study is carried out as a collaborative study supported by National Heart, Lung, and Blood Institute contracts (HHSN268201100005C, HHSN268201100006C, HHSN268201100007C, HHSN268201100008C, HHSN268201100009C, HHSN268201100010C, HHSN268201100011C, and HHSN268201100012C), R01HL087641, R01HL59367 and R01HL086694; National Human Genome Research Institute contract U01HG004402; and National Institutes of Health contract HHSN268200625226C. The authors thank the staff and participants of the ARIC study for their important contributions. Infrastructure was partly supported by Grant Number UL1RR025005, a component of the National Institutes of Health and NIH Roadmap for Medical Research. BioMe: NIH/NHGRI U01HG007417. BMDCS: We thank the investigators Heidi Kalkwarf, Joan Lappe, Sharon Oberfield, Vicente Gilsanz, John Shepherd and Andrea Kelly for their contribution. BMDCS is supported by NIH: R01 HD58886, HD076321, DK094723-01, DK102659-01; NICHD N01-HD-1-3228, -3329, -3330, -3331, -3332, -3333; CTSA program Grant 8 UL1 TR000077. CARDIA: The Coronary Artery Risk Development in Young Adults (CARDIA) study is supported by the NHLBI (contracts HHSN268201300025C, HHSN268201300026C, HHSN268201300027C,

Medicine, University of Pennsylvania, Philadelphia, PA, United States of America, **70** Department of Medicine, University of Pennsylvania, Philadelphia, PA, United States of America, **71** School of Biomedical Informatics, University of Texas Health Science Center at Houston, Houston, TX, United States of America, **72** School of Public Health, University of Kentucky, Lexington, KY, United States of America, **73** Division of Endocrinology, The Children’s Hospital of Philadelphia, Philadelphia, PA, United States of America, **74** Center for Clinical Cancer Genetics, Department of Medicine and Human Genetics, University of Chicago, Chicago, IL, United States of America, **75** Division of Biostatistics, Washington University School of Medicine, St. Louis, MO, United States of America, **76** Department of Internal Medicine, Henry Ford Health System, Detroit, MI, United States of America, **77** Division of Gastroenterology, Hepatology and Nutrition, The Children’s Hospital of Philadelphia, Philadelphia, PA, United States of America, **78** Regeneron Genetics Center, Regeneron Pharmaceuticals, Inc, United States of America, **79** Departments of Genetics and Pediatrics, Harvard Medical School, Boston, MA, United States of America, **80** Department of Biostatistics, University of North Carolina at Chapel Hill, Chapel Hill, NC, United States of America, **81** Department of Computer Science, University of North Carolina at Chapel Hill, Chapel Hill, NC, United States of America, **82** Department of Medicine, University of Pittsburgh, Pittsburgh, PA, United States of America, **83** Cardiovascular Health Research Unit, Departments of Medicine, Epidemiology, and Health Services, University of Washington, Seattle, WA, United States of America, **84** Kaiser Permanente Washington Health Research Institute, Seattle, WA, United States of America, **85** Division of Genomic Outcomes, Departments of Pediatrics and Medicine, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Los Angeles, CA, United States of America, **86** Department of Physiology and Biophysics, University of Mississippi Medical Center, Jackson, MS, United States of America, **87** Department of Biochemistry, Wake Forest School of Medicine, Winston-Salem, NC, United States of America, **88** NHLBI Framingham Heart Study, Framingham, MA, United States of America, **89** The Mindich Child Health and Development Institute, Ichan School of Medicine at Mount Sinai, New York, NY, United States of America

☉ These authors contributed equally to this work.

‡ These authors jointly supervised this work.

¶ Full membership of the Bone Mineral Density in Childhood Study (BMDCS) Group is provided in [S2 Text](#).

* hairman@usc.edu (CAH); Ruth.loos@mssm.edu (RJFL); kari_north@unc.edu (KEN)

Abstract

Genome-wide association studies (GWAS) have identified >300 loci associated with measures of adiposity including body mass index (BMI) and waist-to-hip ratio (adjusted for BMI, WHR_{adjBMI}), but few have been identified through screening of the African ancestry genomes. We performed large scale meta-analyses and replications in up to 52,895 individuals for BMI and up to 23,095 individuals for WHR_{adjBMI} from the African Ancestry Anthropometry Genetics Consortium (AAAGC) using 1000 Genomes phase 1 imputed GWAS to improve coverage of both common and low frequency variants in the low linkage disequilibrium African ancestry genomes. In the sex-combined analyses, we identified one novel locus (*TCF7L2/HABP2*) for WHR_{adjBMI} and eight previously established loci at $P < 5 \times 10^{-8}$: seven for BMI, and one for WHR_{adjBMI} in African ancestry individuals. An additional novel locus (*SPRYD7/DLEU2*) was identified for WHR_{adjBMI} when combined with European GWAS. In the sex-stratified analyses, we identified three novel loci for BMI (*INTS10/LPL* and *MLC1* in men, *IRX4/IRX2* in women) and four for WHR_{adjBMI} (*SSX2IP*, *CASC8*, *PDE3B* and *ZDHHC1/HSD11B2* in women) in individuals of African ancestry or both African and European ancestry. For four of the novel variants, the minor allele frequency was low (<5%). In the trans-ethnic fine mapping of 47 BMI loci and 27 WHR_{adjBMI} loci that were locus-wide significant ($P < 0.05$ adjusted for effective number of variants per locus) from the African ancestry sex-combined and sex-stratified analyses, 26 BMI loci and 17 WHR_{adjBMI} loci contained ≤ 20 variants in the credible sets that jointly account for 99% posterior probability of driving the associations. The lead variants in 13 of these loci had a high probability of being causal. As compared to our previous HapMap imputed GWAS for BMI and WHR_{adjBMI}

HHSN268201300028C, HHSN268201300029C, and HHSN268200900041C); the Intramural Research Program of the National Institute on Aging (NIA); and an intra-agency agreement between the NIA and the NHLBI (grant AG0005). CFS: The Cleveland Family Study (CFS) is supported by the National Institutes of Health: HL046380, HL113338, RR000080. Dr. Cade has been supported by HL007901. CHOP: This research was financially supported by an Institute Development Award from the Children's Hospital of Philadelphia, a Research Development Award from the Cotswold Foundation, NIH grant R01HD056465 and the Daniel B. Burke Endowed Chair for Diabetes Research (SFAG). CHS: This CHS research was supported by NHLBI contracts HHSN268201200036C, HHSN268200800007C, N01HC55222, N01HC85079, N01HC85080, N01HC85081, N01HC85082, N01HC85083, N01HC85086; and NHLBI grants U01HL080295, R01HL085251, R01HL087652, R01HL105756, R01HL103612, R01HL120393 and HL130114 with additional contribution from the National Institute of Neurological Disorders and Stroke (NINDS). Additional support was provided through R01AG023629 from the National Institute on Aging (NIA). A full list of principal CHS investigators and institutions can be found at CHS-NHLBI.org. The provision of genotyping data was supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR000124, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center. Participants provided informed consent for participation in DNA studies. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. FamHS: NIH/NIDDK R01DK089256, and NIH/NHLBI R01HL117078. GeneSTAR: NIH/NHLBI U01 HL72518; NIH/NHLBI HL087698; NIH/NHLBI 58625-01A1; NIH/NHLBI HL071025-01A1; NIH/NINR NR0224103; NIH/NICRR M01-RR000052. HANDLS: Healthy Aging in Neighborhoods of Diversity across the Life Span Study (HANDLS) was supported by the Intramural Research Program of the NIH, National Institute on Aging and the National Center on Minority Health and Health Disparities (project # Z01-AG000513 and human subjects protocol # 2009-149). HealthABC: Health, Aging, and Body Composition Study (Health ABC Study) was supported by NIA contracts N01AG62101, N01AG62103, and N01AG62106. The genome-wide association study was funded by NIA grant 1R01AG032098-01A1 to Wake Forest University Health Sciences and

including up to 71,412 and 27,350 African ancestry individuals, respectively, our results suggest that 1000 Genomes imputation showed modest improvement in identifying GWAS loci including low frequency variants. Trans-ethnic meta-analyses further improved fine mapping of putative causal variants in loci shared between the African and European ancestry populations.

Author summary

Genome-wide association studies (GWAS) have identified >300 genetic regions that influence body size and shape as measured by body mass index (BMI) and waist-to-hip ratio (WHR), respectively, but few have been identified in populations of African ancestry. We conducted large scale high coverage GWAS and replication of these traits in 52,895 and 23,095 individuals of African ancestry, respectively, followed by additional replication in European populations. We identified 10 genome-wide significant loci in all individuals, and an additional seven loci by analyzing men and women separately. We combined African and European ancestry GWAS and were able to narrow down 43 out of 74 African ancestry associated genetic regions to contain small number of putative causal variants. Our results highlight the improvement of applying high density genome coverage and combining multiple ancestries in the identification and refinement of location of genetic regions associated with adiposity traits.

Introduction

Obesity is a worldwide public health epidemic, with current US estimates of 37.9% obese and 7.7% morbidly obese adults [1]. Disparities in obesity rates, as well as rates of comorbidities and mortality, are evident across sex and racial/ethnic groups. Estimates from NHANES for 2013–2014 [1] show that obesity is more prevalent among African Americans (48.5%) than among non-Hispanic Whites (37.1%). In addition, obesity rates are higher among African American women (57.2%) than among African American men (38.2%). For comparison, the obesity rates in non-Hispanic Whites were 38.7% and 35.4%, respectively, for women and men.

Genome-wide association studies (GWAS) in diverse populations have identified > 300 loci associated with measures of adiposity including body mass index (BMI) and waist-to-hip ratio (adjusted for BMI, WHR_{adjBMI}) in populations of European [2–9], African [10–12], and East Asian ancestry [13–15]. The majority of associated variants are common ($MAF > 5\%$) with small effect size, and jointly explain only a fraction of the phenotypic variances [7–8]. It has long been hypothesized that low frequency ($MAF = 0.5\text{--}5\%$) and rare ($MAF < 0.5\%$) variants may also contribute to variability in complex traits. However, these variants are not well captured in previous GWAS imputed to the HapMap reference panel [16–17]. The availability of higher density reference panels such as the 1000 Genomes Project (38M variants in 1092 individuals from phase 1) [18] has demonstrated improved imputation quality in European populations particularly for low frequency variants (aggregate $R^2 \sim 0.6$ for $MAF = 0.5\%$). However its impact is less clear for non-European populations [19]. We took this opportunity to use higher density imputation to reevaluate our previous GWAS for associations with anthropometric traits in individuals of African ancestry (AA) including African Americans and Africans.

genotyping services were provided by the Center for Inherited Disease Research (CIDR). CIDR is fully funded through a federal contract from the National Institutes of Health to The Johns Hopkins University, contract number HHSN268200782096C. This research was supported in part by the Intramural Research Program of the NIH, National Institute on Aging. HRS: HRS is supported by the National Institute on Aging (NIA U01AG009740). The genotyping was funded separately by the National Institute on Aging (RC2 AG036495, RC4 AG039029), and the analysis was funded in part by R03 AG046389. Our genotyping was conducted by the NIH Center for Inherited Disease Research (CIDR) at Johns Hopkins University. Genotyping quality control and final preparation of the data were performed by the Genetics Coordinating Center at the University of Washington. HUFs: The HUFs (Howard University Family Study) was supported by grants S06GM008016-380111 to AA and S06GM008016-320107 to CR, both from the NIGMS/MBRS/SCORE Program. Participant enrollment for the HUFs was carried out at the Howard University General Clinical Research Center (GCRC), which was supported by grant 2M01RR010284 from the National Center for Research Resources (NCR), a component of the National Institutes of Health (NIH). This research was supported in part by the NIH Intramural Research Program in the Center for Research on Genomics and Global Health (CRGGH) with support from the National Human Genome Research Institute, the National Institute of Diabetes and Digestive and Kidney Diseases, the Center for Information Technology, and the Office of the Director at the National Institutes of Health (Z01HG200362). HyperGEN: The HyperGEN network is funded by cooperative agreements (U10) with NHLBI: HL54471, HL54472, HL54473, HL54495, HL54496, HL54497, HL54509, HL54515, and R01 HL55673 from the National Heart, Lung, and Blood Institute. The study involves: University of Utah: (Network Coordinating Center, Field Center, and Molecular Genetics Lab); Univ. of Alabama at Birmingham: (Field Center and Echo Coordinating and Analysis Center); Medical College of Wisconsin: (Echo Genotyping Lab); Boston University: (Field Center); University of Minnesota: (Field Center and Biochemistry Lab); University of North Carolina: (Field Center); Washington University: (Data Coordinating Center); Weil Cornell Medical College: (Echo Reading Center); National Heart, Lung, & Blood Institute. For a complete list of HyperGEN Investigators: <http://www.biostat.wustl.edu/hypergen/Acknowledge.html>. JHS: The Jackson Heart Study is supported by contracts HHSN268201300046C,

The African Ancestry Anthropometry Genetics Consortium (AAAGC) previously identified seven genome-wide significant loci for BMI in up to 71,412 AA individuals, and an additional locus when combined with European ancestry (EA) data from the Genetic Investigation of ANthropometric Traits (GIANT) consortium using GWAS imputed to the HapMap Phase 2 reference panel [11]. No genome-wide significant loci were identified for WHR_{adjBMI} in a GWAS of up to 27,350 AA individuals [12]. The low yield of discovery in AA studies is likely due to their relatively smaller sample sizes in comparison to EA studies [7–8], as well as their lower degree of linkage disequilibrium (LD) and thus poorer imputation quality. Here, we extended our previous work in the AAAGC to perform meta-analyses and replication of GWAS imputed to the 1000 Genomes reference panel in up to 52,895 AA individuals for BMI and up to 23,095 AA individuals for WHR_{adjBMI}. We aimed to 1) discover novel variants, 2) fine map established loci, and 3) evaluate the coverage and contribution of low frequency variants in genetic associations in AA populations.

Results

Study overview

We conducted sex-combined and sex-stratified meta-analyses of GWAS summary statistics across 17 studies for BMI (N = 42,752) and 10 studies for WHR_{adjBMI} (N = 20,384) in AA individuals in stage 1 discovery (S1 and S2 Tables, S1 Fig). Missing genotypes in individual studies were imputed to the 1000 Genomes Project cosmopolitan reference panel (Phase I Integrated Release Version 3, March 2012) [18] using MaCH/minimac [20] or SHAPEIT2/IMPUTEv2 [21–22] (S3 Table). Among all variants with MAF ≥ 0.1% in the largest Women’s Health Initiative (WHI) study, the average info score was 0.81 and 90.5% had imputation info score ≥ 0.3 (S4 Table). Genomic control corrections were applied to each study and after meta-analysis ($\lambda = 1.07$ for BMI, 1.01 for WHR_{adjBMI}) (S3 Table, S2–S5 Figs). Association results for ~18M variants for BMI and ~21M variants for WHR_{adjBMI} were subsequently interrogated further.

From stage 1 meta-analyses, variants associated with BMI (3,241 in all, 1,498 in men, 2,922 in women) and WHR_{adjBMI} (2,496 in all, 1,408 in men, 2,827 in women) at $P < 1 \times 10^{-4}$ were carried forward for replication in AA and EA. Stage 2 included 10,143 AA (2,458 men and 7,685 women) for BMI and 2,711 AA (981 men and 1,730 women) for WHR_{adjBMI} analyses. Stage 3 included 322,154 EA (152,893 men and 171,977 women) for BMI and 210,086 EA (104,079 men and 116,742 women) for WHR_{adjBMI} analyses by imputing HapMap summary statistics results [7–8] to 1000 Genomes [23] (S1 Fig). Meta-analyses were performed to combine either sex-combined or sex-specific results from AA (stages 1+2, $N \leq 57,895$ for BMI, $\leq 23,095$ for WHR_{adjBMI} in sex-combined analyses) and both AA and EA (stages 1+2+3, $N \leq 380,049$ for BMI, $\leq 233,181$ for WHR_{adjBMI} in sex-combined analyses, S6–S9 Figs). Variants that reached genome-wide statistical significance ($P < 5 \times 10^{-8}$) were assessed for generalization of associations with BMI to children in two additional AA cohorts (N = 7,222).

Genome-wide significant loci in meta-analyses

Sex-combined analyses. In the sex-combined meta-analysis of BMI in AA, seven previously established European or African ancestry-derived loci in/near *SEC16B*, *TMEM18*, *GNPDA2*, *GALNT10*, *KLHL32*, *FTO* and *MC4R* reached genome-wide significance ($P < 5 \times 10^{-8}$) (Table 1, S6 and S10A Figs). The rs7708584 variant at *GALNT10* had the lowest P-value ($P = 4.2 \times 10^{-14}$) and was the same lead variant as reported in our previous AA study (S5 Table) [11]. The association at *KLHL32* was specific to the AA population as the lead variant was not statistically significant in EA ($P > 0.05$), consistent with our previous finding (S5 Table) [11]. No additional novel BMI loci were identified after meta-analysis of AA and EA

HHSN268201300047C, HHSN268201300048C, HHSN268201300049C, HHSN268201300050C from the National Heart, Lung, and Blood Institute and the National Institute on Minority Health and Health Disparities. JGW is supported by U54GM115428 from the National Institute of General Medical Sciences. Maywood: NIH: R37-HL045508, R01-HL074166, R01-HL086718, R01-HG003054; analysis also funded through R01-DK075787. MESA: The Multi-Ethnic Study of Atherosclerosis study (MESA) was supported by the Multi-Ethnic Study of Atherosclerosis (MESA) contracts N01-HC-95159, N01-HC-95160, N01-HC-95161, N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169 and by grants UL1-TR-000040 and UL1-RR-025005 from NCCR. Funding for MESA SHARe genotyping was provided by NHLBI Contract N02-HL-6-4278. The provision of genotyping data was supported in part by the National Center for Advancing Translational Sciences, CTSI grant UL1TR001881, and the National Institute of Diabetes and Digestive and Kidney Disease Diabetes Research Center (DRC) grant DK063491 to the Southern California Diabetes Endocrinology Research Center. Nigeria: NIH: R37-HL045508, R01-HL053353, R01-DK075787 and U01-HL054512; analysis also funded through R01-DK075787. ROOT: NIH: R01-CA142996, P50-CA125183, R01-CA89085, and U01-CA161032. American Cancer Society: MRS-13-063-01-TBG and CRP-10-119-01-CCE. SAPPHIRE: NIH/NHLBI R01HL118267; NIH/NIAID R01AI079139; NIH/NIDDK R01DK064695 and R01DK113003; American Asthma Foundation. SIGNET-REGARDS: The Sea Islands Genetics Network (SIGNET) was supported by R01DK084350 (MM Sale). The authors thank the other investigators, the staff, and the participants of the REGARDS study for their valuable contributions. A full list of participating REGARDS investigators and institutions can be found at <http://www.regardsstudy.org>. This study (REGARDS) was supported by a cooperative agreement U01NS041588 from the National Institute of Neurological Disorders and Stroke (NINDS). WFSM: Genotyping services were provided by the Center for Inherited Disease Research (CIDR). CIDR is fully funded through a federal contract from the National Institutes of Health to The Johns Hopkins University, contract number HHSC268200782096C. This work was supported by National Institutes of Health grants R01DK087914, R01DK066358, R01DK053591, R01HL56266, R01DK070941 and by the Wake Forest School of Medicine grant M01RR07122 and Venture Fund 20543. WHI: Funding support for the

data. Two previously reported loci in AA, *ADCY3* and *MIR148A-NFE2L3* [11], did not reach genome-wide significance in the present study. The previously reported lead variant at *ADCY3*, rs7586879, showed weaker effect and association in AA (effect = 0.047, $P = 3.60 \times 10^{-8}$ [11] vs. effect = 0.032, $P = 1.05 \times 10^{-4}$, this study). On the other hand, a moderately correlated ($r^2 = 0.52$ in AFR) variant, rs10203482, show stronger association at $P = 3.35 \times 10^{-7}$ (S5 Table). At *MIR148A/NFE2L3*, the previously reported lead variant identified by meta-analysis of AA and EA, rs10261878, also showed weaker association in the current study primarily due to weak association in EA ($P = 4.10 \times 10^{-5}$ in AA, 4.69×10^{-3} in EA and 2.10×10^{-5} in AA+EA) (S5 Table). For WHR_{adjBMI} , an established locus (*ADAMTS9-AS2*) (S10B Fig) and a novel locus (*TCF7L2/HABP2*) (Fig 1A) showed significant associations, the latter lead variant rs116718588 was low frequency (MAF = 0.045, Table 1). Meta-analyses including both AA and EA individuals revealed an additional novel locus at *SPRYD7/DLEU2* for WHR_{adjBMI} (Table 1, Fig 1A and S7 Fig). Overall, all the BMI associated lead variants were present in HapMap and were in high LD ($r^2 > 0.5$ in 1000 Genomes AFR population) with the lead variants in our previous HapMap imputed data, except for *TMEM18* rs62105306 ($r^2 = 0.17$) which is absent in HapMap. In contrast, all three WHR_{adjBMI} lead variants were absent in HapMap and the lead variants at *ADAMTS9-AS2* and *TCF7L2-HABP2* were in low LD ($r^2 < 0.5$) with the lead variants in our previous HapMap imputed data [11–12]. We used conditional and joint association analyses to examine the genome-wide significant locus for secondary signals, but no additional independent signals were found.

Sex-stratified analyses. In the sex-stratified meta-analysis in AA, four established BMI loci (*SEC16B*, *GALNT10*, *FTO* and *MC4R*) and one established WHR_{adjBMI} locus (*ADAMTS9-AS2*) were genome-wide significant among women (S6 Table, S8 and S9 Figs). *ADAMTS9-AS2* showed a stronger association with WHR_{adjBMI} among women than among men ($P_{het} = 0.02$) (S6 Table), consistent with findings among EA [9]. On the other hand, although our observed *SEC16B* rs543874 effect size differences (0.064 vs. 0.038, $P_{het} = 0.08$) for BMI in women compared to men were similar to those previously observed among EA (0.060 vs. 0.034, $P_{het} = 5.23 \times 10^{-5}$) [7], we did not observe statistically significant differences in effect size, likely due to a much smaller sample size and thus lower statistical power in our study. All these five loci were also genome-wide significant in the sex-combined meta-analyses. They were not further examined in subsequent sex-stratified analyses given their smaller sample sizes compared to the sex-combined analyses. In AA, additional novel loci were observed for association with BMI; these were variants in *IRX4/IRX2* among women, variants in *INTS10/LPL* and *MLC1* among men (Fig 2), and for WHR_{adjBMI} , variants in *SSX2IP* and *PDE3B* among women (Fig 1B, Table 2). In meta-analyses including both AA and EA, two additional novel loci at *CASC8* and *ZDHC1/HSD11B2* were identified for WHR_{adjBMI} in women (Table 2, Fig 1B). Among all loci, the effect sizes of six variants (*IRX4/IRX2*, *INTS10/LPL*, *MLC1*, *ADAMTS9-AS2*, *PDE3B* and *CASC8*) were nominally significant different between men and women in AA ($P_{het} < 0.05$) (S6 Table).

Replication in children. We evaluated the seven sex-combined and three sex-specific genome-wide significant BMI loci for associations in 7,222 AA children (3,552 boys and 3,670 girls). All lead variants displayed directional consistency, and five of these including *SEC16B*, *TMEM18*, *GNPDA2*, *GALNT10* and *MC4R* showed nominal associations with BMI ($P < 0.05$, $P_{binomial} = 4.70 \times 10^{-8}$) (S7 Table), supporting the role of these loci in modulating adiposity in AA children. WHR_{adjBMI} data were not available in the cohorts of children.

Functional characterization of novel loci. We used multiple complementary approaches to elucidate the putative causal genes and/or variants associated with the nine novel BMI and WHR_{adjBMI} loci from the sex-combined and sex-stratified analyses, including annotating nearby coding variants, cis-expression quantitative trait loci (cis-eQTL) analyses, and

“Epidemiology of putative genetic variants: The Women’s Health Initiative” study is provided through the NHGRI PAGE program (U01HG004790 and its NHGRI ARRA supplement). The WHI program is funded by the National Heart, Lung, and Blood Institute; NIH; and U.S. Department of Health and Human Services through contracts N01WH22110, 24152, 32100-2, 32105-6, 32108-9, 32111-13, 32115, 32118-32119, 32122, 42107-26, 42129-32, and 44221. The authors thank the WHI investigators and staff for their dedication, and the study participants for making the program possible. A full listing of WHI investigators can be found at: http://www.whiscience.org/publications/WHI_investigators_shortlist.pdf. CAH: R01 DK101855-01. CL: R01 DK089256. KEN: R01 DK089256, R01 DK101855-01, 15GRNT25880008. KLY: KL2TR001109. LAC: R01 DK089256. RJL: R01 DK101855-01. YLi: R01HG006292 and R01HL129132. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: I have read the journal’s policy and the authors of this manuscript have the following competing interests: BMP serves on the DSMB of a clinical trial funded by the manufacturer and on the Steering Committee for the Yale Open Data Access Project funded by Johnson & Johnson. IB works in Regeneron Pharmaceuticals, Inc. MAN’s participation is supported by a consulting contract between Data Tecnica International and the National Institute on Aging, NIH, Bethesda, MD, USA. As a possible conflict of interest, MAN also consults for Illumina Inc, the Michael J. Fox Foundation and University of California Healthcare.

functional regulatory genomic element analyses. One missense variant in *PLEKHG4*, rs8044843, was in high LD ($r^2 = 0.75$ in AFR) with rs6499129 associated with WHR_{adjBMI} in women (S8 Table). We did not identify any coding variants in high LD ($r^2 > 0.7$) with other lead variants within the flanking 1Mb-regions. Regulatory element analyses using Regulo-meDB [24] and HaploReg [25] revealed that proxies ($r^2 = 0.73–0.84$) to lead variants at three WHR_{adjBMI} loci (*SPRYD7/DLEU2*, *PDE3B*, and *ZDHHC1/HSD11B2*) were associated with transcription factor binding, DNase peak, promoter or enhancer histone marks (S8 Table). In addition, the lead variant rs2472591 at *SPRYD7/DLEU2* was in high LD ($r^2 = 0.85$) with rs790943, a cis-eQTL associated with expression of the nearby gene, *TRIM13*, in blood dendritic cells in tuberculosis patients [26] (S9 Table), suggesting the associations at the *SPRYD7/DLEU2* locus may be involved in the regulation of nearby gene expression at *TRIM13*.

Cross-trait associations of novel loci. We searched the NHGRI-EBI GWAS [27] and Genome-Wide Repository of Associations Between SNPs and Phenotypes (GRASP) [28] catalogs to assess if any of the nine novel lead variants were in high LD with variants that were genome-wide significantly ($P < 5 \times 10^{-8}$) or nominally ($P < 0.05$) associated with related anthropometric and cardiometabolic traits or gene expression in prior studies. Although a few lead variants were physically close (< 500 kb) to GWAS loci for related traits in the NHGRI-EBI GWAS Catalog (Figs 1 and 2), none of our lead variants were in high LD with the previously associated lead variants. Additionally, there were no nearby associations for novel BMI loci in the GRASP Catalog. Of the novel variants associated with WHR_{adjBMI} , rs2472591 at *SPRYD7/DLEU2*, rs378854 near *MYC*, and rs6499129 near *ZDHHC1/HSD11B2* were in high LD ($r^2 > 0.7$) with previously-reported WHR_{adjBMI} variants, but they did not reach genome-wide significance ($P > 2 \times 10^{-5}$) [3] (S9 Table). Other nearby associations with related cardiometabolic traits include chronic kidney disease (CKD), high density lipoprotein cholesterol (HDL-C), anthropometric traits (BMI, height, and birth weight), blood pressure (systolic blood pressure and hypertension), diabetes-related traits (blood glucose and HOMA-IR), and gene expression of several genes (e.g. *ATP6V0D1*, *ZDHHC1*, *DUS2L*, *AGRP*, *GFOD2* and *LRRC29*).

Evaluation of established European loci in African ancestry populations

Conditional analysis in GWAS loci. Among the six BMI (*SEC16B*, *TMEM18*, *GNPDA2*, *GALNT10*, *FTO* and *MC4R*) and one WHR_{adjBMI} (*ADAMTS9-AS2*) genome-wide significant loci in AA that were previously reported in EA [7–8], we tested whether the African derived lead variants were independent of the reported European signals by conditioning on the European lead variants or their surrogates. For three of the BMI loci (*SEC16B*, *GNPDA2* and *MC4R*), our lead variants are the same as those reported in the previous literature [7]. For all other loci, the lead variants demonstrated substantially lower significance upon conditional analysis, suggesting that the African ancestry results represented the same association signals as previously reported in GWAS performed predominantly in EA populations (S10 Table).

SNP transferability. We further examined all sex-combined and sex-stratified BMI and WHR_{adjBMI} loci identified from previous EA studies [7–9] in our AA data. Among 176 EA lead variants from 170 BMI loci, 119 variants displayed directionally consistent associations with BMI in our data, 31 of these were nominally significant at $P < 0.05$ ($P_{binomial} = 2.2 \times 10^{-18}$ among 176 variants). Among 84 EA lead variants from 65 WHR_{adjBMI} loci, 69 variants displayed directionally consistent associations with WHR_{adjBMI} , and 23 of these were nominally significant ($P_{binomial} = 5.3 \times 10^{-19}$ among 84 variants) (S11 Table). EA lead variants in 11 BMI and 3 WHR_{adjBMI} loci showed directional consistency and significant associations after correction for multiple comparisons ($P < 1.92 \times 10^{-4}$). Among the 54 nominally transferable lead

Table 1. Novel and previously identified BMI and WHR_{adjBMI} loci at $P < 5 \times 10^{-8}$ in African ancestry discovery and replication samples, and European ancestry replication samples.

Trait	Lead SNP	Chr	Position (b37/hg19)	Known Locus (if Yes, lead published variant)	Locus	Effect/Other alleles	EAF	AA Discovery			AA Replication			AA Discovery + Replication		EA		AA + EA		
								Effect (SE)	P	HetISq	N	Effect (SE)	P	HetISq	N	Effect (SE)	P	N	P	N
BMI	rs543874	1	177,889,480	Yes rs543874	SEC1B6	G/A	0.248	0.055 (0.008)	5.75E-11	0	42,681	0.057 (0.017)	6.59E-04	28.9	10,143	0.055 (0.008)	1.76E-13	4.36E-35	322,008	6.35E-46
BMI	rs62105306	2	633,660	Yes rs13021737	TMEEM18	T/C	0.751	0.056 (0.01)	1.55E-08	0	41,492	0.04 (0.02)	4.40E-02	49.4	10,143	0.053 (0.009)	2.17E-09	6.10E-21	244,176	3.04E-28
BMI	rs10938397	4	45,182,527	Yes rs10938397	GNPDA2	G/A	0.243	0.053 (0.008)	3.76E-10	3.6	42,752	0.011 (0.017)	5.40E-01	54	10,143	0.044 (0.008)	3.95E-09	1.87E-38	320,955	5.60E-46
BMI	rs7708584	5	153,543,466	Yes rs7715256; rs7708584 ^a	GALNT10	A/G	0.307	0.059 (0.008)	1.05E-13	4.6	42,750	0.034 (0.016)	3.93E-02	6.2	10,143	0.054 (0.007)	4.21E-14	3.80E-07	234,015	4.35E-15
BMI	rs17057164	6	97,410,536	Yes rs974417 ^a	KLHL32	T/C	0.659	0.043 (0.008)	1.75E-08	0	42,751	0.025 (0.015)	9.97E-02	35.2	10,143	0.04 (0.007)	6.08E-09	7.44E-01	233,997	5.43E-03
BMI	rs17817964	16	53,828,066	Yes rs1558902	FTO	T/C	0.117	0.067 (0.011)	5.48E-09	0	42,750	0.08 (0.025)	1.19E-03	49.2	10,143	0.069 (0.01)	2.72E-11	2.40E-139	321,602	1.13E-146
BMI	rs6567160	18	57,829,135	Yes rs6567160	MC4R	C/T	0.197	0.062 (0.009)	2.74E-11	35.3	42,750	0.044 (0.019)	1.99E-02	33.5	10,143	0.059 (0.008)	2.29E-12	8.23E-54	321,958	2.09E-64
WHR _{adjBMI}	rs66815886	3	64,703,394	Yes rs2371767	ADAMTS9-AS2	G/T	0.457	0.07 (0.01)	3.90E-12	0	20,383	0.005 (0.033)	8.75E-01	42.1	2,711	0.064 (0.01)	2.46E-11	5.17E-19	145,257	9.13E-27
WHR _{adjBMI}	rs116718588	10	115,189,239	No	TCF7L2/HABP2	A/G	0.955	0.114 (0.025)	5.88E-06	0	20,384	0.348 (0.084)	3.82E-05	45.6	2,711	0.134 (0.024)	3.22E-08	NA	NA	NA
WHR _{adjBMI}	rs2472591	13	50,536,360	No	SPRYD7/DLEU2	T/A	0.206	0.05 (0.013)	9.72E-05	0	20,371	0.06 (0.049)	2.21E-01	0	2,160	0.05 (0.012)	4.36E-05	1.69E-05	140,431	3.53E-08

AA: African ancestry; BMI: body mass index; Chr: chromosome; EA: European ancestry; EAF: effect allele frequency; HetISq: heterogeneity measured by I-square; SE: standard error; WHR_{adjBMI}: waist-to-hip ratio adjusted for BMI
^a lead published variants reported in African ancestry

<https://doi.org/10.1371/journal.pgen.1006719.t001>




Fig 1. Locuszoom plots of six novel waist-to-hip ratio adjusted for BMI (WHR_{adjBMI}) loci: (A) *TCF7L2/HABP2* and *SPRYD7/DLEU2* in men and women combined; and (B) *SSX2IP*, *PDE3B*, *CASC8*, and *ZDHHC1/HSD11B2* in women only. All plots use AFR LD from the 1000 Genomes phase 1 reference panel. In each plot, the most significant variant within a 1Mb regional locus is highlighted. *P*-values for all variants including the most significant variant are based on the African ancestry discovery phase only (AA-Discovery). In addition, for the most significant variant, *P*-values are annotated and illustrated from the African ancestry discovery and replication phases (AA-Discovery+Replication). SNP rs2472591 was available in the Europeans from the GIANT consortium effort and combined with the African ancestry discovery and replication phases (AA+EA).

<https://doi.org/10.1371/journal.pgen.1006719.g001>




Fig 2. Locuszoom plots for three novel BMI loci: (A) *IRX4/IRX2* in women only; and (B) *INTS10/LPL* and *MLC1* in men only. All plots use AFR LD from the 1000 Genomes phase 1 reference panel. In each plot, the most significant variant within a 1Mb regional locus is highlighted. *P*-values for all variants including the most significant variant are based on the African ancestry discovery phase only (AA-Discovery). In addition, for the most significant variant, *P*-values are annotated and illustrated from the African ancestry discovery and replication phases (AA-Discovery+Replication).

<https://doi.org/10.1371/journal.pgen.1006719.g002>

variants for BMI and WHR_{adjBMI} , 45% and 43% of the effect sizes, respectively, were larger in the EA than the AA populations. In addition, 65% of the frequencies of the trait-raising alleles were higher in the EA populations for both traits. The correlations of both effect sizes and allele frequency of the transferable variants were high (0.74 and 0.79, respectively) for BMI but weak (0.19 and 0.37, respectively) for WHR_{adjBMI} (S11 Fig). The significant but low proportion of lead variants that were transferable from EA to AA (18% for BMI and 27% for WHR_{adjBMI}) suggests either that many loci are not implicated in AA or population differences in LD mask the detection of associated variants in AA. On the other hand, those variants that were transferable explain similar levels of variances for BMI in both populations, but not for WHR_{adjBMI} .

Locus transferability. We further investigated locus transferability in EA loci derived from sex-combined and sex-stratified analyses by considering varying LD between EA and AA populations. S12 Table reports the most significant lead regional variants in our AA sex-combined and sex-stratified data within 0.1cM region of the previously published EA loci (from 176 BMI and 84 WHR_{adjBMI} lead variants) [7–8]. Forty-five (26%) lead regional variants from

Table 2. Additional novel BMI and WHR_{adjBMI} loci at $P < 5 \times 10^{-8}$ in sex-stratified analyses of African ancestry discovery and replication samples.

Trait	Cohort	Lead SNP	Chr	Position (hg19)	Locus	Effect/Other alleles	EAF	AA Discovery				AA Replication				AA Discovery + Replication				EA		AA + EA
								P	Effect (SE)	HetS _q	N	P	Effect (SE)	HetS _q	N	P	Effect (SE)	N	P	N	P	
BMI	Women	rs112778462	5	2,177,693	<i>IRX4/IRX2</i>	A/G	0.023	0.159 (0.035)	35.4	6.06E-06	25,792	0.219 (0.059)	47.6	2.11E-04	6,984	0.175 (0.03)	32,776	7.21E-09	NA	NA	NA	NA
BMI	Men	rs149352150	8	19,736,154	<i>INTS10/LPL</i>	G/A	0.013	0.341 (0.058)	0	4.29E-09	15,179	-0.034 (0.163)	1.0	8.35E-01	2,147	0.299 (0.055)	17,326	4.68E-08	NA	NA	NA	NA
BMI	Men	rs56330886	22	50,493,427	<i>MLC1</i>	G/T	0.189	0.096 (0.016)	3.3	4.81E-09	15,721	0.063 (0.052)	0.2	2.21E-01	2,147	0.093 (0.016)	17,868	2.88E-09	NA	NA	NA	NA
WHR _{adjBMI}	Women	rs140858719	1	85,203,061	<i>SSX2IP</i>	G/A	0.994	0.506 (0.093)	0	5.07E-08	11,314	0.403 (0.487)	0	4.08E-01	834	0.502 (0.091)	12,148	3.69E-08	NA	NA	NA	NA
WHR _{adjBMI}	Women	rs378854	8	128,323,819	<i>CASC8</i>	C/T	0.787	0.062 (0.015)	4.5	3.34E-05	15,600	0.034 (0.054)	73.2	5.29E-01	1,730	0.060 (0.014)	17,330	2.99E-05	3.70E-06	85325	3.26E-09	NA
WHR _{adjBMI}	Women	rs185693786	11	14,804,296	<i>PDE3B</i>	G/A	0.930	0.122 (0.023)	0	2.01E-07	15,601	0.17 (0.065)	21.6	4.50E-02	1,730	0.125 (0.023)	17,331	2.98E-08	NA	NA	NA	NA
WHR _{adjBMI}	Women	rs6499129	16	67,458,251	<i>ZDHC1/HSD11B2</i>	A/C	0.434	0.045 (0.012)	30.9	1.12E-04	15,588	0.071 (0.042)	41.2	9.29E-02	1,730	0.047 (0.011)	17,318	3.07E-05	4.13E-05	86328	4.84E-08	NA

AA: African ancestry; BMI: body mass index; Chr: chromosome; EA: European ancestry; EAF: effect allele frequency; HetS_q: heterogeneity measured by I-square; SE: standard error; WHR_{adjBMI}: waist-to-hip ratio adjusted for BMI

<https://doi.org/10.1371/journal.pgen.1006719.t002>

BMI loci remained significant ($P_{locus} < 0.05$) after adjustment for the number of independent variants tested at each locus. Sixteen (36%) and 22 (49%) of these 45 lead regional variants are in LD ($r^2 > 0.2$) with the EA BMI lead variants using 1000 Genomes AFR and CEU LD, respectively. Twenty-five of these variants are highly correlated with EA lead variants ($r^2 > 0.7$ in CEU) or had ≥ 1 standard error decrease in effect sizes after conditional analyses, representing same association signals as in EA populations. Twenty-one (32%) lead regional variants for WHR_{adjBMI} loci remained significant. Nine (43%) and seven (33%) of the 21 lead regional variants was in LD with the EA lead variant using 1000 Genomes AFR and CEU LD, respectively. Seven of these variants represented the same EA association in conditional analyses (S12 Table).

Fine mapping of novel AA loci and EA-AA transferable established loci

Among the locus-wide significant established loci (44 for BMI given two of 45 lead regional variants were identical in two loci, and 21 for WHR_{adjBMI}), and novel loci (three for BMI and six for WHR_{adjBMI}) derived from the sex-combined and sex-stratified analyses, we performed fine mapping to localize putative causal variants. We constructed 99% credible sets containing variants that jointly account for 99% posterior probability of driving the association in a locus using the corresponding sex-combined or sex-stratified meta-analysis results from AA, EA and combined ancestry (S13 Table). A smaller number of variants in a credible set represent a higher resolution of fine mapping and we considered a credible set containing ≤ 20 variants as ‘tractable’ for follow up. The credible sets in the EA analyses were generally smaller than those in the AA given their larger sample size. As compared to the EA analyses, the number of tractable loci in the meta-analyses of AA and EA increased from 23 to 26 for BMI, and from 14 to 17 for WHR_{adjBMI} .

Among these 43 tractable loci, the lead variants in the combined ancestry analyses had posterior probability ≥ 0.95 in six BMI loci (*SEC16B*, *TLR4*, *STXBP6*, *NLRC3*, *FTO* and *MC4R*) and seven WHR_{adjBMI} loci (*DCST2*, *PPARG*, *ADAMTS9*, *SNX10*, *KLF13*, *CMIP* and *PEMT*) (S13 Table). Functional characterization of variants within the tractable credible sets revealed two loci contain nonsynonymous variants (*ADCY3*: rs11676272 S107P; *SH2B1*: rs7498665 T484A from the *ATP2A1* locus), but they had low posterior probability to drive the respective associations (0.02 and 0.15, respectively) (S14 Table). On the other hand, the *ADCY3* non-coding variants rs10182181 and rs6752378 had higher posterior probability (0.26–0.72) and are cis-eQTLs of *ADCY3* and nearby genes. Several BMI loci including *MTCH2*, *MAP2K5*, *NLRC3* and *ATP2A1*, and WHR_{adjBMI} loci including *TBX15-WARS2* and *FAM13A*, also contained cis-eQTL variants regulating nearby gene expression in subcutaneous and/or visceral adipose tissue (S14 Table).

Discussion

In our large-scale meta-analyses of GWAS in up to 52,895 and 23,095 individuals of African ancestry for BMI and WHR_{adjBMI} , respectively, we identified three novel (*IRX4/IRX2*, *INTS10/LPL* and *MLC1*) and seven established (*SEC16B*, *TMEM18*, *GNPDA2*, *GALNT10*, *KLHL32*, *FTO* and *MC4R*) BMI loci, as well as three novel (*TCF7L2/HABP2*, *SSX2IP* and *PDE3B*) and one established (*ADAMTS9-AS2*) WHR_{adjBMI} loci in either sex-combined or sex-stratified analyses. By employing a recently developed method [23] to impute European GWAS summary statistics to the denser 1000 Genomes reference panel, followed by meta-analyses of both African and European ancestry individuals, we also identified three additional novel loci (*SPRYD7/DLEU2*, *CASC8* and *ZDHC1/HSD11B2*) for WHR_{adjBMI} . While all lead variants from established loci are common ($MAF \geq 5\%$), four of the nine lead variants from novel loci

were low frequency ($0.5\% \leq \text{MAF} < 5\%$). In addition, the lead variants from established loci including *TMEM18* and *ADAMTS9-AS2* were absent in HapMap. Overall, these results suggest the deeper genome coverage and/or improved imputation quality using 1000 Genomes, and complemented with additional sex-stratified analyses, facilitate the discovery of novel loci and identification of variants with stronger effects in established loci.

Among the novel sex-specific BMI loci (*IRX4/IRX2*, *INTS10/LPL* and *MLC1*), we did not identify any putative coding variants or regulatory regions underlying our association signals. Additionally, no associations have been reported with other metabolic traits in these novel BMI-associated signals. The first lead variant rs112778462 is located between the *IRX4* and *IRX2* genes which are members of the Iroquois homeobox gene family. *IRX2* expression has been associated with deposition of fat in the subcutaneous abdominal adipose tissue but no sex difference was observed [29–30]. *Irx4* knock out mice demonstrated cardiomyopathy with compensated increased *Irx2* expression [31]. The second lead variant rs149352150 is located between the *INTS10* and *LPL* genes. *LPL* encoded lipoprotein lipase is expressed in several tissues including adipose to mediate triglyceride hydrolysis and lipoprotein uptake. The serum LPL mass [32] and LPL activity and fat cell size of adipose tissues at gluteus and thigh [33] have been reported to be higher in women than in men. Previous GWAS demonstrated association of *LPL* with triglycerides and HDL cholesterol [34–35]. However, the reported lead variant rs12678919 was not in strong LD with rs149352150 ($r^2 = 0.005$ in AFR and 0.006 in EUR). The third lead variant rs56330886 is located in a gene-rich region on chromosome 22q13 including *MLC1*. No biological candidates are identified in this region, therefore further analyses may be needed to explain the causative mechanism for this association signal.

Among the novel $\text{WHR}_{\text{adjBMI}}$ loci, rs116718588 is located between *TCF7L2* and *HABP2*. *TCF7L2* is the most significant type 2 diabetes locus in African Americans [36] and other populations [37]. However, rs116718588 was not in LD ($r^2 < 0.01$ in AFR) with the reported type 2 diabetes associated variants. The second lead variant rs2472591 is located near *SPRYD7*, *DLEU2* and *TRIM13*. This locus was associated with height in previous GWAS [6], but rs2472591 was not associated with height in our study ($P > 0.05$), suggesting different variants in this locus regulate different measures of body size. In addition, a surrogate of rs2472591, rs790943, is a cis-eQTL for *TRIM13* [26] suggesting it may be the target gene. *TRIM13* encodes an E3 ubiquitin-protein ligase involved in endoplasmic reticulum-associated degradation. The third lead variant rs140858719 is located between *SSX2IP* and *LPAR3*. *LPAR3* is a plausible candidate as it encodes a receptor for lysophosphatidic acid (LPA). The autotaxin/LPA pathway mediates diverse biological actions including activation of preadipocyte proliferation [38], suppression of brown adipose differentiation [39], and promotion of systematic inflammation [40] which lead to increased risk for cardiometabolic diseases including obesity and insulin resistance [41–42]. LPA receptor 1 which is highly expressed in adipocytes and the gut primarily mediates these effects [43]. It has also been reported that LPA, via LPA1 and LPA3 receptors, mediated leukocytes recruitment and pro-inflammatory chemokine secretion during inflammation [44]. The fourth lead variant rs185693786 is located at intron 2 of *PDE3B*. The association signal spanned a large genomic region and harbors GWAS loci for adiponectin and height. Phosphodiesterase 3B is critical for mediating insulin/IGF-1 inhibition of cAMP signaling in adipocytes, liver, hypothalamus and pancreatic β cells [45]. *Pde3b*-knockout mice exhibited multiple alterations in regulation of lipolysis, lipogenesis, and insulin secretion, as well as signs of peripheral insulin resistance [46]. PDE3B expression has been reported to be higher in microvascular endothelial cell culture derived from skeletal muscles from male rats than in female rats [47]. The fifth lead variant rs6499129 is located intergenic between *ZDHHC1* and *HSD11B2*. *HSD11B2* encodes 11 β -hydroxysteroid dehydrogenase type 2 which converts the active glucocorticoids to inactive metabolites. HSD2 activity was elevated in

severe obesity and negatively associated with insulin sensitivity [48]. HSD2 expression is higher in omental than abdominal subcutaneous adipose tissue which may contribute to adipocyte hypertrophy and visceral obesity [49]. The sixth lead variant rs378854 is located at the long non-coding RNA *CASC8*. Associations of variants at *CASC8* have been reported for various cancers [50–52] but no association was reported for cardiometabolic traits.

In our SNP and locus transferability analyses, a moderate number of EA-derived BMI and WHR_{adjBMI} associated variants shared the same trait-raising alleles and displayed nominally significant associations in AA individuals, similar to previous findings [11–12]. While the BMI variants were similar in terms of their effect sizes and frequencies of trait-raising alleles between EA and AA populations, there were more discrepancies for WHR_{adjBMI} variants. In addition, a substantial proportion of lead regional variants in AA were not in strong LD with EA lead variants, suggesting AA populations either have different association signals or the results may be spurious. Taken together, only <30% of EA loci were associated with BMI and WHR_{adjBMI} in AA.

Trans-ethnic fine mapping improved resolution to refine putative causal variant(s) in some loci as compared to using EA studies alone. In the meta-analyses of AA and EA GWAS, four BMI loci (*SEC16B*, *STXBP6*, *FTO* and *MC4R*) and six WHR_{adjBMI} loci (*PPARG*, *ADAMTS9*, *SNX10*, *KLF13*, *CMIP* and *PEMT*) only contained one variant in the 99% credible sets. Among 16 BMI and 3 WHR_{adjBMI} loci that were examined in both the previous trans-ethnic meta-analysis studies using HapMap imputation [7–8] and the present study, the number of variants and the interval of credible sets were either the same or lower in the present study for 13 and 15 loci, respectively. The majority of credible variants are non-coding in those sets containing ≤ 20 variants. Several of them located at the *MTCH2*, *MAP2K5*, *NLRC3*, *ATP2A1*, *TBX15*, *WARS2* and *FAM13A* loci are cis-eQTL variants regulating nearby gene expression in subcutaneous and/or visceral adipose tissue, suggesting the putative causal variants may have a regulatory role instead of directly altering protein structure and function. Despite the low posterior probabilities, the coding changes of credible variants at *ADCY3* and *SH2B1* suggest that they may be the causal genes in the respective loci modulating BMI. Further studies are warranted to delineate putative causal variants including functional annotation in trans-ethnic fine mapping efforts [53].

Our large-scale GWAS meta-analyses in African ancestry individuals imputed to the 1000 Genomes reference panel, complemented by imputation of European GWAS using summary statistics and additional sex-stratified analyses, boosts the study power and improves resolution, leading to the identification of nine novel loci and fine mapping 37 loci with tractable credible sets. We observed significant associations for variants with $MAF \geq 0.5\%$, but rare variants were unlikely to be detected due to limited power and poor imputation quality. Large scale sequencing studies are needed to evaluate the contribution of rare variants in modulating complex traits such as BMI and WHR. Given the substantially larger sample size in European than in African ancestry samples, the trans-ethnic fine mapping results are largely driven by variants showing strong associations in Europeans. Future trans-ethnic studies including additional non-European populations will further improve the fine mapping effort.

Materials and methods

Study design

We used a three-stage design to evaluate genetic associations with BMI and WHR_{adjBMI} in sex-combined and sex-stratified samples (S1 Fig). Stage 1 included GWAS meta-analyses in AA individuals and stage 2 included replication of top associations from stage 1. Stage 3 included meta-analysis of top associations from stages 1 and 2 AA studies and EA meta-analysis results.

In the discovery stage 1 of AAAGC, 17 GWAS of up to 42,752 AA individuals (16,559 men and 26,193 women; 41,696 African Americans and 1,056 Africans) were included for the BMI analyses. A total of 10 GWAS of up to 20,384 AA individuals (4,783 men and 15,601 women; all African Americans) were included for the WHR_{adjBMI} analyses. For variants with $P < 1 \times 10^{-4}$ in either the sex-combined or the sex-stratified meta-analyses, stage 2 replication was performed in additional AA individuals from AAAGC ($N = 10,143$ for BMI, $N = 2,711$ for WHR_{adjBMI}), followed by meta-analysis with EA individuals from the GIANT consortium (322,154 for BMI, 210,086 for WHR_{adjBMI}). Variants that reached genome-wide significance ($P < 5 \times 10^{-8}$) were assessed for associations with BMI in two cohorts of children ($N = 7,222$). All AA participants in these studies provided written informed consent for the research, and approval for the study was obtained from the ethics review boards at all participating institutions. Detailed descriptions of each participating study and measurement and collection of height, weight, waist and hip circumferences are provided in [S1 Text](#), [S1](#) and [S2](#) Tables.

Genotyping, imputation and quality control

Genotyping in each study was performed with Illumina or Affymetrix genome-wide SNP arrays. Pre-phasing and imputation of missing genotypes in each study was performed using MaCH/ minimac [20] or SHAPEIT2/IMPUTEv2 [21–22] using the 1000 Genomes Project cosmopolitan reference panel (Phase I Integrated Release Version 3, March 2012) [18]. The details of the array, genotyping and imputation quality-control procedures and sample exclusions for each study are listed in [S3 Table](#). In general, samples reflecting duplicates, low call rates, gender mismatch, or population outliers were excluded. Variants were excluded by the following criteria: call rate < 0.95 , minor allele count (MAC) ≤ 6 , Hardy-Weinberg Equilibrium (HWE) $P < 1 \times 10^{-4}$, imputation quality score < 0.3 for minimac or < 0.4 for IMPUTE, or absolute allele frequency difference > 0.3 compared with expected allele frequency (calculated as 1000 Genomes frequency of $AFR \times 0.8 + EUR \times 0.2$).

Performance of 1000 Genomes imputation in African ancestry

We evaluated the performance of 1000 Genomes imputation using the largest study, the Women's Health Initiative (WHI) ($N = 8,054$). A total of 25.1 million variants with $MAF \geq 0.1\%$ were imputed to the 1000 Genomes reference panel. Of these, 98.1% (8.8 million) common variants, 95.4% (9.3 million) low frequency variants ($0.5\% \leq MAF < 5\%$), and 72.5% (4.6 million) rare variants ($0.1\% \leq MAF < 0.5\%$) were well imputed with IMPUTE info scores ≥ 0.3 ([S4 Table](#)). Notably, these frequencies are slightly lower than those obtained by imputation using 1000 Genomes phase 1 interim reference panel in Europeans [54]. However, 72.6%, 95.5% and 99.5% of the common, low frequency and rare variants, respectively, from the 1000 Genomes reference panel were not present in the HapMap and therefore demonstrate deeper coverage of the genome, particularly for the low frequency and rare variants.

Study-level association analyses

At all stages, genome-wide association analyses were performed by each of the participating studies. BMI was regressed on age, age squared, principal components and study site (if needed) to obtain residuals, separately by sex and case-control status, if needed. WHR was regressed on age, age squared, principal components, BMI and study site to obtain residuals, separately by sex and case-control status. Principal components were included to adjust for admixture proportion and population structure within each study. Residuals were inverse-normally transformed to obtain a standard normal distribution with mean of zero and standard deviation of one. For studies with unrelated subjects, each variant was tested assuming an

additive genetic model with each trait by regressing the transformed residuals on the number of copies of the variant effect allele. The analyses were stratified by sex and case-control status (if needed). For studies that included related individuals, family based association tests were conducted that took into consideration the genetic relationships among the individuals. Sex stratified, case-control stratified and combined analyses were performed. Association results with extreme values (absolute beta coefficient or standard error ≥ 10), primarily due to small sample sizes and/or low minor allele count, were excluded for meta-analysis.

Imputation of European GWAS summary statistics to 1000 Genomes

The latest summary statistics of sex-combined and sex-stratified meta-analyses of BMI and WHR_{adjBMI} imputed to the HapMap reference panel in EA from the Genetic Investigation of ANthropometric Traits (GIANT) consortium were obtained from http://www.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files [7–8]. These association summary statistics were used to impute z-scores of unobserved variants at the 1000 Genomes Project EUR reference panel (Phase I Integrated Release Version 3) using the ImpG program [23]. In brief, palindromic variants (AT/CG) and variants with allele mismatch with the reference were removed from the data. Using the ImpG-Summary method, the z-score of an unobserved variant was calculated as a linear combination of observed z-scores weighted by the variance-covariance matrix between variants induced by LD within a 1 Mb window from the reference haplotypes. The sample size of each unobserved variant was also interpolated from the sample sizes of observed variants using the same weighting method for z-score as $N_i = \sum_{t=1}^{t=T} \frac{|w_{i,t}|}{\sum |w_{i,t}|} N_t$. Here, $t = 1, 2, \dots, T$,

where T is the number of observed variants, $w_{i,t}$ is the element of the covariance matrix $\Sigma_{i,t}$ for the unobserved variant i and the observed variant t within window. The performance of imputation was assessed by r^2_{pred} , with similar characteristics as the standard imputation accuracy metric r^2_{hat} [20]. Results of variants with $r^2_{pred} \geq 0.6$ were used in subsequent analyses.

Meta-analysis

In the discovery stage 1, association results were combined across studies in sex-combined and sex-stratified samples using inverse-variance weighted fixed-effect meta-analysis implemented in the program METAL [55]. The study-specific λ values of association ranged from 0.97 to 1.05 for BMI, and 0.98 to 1.05 for WHR_{adjBMI} (S3 Table). Genomic control correction [56] was applied to each study before meta-analysis, and to the overall results after meta-analysis ($\lambda = 1.07$ for BMI, 1.01 for WHR_{adjBMI}). Variants with results generated from $< 50\%$ of the total sample size for each trait were excluded. After filtering, the numbers of variants reported in the meta-analyses were 17,972,087 for BMI, and 20,502,658 for WHR_{adjBMI} .

Variants with $P < 1 \times 10^{-4}$ in stage 1 sex-combined or sex-stratified meta-analyses were carried forward for replication in additional AA individuals (stage 2) and EA individuals (stage 3). For each of the replication AA studies, trait transformation and association were performed as in stage 1 and results were meta-analyzed using the inverse-variance method in METAL. For the replication study in EA, HapMap imputed summary statistics of each trait from the GIANT consortium were used to impute z-scores of unobserved variants at the 1000 Genomes.

In stages 1 and 2, meta-analysis results of AA studies were combined using the inverse-variance weighted method. In all stages including both AA and EA studies, meta-analysis results expressed as signed z-scores were combined using the fixed effect sample size weighted method in METAL due to the lack of beta and standard error estimates from the ImpG program [23]. Evidence of heterogeneity of allelic effects between males and females, within and across stages were assessed by the I^2 statistic in METAL. Genome-wide significance was

declared at $P < 5 \times 10^{-8}$ from each of the sex-combined and sex-stratified meta-analysis including AA and/or combined AA and EA individuals. Difference in effects between men and women was assessed using Cochran's Q test and nominal $P_{het} < 0.05$ declared as significant. A lead variant in a locus was defined as the most significant variant within a 1 Mb region. A novel locus was defined as a lead variant with distance > 500 kb from any established lead variants reported in previous studies. By convention, a locus was named by the closest gene(s) to the lead variant.

Conditional and joint analyses of summary statistics

For the genome-wide significant loci identified in sex-combined and sex-stratified analyses in AA (stages 1+2), we used the program GCTA [57–58] to select the top independent associated variants from summary statistics of the meta-analyses. This method uses the LD correlations between variants estimated from a reference sample to perform an approximate conditional association analysis. We used 8,054 unrelated individuals of African ancestry from the WHI cohort with ~ 15.7 M variants available as the reference sample for LD estimation. To select the top independent variants in the discovery and replication meta-analysis results, we first selected all variants that had $P < 5 \times 10^{-8}$ and conducted analysis conditioning on the selected variants to search for the top variants iteratively via a stepwise model to select the independent variants from this list. Then we proceeded to condition the rest of the variants that had $P > 5 \times 10^{-8}$ on the list of independent variants in the same fashion until no variant had conditional P that passed the significance level $P < 5 \times 10^{-8}$. Finally, all the selected variants were fitted jointly in the model for effect size estimation.

We also tested if the genome-wide significant variants identified from sex-combined GWAS in AA and the locus-wide significant variants identified from sex-combined and sex-specific locus transferability studies in AA were independent from nearby established loci identified from EA studies [7–8]. First, the published lead variants from EA studies were used to search for all surrogate variants that were in high LD ($r^2 > 0.8$ in 1000 Genomes Project EUR population). Second, these variants were pruned to select only variants in low LD in AA ($r^2 < 0.3$ in the 1000 Genomes Project AFR population) to avoid collinearity in conditional analysis. Third, association analysis was conducted on the AA significant variants conditioned on the selected EA lead and surrogate variants, using the program GCTA and estimated LD correlation from the WHI cohort. For genome-wide significant loci, an AA derived association signal is considered as independent from the established EA signals when the difference in $-\log P < 3$ and difference in effect size < 1 standard error after conditional analysis. For locus-wide significant loci, given the lower level of significance, independence is only considered as difference in effect size < 1 standard error after conditional analysis.

SNP and locus transferability analyses

We investigated the transferability of EA BMI and WHR associated variants and loci in AA individuals from stage 1 sex-combined and sex-stratified meta-analyses. First, we tested for replication of lead variants previously reported to be associated with BMI (176 variants from 170 loci) and $\text{WHR}_{\text{adjBMI}}$ (84 variants from 65 loci) at genome-wide significance in sex-combined and sex-stratified analyses from the GIANT consortium studies [7–9]. We defined SNP transferability as an EA lead variant sharing the same trait-raising allele at nominal $P < 0.05$ in AA individuals. To account for differences in local LD structure across populations, we also interrogated the flanking 0.1cM regions of the lead variants to search for the best variants with the smallest association P in AA individuals. Locus-wide significance was declared as $P_{locus} <$

0.05 by Bonferroni correction for the effective number of tests within a locus, estimated using the Li and Ji approach [59].

Fine mapping analyses

We compared the credible set intervals of established loci that showed locus-wide significance ($P_{locus} < 0.05$) in the sex-combined or sex-specific analyses from this study in summary statistics datasets including the 1000 Genomes imputed results from GIANT, AAAGC and meta-analysis of GIANT and AAAGC. In each dataset, a candidate region is defined as the flanking 0.1cM region of the lead variant reported by the GIANT consortium. Under the assumption of one causal variant in a region of M variants, the posterior probability of a variant j with association statistics Z driving the association, $P(C_j|Z)$, was calculated using the formula

$$P(C_j|Z) = \frac{\exp(\frac{1}{2}Z_j^2)}{\sum_{j=1}^M \exp(\frac{1}{2}Z_j^2)}. \text{ A 99\% credible set was constructed by ranking all variants by their}$$

posterior probability, followed by adding variants until the credible set has a cumulative posterior probability > 0.99 [53].

Bioinformatics

Functional annotation of novel variants. To determine whether any of our nine novel GWAS lead variants identified in the sex-combined and sex-specific analyses might be tagging potentially functional variants, we identified all variants within 1 Mb and in LD ($r^2 > 0.7$, 1000 Genomes AFR) with our lead variants. As such, we identified 137 variants and annotated each of them using ANNOVAR [60]. The predicted functional impact for coding variants were assessed via the Exome Variant Server (<http://evs.gs.washington.edu/EVS/>) for PhastCon, GERP [61], and PolyPhen [62], as well as SIFT [63].

We further characterized the variants that were in LD with the novel variants using the web-based tool RegulomeDB (<http://regulomedb.org/>) [24]. The variants that were likely to affect binding and linked to expression of a gene target (scores 1a-1f) based on “eQTL, transcription factor (TF) binding, matched TF motif, matched DNase footprint and DNase peak” or were only likely to affect binding (scores 2a-2c) based on “TF binding, matched TF motif, matched DNase footprint and DNase peak” were selected. For these variants, the sequence conservation (GERP and SiPhy [64]), the epigenomic data from the Roadmap Epigenomic project (ChromHMM states corresponding to enhancer or promoter elements, histone modification ChIP-seq peaks, and DNase hypersensitivity data peaks), the regulatory protein binding from the ENCODE project, the regulatory motifs based on commercial, literature and motif-finding analysis of the ENCODE project, and the eQTLs from Genotype-Tissue Expression (GTEx) project [65] were extract from web-based HaploReg v4 [25]. For variants within the tractable credible sets in the fine mapping analyses, similar analyses were also conducted.

Cross-trait associations. To assess whether the novel loci identified in the sex-combined and sex-specific analyses were associated with any related cardiometabolic and anthropometric traits, or may be in high LD with known eQTLs, we examined the NHGRI-EBI GWAS Catalog [27] and the GRASP (Genome-Wide Repository of Associations Between SNPs and Phenotypes) catalog [28] for reported variant-trait associations near our lead variants. We supplemented the catalogs with additional genome-wide significant associations of interest from the literature [7–9,66]. We used PLINK to identify variants within 1 Mb of lead variants. All variants within the specified regions with $r^2 > 0.7$ (1000 Genomes AFR) were retained from the catalogs for further evaluation.

Power analysis

Given our sample sizes in the discovery and replication stages in our African ancestry populations, we have >80% power to detect variants explaining 0.08% variance for BMI that corresponds to effect sizes of 0.09 and 0.20 SD units for MAF of 0.05 and 0.01, respectively. For WHR_{adjBMI} , we have >80% power to detect variants explaining 0.18% variance that corresponds to effect sizes of 0.14 and 0.30 SD units for MAF of 0.05 and 0.01, respectively.

Supporting information

S1 Fig. Study design of GWAS meta-analyses and replications for BMI and WHR_{adjBMI} .
(PDF)

S2 Fig. Quantile-quantile plot of 1000 genomes phase 1 imputed discovery results and their associations to adult BMI in men and women of African ancestry using all variants and only variants outside of known GWAS loci.
(PDF)

S3 Fig. Quantile-quantile plot of 1000 genomes phase 1 imputed discovery results and their associations to adult BMI in women only and men only of African ancestry using all variants and only variants outside of known GWAS loci.
(PDF)

S4 Fig. Quantile-quantile plot of 1000 genomes phase 1 imputed discovery results and their associations to adult waist-to-hip ratio adjusted for BMI (WHR_{adjBMI}) in men and women of African ancestry using all variants and only variants outside of known GWAS loci.
(PDF)

S5 Fig. Quantile-quantile plot of 1000 genomes phase 1 imputed discovery results and their associations to adult waist-to-hip ratio adjusted for BMI (WHR_{adjBMI}) in women only and men only of African ancestry using all variants and only variants outside of known GWAS loci.
(PDF)

S6 Fig. Manhattan plot of 1000 genomes phase 1 imputed discovery results and their associations to adult BMI in men and women of African ancestry.
(PDF)

S7 Fig. Manhattan plot of 1000 genomes phase 1 imputed discovery results and their associations to waist-to-hip ratio adjusted for BMI (WHR_{adjBMI}) in men and women of African ancestry.
(PDF)

S8 Fig. Miami plot of 1000 genomes phase 1 imputed discovery results and their associations to adult BMI in women only (top) and men only (bottom) of African ancestry.
(PDF)

S9 Fig. Miami plot of 1000 genomes phase 1 imputed discovery results and their associations to adult waist-to-hip ratio adjusted for BMI (WHR_{adjBMI}) in women only (top) and men only (bottom) of African ancestry.
(PDF)

S10 Fig. Locuszoom plots using discovery results for established loci that reached genome-wide significance: (A) *SEC16B*, *TMEM18*, *GNPDA2*, *GALNT10*, *KLHL32*, *FTO* and *MC4R* for BMI in men and women combined; and (B) *ADAMTS9-AS2* for waist-to-hip ratio adjusted

for BMI (WHR_{adjBMI}) in men and women combined. All plots use AFR LD from the 1000 Genomes phase 1 reference panel. In each plot, the most significant variant within a 1Mb regional locus is highlighted. *P*-values for all variants including the most significant variant are based on the African ancestry discovery phase only (AA-Discovery). In addition, for the most significant variant, *P*-values are annotated and illustrated from the African ancestry discovery and replication phases (AA-Discovery+Replication).

(PDF)

S11 Fig. Correlation of effect sizes for (A) BMI and (B) WHR_{adjBMI} , and effect allele frequencies for (C) BMI and (D) WHR_{adjBMI} in European and African ancestry studies in SNP transferability analyses.

(PDF)

S1 Table. Study design and sample quality control of discovery and replication studies.

(XLSX)

S2 Table. Study-specific descriptive statistics of discovery and replication studies.

(XLSX)

S3 Table. Genotyping methods, quality control of variants, imputation, and statistical analysis in discovery and replication studies.

(XLSX)

S4 Table. Comparison of coverage of variants using the 1000 Genomes and HapMap reference panels for imputation in the Women's Health Initiative study.

(XLSX)

S5 Table. Comparison of lead variants between 1000 Genomes and HapMap imputed meta-analysis in AA in previously identified BMI loci in discovery and replication studies.

(XLSX)

S6 Table. Associations of lead variants from novel and previously identified BMI and WHR_{adjBMI} loci in combined and sex-stratified analyses of African ancestry discovery and replication samples.

(XLSX)

S7 Table. Association of African Ancestry sex-combined genome-wide significant variants in children of African ancestry.

(XLSX)

S8 Table. Putative coding or regulatory variants in linkage disequilibrium ($r^2 > 0.7$) with WHR_{adjBMI} loci.

(XLSX)

S9 Table. Previously-reported associations of novel BMI and WHR_{adjBMI} loci with other traits in the GRASP Catalog. This table lists all previously-reported associations within 1 Mb (+/- 500 kb) and in high LD ($r^2 > 0.7$) with our lead novel SNPs along with relevant annotation (e.g. miRNA target binding site, variant location relevant to nearest gene, gene function prediction) reported in the GRASP Catalog.

(XLSX)

S10 Table. Conditional analysis of African ancestry primary and secondary lead SNPs with European lead SNPs in previously identified BMI and WHR_{adjBMI} loci.

(XLSX)

S11 Table. SNP transferability of BMI and WHR_{adjBMI} lead SNPs from European sex combined and sex stratified GWAS in African ancestry individuals.

(XLSX)

S12 Table. Locus transferability and conditional analyses of European BMI and WHR_{adjBMI} loci in African ancestry individuals.

(XLSX)

S13 Table. Fine mapping of novel loci and previously identified loci with locus-wide significance in African ancestry individuals using 1000 Genomes imputed results from African, European and combined ancestries.

(XLSX)

S14 Table. Functional characterization of variants in tractable credible sets in meta-analysis of African and European ancestry GWAS.

(XLSX)

S1 Text. Supplementary note.

(DOCX)

S2 Text. Members of the BMDCS Group.

(DOCX)

Acknowledgments

We gratefully acknowledge the contribution of all participants from the various studies that contributed to this work.

Author Contributions

Conceptualization: CAH KEN MCYN RJFL.

Formal analysis: AC AEJ AML AS BEC BOT BP CTL DH DV DZ EBW GC GL HO JAB JAS JPB JDF JL JY KR KY LAL LD LRY MCYN MFF MGr MKW MAN MRI PM QD RR SMT SV TMB WMC WZha XG YHHH YLi YLu YS.

Project administration: AA ABZ AOI BAR BIF BM BMP BN BOT BSZ CAH CBA CDH CNR DCR DH DKA DMB DRW DSS DVC DWB EK EMJ EPB EVB HH IBB JC JGW JH JIR JLS JNH JSW KEN KLW LAC LB MCYN MF MKE MMS MFP MS OIO PJG RGZ RJFL RK RSC SAI SFAG SIB SJC SLRK SRP SSS TBH VLS WJB WZhe XZ.

Resources: AGF AOg BIF BMP BN BS BSZ DRW HH JAS JDF JIR KLN MAA MC MGa OO SA SFAG SLRK TOO UN WZha WZhe YDIC.

Supervision: CAH KEN MCYN RJFL.

Writing – original draft: AEJ CTL LAC LRY MCYN MFF MGr MKW YLu.

Writing – review & editing: AEJ BEC BMP CAH CTL DCR DKA DMB JPB KEN KR KY LAC LRY MC MCYN MFF MGr MKW MAN MRI RJFL WMC XG YLu.

References

1. Flegal KM, Kruszon-Moran D, Carroll MD, Fryar CD, Ogden CL (2016) Trends in obesity among adults in the United States, 2005 to 2014. *JAMA* 315: 2284–2291. <https://doi.org/10.1001/jama.2016.6458> PMID: 27272580

2. Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, et al. (2010) Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. *Nat Genet* 42: 937–948. <https://doi.org/10.1038/ng.686> PMID: 20935630
3. Heid IM, Jackson AU, Randall JC, Winkler TW, Qi L, et al. (2010) Meta-analysis identifies 13 new loci associated with waist-hip ratio and reveals sexual dimorphism in the genetic basis of fat distribution. *Nat Genet* 42: 949–960. <https://doi.org/10.1038/ng.685> PMID: 20935629
4. Scherag A, Dina C, Hinney A, Vatin V, Scherag S, et al. (2010) Two new loci for body-weight regulation identified in a joint analysis of genome-wide association studies for early-onset extreme obesity in French and German study groups. *PLoS Genet* 6: e1000916. <https://doi.org/10.1371/journal.pgen.1000916> PMID: 20421936
5. Bradfield JP, Taal HR, Timpson NJ, Scherag A, Lecocoeur C, et al. (2012) A genome-wide association meta-analysis identifies new childhood obesity loci. *Nat Genet* 44: 526–531. <https://doi.org/10.1038/ng.2247> PMID: 22484627
6. Berndt SI, Gustafsson S, Magi R, Ganna A, Wheeler E, et al. (2013) Genome-wide meta-analysis identifies 11 new loci for anthropometric traits and provides insights into genetic architecture. *Nat Genet* 45: 501–512. <https://doi.org/10.1038/ng.2606> PMID: 23563607
7. Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, et al. (2015) Genetic studies of body mass index yield new insights for obesity biology. *Nature* 518: 197–206. <https://doi.org/10.1038/nature14177> PMID: 25673413
8. Shungin D, Winkler TW, Croteau-Chonka DC, Ferreira T, Locke AE, et al. (2015) New genetic loci link adipose and insulin biology to body fat distribution. *Nature* 518: 187–196. <https://doi.org/10.1038/nature14132> PMID: 25673412
9. Winkler TW, Justice AE, Graff M, Barata L, Feitosa MF, et al. (2015) The influence of age and sex on genetic associations with adult body size and shape: a large-scale genome-wide interaction study. *PLoS Genet* 11: e1005378. <https://doi.org/10.1371/journal.pgen.1005378> PMID: 26426971
10. Ng MC, Hester JM, Wing MR, Li J, Xu J, et al. (2012) Genome-wide association of BMI in African Americans. *Obesity (Silver Spring)* 20: 622–627.
11. Monda KL, Chen GK, Taylor KC, Palmer C, Edwards TL, et al. (2013) A meta-analysis identifies new loci associated with body mass index in individuals of African ancestry. *Nat Genet* 45: 690–696. <https://doi.org/10.1038/ng.2608> PMID: 23583978
12. Liu CT, Monda KL, Taylor KC, Lange L, Demerath EW, et al. (2013) Genome-wide association of body fat distribution in African ancestry populations suggests new loci. *PLoS Genet* 9: e1003681. <https://doi.org/10.1371/journal.pgen.1003681> PMID: 23966867
13. Okada Y, Kubo M, Ohmiya H, Takahashi A, Kumasaka N, et al. (2012) Common variants at CDKAL1 and KLF9 are associated with body mass index in East Asian populations. *Nat Genet* 44: 302–306. <https://doi.org/10.1038/ng.1086> PMID: 22344221
14. Wen W, Cho YS, Zheng W, Dorajoo R, Kato N, et al. (2012) Meta-analysis identifies common variants associated with body mass index in East Asians. *Nat Genet* 44: 307–311. <https://doi.org/10.1038/ng.1087> PMID: 22344219
15. Wen W, Zheng W, Okada Y, Takeuchi F, Tabara Y, et al. (2014) Meta-analysis of genome-wide association studies in East Asian-ancestry populations identifies four new loci for body mass index. *Hum Mol Genet* 23: 5492–5504. <https://doi.org/10.1093/hmg/ddu248> PMID: 24861553
16. Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, et al. (2007) A second generation human haplotype map of over 3.1 million SNPs. *Nature* 449: 851–861. <https://doi.org/10.1038/nature06258> PMID: 17943122
17. Altshuler DM, Gibbs RA, Peltonen L, Dermitzakis E, Schaffner SF, et al. (2010) Integrating common and rare genetic variation in diverse human populations. *Nature* 467: 52–58. <https://doi.org/10.1038/nature09298> PMID: 20811451
18. Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM, et al. (2012) An integrated map of genetic variation from 1,092 human genomes. *Nature* 491: 56–65. <https://doi.org/10.1038/nature11632> PMID: 23128226
19. Delaneau O, Marchini J (2014) Integrating sequence and array data to create an improved 1000 Genomes Project haplotype reference panel. *Nat Commun* 5: 3934. <https://doi.org/10.1038/ncomms4934> PMID: 25653097
20. Li Y, Willer CJ, Ding J, Scheet P, Abecasis GR (2010) MaCH: using sequence and genotype data to estimate haplotypes and unobserved genotypes. *Genet Epidemiol* 34: 816–834. <https://doi.org/10.1002/gepi.20533> PMID: 21058334
21. Delaneau O, Marchini J, Zagury JF (2011) A linear complexity phasing method for thousands of genomes. *Nat Methods* 9: 179–181. <https://doi.org/10.1038/nmeth.1785> PMID: 22138821

22. Howie BN, Donnelly P, Marchini J (2009) A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet* 5: e1000529. <https://doi.org/10.1371/journal.pgen.1000529> PMID: 19543373
23. Pasaniuc B, Zaitlen N, Shi H, Bhatia G, Gusev A, et al. (2014) Fast and accurate imputation of summary statistics enhances evidence of functional enrichment. *Bioinformatics* 30: 2906–2914. <https://doi.org/10.1093/bioinformatics/btu416> PMID: 24990607
24. Boyle AP, Hong EL, Hariharan M, Cheng Y, Schaub MA, et al. (2012) Annotation of functional variation in personal genomes using RegulomeDB. *Genome Res* 22: 1790–1797. <https://doi.org/10.1101/gr.137323.112> PMID: 22955989
25. Ward LD, Kellis M (2016) HaploReg v4: systematic mining of putative causal variants, cell types, regulators and target genes for human complex traits and disease. *Nucleic Acids Res* 44: D877–881. <https://doi.org/10.1093/nar/gkv1340> PMID: 26657631
26. Barreiro LB, Tailleux L, Pai AA, Gicquel B, Marioni JC, et al. (2012) Deciphering the genetic architecture of variation in the immune response to *Mycobacterium tuberculosis* infection. *Proc Natl Acad Sci U S A* 109: 1204–1209. <https://doi.org/10.1073/pnas.1115761109> PMID: 22233810
27. Welter D, MacArthur J, Morales J, Burdett T, Hall P, et al. (2014) The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. *Nucleic Acids Res* 42: D1001–1006. <https://doi.org/10.1093/nar/gkt1229> PMID: 24316577
28. Leslie R, O'Donnell CJ, Johnson AD (2014) GRASP: analysis of genotype-phenotype results from 1390 genome-wide association studies and corresponding open access database. *Bioinformatics* 30: i185–194. <https://doi.org/10.1093/bioinformatics/btu273> PMID: 24931982
29. Karastergiou K, Fried SK, Xie H, Lee MJ, Divoux A, et al. (2013) Distinct developmental signatures of human abdominal and gluteal subcutaneous adipose tissue depots. *J Clin Endocrinol Metab* 98: 362–371. <https://doi.org/10.1210/jc.2012-2953> PMID: 23150689
30. Pinnick KE, Nicholson G, Manolopoulos KN, McQuaid SE, Valet P, et al. (2014) Distinct developmental profile of lower-body adipose tissue defines resistance against obesity-associated metabolic complications. *Diabetes* 63: 3785–3797. <https://doi.org/10.2337/db14-0385> PMID: 24947352
31. Bruneau BG, Bao ZZ, Fatkin D, Xavier-Neto J, Georgakopoulos D, et al. (2001) Cardiomyopathy in *Irf4*-deficient mice is preceded by abnormal ventricular gene expression. *Mol Cell Biol* 21: 1730–1736. <https://doi.org/10.1128/MCB.21.5.1730-1736.2001> PMID: 11238910
32. Onat A, Hergenc G, Agirbasli M, Kaya Z, Can G, et al. (2009) Preheparin serum lipoprotein lipase mass interacts with gender, gene polymorphism and, positively, with smoking. *Clin Chem Lab Med* 47: 208–215. <https://doi.org/10.1515/CCLM.2009.018> PMID: 19191728
33. Votruba SB, Jensen MD (2007) Sex differences in abdominal, gluteal, and thigh LPL activity. *Am J Physiol Endocrinol Metab* 292: E1823–1828. <https://doi.org/10.1152/ajpendo.00601.2006> PMID: 17311894
34. Teslovich TM, Musunuru K, Smith AV, Edmondson AC, Stylianou IM, et al. (2010) Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* 466: 707–713. <https://doi.org/10.1038/nature09270> PMID: 20686565
35. Willer CJ, Schmidt EM, Sengupta S, Peloso GM, Gustafsson S, et al. (2013) Discovery and refinement of loci associated with lipid levels. *Nat Genet* 45: 1274–1283. <https://doi.org/10.1038/ng.2797> PMID: 24097068
36. Ng MC, Shriner D, Chen BH, Li J, Chen WM, et al. (2014) Meta-analysis of genome-wide association studies in African Americans provides insights into the genetic architecture of type 2 diabetes. *PLoS Genet* 10: e1004517. <https://doi.org/10.1371/journal.pgen.1004517> PMID: 25102180
37. Mahajan A, Go MJ, Zhang W, Below JE, Gaulton KJ, et al. (2014) Genome-wide trans-ancestry meta-analysis provides insight into the genetic architecture of type 2 diabetes susceptibility. *Nat Genet* 46: 234–244. <https://doi.org/10.1038/ng.2897> PMID: 24509480
38. Ferry G, Tellier E, Try A, Gres S, Naime I, et al. (2003) Autotaxin is released from adipocytes, catalyzes lysophosphatidic acid synthesis, and activates preadipocyte proliferation. Up-regulated expression with adipocyte differentiation and obesity. *J Biol Chem* 278: 18162–18169. <https://doi.org/10.1074/jbc.M301158200> PMID: 12642576
39. Federico L, Ren H, Mueller PA, Wu T, Liu S, et al. (2012) Autotaxin and its product lysophosphatidic acid suppress brown adipose differentiation and promote diet-induced obesity in mice. *Mol Endocrinol* 26: 786–797. <https://doi.org/10.1210/me.2011-1229> PMID: 22474126
40. Hui DY (2016) Intestinal phospholipid and lysophospholipid metabolism in cardiometabolic disease. *Curr Opin Lipidol* 27: 507–512. <https://doi.org/10.1097/MOL.0000000000000334> PMID: 27438680

41. Rancoule C, Dusaulcy R, Treguer K, Gres S, Attane C, et al. (2014) Involvement of autotaxin/lysophosphatidic acid signaling in obesity and impaired glucose homeostasis. *Biochimie* 96: 140–143. <https://doi.org/10.1016/j.biochi.2013.04.010> PMID: 23639740
42. Reeves VL, Trybula JS, Wills RC, Goodpaster BH, Dube JJ, et al. (2015) Serum Autotaxin/ENPP2 correlates with insulin resistance in older humans with obesity. *Obesity (Silver Spring)* 23: 2371–2376.
43. Yung YC, Stoddard NC, Chun J (2014) LPA receptor signaling: pharmacology, physiology, and pathophysiology. *J Lipid Res* 55: 1192–1214. <https://doi.org/10.1194/jlr.R046458> PMID: 24643338
44. Zhao C, Sardella A, Chun J, Poubelle PE, Fernandes MJ, et al. (2011) TNF-alpha promotes LPA1- and LPA3-mediated recruitment of leukocytes in vivo through CXCR2 ligand chemokines. *J Lipid Res* 52: 1307–1318. <https://doi.org/10.1194/jlr.M008045> PMID: 21521824
45. Degerman E, Ahmad F, Chung YW, Guirguis E, Omar B, et al. (2011) From PDE3B to the regulation of energy homeostasis. *Curr Opin Pharmacol* 11: 676–682. <https://doi.org/10.1016/j.coph.2011.09.015> PMID: 22001403
46. Choi YH, Park S, Hockman S, Zmuda-Trzebiatowska E, Svennelid F, et al. (2006) Alterations in regulation of energy homeostasis in cyclic nucleotide phosphodiesterase 3B-null mice. *J Clin Invest* 116: 3240–3251. <https://doi.org/10.1172/JCI24867> PMID: 17143332
47. Wang J, Bingaman S, Huxley VH (2010) Intrinsic sex-specific differences in microvascular endothelial cell phosphodiesterases. *Am J Physiol Heart Circ Physiol* 298: H1146–1154. <https://doi.org/10.1152/ajpheart.00252.2009> PMID: 20139324
48. Mussig K, Remer T, Haupt A, Gallwitz B, Fritsche A, et al. (2008) 11beta-hydroxysteroid dehydrogenase 2 activity is elevated in severe obesity and negatively associated with insulin sensitivity. *Obesity (Silver Spring)* 16: 1256–1260.
49. Lee MJ, Fried SK, Mundt SS, Wang Y, Sullivan S, et al. (2008) Depot-specific regulation of the conversion of cortisone to cortisol in human adipose tissue. *Obesity (Silver Spring)* 16: 1178–1185.
50. Gudmundsson J, Sulem P, Gudbjartsson DF, Blondal T, Gylfason A, et al. (2009) Genome-wide association and replication studies identify four variants associated with prostate cancer susceptibility. *Nat Genet* 41: 1122–1126. <https://doi.org/10.1038/ng.448> PMID: 19767754
51. Yao K, Hua L, Wei L, Meng J, Hu J (2015) Correlation Between CASC8, SMAD7 Polymorphisms and the Susceptibility to Colorectal Cancer: An updated meta-analysis based on GWAS results. *Medicine (Baltimore)* 94: e1884.
52. Ma G, Gu D, Lv C, Chu H, Xu Z, et al. (2015) Genetic variant in 8q24 is associated with prognosis for gastric cancer in a Chinese population. *J Gastroenterol Hepatol* 30: 689–695. <https://doi.org/10.1111/jgh.12801> PMID: 25302443
53. Kichaev G, Pasaniuc B (2015) Leveraging functional-annotation data in trans-ethnic fine-mapping studies. *Am J Hum Genet* 97: 260–271. <https://doi.org/10.1016/j.ajhg.2015.06.007> PMID: 26189819
54. Horikoshi M, Mgi R, van de Bunt M, Surakka I, Sarin AP, et al. (2015) Discovery and fine-mapping of glycaemic and obesity-related trait loci using high-density imputation. *PLoS Genet* 11: e1005230. <https://doi.org/10.1371/journal.pgen.1005230> PMID: 26132169
55. Willer CJ, Li Y, Abecasis GR (2010) METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* 26: 2190–2191. <https://doi.org/10.1093/bioinformatics/btq340> PMID: 20616382
56. Devlin B, Roeder K, Wasserman L (2001) Genomic control, a new approach to genetic-based association studies. *Theor Popul Biol* 60: 155–166. <https://doi.org/10.1006/tpbi.2001.1542> PMID: 11855950
57. Yang J, Lee SH, Goddard ME, Visscher PM (2011) GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet* 88: 76–82. <https://doi.org/10.1016/j.ajhg.2010.11.011> PMID: 21167468
58. Yang J, Ferreira T, Morris AP, Medland SE, Madden PA, et al. (2012) Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nat Genet* 44: 369–375, S361–363. <https://doi.org/10.1038/ng.2213> PMID: 22426310
59. Li J, Ji L (2005) Adjusting multiple testing in multilocus analyses using the eigenvalues of a correlation matrix. *Heredity (Edinb)* 95: 221–227.
60. Wang K, Li M, Hakonarson H (2010) ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res* 38: e164. <https://doi.org/10.1093/nar/gkq603> PMID: 20601685
61. Cooper GM, Stone EA, Asimenos G, Green ED, Batzoglou S, et al. (2005) Distribution and intensity of constraint in mammalian genomic sequence. *Genome Res* 15: 901–913. <https://doi.org/10.1101/gr.3577405> PMID: 15965027
62. Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, et al. (2010) A method and server for predicting damaging missense mutations. *Nat Methods* 7: 248–249. <https://doi.org/10.1038/nmeth0410-248> PMID: 20354512

63. Sim NL, Kumar P, Hu J, Henikoff S, Schneider G, et al. (2012) SIFT web server: predicting effects of amino acid substitutions on proteins. *Nucleic Acids Res* 40: W452–457. <https://doi.org/10.1093/nar/gks539> PMID: 22689647
64. Garber M, Guttman M, Clamp M, Zody MC, Friedman N, et al. (2009) Identifying novel constrained elements by exploiting biased substitution patterns. *Bioinformatics* 25: i54–62. <https://doi.org/10.1093/bioinformatics/btp190> PMID: 19478016
65. Consortium GTEx (2015) Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* 348: 648–660. <https://doi.org/10.1126/science.1262110> PMID: 25954001
66. Wood AR, Esko T, Yang J, Vedantam S, Pers TH, et al. (2014) Defining the role of common variation in the genomic and biological architecture of adult human height. *Nat Genet* 46: 1173–1186. <https://doi.org/10.1038/ng.3097> PMID: 25282103