



University of Kentucky
UKnowledge

Psychology Faculty Publications

Psychology

6-1-2001

Adolescent and Young Adult Substance Use: Association with Sensation Seeking, Self Esteem and Retrospective Report of Early Pubertal Onset. A Preliminary Examination

Catherine A. Martin

University of Kentucky, catherine.martin@uky.edu

T. K. Logan

University of Kentucky, tklogan@email.uky.edu

Carl Leukefeld

University of Kentucky, cleukef@uky.edu

Rich Milich

University of Kentucky, richard.milich@uky.edu

Hatim Omar

University of Kentucky, hatim.omar@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/psychology_facpub



Part of the [Child Psychology Commons](#), [Clinical and Medical Social Work Commons](#), and the [Substance Abuse and Addiction Commons](#).
Right click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Martin, Catherine A.; Logan, T. K.; Leukefeld, Carl; Milich, Rich; Omar, Hatim; and Clayton, Richard, "Adolescent and Young Adult Substance Use: Association with Sensation Seeking, Self Esteem and Retrospective Report of Early Pubertal Onset. A Preliminary Examination" (2001). *Psychology Faculty Publications*. 60.
https://uknowledge.uky.edu/psychology_facpub/60

This Article is brought to you for free and open access by the Psychology at UKnowledge. It has been accepted for inclusion in Psychology Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Authors

Catherine A. Martin, T. K. Logan, Carl Leukefeld, Rich Milich, Hatim Omar, and Richard Clayton

Adolescent and Young Adult Substance Use: Association with Sensation Seeking, Self Esteem and Retrospective Report of Early Pubertal Onset. A Preliminary Examination**Notes/Citation Information**

Published in *International Journal of Adolescent Medicine and Health*, v. 13, issue 3, p. 211-219.

© Freund Publishing House Ltd.

The copyright holder has granted permission for posting the article here.

Digital Object Identifier (DOI)

<https://doi.org/10.1515/IJAMH.2001.13.3.211>

Adolescent and young adult substance use: Association with sensation seeking, self esteem and retrospective report of early pubertal onset. A preliminary examination

Catherine Martin, MD, TK Logan, PhD, Carl Leukefeld, DSW, Rich Milich, PhD, Hatim Omar, MD and Richard Clayton, PhD

Department of Psychiatry, Center on Drug and Alcohol Research and Department of Pediatrics, University of Kentucky College of Medicine, Department of Psychology, and Center for Prevention Research, University of Kentucky, Lexington, KY, USA

Abstract: Structured questionnaires were administered to investigate the relationship between early pubertal onset, substance use, sensation seeking, and self-esteem. The current study presents data from 1,002 subjects, who were followed from the 6th to the 10th grades and again at the age of 20. In females, early pubertal onset was associated with greater cigarette use and lower self-esteem. Further the interaction of early pubertal onset and low self-esteem was predictive for cigarette use in females, as was early pubertal onset and high sensation seeking. Late pubertal onset was associated with decreased alcohol use and lower sensation seeking in males, with the opposite trends for early pubertal onset. These interrelationships must be considered when attempting to understand and identify early adolescent initiation of substance use.

Keywords: Pubertal onset, substance use, sensation seeking, self-esteem, USA

Correspondence: Hatim A. Omar, M.D., Director, Adolescent Medicine, University of Kentucky, Department of Pediatrics, Kentucky Clinic, Lexington, KY 40536-0284, USA. Tel office (859) 323-6426 ext. 307; Fax (859) 257-7706; e-mail: haomar2@pop.uky.edu

Submitted: December 25, 2000. **Revised:** January 30, 2001; **Accepted:** February 3, 2001.

INTRODUCTION

The relationship between substance use and pubertal maturation is complicated, in part because of the association between pubertal maturation and age, hormonal change (1), and peer associations (2-4). All these factors can affect substance use. Despite these complexities, there is evidence for the association of early pubertal onset with substance use. For example, early maturing girls use marijuana more than girls who mature later, and this difference becomes more pronounced over time (5). Similarly, early maturing girls are more likely to report frequent intoxication (5). Adult females who report maturing early also report earlier substance use, while late-maturing adult females are more likely to report never having started smoking (5).

Early and late-maturing 14-year-old boys are more likely to report having been drunk than "on-time" boys (6).

There are several other important variables that have been associated with both pubertal maturation and substance use. Lower self-esteem and sensation seeking have been associated with earlier pubertal maturation in girls (7) and later maturation in boys (8), as well as substance use (9-12). Although the relationship of sensation seeking with early pubertal onset has not been systematically examined, an association between sexual hormones (Hypothalamic Pituitary Gonadal Axis hormones) and sensation seeking has been reported (13).

The purpose of this study was to examine the relationship between early

pubertal onset and substance use. The following questions were examined: 1) Was early pubertal onset related to increased substance use? 2) Was the association of pubertal onset with substance use related to differences in sensation seeking, and self-esteem? and 3) Would the interaction of pubertal onset, sensation seeking, and self-esteem, account for more variance in substance use than early pubertal onset or sensation seeking or self-esteem alone?

METHODS

Participants

Participants were part of a longitudinal study designed to examine drug use over time. The initial sample was composed of 2,071 6th graders who participated in a citywide school system drug prevention program. Students completed questionnaires, which measured drug use and other variables in the 6th – 10th grades. Completed questionnaires were obtained from original participants on at least three occasions, once in the 6th grade, once in the 7th or 8th grade, and once in the 9th or 10th grade. These subjects were surveyed again at the age of 20 (n=1,429). Findings reported here are based on the data collected from the final sample of 1,002. The sample consisted of 571 females (57%) and 431 males (43%). In addition, 748 (75.1%) were White, 204 (20.4%) were African American, and 44 (4.5%) were of other race or ethnicity.

An attrition analysis was conducted to test the possibility that the 1,002 respondents were different from the 427 who were mailed a survey, but did not respond. A dummy variable representing present-missing status was simultaneously regressed using a paired correlation matrix onto 15 variables from the original assessment: sex; ethnicity; age; peer-pressure resistance; self-esteem; DARE status; and use of, and positive and negative

expectancies toward, cigarettes, alcohol, and marijuana. Missing status accounted for a small but significant proportion of variance in the linear combination of the 15 study variables ($R^2 = .06$), $F(15, 1339) = 6.08$, $p < .001$. Only 3 variables were independently linked to missing status. Specifically, participants who were missing completed surveys tended to be older males, who reported using cigarettes in the 6th grade (14).

Questionnaires:

Figure 1 presents the sample questions used for the current study from each of the following areas.

Onset of Puberty: Subjects were asked to endorse retrospectively one gender-specific pubertal marker at 6-month intervals between ages 10 and 17½. These questions were asked at age 20 only. Correlations of self-reported recalled age of menarche and actual age of menarche are quite high (.6 to .9) (15,16). Self-reports of menarche to measure pubertal development are commonly used (17). Similarly, self-reports of male sexual maturation demonstrate test-retest reliability and the magnitude of the correlation of self reported onset of sexual maturation increases with the degree of genetic similarity between subjects (18).

Drug Use: Self-reported drug use items were asked on 5 occasions. For this study, past month use of cigarettes and alcohol was included.

Self-Esteem: The Rosenberg Self-Esteem Scale (19) is a 10 item scale designed to measure an individuals' perceived self worth. Responses are recorded on a 5 point Likert scale. Test-retest reliability has been shown to be as high as .85 for college students and internal consistency and validity for this scale have been reported previously (19). The Alpha coefficient for this study was .87.

Figure 1. *Questions*Onset of Puberty*If you are a male, when was your first nocturnal ejaculation (e.g., wet dream)?*

10 years old	12	14	16	11	13	15	17
10 ½	12 ½	14 ½	16 ½	11 ½	13 ½	15 ½	17 ½

If you are a female, when was your first menstrual period?

10 years old	12	14	16	11	13	15	17
10 ½	12 ½	14 ½	16 ½	11 ½	13 ½	15 ½	17 ½

Drug Use

Substance Use Asked at Age 21.

How frequently have you smoked cigarettes during the past 30 days?

- | | |
|------------------------------------|---|
| 1. Not at all | 5. About one pack per day |
| 2. Less than one cigarette per day | 6. About one and one-half packs per day |
| 3. One to five cigarettes per day | 7. Two packs or more per day |
| 4. About one-half pack per day | |

How many times have you had a drink of alcohol (a drink is a glass of wine, a wine cooler, a bottle of beer, a shot glass of liquor or a mixed drink) in the past month?

- | | | | |
|---------|----------|--------|---------------|
| A. None | E. 10-19 | C. 3-5 | G. 40 or more |
| B. 1-2 | F. 20-39 | D. 6-9 | |

Substance Use Asked at 7th through 10th Grades.*How many cigarettes have you smoked in THE PAST MONTH?*

- | | |
|----------------------------|--|
| 1. None | 5. 11-20 cigarettes (1/2 pack to 1 pack) |
| 2. Less than one cigarette | 6. 21-30 cigarettes (1 pack to 1½ pack) |
| 3. 1-5 cigarettes | 7. 31 or more cigarettes (over 1½ pack) |
| 4. 6-10 cigarettes | |

IN THE PAST MONTH, how many times have you drunk a FULL GLASS of alcohol (beer, wine, liquor)?

- | | | | |
|------------|--------|----------|---------------|
| 1. 0 times | 3. 3-5 | 5. 10-19 | 7. 40 or more |
| 2. 1-2 | 4. 6-9 | 6. 20-39 | |

Sensation Seeking*Likert**Someday I would like to try sky-diving or parachute jumping. (Zuckerman, 1979)*

I agree strongly	I agree but not strongly	I'm not sure	I disagree	I disagree but not strongly
------------------	--------------------------	--------------	------------	-----------------------------

Forced choice

I would like to try parachute jumping.

I would never want to try jumping out of a plane, with or without a parachute.

Self-Esteem

I feel that I am an important person

Sensation Seeking: Sensation seeking was measured using the Zuckerman Sensation Seeking Scales form V (20). This scale measured the respondents' need to experience varied, novel, and complex sensations (20). From the 7th to the 10th grades, responses were recorded on a five point scale ranging from: I agree strongly to I disagree strongly (20). The scale was revised when the subjects were 21 to reflect recent modifications in the scale (21). In this version, subjects were asked to choose between two alternative statements. The Alpha coefficient for the most recent survey was .86

Data Analysis

First, early pubertal onset was defined as one standard deviation below the mean, which was age 10-11 for females and age 10-11½ for males. Late pubertal onset was determined as one standard deviation above the mean, which was age 14-17½ for females and age 14½ -17 for males. Mid onset was age 11½ -13½ for females and age 12-14 for males.

Then, two different analyses were carried out to examine the relationship of substance use with pubertal onset, sensation seeking, and self-esteem. First, post hoc tests were done using ANOVA with a Tukey HSD to compare group differences across substance use, sensation seeking, and self-esteem scores between females and males with early, mid and late pubertal onset. Reported differences are at the $p \leq .05$ level.

The interactions of pubertal onset, self-esteem, and sensation seeking were analyzed in those situations where substance use and either self-esteem or sensation seeking were associated with early pubertal onset. Specifically, in 7th grade females, early pubertal onset was associated with increased cigarette use and low self-esteem. Therefore, early pubertal onset, self-esteem, and the interaction of these two measures were entered as

independent variables in a backward stepwise regression. For 8th and 9th grade females, early pubertal onset was associated with cigarette use. Late pubertal onset was associated with lower sensation seeking levels so early pubertal onset, sensation seeking, and the interaction of these two measures were entered as an independent variables in a backward stepwise regression. For 9th grade males, early pubertal onset was associated with alcohol use and increased sensation seeking so sensation seeking, pubertal onset, and the interaction of these two measures were entered in a backward stepwise regression. Sensation seeking was coded so greater sensation seeking had a greater value than lower sensation seeking scores. Self-esteem was coded so lower self-esteem had a greater value, and pubertal onset was coded so that early pubertal onset had a greater value than late pubertal onset. This enabled the interaction terms to have the greatest value if puberty was early, sensation seeking was high, and self-esteem was low. Each analysis examined gender and grade separately.

RESULTS

Table 1 presents the ANOVA findings with a Tukey HSD comparison between retrospectively reported early, mid, and late pubertal onset in females and males for previous month cigarette and alcohol use, sensation seeking, and self-esteem. Early pubertal onset was associated with greater cigarette use in 7th, 8th, and 9th grade females and greater alcohol use in 9th and 10th grade males. Sensation seeking was significantly lower in late pubertal onset 8th and 9th grade females and 7th, 8th, and 9th grade and young adult males. In contrast, young adult females with early pubertal onset reported lower sensation seeking scores. In addition, self-esteem scores were significantly lower in early pubertal onset 7th grade females.

Table 1: ANOVA - Tukey HSD Comparisons of Reports of Substance Use, Self Esteem, and Sensation Seeking in Females and Males between Early, Mid and Late Pubertal Onset: Significant Mean Differences ($p \leq 0.05$).

		Early	Mid	Late	F	df	Group Differences*
7th Grade							
Cigarette/Month	Females	1.38	1.14	1.20	3.38 ^a	538	Early>Mid
Self Esteem	Females	38.73	40.78	42.25	9.34 ^c	538	Early<Mid&Late
Sensation Seeking	Males	54.08	55.14	50.74	3.55 ^a	402	Mid>Late
8th Grade							
Cigarette/Month	Females	1.81	1.28	1.23	8.91 ^c	532	Early>Mid&Late
Sensation Seeking	Females	49.96	50.61	47.20	4.01 ^a	533	Mid>Late
Sensation Seeking	Males	54.08	55.14	50.74	5.02 ^b	396	Mid>Late
9th Grade							
Cigarette/Month	Females	1.38	1.14	1.20	3.21 ^a	525	Early>Mid
Alcohol/Month	Males	1.88	1.57	1.28	4.40 ^b	387	Early>Late
Sensation Seeking	Females	50.99	53.23	49.77	5.25 ^b	528	Mid>Late
Sensation Seeking	Males	57.64	57.14	52.03	7.02 ^c	387	Early&Mid>Late
10th Grade							
Alcohol/Month	Males	2.13	2.01	1.48	3.27 ^a	341	Early>Late
Age 20							
Sensation Seeking	Females	12.87	15.52	15.30	6.74 ^c	564	Early<Mid&Late
Sensation Seeking	Males	19.58	20.03	16.88	6.31 ^c	419	Early&Mid>Late

a = $p \leq 0.05$

b = $p \leq 0.01$

c = $p \leq 0.001$

* = Tukey significant at ≤ 0.05

Table 2 presents the significant predictors of substance use that are associated with early pubertal onset using backward stepwise regression. The interaction of pubertal onset and self-esteem was predictive of cigarette use in 7th grade females while the interaction of pubertal onset and sensation seeking predicted cigarette use in 8th and 9th grade females and alcohol use in 9th grade males.

DISCUSSION

Early pubertal onset and substance use

This and other studies support the premise that adolescent substance use is associated with early pubertal onset (5, 6). Specifically, early pubertal onset is associated with cigarette use in adolescent females (4, 22). Similarly, the association of early pubertal onset with alcohol use in adolescent males (6) was confirmed in the current study. However, in the current study late pubertal onset in males is associated with decreased alcohol use, which is in contrast to the observations by Andersson and Magnusson (6) that late pubertal onset (like early pubertal onset), is associated with drinking to intoxication.

Early pubertal onset, sensation seeking and self-esteem

In this and other studies, early pubertal onset is associated with decreases in self-esteem (7) in adolescent females. While early pubertal onset is associated with decreases in self-esteem (7) in females, males report their pubertal changes more positively (23). Low self-esteem has been associated with smoking in females grades 6-8 but not males (24) and predicts alcohol use disorder in college aged women but not men (10). The association of early pubertal onset with substance use risk in females may be in part related to the association with low self-esteem.

Late pubertal onset is associated with decreased sensation seeking in males and females. Although sensation seeking has

been correlated with pubertal hormones [testosterone and estradiol (13)], the investigation of sensation seeking as it relates to pubertal onset has received limited attention. It is possible that late pubertal onset adolescent males and females are less biologically prone to be sensation seekers.

The lower sensation seeking scores in 20-year-old females reporting early pubertal onset may reflect a maturational process in which females with early pubertal onset may not have elevated sensation seeking in young adulthood. For example, Martin *et al.* (25) reported that adolescent females did not have increases in Conduct Disordered behaviors with age while adolescent males did have increased Conduct Disordered behaviors with age. This may imply a maturational process in females, which was not observed in males.

The interaction of early pubertal onset with self-esteem or sensation seeking

The prediction of cigarette and alcohol use by the interaction of pubertal onset with either self-esteem or sensation seeking highlights the importance of understanding why early pubertal onset is predictive of substance use. The observed interactions suggest the need for further investigation to determine whether there are independent variables such as pubertal hormones (i.e. testosterone, estradiol) that are related to early pubertal onset, sensation seeking, self-esteem, and substance use. Further investigation of these interactions should include youth as they enter and progress through puberty.

Examining the relationship between pubertal maturation and substance use is difficult because of the multiple complex variables (2,3,26). In addition, pubertal development measures are not precise and include current and retrospective self-reports, examination of secondary sexual characteristics, hormonal assays and laboratory measures of growth such as bone

Table 2: Significant Predictors of Substance Use Associated with Early Puberty: Backward Stepwise Regression of Pubertal Onset, Self Esteem, Sensation Seeking, (Pubertal Onset) X (Self Esteem), and (Pubertal Onset) X (Sensation Seeking)

Grade	Sex	Dependent Variables	Independent Variables	Regression Coefficient	Beta	R Square	P
8 th	F	Cigarette Use/Mon.h	Sensation Seeking	-.04	-.35 ^a	.12	.000
			Pubertal Onset	-.27	-.64 ^c		
			(Pubertal Onset) X (Sensation Seeking)	.007	1.04 ^b		
9 th	F	Cigarette Use/Month	Pubertal Onset	.09	-.16 ^a	.04	.000
9 th	M	Alcohol Use/Month	Pubertal Onset	.21	.54 ^a	.08	.000
			(Pubertal Onset) X (Sensation Seeking)	.005	.92 ^b		

a = p ≤ 0.05
 b = p ≤ 0.01
 c = p ≤ 0.001

age. The current study used retrospective self-report; although this approach is considered to be valid (15,16,18), it is limited when compared to current measure of pubertal onset.

Despite these limitations this study highlights the importance of increased monitoring of youth with early pubertal onset. The importance of understanding and monitoring youth with early pubertal onset is emphasized by recent observations regarding increased early adolescent substance use (27). The importance of identification of both youth at increased risk and developmental periods of increased risk for substance use is evident. Monitoring of youth should include measures of drug use, self-esteem, and sensation seeking.

ACKNOWLEDGEMENTS

This study was supported by NIDA grants number DA 05312 and 1 K08 DA00333

REFERENCES

1. Petersen AC, Taylor B. The biological approach to adolescence: Biological change and psychological adaptation. In: Adelson J, ed. Handbook of adolescent psychology. New York: Wiley, 1980:117-55.
2. Magnusson D, Stattin H, Allen VL. Biological maturation and social development: a longitudinal study of some adjustment processes from mid-adolescence to adulthood. *J Youth Adolesc* 1985;14:267-83.
3. Magnusson D. Individual development from an interactional perspective: a longitudinal study. Hillsdale, NJ: Earlbaum, 1988.
4. Caspi A, Lynam D, Moffitt TE, et al. Unraveling girls' delinquency: biological, dispositional, and contextual contributions to adolescent misbehavior. *Dev Psychol* 1993; 29:19-30.
5. Stattin H, Magnusson D. Pubertal Maturation in female development. Hillsdale, NJ: Earlbaum, 1990
6. Andersson T, Magnusson D. Biological maturation in adolescence and the development of drinking habits and alcohol abuse among young males. *J Youth Adolesc* 1990; 19:33-41.
7. Lackovic-Grgin K, Dekovic M, Opacoc G. Pubertal status, interaction with significant others, and self-esteem of adolescent girls. *Adolescence* 1994; 115:691-700.
8. Crockett LJ, Petersen AC. Pubertal status and psychosocial development: Findings from the early adolescence study. In: Lerner RM, Foch TT, eds. Biological-Psychosocial interactions in early adolescence: A life-span perspective. Hillsdale, NJ: Earlbaum; 1987.
9. Gordon WR, Caltabiano ML. Urban-rural differences in adolescent self-esteem, leisure boredom, and sensation seeking as predictors of leisure-time usage and satisfaction. *Adolescence* 1996; 31:883-901.
10. Walitzer KS, Sher KJ. A prospective study of self-esteem and alcohol use disorders in early adulthood: evidence for gender differences. *Alcohol Clin Exp Res* 1996; 20:1118-24.
11. Zuckerman M. Biological bases of sensation seeking, impulsivity, and anxiety. Hillsdale, NJ: Earlbaum, 1983.
12. Jaffe LT, Archer RP. The prediction of drug use among college students from MMPI, MCMC, and sensation seeking scales. *J Personality Assessment* 1987; 51:243-253.
13. Daitzman RJ, Zuckerman M, Sammelwitz P et al. Sensation seeking and gonadal hormones. *J Biosoc Sci* 1978;10:401-8.
14. Lynam DR, Milich R, Zimmerman R et al. *J Consult Clin Psychol* 1999; 67:590-93.

15. Bean JA, Leeper JD, Wallace RB, Sherman BM, Jagger H. Variations in the reporting of menstrual histories. *Am J Epidemiology* 1979;109:181-5.
16. Damon A, Bajema CJ. Age at menarche: accuracy of recall after thirty-nine years. *Hum Biol* 1974; 46:381-4.
17. Brooks-Gunn J, Warren MP, Rosso J et al. Validity of self-report measures of girls' pubertal status. *Child Dev* 1987; 58:829-41.
18. Gilger JW, Geary DC, Eisele LM. Reliability and validity of retrospective self-reports of the age of pubertal onset using twin, sibling, and college student data. *Adolescence* 1991; 26:41-53.
19. Rosenberg M. *Society and the adolescent self-image*. Princeton, NJ: Princeton Univ Press, 1965.
20. Zuckerman M. *Sensation Seeking: Beyond the Optimal Level of Arousal*. Hillsdale, NJ: Earlbaum, 1979.
21. Zuckerman M, Como P. Sensation seeking and arousal systems. *Pers Individ Differ* 1983; 4:381-86.
22. Wilson DM, Killen JD, Hayward C, et al. Timing and rate of sexual maturation and the onset of cigarette and alcohol use among teenage girls. *Arch Pediatr Adolesc Med* 1994;148:789-95.
23. Simmons RG, Blyth DA, Van Cleave EF et al. Entry into early adolescence: the impact of school structure, puberty, and early dating on self-esteem. *Am Soc Rev* 1979; 44: 948-67.
24. Abernathy TJ, Massad L, Romamo-Dwyer L. The relationship between smoking and self-esteem. *Adolescence* 1995;30:899-907.
25. Martin CA, Milich R, Martin WR et al. Gender differences in adolescent psychiatric outpatient substance use: associated behaviors and feelings. *J Am Acad Child Adolesc Psychiatr* 1997; 36:486-94.
26. Tschann JM, Adler NE, Irwin CE et al. Initiation of substance use in early adolescence: the roles of pubertal timing and emotional distress. *Health Psychol* 1994;13:326-33.
27. Johnston LD, O'Malley P, Bachman JG. National Survey results on drug use from The Monitoring the Future Study, 1975-1995, Vol. 1, Secondary School Students. Washington, D.C.: National Institute on Drug Abuse, 1996.

BITS 'N PIECES

In China, suicide in young women is a problem too

EDITOR—Mayor reports that the Men's Health Forum recommends that suicide in young British men needs multiagency solutions rather than efforts by psychiatrists or general practitioners alone. This is true in China too, except that it is young women who should be the target of interventions.

China's national suicide rate has been estimated to be about 30/100 000 annually, about three times the global average. Consequently, China accounts for more than four tenths of suicides reported worldwide, including more than half of suicides among women (although reliable data on suicide from Central and South America, Africa, the Middle East, and huge Asian countries such as India and Indonesia are still lacking). Among women, completed and attempted suicide is a greater source of lost working days than diabetes, heart disease, or cancer.

The rate is astonishingly high in young rural women aged 15-24, which casts doubt on the view of experts such as Baechler that "women endure misfortune better than do men. Their social roles require them to face unbearable problems less frequently. . . . As daughters, wives, and mistresses, and conforming to the dependency which nature and culture encourage, women have a greater tendency to reach their ends by the threat of trying to kill themselves. . . . Dangerous and aggressive behavior generally is not characteristic of women."

Evidence suggests that high rates of suicide in young Chinese rural women have multiple causes: low social status, forced marriage, domestic abuse, birth control policy, harassment by the husband's family, frustration over rural life, easy availability of pesticides, and greatly limited access to medical resuscitation facilities. Since most rural Chinese people do not have medical insurance and psychiatrists and even general practitioners are barely available in many rural areas, the prevention of suicide must rest, provided political will is available, on the intersectoral collaboration of multiple local agencies.

Sing Lee S. *BMJ* 2000;321:636.