



University of Kentucky
UKnowledge

Pharmaceutical Sciences Faculty Publications

Pharmaceutical Sciences

8-9-2017

A New Approach to Synthesize of 4-Phenacylidene-flavene Derivatives and to Evaluate Their Cytotoxic Effects on HepG2 Cell Line

Hongbin Chen
Quanzhou Normal University, China

Yang Xu
Xiamen University, China

Yinan Zhang
University of Kentucky, yinan.zhang@uky.edu

Zongping Zheng
Quanzhou Normal University, China

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/ps_facpub

 Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

A New Approach to Synthesize of 4-Phenacylidene-flavone Derivatives and to Evaluate Their Cytotoxic Effects on HepG2 Cell Line

Notes/Citation Information

Published in *Molecules*, v. 22, issue 8, p. 1-7.

© 2017 by the authors.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Digital Object Identifier (DOI)

<https://doi.org/10.3390/molecules22081296>

Article

A New Approach to Synthesize of 4-Phenacylidene-flavene Derivatives and to Evaluate Their Cytotoxic Effects on HepG2 Cell Line

Hongbin Chen ¹, Yang Xu ², Yinan Zhang ³ and Zongping Zheng ^{1,*} 

¹ Fujian Province Key Laboratory for the Development of Bioactive Material from Marine Alge, College of Oceanology and Food Science, Quanzhou Normal University, Quanzhou 362000, China; chenhongbin@qztc.edu.cn

² School of Pharmaceutical Sciences, Xiamen University, Xiamen 361005, China; xu_yang@xmu.edu.cn

³ College of pharmacy, University of Kentucky, Lexington, KY 40536, USA; noahzhang@gmail.com

* Correspondence: zzpsea@qztc.edu.cn; Tel./Fax: +86-595-2291-9563

Received: 29 June 2017; Accepted: 2 August 2017; Published: 9 August 2017

Abstract: In this study, a convenient approach and green procedure for the synthesis of 4-phenacylidene-flavenes has been developed from the reaction between 2,4-dihydroxybenzaldehyde and substituted acetophenones using boric acid as a catalyst in polyethylene glycol 400. Seven 4-phenacylidene-flavenes were synthesized and their structures were confirmed by NMR and mass spectral analyses. Meanwhile, their possible mechanism of formation was also discussed. These products were found to have potential cytotoxic effect on HepG2 cell line with IC₅₀ values from 12.5 to 50 μM.

Keywords: substituted acetophenones; 2,4-dihydroxybenzaldehyde; 4-phenacylidene-flavenes; cytotoxic effect; HepG2 cell line


1. Introduction

Chromene moieties are widely found in flavonoid derivatives. Their derivatives have become of much interest due to a variety of biological activities such as anticancer, antimicrobial, anti-inflammatory, and anti-HIV activities [1–4]. 4-Phenacylidene-flavene derivatives are one type of chromene moiety derivatives. However, the methods for the synthesis of phenacylidene-flavene derivatives were limited to only a few until now. Bhattacharjee and co-author recently reported one-pot synthesis of 4-phenacylidene-flavene derivatives with a simple reaction procedure, high bond-forming efficiency, good yields, and environmentally benign reaction conditions using bromodimethylsulfonium bromide as catalyst in [5]. Sashidhara et al. also reported one-pot synthesis of 4-phenacylidene-flavene derivatives through the reaction of salicylaldehydes and acetophenones using I₂ as catalyst under reflux conditions [6]. Other methods available for the synthesis of phenacylidene-flavene derivatives were as follows: Fichtner et al. reported to obtain 4-phenacylidene-flavene derivatives from the reaction of flavylum salt with 1-phenyl-1-(trimethylsiloxy)ethene in the presence of HBF₄·OEt₂ or TfOH, Vanallan et al. and Hill used multi-step condensation reaction of salicylaldehyde and acetophenone to prepare 4-phenacylidene-flavene derivatives in hot HCl or AcOH [7–9]. Although some of these methods are quite useful, there is still further need to develop the simple, efficient, inexpensive, and environment-friendly procedures for the synthesis of 4-phenacylidene-flavene.

With an increasing ecological pressure, new green synthetic strategies are of continuously growing interest to synthesize valuable organic compounds due to their environment-friendliness, but this depends on the discovery and usage of mild, low-cost, and high-performance catalysts. Boric acid is a commercially available, environmentally benign, and inexpensive catalyst. Boric acid is a weak solid

acid and slightly soluble in water. It forms tetrahydroxyborate in water and serves as a Lewis acid to play catalytic role in various reactions, such as aza-Michael reaction, Biginelli reaction, Mannich reaction, and transamidation of carboxamides [10–14].

Recently, in the course of our synthesizing 2',4'-dihydroxychalcone derivatives, our research group accidentally discovered a new way to synthesize 4-phenacylidene flavene derivatives. To the best of our knowledge, no reports for the use of boric acid for condensation of 2,4-dihydroxybenzaldehyde (**1**) and substituted acetophenones (**2a–2g**) to form 4-phenacylidene flavenes were reported. Herein, we report a one-pot procedure for the green synthesis of 4-phenacylidene flavenes (**4a–4g**) by the condensation of 2,4-dihydroxybenzaldehyde with various substituted acetophenones using boric acid as a green catalyst in polyethylene glycol 400 (Scheme 1). Meanwhile, the products (**4a–4g**) were also subjected to cytotoxic tests on liver cancer cell line (HepG2).



Scheme 1. One-pot synthesis of 4-phenacylidene flavene derivatives.


2. Results and Discussion

2.1. Chemistry

The desired seven 4-phenacylidene flavene derivatives (**4a–4g**) were prepared using a straightforward one step reaction, as presented in Scheme 1. For the present study, the mixtures of 2,4-dihydroxybenzaldehyde (**1**, 1 mmol) and various substituted acetophenones (**2a–2g**, 1 mmol) were stirred in the presence of 0.5 mmol of boric acid in PEG-400 (5 mL) for 6 h at 130 °C. Seven 4-phenacylidene flavene products **4a–4g** were isolated by chromatographic purification. The yields of products **4a–4g** ranging from 3.06% to 25.28% were determined by HPLC analysis. The yields of products were maybe affected by the molar ratios between 2,4-dihydroxybenzaldehyde and acetophenones and solvent conditions. Taking the product **4e** as example, when the ratio of 2,4-dihydroxybenzaldehyde and 4-methylacetophenone was 1:2, the yield of **4e** had great improvement—up to 58.14%—whereas when the reaction was taken in methanol, no product of **4e** was found. On the other hand, the different substituent groups and substituent positions also affected the yields of products. The results indicated that the conversion to acylidene flavenes preferred electron-rich substituents (R: OH, OMe, etc.) on the aromatic ring than electron-neutral one (R: H). However, electron withdraw groups (R: CN, NO₂) did not give any desired products under this condition (data not shown). The low yield of **4b** was probably due to the formation of intramolecular hydrogen bond between 2-OH and acetophenone, which inhibited the Aldol reaction and Michael addition afterwards. The structures of the synthesized compounds were characterized by ¹H-NMR, ¹³C-NMR, and ESI mass spectra. The structures of compounds **4b–4d** were finally determined on the base of the 2D-NMR (¹H-¹H COSY, HSQC, HMBC, Supplementary materials), whereas the structures of **4a**, **4e–4g** were determined on the base of the ¹H-NMR and ¹³C-NMR and the structure of **4d**. ¹H-NMR, ¹³C-NMR and mass spectrometric data of synthesized compounds are summarized in Materials and Methods.

A plausible mechanism for the formation of products **4a–4g** was derived and shown in Scheme 2 on the basis of the reported literature [6]. The initial step was Aldol reaction catalyzed by boric acid between 2,4-dihydroxybenzaldehyde (**1**) and substituted acetophenones (**2a–2g**) and firstly led to the products of 2',4'-dihydroxychalcones (**3a–3g**). The mechanism in path 1 involved in a Michael

addition between substituted acetophenone (**2a–2g**) and 2',4'-dihydroxychalcones (**3a–3g**) to offer the products **5a–5g**. The products **5a–5g** further underwent intramolecular cyclization, followed by dehydration to form adducts (**6a–6g**) and then were oxidized to form the product **4a–4g**. The other plausible mechanism in path 2 involved in intramolecular cyclization of 2',4'-dihydroxychalcones (**3a–3g**) to form their hemiacetal species **5a'–5g'**, which were then converted to more reactive flavylum ions (**6a'–6g'**). The flavylum ions (**6a'–6g'**) further reacted with substituted acetophenone (**2a–2g**) to form adducts (**6a–6g**). Finally, adducts were oxidized to form the products **4a–4g**, probably through a disproportionation mechanism suggested by VanAllen [8], which also accounted for the low yield of 4-phenacylidene flavene.



Scheme 2. Proposed mechanism for the synthesis of 4-phenacylidene flavene derivatives.

2.2. Cytotoxic Effects on HepG2 Cell Line

In vitro cytotoxicities of seven 4-phenacylidene flavene derivatives (**4a–4g**) were preliminarily evaluated against the selected cancer cell line (HepG2). The results of anticancer activity are expressed as IC_{50} values and are shown in Table 1. As presented in Table 1, among the seven compounds,

compound **4e** showed the greatest cytotoxic activity against HepG2 cell. Substitution at the 4 position with methyl group showed significant activity against HepG2, while substitutions with the hydroxyl, methoxyl, and chlorine groups caused substantial decrease in activities, suggesting that the substitution with different groups will cause substantial difference in activities. Thus, more research is worth to carrying out to figure out what group leads to more potential cytotoxicity against HepG2 cell.

Table 1. Inhibitory effects of the products against HepG2 cell line

Products	R	HepG2 (IC ₅₀ μM)
4a	H	20–50
4b	2-OH	20–50
4c	3-OH	>50
4d	4-OH	>50
4e	4-CH ₃	≈12.5 ^a
4f	4-OCH ₃	20–50
4g	4-Cl	>50

^a Values are means ± SD (*n* = 3).

3. Materials and Methods

3.1. General Information

Analytical HPLC was carried out on a Waters 1525 system (Waters, Milford, MA, USA) equipped with a 2487 dual-wavelength detector and the Empower 2 Pro software (Waters, Milford, MA, USA). Alltima C₁₈ column (250 × 4.6 mm, 5 μm, Delta Technical Products Co., Des Plaines, IL, USA) was used for analytical HPLC. ¹H-NMR, ¹³C-NMR, HSQC, and HMBC data were acquired on a Bruker 400 DRX NMR spectrometer (Bruker, Colorado Springs, CO, USA). Molecular weights of compounds were analyzed on Waters Maldi Syapt Q-ToF mass spectrometer (Waters, Milford, MA, USA). Spectrophotometric measurements for the tyrosinase inhibition assay were taken on a UV-5300PC Spectro-photometer (Metash Instrument Co., Ltd., Shanghai, China). 2,4-Dihydroxybenzaldehyde (**1**), acetophenone (**2a**), 2-hydroxyacetophenone (**2b**), 3-hydroxyacetophenone (**2c**), 4-hydroxyacetophenone (**2d**), 4-methylacetophenone (**2e**), 4-methoxyacetophenone (**2f**), and 4-chloroacetophenone (**2g**) were purchased from Shanghai Darui Company (Shanghai, China). Polyethylene glycol 400 (PEG 400), ethanol (EtOH), methanol (MeOH), boric acid, and dichloromethane (CH₂Cl₂) were purchased from Sinopharm Chemical Reagent Co., Ltd. (Suzhou, China). Silica gel (200–300 mesh) for column chromatography and TLC plates (HSGF254) were purchased from Yantai Jiangyou Silicone Development Co. (Yantai, China). Dichloromethane (CH₂Cl₂), dimethyl sulfoxide (DMSO), 95% ethanol (EtOH), methanol (MeOH), sodium dihydrogen orthophosphate (NaH₂PO₄·2H₂O), formic acid, and anhydrous di-sodium hydrogen phosphate (Na₂HPO₄) were purchased from Sinopharm Chemical Reagent Co., Ltd. (Suzhou, China). HPLC grade solvents were purchased from J&K Scientific Ltd. (Beijing, China).

3.2. General Procedure for the Synthesis of Compounds **4a–4g**

The mixture of 2,4-dihydroxybenzaldehyde (**1**, 1 mmol), various substituted acetophenone (**2a–2g**, 1 mmol), and boric acid (0.5 mmol) were dissolved in PEG-400 and then were stirred to react for 6 h at 130 °C. After the completion of the reaction, the reaction mixture was extracted with ethyl acetate three times. The collected organic layers were combined and concentrated in vacuum, and the residue was subjected to column chromatography on silica gel using CH₂Cl₂ (**4a**, **4e–4g**) and CH₂Cl₂-MeOH (30:1, **4b–4d**) as eluent to generate the pure product.

2-(7-Hydroxy-2-phenyl-chromen-4-ylidene)-1-phenyl-ethanone (**4a**): Yellow powder; m.p. 260 °C; Yield 4.79%; ESI-MS *m/z* 339.1 [M – H][−]; ¹H-NMR (400 MHz, DMSO-*d*₆) δ ppm: 10.58 (1H, s, OH-4'), 8.88 (1H, s, H-10), 8.24 (1H, d, *J* = 9.6 Hz, H-6'), 8.10 (2H, d, *J* = 7.6 Hz, H-2, 6), 7.95 (2H, d, *J* = 7.2 Hz, H-2',

6'), 7.58 (2H, d, $J = \text{Hz}$, H-3', 5'), 7.58 (1H, s, H-8), 7.56 (1H, m, H-4), 7.53 (2H, d, $J = 7.6 \text{ Hz}$, H-3, 5), 7.19 (1H, s, H-4'), 6.90 (2H, overlapped, H-3'', 5''); $^{13}\text{C-NMR}$ (100 MHz, DMSO- d_6) δ ppm: 188.5 (C=O, C-7), 161.3 (C, C-4''), 155.1 (C, C-11), 153.9 (C, C-2''), 142.1 (C, C-1''), 140.7 (C, C-1'), 132.0 (C, C-1), 131.5 (CH, C-4), 130.7 (CH, C-4'), 129.1 (CH, C-2, 6), 128.37 (CH, C-2', 6'), 127.6 (CH, C-3, 5), 125.6 (CH, C-6''), 125.4 (CH, C-3', 5'), 114.8 (CH, C-10), 111.5 (C, C-9), 103.0 (CH, C-5''), 101.8 (CH, C-3''), 101.0 (CH, C-8).

2-[7-Hydroxy-2-(2-hydroxy-phenyl)-chromen-4-ylidene]-1-(2-hydroxy-phenyl)-ethanone (4b): Red power; m.p. 249 °C; Yield 6.46%; ESI-MS m/z 371.1 [M – H] $^-$; $^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ ppm: 13.82 (1H, s, OH-2), 10.65 (2H, s, OH-2', 4''), 9.20 (1H, s, H-10), 8.32 (1H, d, $J = 9.2 \text{ Hz}$, H-6''), 8.26 (1H, d, $J = 8.0 \text{ Hz}$, H-6), 7.84 (1H, d, $J = 7.6 \text{ Hz}$, H-6'), 7.46 (1H, t, $J = 7.6 \text{ Hz}$, H-4), 7.35 (1H, t, $J = 8.0 \text{ Hz}$, H-4'), 7.21 (1H, s, H-8), 7.02 (1H, d, $J = 8.8 \text{ Hz}$, H-3'), 6.98 (1H, d, $J = 7.6 \text{ Hz}$, H-3), 6.93 (1H, overlapped, H-5'), 6.92 (1H, overlapped, H-5), 6.90 (1H, overlapped, H-3''), 6.88 (1H, overlapped, H-5''); $^{13}\text{C-NMR}$ (100 MHz, DMSO- d_6) δ ppm: 192.3 (C=O, C-7), 162.3 (C, C-2), 161.6 (C, C-4''), 156.2 (C, C-2'), 154.4 (C, C-11), 154.3 (C, C-2''), 145.0 (C, C-1''), 134.6 (CH, C-4), 131.7 (CH, C-4'), 129.4 (CH, C-6), 128.3 (CH, C-6'), 126.0 (CH, C-6''), 121.8 (C, C-1), 119.3 (CH, C-3), 118.6 (C, C-1'), 118.4 (CH, C-5'), 117.7 (CH, C-5), 116.8 (CH, C-3'), 115.0 (CH, C-5''), 111.6 (C, C-9), 106.6 (CH, C-10), 102.8 (CH, C-3''), 98.6 (CH, C-8).

2-[7-Hydroxy-2-(3-hydroxy-phenyl)-chromen-4-ylidene]-1-(3-hydroxy-phenyl)-ethanone (4c): Red power; m.p. 310 °C; Yield 19.27%; ESI-MS m/z 371.1 [M – H] $^-$; $^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ ppm: 10.56 (1H, s, OH-4''), 9.82 (1H, s, OH-3), 9.62 (1H, s, OH-3'), 8.79 (1H, s, H-10), 8.18 (1H, d, $J = 9.2 \text{ Hz}$, H-6''), 7.55 (1H, d, $J = 8.0 \text{ Hz}$, H-6), 7.41 (1H, H-4), 7.38 (2H, overlapped, H-5', 6'), 7.35 (1H, H-4'), 7.31 (1H, t, $J = 8.0 \text{ Hz}$, H-5), 7.10 (1H, s, H-8), 6.97 (1H, H-2), 6.95 (1H, H-2'), 6.88 (1H, dd, $J = 8.8, 2.4 \text{ Hz}$, H-5''), 6.85 (1H, d, $J = 2.4 \text{ Hz}$, H-3''); $^{13}\text{C-NMR}$ (100 MHz, DMSO- d_6) δ ppm: 188.5 (C=O, C-7), 161.2 (C, C-4''), 157.8 (C, C-3'), 157.4 (C, C-3), 155.0 (C, C-11), 153.8 (C, C-2''), 142.2 (C, C-1), 141.9 (C, C-1''), 133.2 (C, C-1'), 130.2 (CH, C-5'), 129.3 (CH, C-5), 125.5 (CH, C-6''), 118.6 (CH, C-6), 118.4 (CH, C-2), 117.9 (CH, C-2'), 116.2 (CH, C-6'), 114.8 (CH, C-5''), 114.0 (CH, C-4), 111.9 (CH, C-4'), 111.5 (C, C-9), 102.9 (CH, C-3''), 101.7 (CH, C-10), 101.0 (CH, C-8).

2-[7-Hydroxy-2-(4-hydroxy-phenyl)-chromen-4-ylidene]-1-(4-hydroxy-phenyl)-ethanone (4d): Red power; m.p. 197 °C; Yield 3.06%; ESI-MS m/z 371.2 [M – H] $^-$; $^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ ppm: 10.15 (3H, br s, OH-4, 4', 4''), 8.74 (1H, s, H-10), 8.17 (1H, d, $J = 8.8 \text{ Hz}$, H-6''), 8.00 (2H, d, $J = 8.8 \text{ Hz}$, H-2, 6), 7.79 (2H, d, $J = 8.4 \text{ Hz}$, H-2', 6'), 7.09 (1H, s, H-8), 6.96 (2H, d, $J = 8.8 \text{ Hz}$, H-3', 5'), 6.86 (2H, d, $J = 8.8 \text{ Hz}$, H-3, 5), 6.85 (2H, overlapped, H-3'', 5''); $^{13}\text{C-NMR}$ (100 MHz, DMSO- d_6) δ ppm: 187.2 (C=O, C-7), 160.9 (C, C-4''), 160.7 (C, C-4), 159.9 (C, C-4'), 155.0 (C, C-11), 153.7 (C, C-2''), 141.4 (C, C-1''), 132.2 (C, C-1), 129.9 (CH, C-2, 6), 127.2, 127.1 (CH, C-2', 6'), 125.4 (CH, C-6''), 122.7 (C, C-1'), 116.0, 115.8 (CH, C-3', 5'), 115.0, 114.8 (CH, C-3, 5), 111.7 (C, C-9), 103.1, 102.8 (CH, C-3'', 5''), 99.8 (CH, C-8, 10).

2-(7-Hydroxy-2-p-tolyl-chromen-4-ylidene)-1-p-tolyl-ethanone (4e): Yellow power; m.p. 258 °C; Yield 25.28%; ESI-MS m/z 367.2 [M – H] $^-$; $^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ ppm: 10.52 (1H, s, OH-4''), 8.84 (1H, s, H-10), 8.20 (1H, d, $J = 8.8 \text{ Hz}$, H-6''), 8.00 (2H, d, $J = 8.0 \text{ Hz}$, H-2, 6), 7.82 (2H, d, $J = 8.4 \text{ Hz}$, H-2', 6'), 7.37 (2H, d, $J = 8.4 \text{ Hz}$, H-3', 5'), 7.31 (2H, d, $J = 8.0 \text{ Hz}$, H-3, 5), 7.15 (1H, s, H-8), 6.88 (1H, dd, $J = 8.0, 2.4 \text{ Hz}$, H-5''), 6.87 (1H, d, $J = 2.4 \text{ Hz}$, H-3''); $^{13}\text{C-NMR}$ (100 MHz, DMSO- d_6) δ ppm: 188.7 (C=O, C-7), 161.2 (C, C-4''), 155.0 (C, C-11), 153.8 (C, C-2''), 141.9 (C, C-1''), 141.6 (C, C-4), 140.7 (C, C-4'), 138.1 (C, C-1), 129.7 (CH, C-2, 6), 129.2 (C, C-1'), 129.0 (CH, C-2', 6'), 127.7 (CH, C-3, 5), 125.6 (CH, C-6''), 125.3 (CH, C-3', 5'), 114.7 (CH, C-10), 111.6 (C, C-9), 103.0 (CH, CH-5''), 101.2 (CH, C-3''), 100.6 (CH, C-8), 21.0 (CH₃-4, 4').

2-[7-Hydroxy-2-(4-methoxy-phenyl)-chromen-4-ylidene]-1-(4-methoxy-phenyl)-ethanone (4f): Yellow power; m.p. 260 °C; Yield 19.37%; ESI-MS m/z 399.2 [M – H] $^-$; $^1\text{H-NMR}$ (400 MHz, DMSO- d_6) δ ppm: 8.78 (1H, s, H-10), 8.20 (1H, d, $J = 9.6 \text{ Hz}$, H-6''), 8.09 (2H, d, $J = 8.8 \text{ Hz}$, H-2, 6), 7.88 (2H, d, $J = 8.4 \text{ Hz}$, H-2', 6'), 7.13 (2H, d, $J = 8.8 \text{ Hz}$, H-3', 5'), 7.121 (1H, s, H-8), 7.03 (2H, d, $J = 8.8 \text{ Hz}$, H-3, 5), 6.87 (1H, overlapped, H-3''), 6.86 (1H, overlapped, H-5''); $^{13}\text{C-NMR}$ (100 MHz, DMSO- d_6) δ ppm: 187.3 (C=O,

C-7), 162.0 (C, C-4''), 161.3 (C, C-4), 161.1 (C, C-4'), 154.9 (C, C-11), 153.8 (C, C-2''), 141.7 (C, C-1''), 133.6 (C, C-1), 129.8 (CH, C-2, 6), 127.2 (CH, C-2', 6'), 125.5 (CH, C-6''), 124.3 (C, C-1'), 114.6 (CH, C-3', 5'), 114.6 (CH, C-10), 113.6 (CH, C-3, 5), 111.7 (C, C-9), 103.0 (CH, CH-5''), 100.4 (CH, C-3''), 100.2 (CH, C-8), 55.5 (CH₃O-4), 55.4 (CH₃O-4').

1-(4-Chloro-phenyl)-2-[2-(4-chloro-phenyl)-7-hydroxy-chromen-4-ylidene]-ethanone (**4g**): Yellow powder; m.p. 298 °C; Yield 20.45%; ESI-MS m/z 407.1 [M – H][–]; ¹H-NMR (400 MHz, DMSO-*d*₆) δ ppm: 8.79 (1H, s, H-10), 8.16 (1H, d, *J* = 8.8 Hz, H-6''), 8.08 (2H, d, *J* = 8.4 Hz, H-2, 6), 7.92 (2H, d, *J* = 8.4 Hz, H-2', 6'), 7.60 (2H, d, *J* = 8.4 Hz, H-3', 5'), 7.53 (2H, d, *J* = 8.4 Hz, H-3, 5), 7.12 (1H, s, H-8), 6.89 (1H, dd, *J* = 8.8, 2.0 Hz, H-5''), 6.86 (1H, d, *J* = 2.4 Hz, H-3''); ¹³C-NMR (100 MHz, DMSO-*d*₆) δ ppm: 187.0 (C=O, C-7), 161.3 (C, C-4''), 153.6 (C, C-11), 153.7 (C, C-2''), 142.2 (C, C-1''), 139.2 (C, C-1), 136.3 (C, C-4), 135.3 (C, C-4'), 130.7 (C, C-1'), 129.2 (CH, C-2, 6), 128.2 (CH, C-2', 6'), 127.0 (CH, C-3, 5), 125.3 (CH, C-6''), 114.8 (CH, C-10), 111.3 (C, C-9), 102.9 (CH, CH-5''), 102.1 (CH, C-3''), 100.8 (CH, C-8).

3.3. HPLC Analysis of the Products

The yields of products were determined by HPLC analysis on the crude reaction mixture after having made a calibration curve with pure compounds isolated by chromatography. The analytical HPLC system consisted of a Shimadzu LC-20AT series pumping system, an SIL-20A automatic injector, an SPD-M20A UV-visible detector, and Class-Vp chromatography data station software. All samples were analyzed by HPLC using a reverse-phase GraceSmart column (4.6 μm, 2.1 × 250 mm, Ryss Tech Ltd., Shanghai, China) at 30 °C with a flow rate of 1.0 mL/min. The mobile phases consisted of solvent A (0.1% formic acid in water, *v/v*) and solvent B (methanol). The gradient elution was as follows: initially, 20% B; 0–10 min, 50% B; 10–30 min, 80% B; 30–32 min, 100% B; 32–35 min, 100% B; 35–40 min, 20% B; 40–50 min, 20% B. Flow rate was set at 1.0 mL/min. The sample injection volume was 10 μL. The UV detector was set at 369 nm.

3.4. Biological Evaluation

3.4.1. Cell Culture

HepG2 cell was cultured in Dulbecco's Modified Eagle Medium (high glucose) and RPMI-1640, respectively, with 10% fetal bovine serum under standard culture conditions. When the cells grew to about 80% confluence, they were sub-cultured or treated with products.

3.4.2. Growth Inhibition Study

The proliferation of cells was assessed by using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. Cells were plated on 96-well microplates in 100 μL (1 × 10⁴/well). After 12 h, the cells were treated with the medium (without FBS) containing compounds (12.5, 25, and 50 μM) for 24 h. At the end of experiments, 10 μL of 5 μg/mL MTT was directly added to each well. Cells were then incubated at 37 °C for 4 h. After removing medium, formazan was solubilized by 100 μL of DMSO and measured at 560 nm. All experiments were repeated three times.

4. Conclusions

In conclusion, a simple one pot synthesis of 4-phenacylidene flavene derivatives from the reaction between 2,4-dihydroxybenzaldehyde and substituted acetophenones had been developed in the presence of boric acid as catalyst in PEG 400. The method provided an inexpensive, safe, simple, and eco-friendly way to synthesize 4-phenacylidene flavene derivatives. On the other hand, the synthesized 4-phenacylidene flavene products were found to have potential inhibitory activities against HepG2 cell line. To the best of our knowledge, this is the first report for the use of boric acid for condensation of 2,4-dihydroxybenzaldehyde and substituted acetophenones to form 4-phenacylidene flavenes and

preliminarily evaluated their anticancer activities against HepG2 cell line, which will contribute to the development of green strategy for synthesizing various biologically flavene products.

Supplementary Materials: The 1D- (^1H - and ^{13}C -NMR), 2D-NMR (^1H - ^1H COSY, HSQC and HMBC), ESI-MS spectra of compounds are available in the supplementary materials.

Acknowledgments: This work was supported by grants from the Young Teacher Education Research Project of Fujian Province of China (JA15390) and the Science and Technology Project of Quanzhou of China (2015Z135). We would like to express our sincere appreciation to Shuang Zhang and Xiaohong Gu from state Key Laboratory of Food Science and Technology, Jiangnan University for measuring the mass data and NMR data of all compounds.

Author Contributions: Hongbin Chen contributed to synthesizing some of the compounds and analyzing the ESI-MS data of all compounds. Yang Xu contributed to the activity test and drafted the activity test section of the paper. Yanan Zhang contributed the plausible mechanism of synthesized compounds. Zongping Zheng initiated and coordinated the project and drafted other sections of the paper. All authors read and approved the final manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Korobkova, E.A. Flavonoids and mechanisms of their anticancer action. *Chem. Biol. Interface* **2013**, *3*, 346–365.
2. Ritter, M.; Martins, R.M.; Dias, D.; Pereira, C.M.P. Recent advances on the synthesis of chalcones with antimicrobial activities: A brief review. *Lett. Org. Chem.* **2014**, *11*, 498–508. [[CrossRef](#)]
3. Yasuda, M.; Kawabata, K.; Miyashita, M.; Okumura, M.; Yamamoto, N.; Takahashi, M.; Ashida, H.; Ohigashi, H. Inhibitory effects of 4-hydroxyderricin and xanthoangelol on lipopolysaccharide-induced inflammatory responses in RAW264 macrophages. *J. Agric. Food Chem.* **2014**, *62*, 462–467. [[CrossRef](#)] [[PubMed](#)]
4. Casano, G.; Dumetre, A.; Pannecouque, C.; Hutter, S.; Azas, N.; Robin, M. Anti-HIV and antiplasmodial activity of original flavonoid derivatives. *Bioorgan. Med. Chem.* **2010**, *18*, 6012–6023. [[CrossRef](#)] [[PubMed](#)]
5. Bhattacharjee, S.; Das, D.K.; Khan, A.T. Bromodimethylsulfonium bromide: An efficient catalyst for one-pot synthesis of 4-phenacylidene flavene derivatives. *Tetrahedron Lett.* **2015**, *56*, 2412–2415. [[CrossRef](#)]
6. Sashidhara, K.V.; Kumar, A.; Agarwal, S.; Kumar, M.; Kumar, B.; Sridhar, B. A simple and efficient access to new functionalized 4-phenacylidene flavenes. *Adv. Synth. Catal.* **2012**, *354*, 1129–1140. [[CrossRef](#)]
7. Fichtner, C.; Remennikov, G.; Mayr, H. Kinetics of the reactions of flavylum ions with π -nucleophiles. *Eur. J. Org. Chem.* **2001**, *20*, 4451–4456. [[CrossRef](#)]
8. Vanallan, J.A.; Reynolds, G.A.; Regan, T.H. Formation of 4-phenacylidene flavene from 4-phenacylflavene. *J. Org. Chem.* **1967**, *32*, 1897–1899. [[CrossRef](#)]
9. Hill, D.W. Reactions of *o*-hydroxybenzylidenediacetophenones. Part I. Reaction with acids. *J. Chem. Soc.* **1934**, *9*, 1255–1258. [[CrossRef](#)]
10. Hwang, H.T.; Varma, A. Effect of boric acid on thermal dehydrogenation of ammonia borane: Mechanistic studies. *Int. J. Hydrogen Energy* **2013**, *38*, 1925–1931. [[CrossRef](#)]
11. Chaudhuri, M.K.; Hussain, S.; Kantam, M.L.; Neelima, B. Boric acid: A novel and safe catalyst for aza-Michael reactions in water. *Tetrahedron Lett.* **2005**, *46*, 8329–8331. [[CrossRef](#)]
12. Tu, S.; Fang, F.; Miao, C.; Jiang, H.; Feng, Y.; Shi, Y.; Wang, X. One-pot synthesis of 3,4-dihydropyrimidin-2(1H)-ones using boric acid as catalyst. *Tetrahedron Lett.* **2003**, *44*, 6153–6155. [[CrossRef](#)]
13. Mukhopadhyay, C.; Datta, A.; Butcher, R.J. Highly efficient one-pot, three-component Mannich reaction catalysed by boric acid and glycerol in water with major 'syn' diastereoselectivity. *Tetrahedron Lett.* **2009**, *50*, 4246–4250. [[CrossRef](#)]
14. Nguyen, T.B.; Sorres, J.; Tran, M.Q.; Ermolenko, L.A. Boric acid: A highly efficient catalyst for transamidation of carboxamides with amines. *Org. Lett.* **2012**, *14*, 3202–3205. [[CrossRef](#)] [[PubMed](#)]

Sample Availability: Samples of the compounds **4a–4g** are available from the authors.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).