



University of Kentucky  
UKnowledge

Molecular and Cellular Biochemistry Faculty  
Publications

Molecular and Cellular Biochemistry

6-10-2013

# Archaeal Nucleosome Positioning In Vivo and In Vitro is Directed by Primary Sequence Motifs

Narasimharao Nalabothula  
*University of Kentucky*, [nna226@uky.edu](mailto:nna226@uky.edu)

Liqun Xi  
*Northwestern University*

Sucharita Bhattacharyya  
*Northwestern University*

Jonathan Widom  
*Northwestern University*

Ji-Ping Wang  
*Northwestern University*

*See next page for additional authors*

**Right click to open a feedback form in a new tab to let us know how this document benefits you.**

Follow this and additional works at: [https://uknowledge.uky.edu/biochem\\_facpub](https://uknowledge.uky.edu/biochem_facpub)

 Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#), and the [Genetics and Genomics Commons](#)

## Repository Citation

Nalabothula, Narasimharao; Xi, Liqun; Bhattacharyya, Sucharita; Widom, Jonathan; Wang, Ji-Ping; Reeve, John N.; Santangelo, Thomas J.; and Fondufe-Mittendorf, Yvonne N., "Archaeal Nucleosome Positioning In Vivo and In Vitro is Directed by Primary Sequence Motifs" (2013). *Molecular and Cellular Biochemistry Faculty Publications*. 24.  
[https://uknowledge.uky.edu/biochem\\_facpub/24](https://uknowledge.uky.edu/biochem_facpub/24)

This Article is brought to you for free and open access by the Molecular and Cellular Biochemistry at UKnowledge. It has been accepted for inclusion in Molecular and Cellular Biochemistry Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**Authors**

Narasimharao Nalabothula, Liqun Xi, Sucharita Bhattacharyya, Jonathan Widom, Ji-Ping Wang, John N. Reeve, Thomas J. Santangelo, and Yvonne N. Fondufe-Mittendorf

**Archaeal Nucleosome Positioning In Vivo and In Vitro is Directed by Primary Sequence Motifs****Notes/Citation Information**

Published in *BMC Genomics*, v. 14, 391, p. 1-13 .

© 2013 Nalabothula et al.; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Digital Object Identifier (DOI)**

<https://doi.org/10.1186/1471-2164-14-391>

RESEARCH ARTICLE

Open Access

# Archaeal nucleosome positioning in vivo and in vitro is directed by primary sequence motifs

Narasimharao Nalabothula<sup>1†</sup>, Liqun Xi<sup>2†</sup>, Sucharita Bhattacharyya<sup>3</sup>, Jonathan Widom<sup>3^</sup>, Ji-Ping Wang<sup>2</sup>, John N Reeve<sup>4</sup>, Thomas J Santangelo<sup>4,5</sup> and Yvonne N Fondufe-Mittendorf<sup>1\*</sup>

## Abstract

**Background:** Histone wrapping of DNA into nucleosomes almost certainly evolved in the Archaea, and predates Eukaryotes. In Eukaryotes, nucleosome positioning plays a central role in regulating gene expression and is directed by primary sequence motifs that together form a nucleosome positioning code. The experiments reported were undertaken to determine if archaeal histone assembly conforms to the nucleosome positioning code.

**Results:** Eukaryotic nucleosome positioning is favored and directed by phased helical repeats of AA/TT/AT/TA and CC/GG/CG/GC dinucleotides, and disfavored by longer AT-rich oligonucleotides. Deep sequencing of genomic DNA protected from micrococcal nuclease digestion by assembly into archaeal nucleosomes has established that archaeal nucleosome assembly is also directed and positioned by these sequence motifs, both in vivo in *Methanothermobacter thermoautotrophicus* and *Thermococcus kodakarensis* and in vitro in reaction mixtures containing only one purified archaeal histone and genomic DNA. Archaeal nucleosomes assembled at the same locations in vivo and in vitro, with much reduced assembly immediately upstream of open reading frames and throughout the ribosomal rDNA operons. Providing further support for a common positioning code, archaeal histones assembled into nucleosomes on eukaryotic DNA and eukaryotic histones into nucleosomes on archaeal DNA at the same locations. *T. kodakarensis* has two histones, designated HTkA and HTkB, and strains with either but not both histones deleted grow normally but do exhibit transcriptome differences. Comparisons of the archaeal nucleosome profiles in the intergenic regions immediately upstream of genes that exhibited increased or decreased transcription in the absence of HTkA or HTkB revealed substantial differences but no consistent pattern of changes that would correlate directly with archaeal nucleosome positioning inhibiting or stimulating transcription.

**Conclusions:** The results obtained establish that an archaeal histone and a genome sequence together are sufficient to determine where archaeal nucleosomes preferentially assemble and where they avoid assembly. We confirm that the same nucleosome positioning code operates in Archaea as in Eukaryotes and presumably therefore evolved with the histone-fold mechanism of DNA binding and compaction early in the archaeal lineage, before the divergence of Eukaryotes.

**Keywords:** Archaea, Nucleosome positioning, Dinucleotide repeats, Histone deletions, rDNA expression, Chromatin evolution

\* Correspondence: y.fondufe-mittendorf@uky.edu

†Equal contributors

^Deceased

<sup>1</sup>Department of Molecular and Cellular Biochemistry, College of Medicine, University of Kentucky, Lexington, KY 40536, USA

Full list of author information is available at the end of the article

## Background

Histone wrapping of nuclear DNA generates nucleosomes, the basic unit of chromatin in virtually all Eukaryotes. Nucleosomes are dynamically associated with the genome and their distribution is not random, but often plays a major role in regulating gene expression [1-4]. Nucleosome assembly is favored or deterred by differences in the enthalpic and entropic costs inherent in wrapping and maintaining different sequences in the rigid nucleosome toroid, and this has resulted in a eukaryotic nucleosome positioning code [1,3,5-8]. Histones, and presumably histone-DNA interactions, evolved before the divergence of the archaeal and eukaryotic lineages with histones now distributed throughout the *Euryarchaea*, *Nanoarchaea* and *Thaumarchaea* and also present in some *Crenarchaea* [9-11].

Sequencing DNA molecules selected from a large random population by repetitive selection (SELEX) and PCR-amplification, based on increased archaeal histone-DNA affinity, revealed that DNA molecules preferentially assembled into archaeal nucleosomes in vitro had sequences that conformed to the eukaryotic nucleosome positioning code [12]. However, the technology was not then available to determine if this was also the case in vivo and so confirm in vivo these results obtained in vitro. But, with the advent of large-scale DNA sequencing technologies, such comparisons became possible, and here we document that archaeal nucleosome assembly in vivo is directed by the nucleosome positioning code in both *Methanothermobacter thermautotrophicus* and *Thermococcus kodakarensis*, and that this is reproduced in vitro with archaeal genomic DNA and purified archaeal or eukaryotic histones. With recently developed genetic technologies, it was possible to delete either, but not both of the two archaeal histone-encoding genes present in *T. kodakarensis* generating strains that grow normally but exhibit transcriptome changes when compared with the parental strain [13]. Here we show that the histone deletions also result in changes in the archaeal nucleosome profiles upstream of the genes that have increased or decreased transcription.

## Results

### Nucleosome positioning motifs in vivo


The discovery of archaeal histones [14] and most subsequent studies [9,15], have investigated histones from methanogens for which genetic procedures are not available. We therefore first determined, and confirmed, that archaeal nucleosome positioning in vivo was directed by same sequence motifs in *M. thermautotrophicus* and *T. kodakarensis* and then focused on *T. kodakarensis*, as the experimental system, for which genetic procedures and strains with histone genes deleted were available [13]. As previously documented for *M. thermautotrophicus* [16],

MN digestion of chromatin from *T. kodakarensis* generated a ladder of discrete-length DNA molecules, with a predominant population of ~90 bp molecule first accumulating, consistent with the length of a DNA molecule required to completely encircle an archaeal histone tetramer core [17,18]. With further MN digestion, the number of ~90 bp molecules decreased and ~60 bp molecules accumulated (Figure 1a), the length of DNA directly bound by the histone folds of a histone tetramer. Deep-sequencing of the ~60 bp fragments resulted in 7 and 9 million unique reads, respectively, of molecules with lengths ranging from 55 to 65 nucleotides from *M. thermautotrophicus* and *T. kodakarensis* (Figure 1b). As established for DNA molecules preferentially packaged into eukaryotic nucleosomes [1,3,5-8,19-21], these sequences were dominated by helical repeats (10 bp) of AA/AT/TA/TT dinucleotides offset by 5 bp from repeats of GG/GC/CG/CC dinucleotides with the centers preferentially filled by GC-rich sequences (Figure 2a and 2b).

In contrast to A/T-rich dinucleotides that, in a 10 bp periodicity, offer flexibility and so facilitate DNA incorporation, poly (dA:dT) tracts are relatively rigid. This deters their incorporation into nucleosomes [1-3,6,8,22,23] and poly (dA:dT) tracts are significantly underrepresented in DNA incorporated into eukaryotic nucleosomes [1,6-8,23-33]. Analyses of the archaeal ~60 bp nucleosomal DNA fragments revealed that A/T-rich oligonucleotides were also excluded from incorporation into archaeal nucleosomes assembled in *M. thermautotrophicus* and *T. kodakarensis*. For example, as illustrated in Figures 2c and 2d, when the frequencies of occurrence of each of the 1064 pentamers was determined, all of the 32 pentamers that contain only A and/or T were underrepresented relative to the presence in the genome sequences with ATATA (=TATAT) being the most disfavored pentamer in both *Archaea*. Oligo A/T-rich sequences are consequently located preferentially in nucleosome-depleted regions. In contrast, all of the 32 pentamers that contain only G and/or C were enriched in the ~60 bp MN-protected fragments generated from both *Archaea* relative to their abundances in the genome sequences (Figure 2c and 2d).

### Exclusion of nucleosomes from intergenic regions

In *T. kodakarensis*, at least 92% of the genome is coding sequence [34], and most intergenic regions are <100 bp, with many <50 bp. Within these regions, both transcription initiation (TATA-box sequences) and termination (oligo A/T-rich sequences) are directed by A/T-rich sequences [35,36]. Given compliance with the nucleosome positioning code [5-8,19,37,38], archaeal nucleosome assembly should avoid intergenic regions and this was confirmed by aligning the ~60 bp nucleosomal sequences with the genome sequence. A transcriptome map has not been established for *T. kodakarensis* but, based on bioinformatic predictions [39],



**Figure 1** Archaeal nucleosomes protect ~60 bp chromosomal DNA fragments from micrococcal nuclease (MN) digestion. (a). Ethidium bromide stained electrophoretic separation of DNA molecules protected from MN digestion of *T. kodakarensis* TSS17 chromatin. The control lane (C) contained double-stranded DNA size standards (bp). As indicated, MN digestion generated DNA molecules that migrated to form a band consistent with ~60 bp molecules. These were isolated and sequenced. (b). The number and size profile of the sequences of the ~60 bp DNA fragment generated by ABI-SolID deep sequencing.

non-transcribed intergenic regions are preferentially depleted of nucleosomes. Genome-wide, there was a substantial underrepresentation of nucleosomes immediately upstream of translation initiation codons (Figure 3a) and, in multigene operons, this was predominantly upstream of the promoter proximal gene. For example, TK1761-TK1762-TK1763 constitutes an operon transcribed divergently from TK1760 [40]. The intergenic region separating TK1760 and TK1761 contains several oligo A/T-rich sequences and had minimal nucleosome occupancy in vivo (Figure 3b). There was no similar region of nucleosome exclusion downstream from the promoter within the TK1761-1763 operon.

#### Conservation of archaeal nucleosome positioning in vivo and in vitro

Most archaeal histones are single histone folds that form homodimers in solution [9]. However, when mixed with other archaeal histone homodimers, there is rapid and spontaneous reassembly that generates an equilibrium mixture of the homodimers and all possible heterodimers [18]. *M. thermoautotrophicus* has three histones [41] and *T. kodakarensis* has two histones [34] and, practically, it is impossible to know their homo- versus heterodimer configurations in vivo, a concern that limits the reproducibility in vitro of results obtained in vivo. The two histones in *T. kodakarensis*, HTkA and HTkB are encoded by TK1413 and TK2289, respectively, and although constructing a strain with both genes deleted proved impossible, *T. kodakarensis* LC124 ( $\Delta$ TK1413) and LC125 ( $\Delta$ TK2289) were constructed [13]. These strains therefore contain only one archaeal histone and, by heterologous expression of TK1413 and TK2289, preparations of recombinant HTkA and HTkB homodimers were obtained. Direct comparisons could therefore be made of the locations of archaeal


nucleosomes assembled by only HTkA or HTkB, in vivo and in vitro, on *T. kodakarensis* genomic DNA. The archaeal nucleosome profiles were very similar in vivo and in vitro throughout the length of the *T. kodakarensis* genome (Figure 4a). The pattern of nucleosome depletion immediately upstream of translation codons was conserved (Figure 4b) and there was only minimal archaeal nucleosome assembly on the rDNA operon both in vivo and in vitro (Figure 5).

As established for the parental strain (Figure 1a), the ~60 bp fragments of the *T. kodakarensis* genome protected from MN digestion by only HTkA or HTkB assembly in vivo and in vitro also contained 10 bp helical-periodicity repeats of AA/AT/TA/TT and GG/GC/CG/CC dinucleotides, offset by 5 bp, and pentamers containing only A and/or T were under-represented, and those containing only G and/or C were over-represented, relative to their occurrences in the *T. kodakarensis* genome (Additional file 1: Figure S1). Together these results confirm that the positions at which HTkA and HTkB assemble to form archaeal nucleosomes are predominantly determined by the *T. kodakarensis* genome sequence and, as concluded from eukaryotic nucleosome studies [7,33,38,43-46], from an archaeal genome sequence [39], it should be possible to predict where archaeal nucleosomes will preferentially assemble in vivo.

#### Evolutionary conservation of the nucleosome positioning code

source (eukaryotic versus archaeal), we assembled nucleosomes using eukaryotic histones with archaeal DNA and archaeal histones with eukaryotic DNA. As expected, based on many previous studies, chicken histone octamers bound *M. thermoautotrophicus* and *T. kodakarensis* genomic DNAs into nucleosomes that protected ~147 bp





**Figure 3 Depletion of archaeal nucleosomes in intergenic regions. (a)** The occurrence of nucleosomes relative to the start codons of all open reading frames (ORFs), as documented in the Archaeal Genome Browser [39]. The frequency of occurrence of each nucleosome read is plotted relative to the average value of occurrence of all nucleosomal reads sequenced from the *T. kodakarensis* TS517 genome. **(b)** The sequence of the intergenic region separating TK1760 and TK1761 is positioned above the profiles of nucleosomes assembled in vivo at this locus and downstream, within the well-established TK1761-TK1762-TK1763 operon [40].


nucleosome assembly upstream of TK2196, there was increased assembly upstream of TK1927 (Figure 7a). Similarly, in the absence of HTkA, transcripts of TK0166 and TK0982 decreased 5.3- and 3.5-fold, respectively, and there was increased nucleosome assembly upstream of TK0166, but there were both regions with decreased and increased nucleosome assembly upstream of TK0982 (Figure 7b).

## Discussion

### Conservation and ancestral origin of the nucleosome positioning code

Whether a sequence will favor or disfavor assemble into a nucleosome can be predicted from the below- or

above-average energy requirements needed to distort that sequence into the circular configuration of DNA wrapped around the nucleosome core [19,47,48]. To a large extent, these predictions have been confirmed experimentally by the sequences of DNA isolated from eukaryotic nucleosomes and so in the definition of a nucleosome positioning code [5-8,19,21,33,38,43]. The results reported here confirm that two basic features of this code, namely a 10 bp periodicity of AA/AT/TA/TT dinucleotides offset by 5 bp from GG/GC/CG/CC dinucleotides, and the exclusion of A/T-rich oligonucleotides also direct archaeal nucleosome assembly in vivo and in vitro. In the structures established for the eukaryotic



**Figure 4 Archaeal nucleosome profiles in vivo are reproduced in vitro.** (a) Profiles of the archaeal nucleosomes assembled in vivo and in vitro by HTkA and HTkB between nucleotide positions 5,000 and 10,000 on the *T. kodakarensis* genome. The organization of *T. kodakarensis* genes in this region is shown between upper and lower panels [34,39,42]. (b) The occurrence of nucleosome positions relative to the start codon of all open reading frames (ORFs), assembled by HTkA in vivo (blue line) and in vitro (green line), and by HTkB in vivo (red line) and in vitro (black line) normalized to the total number of nucleosomal reads from each sequencing experiment.


nucleosome [49,50], the AA/AT/TA/TT dinucleotide repeats facilitate DNA wrapping as the dinucleotides that most readily accept the distortion needed [48], where the minor groove faces inwards towards the histone octamer. The GG/GC/CG/CC dinucleotide repeats, offset by half a helical turn, facilitate the distortion needed at each site where the minor groove faces outwards and so away from the nucleosome core [3,49,50]. To date, despite a significant effort, there is no high resolution structure of an archaeal nucleosome available but, given that the archaeal and eukaryotic histone folds are virtually identical [9,51] and that DNA-histone interactions are almost entirely mediated by histone fold residues [52], it seems highly likely that the same wrapping mechanism is employed and so DNA distortion is required to assemble archaeal nucleosomes. Given this conservation, the

nucleosome positioning code almost certainly predates the Eukaryotes, and co-evolved with histones and the histone-fold based mechanism of DNA compaction early in the archaeal lineage, before the divergence of *Euryarchaea*, *Nanoarchaea* and *Thaumarchaea* [9-11].

#### Archaeal nucleosome positioning and gene expression

As in *Bacteria*, many archaeal genes are organized into operons and so are co-transcribed from a single promoter and, in the absence of a nuclear membrane; transcription and translation are coupled [53]. However, archaeal promoters and the transcription machinery are substantially more similar to their eukaryotic than bacterial counterparts. Archaeal transcription initiation requires that both transcription factors and RNA polymerase be recruited to the promoter [35] and archaeal histone binding







**Figure 5** Archaeal nucleosomes do not assemble on the ribosomal DNA operon. Profiles of the archaeal nucleosomes assembled by HTkA and HTkB in vivo in *T. kodakarensis* LC125 and LC124, respectively, and in vitro on a 10 Kbp region of the *T. kodakarensis* genome. As illustrated, this region encodes the 16S and 23S rRNA (rDNA operon; red arrows) and several protein-encoding flanking genes [34,39].

to promoter regions has been shown, alternatively, to inhibit or to stimulate transcription in vitro [54,55]. Without a transcriptome, we chose to determine the locations at which HTkA and HTkB assembled into nucleosomes on *T. kodakarensis* genomic DNA, both in vivo and in vitro,

relative to translation start codons. This is a reasonable proxy for transcription initiation sites as most archaeal promoters are located within short intergenic regions and so are close to a start codon. As the results revealed, on a genome wide basis, archaeal nucleosomes are excluded from



**Figure 6** Archaeal and eukaryotic nucleosomes are positioned by conserved sequences. The frequencies of occurrence of AA/AT/TA/TT and CC/GG/GC/CG dinucleotides, at each position relative to the center of (a) eukaryotic nucleosomes assembled by chicken histones on *T. kodakarensis* genomic DNA, and (b) of archaeal nucleosomes assembled by *T. kodakarensis* histones on yeast genomic DNA. (c) The ratio of the presence and absence of all pentamers in the *T. kodakarensis* DNA protected from MN digestion by assembly into eukaryotic nucleosomes, and (d) in yeast DNA assembled into archaeal nucleosomes. As indicated, all pentamers that contained only G and/or C were preferentially incorporated into both eukaryotic and archaeal nucleosomes (blue regions) whereas all pentamers that contained only A and/or T (red region) were preferentially excluded nucleosomes.



the DNA immediately upstream of open reading frames, and these nucleosome depleted regions (NDRs) are flanked by regions with above average nucleosome occupancy (Figure 4b). This is very similar to the pattern of nucleosome occupancy established in promoter regions and around the downstream transcription start sites in the yeast genome [1,3,26,56]. The conservation of this nucleosome organization argues strongly for a participatory role in gene expression. Possibly, avoiding nucleosome assembly in the promoter region coupled with the presence of a nucleosome at the transcription start site evolved as a generic system to facilitate pre-transcription complex assembly while preventing immediate transcription initiation. If so, a mechanism would then also be needed to remove the inhibitory nucleosome when transcription should occur. In Eukaryotes, many complexes have evolved that catalyze histone modifications, nucleosome remodeling, repositioning and/or eviction that relieve transcription inhibition [3] but, to date, there is no evidence for archaeal histone modifications nor for archaeal nucleosome remodeling complexes.

Consistent with the presence of an archaeal nucleosome impeding transcription [54], presumably to maximize

rRNA synthesis, archaeal nucleosome assembly was strongly deterred by the rDNA operon sequences, in vivo and in vitro, in both *T. kodakarensis* (Figure 6) and in *M. thermotrophicus* (Additional file 3: Figure S3). To test the prediction that the presence of an archaeal nucleosome in a promoter region was sufficient to prevent downstream transcription, we compared the archaeal nucleosomes profiles upstream of genes whose transcription was known to increase or decrease in response to the absence of HTkA or HTkB [13]. There were substantial differences in these profiles, in both the abundance and positioning of archaeal nucleosomes, when compared with the parental strain with both histones (Figure 7) but there was no consistent correlation. An increase or decrease in transcription did not simply result from the absence or presence of an archaeal nucleosome in a promoter region.

#### Supporting report

The results of a similar and complementary investigation were published [57] while this report was being finalized. Ammar et al. [57] determined the locations of archaeal nucleosomes assembled in vivo in *Haloferax volcanii*,

also a euryarchaeon, but a species with a relatively high 65% G + C content, and one with a single atypical archaeal histone that has two histone folds fused into one polypeptide [9,15,58]. Both positioning investigations established that G/C-rich sequences predominate at the center of an archaeal nucleosome but a 10 bp periodicity of AA/AT/TA/TT dinucleotides was not detected in the DNA incorporated into nucleosomes in *Hlf. volcanii*. In both studies, a NDR flanked by increased nucleosome assembly was documented in intergenic regions but, with a transcriptome available, Ammar et al. [57] were able to define the location of the NDR relative to the sites of transcription initiation rather than, as here, to translation start codons. They did not report confirmatory in vitro positioning studies, but the nucleosome positioning in vivo in *Hlf. volcanii* was, as established here both in vivo and in vitro, almost certainly directed primarily by the *Hlf. volcanii* genome sequence.

## Conclusions

The results reported establish that an archaeal histone and genome sequence are sufficient for positioned archaeal nucleosome assembly. They confirm that the primary sequence motifs known to facilitate and direct histone assembly into eukaryotic nucleosomes [5-8] also direct nucleosome assembly in Archaea and that this positioning mechanism therefore almost certainly originated in a common ancestor of *Archaea* and Eukaryotes. DNA compaction is often described as the primary function of nucleosomes but positioned nucleosomes clearly also participate in regulating eukaryotic gene expression [3,59,60] and nucleosome positioning, as a regulatory mechanism, likely predates nucleosome assembly for DNA compaction and archiving. When compared with eukaryotic genomes, archaeal (and bacterial) genomes are very small, and many different proteins have been described that participate both in gene expression and prokaryotic genome compaction [15]. The first histone may have evolved as a protein that bound preferentially to sequences that encoded amphipathic peptide helices [61] with this sequence-directed binding participating in gene regulation. But, given that the histone fold mechanism of DNA binding results in DNA wrapping, this would have also inherently resulted in DNA compaction. When evaluated in terms of the length of DNA compacted per unit of protein, histone wrapping is very efficient, and it is possible that it was the availability of this mechanism, employed by all Eukaryotes, that allowed the massive genome expansion needed for eukaryotic evolution [62]. It remains to be determined if archaeal histones still function primarily as regulators, as suggested by their depletion in intergenic regions [57], or if genome compaction is their primary function consistent with their observed assembly here throughout the full

length of the *T. kodakarensis* and *M. thermautotrophicus* genomes.

## Methods

### Isolation of genomic DNA and archaeal nucleosomes assembled in vivo

Cells from exponentially growing cultures of *M. thermautotrophicus*, *T. kodakarensis* TS517 ( $\Delta pyrF$ ;  $\Delta trpE::pyrF$ ;  $\Delta TK0664$ ), LC124 ( $\Delta pyrF$ ;  $\Delta trpE::pyrF$ ;  $\Delta TK0664$ ;  $\Delta TK1413$ ) and LC125 ( $\Delta pyrF$ ;  $\Delta trpE::pyrF$ ;  $\Delta TK0664$ ;  $\Delta TK2289$ ) [13] were harvested by centrifugation, flash frozen and genomic DNA preparations isolated from aliquots of these cells as previously described [36,40]. The remainder were ruptured by grinding in frozen micrococcal nuclease (MN) buffer [50 mM Tris (pH 8), 1 mM  $CaCl_2$ , 100 mM NaCl], and the lysates allowed to thaw at 4°C. Aliquots were incubated with MN (1 U/ $\mu$ l) at 37°C, and the nuclease digestion then terminated, after increasing periods of digestion, by addition of 250 mM EDTA, 1% SDS, 200 mM NaCl. Following incubation with RNase A (10 mg/ml) for 60 min at 42°C, the DNA molecules that remained were purified by phenol:chloroform extraction, concentrated by ethanol precipitation, and separated by electrophoresis through 3.5% NuSieve agarose gels (Fisher Molecular Biology, Trevose, PA) or 6% polyacrylamide gels. Gel fragments that contained DNA molecules with ~60 bp lengths were excised, crushed and the DNA molecules eluted by incubation overnight at 37°C in 300 mM sodium acetate, 1 mM EDTA (pH 8), 0.1% SDS. The DNA molecules were concentrated by ethanol precipitation, and prepared for sequencing (see below).

### Archaeal histone gene cloning, expression and purification of recombinant HTkA and HTkB

The genes, TK1413 and TK2289, that encode HTkA and HTkB respectively in *T. kodakarensis* TS517 [34], were PCR-amplified and cloned into plasmid pQE-80 (Qiagen, Valencia, CA) generating plasmids pTS600 (TK1413) and pTS601 (TK2289) that were transformed into *Escherichia coli* Rosetta 2 (EMD-Millipore, Billerica, MA). Cultures of the transformants were grown to the late exponential phase in LB medium that contained 100  $\mu$ g ampicillin/ml and 30  $\mu$ g chloramphenicol/ml at 37°C, and recombinant HTkA or HTkB synthesis was then induced by adding isopropyl  $\beta$ -D-1-thiogalactopyranoside (500  $\mu$ M final concentration) and continued incubation for 3 h at 37°C. The cells were harvested by centrifugation, resuspended (0.33 g wet cell pellet/ml) in 25 mM Tris-HCl (pH 7), 0.1 mM EDTA, 50 mM NaCl, lysozyme (100  $\mu$ g/ml) added and the mixtures held ice for 30 min. Phenylmethanesulfonyl fluoride (Sigma, St. Louis, MO) was added (100  $\mu$ g/ml) and cells were ruptured by repeated passage through a French press. The lysates were clarified by centrifugation at 4°C (60,000 g,

20 min), MgCl<sub>2</sub> (5 mM) and DNase I (40 µg/ml) added, the mixtures incubated for 1 h at 37°C and then at 85°C for 20 min. Following further centrifugation (60,000 g, 30 min, 4°C), the supernatants generated were loaded onto 5 ml Hi-Trap heparin columns (GE Healthcare; Pataskala, OH). Recombinant HTkA and HTkB were eluted by passage of 10 column volumes of linear 50 to 500 mM, and 200 to 700 mM gradients of NaCl, respectively, dissolved in 25 mM Tris-HCl (pH 7). The eluate fractions that contained HTkA or HTkB were identified by Commassie-staining of the proteins in samples of the fractions separated by electrophoresis through 22% (w/v) denaturing polyacrylamide gels. These fractions were combined and the protein solution concentrated (final volume of ~0.5 ml) by centrifugation through a pre-rinsed Vivaspin 6 centrifugal concentrator (5 K molecular weight cut off; Sartorius AG, Bohemia, NY). The solutions were adjusted to contain 600 mM NaCl in 25 mM Tris-HCl (pH 7) and then passaged through Sephacryl S-100 HR 16/40 column (GE Healthcare) at a flow rate of 0.5 ml/min. Fractions that contained HTkA or HTkB, identified by Commassie-blue staining after electrophoresis of aliquots through 22% denaturing polyacrylamide gels, were pooled and concentrated (final volume of ~2 ml) by centrifugation again through pre-rinsed Vivaspin 6 centrifugal concentrators (5 K molecular weight cut off). These proteins solution, >99% purified archaeal histone, were dialyzed against in 25 mM Tris-HCl (pH 7), 500 mM NaCl, 50% (v/v) glycerol, and stored at -20°C.

#### Purification of eukaryotic histones

Chicken histone octamers were purified from erythrocytes by salt extraction and by hydroxyapatite column chromatography as previously described [63].

#### Archaeal and eukaryotic nucleosome assembly *in vitro*

Eukaryotic nucleosomes were assembled *in vitro* by salt dialysis in 200 µl reaction mixtures that contained 50 µg of genomic DNA and 30 µg of chicken histone octamers [6]. Archaeal nucleosomes were reconstituted by mixing 50 µg of genomic DNA with 30 µg archaeal histone tetramers. The complexes formed were dialyzed into MN digestion buffer, and aliquots containing ~2.5 µg of DNA were incubated with 0.1 U MN/ µl for 5 min at 37°C. The MN digestions were stopped by addition of 125 mM EDTA, 200 mM NaCl, and the DNA molecules remaining were isolated by phenol:chloroform extraction, concentrated by ethanol precipitation and separated by electrophoresis through 6% polyacrylamide or 3.5% NuSieve agarose gels. Gel fragments that contained the ~60 bp, or ~147 bp, DNA molecules protected from MN digestion by incorporation into archaeal or eukaryotic nucleosomes, respectively, were excised and the DNA

molecules extracted, purified and prepared for DNA sequencing as described above.

#### ABI SOLiD sequencing of DNA fragments

The ends of the ~60 bp and ~147 bp DNA fragments were repaired and 5'-phosphorylated by incubation in DNATerminator end-repair kits, as recommended by the manufacturer (Lucigen Corp., Middleton, WI). SOLiD adapters were ligated and the DNA molecules PCR amplified (very low cycle number) and sequenced by using the Applied Biosystems protocol for SOLiD fragment paired-end sequencing [64]. Sequencing generated from 2 to 12 million unique reads which, depending on the experiment, equated to 60- to 800-fold coverage per 60 bp or 147 bp nucleosome footprint.

#### Analysis of DNA reads generated by pair-end sequencing

We first selected reads of length 55–65 bp (nucleosome of 60 bp lengths) to construct the center-weighted nucleosome occupancy scores. If a read length was odd, a Gaussian weight of  $\exp(-0.5 * (d/10)^2)$  was assigned to a position  $d$  bp away from the center of the read for  $d \leq 25$ . If a read length was even, then positions  $i - 1$  and  $i$  were treated as the possible nucleosome centers. For example, for a 60 nucleotide sequence  $i = 31$ , and so the two potential centers were at positions 30 and 31. Each center in an even read was, in turn, assigned a weight of  $0.5 * \exp(-0.5 * (d/10)^2)$  for a position  $d$  bp away from the center and the values for both positions were then divided by 2. The center-weighted occupancy score for any given position was defined as the aggregation of the weighted scores from all reads. We identified well-defined peaks on the reads occupancy-curve as putative nucleosome centers by controlling the peak height and steepness simultaneously. To generate AA/AT/TA/TT frequency plots, after defining the nucleosome center positions based on the peaks of center-weighted occupancy score, dinucleotide frequency scores were computed as described by Segal *et al.* [5]. We searched for a sequencing tag of length 60 bp nearest to the peak position in the +/-5 bp region. If no such read existed, we further searched for reads of lengths 61, 59, 62 and 58 bp sequentially within +/- 5 bp region of the peak until the first read was identified. The center of the identified read was treated as the nucleosome center to generate the AA/TT/TA/AT frequency plot. If no such read was identified in the +/-5 bp region, the peak position was treated as the true nucleosome center to generate the alignment. For paired-end MNase sequencing data for 147 bp long nucleosomes, read lengths of 137–157 bp were used. We followed a similar approach as described above and also employed by Brogaard *et al.* [21,65] to identify the nucleosome centers.

### Analyses of the DNA reads generated by single-end sequencing

For the single-end reads with known start position on the Watson strand, their end positions are unknown. However, since the DNA inserts are mapping nucleosomes, their length must be subject to the constraint of being around one nucleosome repeat length. Thus, if we observe a single-end read on the Watson strand at position  $i$ , we could practically assume that its end position should be within a region, say  $[i + a, i + b]$ , and follows some distribution. For practical purpose, we let  $a = 51, b = 68$ . We further assumed that the start and end positions of the DNA inserts are independently distributed around the two edges of the nucleosome they map. Let  $c_{i+51}, \dots, c_{i+68}$  be the Crick strand tag numbers in this region. Then the rela-

tive frequency defined as  $c_{i+k} / \sum_{j=51}^{68} c_{i+j}$  can be used to esti-

mate the probability of a DNA insert ending at position  $i + k$  for  $k = 51, \dots, 68$ . Thus, if we observe  $w_i$  single-end tags at position  $i$  from the Watson strand, then we could regard that we had observed  $w_i$  paired-end tags ending at  $i + k$  for  $k = 51, \dots, 68$  with respective frequency  $w_i c_{i+k} / \sum_{j=51}^{68} c_{i+j}$ .

Likewise, if we observe  $c_i$  single-end tags at position  $i$  from the Crick strand, we would regard that there were  $c_i$  paired-end tags ending at  $i - k$  for  $k = 51, \dots, 68$  with respective frequency  $c_i w_{i-k} / \sum_{j=51}^{68} w_{i-j}$ . By this calculation the observed data with  $\sum_i (w_i + c_i)$  single-end tags are converted approximately to a pseudo data set consisting of  $\sum_i (w_i + c_i)$  paired-end tags. The approach defined above for paired-end data was then used to define the center-weighted reads occupancy score and the nucleosome centers.

### Primary data

The sequences obtained and detailed descriptions of the computational analyses are available [66]. The *M. thermoautotrophicus* and *T. kodakarensis* genome coordinates and RefSeq transcript annotations used were from the *methTher1* [67] and *therkoda1* [42] genome assemblies available on the Archaeal Genome Browser web site [39].

### Additional files

**Additional file 1: Figure S1.** Shows data that document that archaeal nucleosomes assembled by HtkA and HtkB, in vivo and in vitro, contain 5 bp offset helical repeats of AA/AT/TA/TT and CC/GG/GC/CG dinucleotides and preferentially exclude oligo A/T-rich sequences.

**Additional file 2: Figure S2.** Documents the conserved positioning of archaeal and eukaryotic histone assembly into nucleosomes on *Methanothermobacter thermautotrophicus* genomic DNA.

**Additional file 3: Figure S3.** Documents the absence of archaeal nucleosome assembly in vivo and in vitro on the two rDNA operons present in the *Methanothermobacter thermautotrophicus* genome.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

Designed and directed the project: TJS, JNR, YNF-M. Performed experiments: NN, SB, TJS, YNF-M. Analyzed the data: LX, WJ-P, TJS, JNR, YNF-M. Wrote the paper: NN, SB, WJ-P, TJS, JNR, YNF-M. All authors read and approved the final manuscript.

### Acknowledgments

We gratefully acknowledge the life and achievements of Jonathan Widom. He was a mentor, colleague and friend who played a major role in the initial stages of this project. Sadly, he passed away before its completion. We thank the staff of the Northwestern University Genomic Core for all the DNA sequencing. This research was supported by NIH grants GM075313 (WJ-P), GM098176 (JNR; TJS), GM100329 (TJS) and 2P20 RR020171 (YNF-M).

### Author details

<sup>1</sup>Department of Molecular and Cellular Biochemistry, College of Medicine, University of Kentucky, Lexington, KY 40536, USA. <sup>2</sup>Department of Statistics, Northwestern University, Chicago, IL 60208, USA. <sup>3</sup>Department of Molecular Biosciences, Northwestern University, Evanston, IL 60208, USA. <sup>4</sup>Department of Microbiology, Ohio State University, Columbus, OH 43210, USA. <sup>5</sup>Center for RNA Biology, Ohio State University, Columbus, OH 43210, USA.

Received: 19 February 2013 Accepted: 31 May 2013

Published: 10 June 2013

### References

1. Field Y, Kaplan N, Fondufe-Mittendorf Y, Moore IK, Sharon E, Lubling Y, Widom J, Segal E: **Distinct modes of regulation by chromatin encoded through nucleosome positioning signals.** *PLoS Comput Biol* 2008, **4**:e1000216.
2. Shivaswamy S, Bhinge A, Zhao Y, Jones S, Hirst M, Iyer VR: **Dynamic remodeling of individual nucleosomes across a eukaryotic genome in response to transcriptional perturbation.** *PLoS Biol* 2008, **6**:e65.
3. Jiang C, Pugh BF: **Nucleosome positioning and gene regulation: advances through genomics.** *Nature Rev Genet* 2009, **10**:161–172.
4. Guan Y, Yao V, Tsui K, Gebbia M, Dunham MJ, Nislow C, Troyanskaya OG: **Nucleosome-coupled expression differences in closely-related species.** *BMC Genomics* 2011, **12**:466.
5. Segal E, Fondufe-Mittendorf Y, Chen L, Thastrom A, Field Y, Moore IK, Wang JP, Widom J: **A genomic code for nucleosome positioning.** *Nature* 2006, **442**:772–778.
6. Kaplan N, Moore IK, Fondufe-Mittendorf Y, Gossett AJ, Tillo D, Field Y, LeProust EM, Hughes TR, Lieb JD, Widom J, Segal E: **The DNA-encoded nucleosome organization of a eukaryotic genome.** *Nature* 2009, **458**:362–366.
7. Yuan GC, Liu JS: **Genomic sequence is highly predictive of local nucleosome depletion.** *PLoS Comput Biol* 2008, **4**:e13.
8. Rapoport AE, Frenkel ZM, Trifonov EN: **Nucleosome positioning pattern derived from oligonucleotide compositions of genomic sequences.** *J Biomol Struct Dynam* 2011, **28**:567–574.
9. Sandman K, Reeve JN: **Archaeal histones and the origin of the histone fold.** *Curr Opin Microbiol* 2006, **9**:520–525.
10. Cubonova L, Sandman K, Hallam SJ, Delong EF, Reeve JN: **Histones in Crenarchaea.** *J Bacteriol* 2005, **187**:5482–5485.
11. Friedrich-Jahn U, Aigner J, Langst G, Reeve JN, Huber H: **Nanoarchaeal origin of histone H3?** *J Bacteriol* 2009, **191**:1092–1096.
12. Bailey KA, Pereira SL, Widom J, Reeve JN: **Archaeal histone selection of nucleosome positioning sequences and the prokaryotic origin of histone-dependent genome evolution.** *J Mol Biol* 2000, **303**:25–34.
13. Cubonova L, Katano M, Kanai T, Atomi H, Reeve JN, Santangelo TJ: **An archaeal histone is required for transformation of Thermococcus kodakarensis.** *J Bacteriol* 2012, **194**:6864–6874.

14. Sandman K, Krzycki JA, Dobrinski B, Lurz R, Reeve JN: **Hmf, a DNA-binding protein isolated from the hyperthermophilic archaeon *Methanothermus fervidus*, is most closely related to histones.** *Proc Natl Acad Sci USA* 1990, **87**:5788–5791.
15. Samson R, Reeve JN: **DNA binding proteins and chromatin.** In *Archaea: Molecular and Cellular Biology*. Edited by Cavicchioli R. Washington, DC: ASM Press; 2007:110–119.
16. Pereira SL, Grayling RA, Lurz R, Reeve JN: **Archaeal nucleosomes.** *Proc Natl Acad Sci USA* 1997, **94**:12633–12637.
17. Bailey KA, Marc F, Sandman K, Reeve JN: **Both DNA and histone fold sequences contribute to archaeal nucleosome stability.** *J Biol Chem* 2002, **277**:9293–9301.
18. Marc F, Sandman K, Lurz R, Reeve JN: **Archaeal histone tetramerization determines DNA affinity and the direction of DNA supercoiling.** *J Biol Chem* 2002, **277**:30879–30886.
19. Kaplan N, Moore I, Fondufe-Mittendorf Y, Gossett AJ, Tillo D, Field Y, Hughes TR, Lieb JD, Widom J, Segal E: **Nucleosome sequence preferences influence in vivo nucleosome organization.** *Nat Struct Mol Biol* 2010, **17**:918–920. author reply 920–922.
20. Gaffney DJ, McVicker G, Pai AA, Fondufe-Mittendorf YN, Lewellen N, Michelini K, Widom J, Gilad Y, Pritchard JK: **Controls of nucleosome positioning in the human genome.** *PLoS Genet* 2012, **8**:e1003036.
21. Brogaard K, Xi L, Wang JP, Widom J: **A map of nucleosome positions in yeast at base-pair resolution.** *Nature* 2012, **486**:496–501.
22. Lee W, Tillo D, Bray N, Morse RH, Davis RW, Hughes TR, Nislow C: **A high-resolution atlas of nucleosome occupancy in yeast.** *Nat Genet* 2007, **39**:1235–1244.
23. Segal E, Widom J: **Poly(dA:dT) tracts: major determinants of nucleosome organization.** *Curr Opin Struct Biol* 2009, **19**:65–71.
24. Iyer V, Struhl K: **Poly(dA:dT), a ubiquitous promoter element that stimulates transcription via its intrinsic DNA structure.** *EMBO J* 1995, **14**:2570–2579.
25. Lascaris RF, Groot E, Hoen PB, Mager WH, Planta RJ: **Different roles for abf1p and a T-rich promoter element in nucleosome organization of the yeast RPS28A gene.** *Nucl Acids Res* 2000, **28**:1390–1396.
26. Morohashi N, Yamamoto Y, Kuwana S, Morita W, Shindo H, Mitchell AP, Shimizu M: **Effect of sequence-directed nucleosome disruption on cell-type-specific repression by alpha2/Mcm1 in the yeast genome.** *Euk Cell* 2006, **5**:1925–1933.
27. Bao Y, White CL, Luger K: **Nucleosome core particles containing a poly(dA:dT) sequence element exhibit a locally distorted DNA structure.** *J Mol Biol* 2006, **361**:617–624.
28. Struhl K: **Naturally occurring poly(dA-dT) sequences are upstream promoter elements for constitutive transcription in yeast.** *Proc Natl Acad Sci USA* 1985, **82**:8419–8423.
29. Nelson HC, Finch JT, Luisi BF, Klug A: **The structure of an oligo(dA)-oligo(dT) tract and its biological implications.** *Nature* 1987, **330**:221–226.
30. Suter B, Schnappauf G, Thoma F: **Poly(dA:dT) sequences exist as rigid DNA structures in nucleosome-free yeast promoters in vivo.** *Nucl Acids Res* 2000, **28**:4083–4089.
31. Anderson JD, Widom J: **Poly(dA-dT) promoter elements increase the equilibrium accessibility of nucleosomal DNA target sites.** *Mol Cell Biol* 2001, **21**:3830–3839.
32. Sekinger EA, Moqtaderi Z, Struhl K: **Intrinsic histone-DNA interactions and low nucleosome density are important for preferential accessibility of promoter regions in yeast.** *Mol Cell* 2005, **18**:735–748.
33. Ioshikhes IP, Albert I, Zanton SJ, Pugh BF: **Nucleosome positions predicted through comparative genomics.** *Nat Genet* 2006, **38**:1210–1215.
34. Fukui T, Atomi H, Kanai T, Matsumi R, Fujiwara S, Imanaka T: **Complete genome sequence of the hyperthermophilic archaeon *Thermococcus kodakaraensis* KOD1 and comparison with *Pyrococcus* genomes.** *Genome Res* 2005, **15**:352–363.
35. Grohmann D, Werner F: **Recent advances in the understanding of archaeal transcription.** *Curr Opin Microbiol* 2011, **14**:328–334.
36. Santangelo TJ, Reeve JN: **Archaeal RNA polymerase is sensitive to intrinsic termination directed by transcribed and remote sequences.** *J Mol Biol* 2006, **355**:196–210.
37. Nishida H, Kondo S, Matsumoto T, Suzuki Y, Yoshikawa H, Taylor TD, Sugiyama J: **Characteristics of nucleosomes and linker DNA regions on the genome of the basidiomycete *Mixia osmundae* revealed by mono- and dinucleosome mapping.** *Open Biol* 2012, **2**:120043.
38. Peckham HE, Thurman RE, Fu Y, Stamatoyannopoulos JA, Noble WS, Struhl K, Weng Z: **Nucleosome positioning signals in genomic DNA.** *Genome Res* 2007, **17**:1170–1177.
39. *Archaeal Genome Browser*. <http://archaea.ucsc.edu>.
40. Santangelo TJ, Cubonova L, Matsumi R, Atomi H, Imanaka T, Reeve JN: **Polarity in archaeal operon transcription in *Thermococcus kodakaraensis*.** *J Bacteriol* 2008, **190**:2244–2248.
41. Smith DR, Doucette-Stamm LA, Deloughery C, Lee H, Dubois J, Aldredge T, Bashirzadeh R, Blakely D, Cook R, Gilbert K, Harrison D, Hoang L, Keagle P, Lum W, Pothier B, Qiu D, Spadafora R, Vicaire R, Wang Y, Wierzbowski J, Gibson R, Jiwani N, Caruso A, Bush D, Reeve JN: **Complete genome sequence of *Methanobacterium thermoautotrophicum* ΔH: functional analysis and comparative genomics.** *J Bacteriol* 1997, **179**:7135–7155.
42. *UCSC Genome Browser on Thermococcus kodakaraensis May 2004 Assembly (therKoda1)*. <http://archaea.ucsc.edu/lists/therKoda1/refSeq-list.html>.
43. Gupta S, Dennis J, Thurman RE, Kingston R, Stamatoyannopoulos JA, Noble WS: **Predicting human nucleosome occupancy from primary sequence.** *PLoS Comput Biol* 2008, **4**:e1000134.
44. Zhang Y, Moqtaderi Z, Rattner BP, Euskirchen G, Snyder M, Kadonaga JT, Liu XS, Struhl K: **Intrinsic histone-DNA interactions are not the major determinant of nucleosome positions in vivo.** *Nat Struct Mol Biol* 2009, **16**:847–852.
45. Ioshikhes I, Hosid S, Pugh BF: **Variety of genomic DNA patterns for nucleosome positioning.** *Genome Res* 2011, **21**:1863–1871.
46. Tillo D, Hughes TR: **G + C content dominates intrinsic nucleosome occupancy.** *BMC Bioinforma* 2009, **10**:442.
47. Visnapuu ML, Greene EC: **Single-molecule imaging of DNA curtains reveals intrinsic energy landscapes for nucleosome deposition.** *Nat Struct Mol Biol* 2009, **16**:1056–1062.
48. Lankas F, Sponer J, Langowski J, Cheatham TE: **DNA basepair step deformability inferred from molecular dynamics simulations.** *Biophys J* 2003, **85**:2872–2883.
49. Luger K, Mäder AW, Richmond RK, Sargent DF, Richmond TJ: **Crystal structure of the nucleosome core particle at 2.8 Å resolution.** *Nature* 1997, **389**:251–260.
50. Luger K, Mader A, Sargent DF, Richmond TJ: **The atomic structure of the nucleosome core particle.** *J Biomol Struct Dyn* 2000, **17**(Suppl 1):185–188.
51. Decanniere K, Babu AM, Sandman K, Reeve JN, Heinemann U: **Crystal structures of recombinant histones HmfA and HmfB from the hyperthermophilic archaeon *Methanothermus fervidus*.** *J Mol Biol* 2000, **303**:35–47.
52. Soares DJ, Sandman K, Reeve JN: **Mutational analysis of archaeal histone-DNA interactions.** *J Mol Biol* 2000, **297**:39–47.
53. French SL, Santangelo TJ, Beyer AL, Reeve JN: **Transcription and translation are coupled in Archaea.** *Mol Biol Evol* 2007, **24**:893–895.
54. Wilkinson SP, Ouhammouch M, Geiduschek EP: **Transcriptional activation in the context of repression mediated by archaeal histones.** *Proc Natl Acad Sci USA* 2010, **107**:6777–6781.
55. Xie Y, Reeve JN: **Transcription by an archaeal RNA polymerase is slowed but not blocked by an archaeal nucleosome.** *J Bacteriol* 2004, **186**:3492–3498.
56. Yuan GC, Liu YJ, Dion MF, Slack MD, Wu LF, Altschuler SJ, Rando OJ: **Genome-scale identification of nucleosome positions in *S. cerevisiae*.** *Science* 2005, **309**:626–630.
57. Ammar R, Torti D, Tsui K, Gebbia M, Durbin T, Bader GD, Giaever G, Nislow C: **Chromatin is an ancient innovation conserved between Archaea and Eukarya.** *Life* 2012, **1**:e00078.
58. Hartman AL, Norais C, Badger JH, Delmas S, Haldenby S, Madupu R, Robinson J, Khouri H, Ren Q, Lowe TM, Maupin-Furlow J, Pohlschroder M, Daniels C, Pfeiffer F, Allers T, Eisen JA: **The complete genome sequence of *Haloflex volcanii* DS2, a model archaeon.** *PLoS One* 2010, **5**:e9605.
59. Mavrich TN, Jiang C, Ioshikhes IP, Li X, Venters BJ, Zanton SJ, Tomsho LP, Qi J, Glaser RL, Schuster SC, Gilmour DS, Albert I, Pugh BF: **Nucleosome organization in the *Drosophila* genome.** *Nature* 2008, **453**:358–362.
60. Field Y, Fondufe-Mittendorf Y, Moore IK, Mieczkowski P, Kaplan N, Lubling Y, Lieb JD, Widom J, Segal E: **Gene expression divergence in yeast is coupled to evolution of DNA-encoded nucleosome organization.** *Nat Genet* 2009, **41**:438–445.
61. Rapoport AE, Trivonof EF: **“Anticipated” nucleosome positioning pattern in prokaryotes.** *Gene* 2011, **488**:41–45.

62. Reeve JN: Archaeal chromatin and transcription. *Mol Microbiol* 2003, **48**:587–598.
63. Feng HP, Scherl DS, Widom J: Lifetime of the histone octamer studied by continuous-flow quasielastic light scattering: test of a model for nucleosome transcription. *Biochemistry* 1993, **32**:7824–7831.
64. *Applied Biosystems protocol for SOLiD sequencing.*  
[http://tools.invitrogen.com/content/sfs/manuals/SOLiD4\\_Library\\_Preparation\\_man.pdf](http://tools.invitrogen.com/content/sfs/manuals/SOLiD4_Library_Preparation_man.pdf).
65. Brogaard KR, Xi L, Wang JP, Widom J: A chemical approach to mapping nucleosomes at base pair resolution in yeast. *Meth Enzymol* 2012, **513**:315–334.
66. *Genome-database of Methanobacterium thermoautotrophicum and Thermococcus kodakarensis sequenced nucleosomal reads 2012.*  
<http://ncode.northwestern.edu>.
67. *UCSC Genome Browser on Methanobacterium thermoautotrophicum Aug 1997 Assembly (methTher1).* <http://archaea.ucsc.edu/lists/methTher1/refSeq-list.html>.

doi:10.1186/1471-2164-14-391

**Cite this article as:** Nalabothula *et al.*: Archaeal nucleosome positioning in vivo and in vitro is directed by primary sequence motifs. *BMC Genomics* 2013 **14**:391.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

