



University of Kentucky
UKnowledge

Center for Health Services Research Faculty
Publications

Center for Health Services Research

8-2017

Substance Use Disorders, Violence, Mental Health, and HIV: Differentiating a Syndemic Factor by Gender and Sexuality

Kiyomi Tsuyuki

University of California - San Diego

Eileen V. Pitpitan

University of California - San Diego

Maria A. Levi-Minzi

Nova Southeastern University

Lianne A. Urada

San Diego State University

Steven P. Kurtz

Nova Southeastern University

Follow this and additional works at: https://uknowledge.uky.edu/chsr_facpub

See next page for additional authors



Part of the [Feminist, Gender, and Sexuality Studies Commons](#), [Health Services Research Commons](#),
and the [Substance Abuse and Addiction Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Tsuyuki, Kiyomi; Pitpitan, Eileen V.; Levi-Minzi, Maria A.; Urada, Lianne A.; Kurtz, Steven P.; Stockman, Jamila K.; and Surratt, Hilary L., "Substance Use Disorders, Violence, Mental Health, and HIV: Differentiating a Syndemic Factor by Gender and Sexuality" (2017). *Center for Health Services Research Faculty Publications*. 6.

https://uknowledge.uky.edu/chsr_facpub/6

This Article is brought to you for free and open access by the Center for Health Services Research at UKnowledge. It has been accepted for inclusion in Center for Health Services Research Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Substance Use Disorders, Violence, Mental Health, and HIV: Differentiating a Syndemic Factor by Gender and Sexuality

Digital Object Identifier (DOI)

<https://doi.org/10.1007/s10461-017-1841-3>

Notes/Citation Information

Published in *AIDS and Behavior*, v. 21, issue 8, p. 2270-2282.

© Springer Science+Business Media, LLC 2017

The copyright holder has granted the permission for posting the article here.

This is a post-peer-review, pre-copyedit version of an article published in *AIDS and Behavior*. The final authenticated version is available online at: <https://doi.org/10.1007/s10461-017-1841-3>.

Authors

Kiyomi Tsuyuki, Eileen V. Pitpitan, Maria A. Levi-Minzi, Lianne A. Urada, Steven P. Kurtz, Jamila K. Stockman, and Hilary L. Surratt



Published in final edited form as:

AIDS Behav. 2017 August ; 21(8): 2270–2282. doi:10.1007/s10461-017-1841-3.

Substance Use Disorders, Violence, Mental Health, and HIV: Differentiating a Syndemic Factor by Gender and Sexuality

Kiyomi Tsuyuki¹, Eileen V. Pitpitan¹, Maria A. Levi-Minzi², Lianne A. Urada³, Steve Kurtz²,
Jamila K. Stockman¹, and Hilary L. Surratt^{2,4}

¹Division of Global Public Health, School of Medicine, University of California, San Diego (UCSD)

²Center for Applied Research on Substance Use and Health Disparities, Nova Southeastern University, Miami FL

³School of Social Work, San Diego State University (SDSU)

⁴Center for Health Services Research, University of Kentucky, Lexington KY

Abstract

This paper measures syndemic substance use disorder, violence, and mental health and compares the syndemic among HIV-infected heterosexual men, heterosexual women, and men who have sex with men (MSM). Data were from a sample of vulnerable, substance-using, HIV-infected people in South Florida between 2010 and 2012 (n=481). We used confirmatory factor analysis to measure a syndemic latent variable and applied a measurement invariance model to identify group differences in the data structure of syndemic co-morbidities among heterosexual men, heterosexual women, and MSM. We found that variables used to measure the syndemic fit each sub-group, supporting that substance use disorder, violence, and mental health coincide in HIV-infected individuals. Heterosexual men and MSM demonstrated similar syndemic latent variable factor loadings, but significantly different item intercepts, indicating that heterosexual men had larger mean values on substance use disorder, anxiety, and depression than MSM. Heterosexual men and heterosexual women demonstrated significantly different syndemic variable factor loadings, indicating that anxiety and depression contribute more (and substance use contributes less) to the syndemic in heterosexual men compared to heterosexual women. MSM and heterosexual women demonstrated similar syndemic latent variable factor loadings and intercepts, but had significantly different factor residual variances indicating more variance in violent victimization and depression for MSM and more variance in stress for heterosexual women than what is captured by the observed syndemic indicators. Furthermore, heterosexual women had a

Corresponding Author: Kiyomi Tsuyuki, Ph.D., M.P.H., Assistant Professor, Division of Global Public Health, School of Medicine, Center on Gender Equity and Health, University of California, San Diego (UCSD), 9500 Gilman Drive, MC 0507, La Jolla, CA 92093-0507, Tel: (858)822-6468, ktsuyuki@ucsd.edu.

Conflict of Interest: No conflict exists. Kiyomi Tsuyuki declares that she has no conflict of interest. Eileen V. Pitpitan declares that she has no conflict of interest. Maria A. Levi-Minzi declares that she has no conflict of interest. Lianne A. Urada declares that she has no conflict of interest. Steve Kurtz declares that he has no conflict of interest. Jamila K. Stockman declares that she has no conflict of interest. Hilary L. Surratt declares that she has no conflict of interest.

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the Institutional Review Boards of the University of Delaware (predecessor institution) and Nova Southeastern University and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. An NIH Certificate of Confidentiality was acquired and informed consent was obtained from every participant prior to interview.

larger syndemic factor mean than MSM, indicating that the syndemic burden is greater among heterosexual women than MSM. Our findings support that measurement invariance can elucidate syndemic differences to tailor interventions to sub-group needs.

INTRODUCTION

Antiretroviral therapy (ART) has lengthened the lifespan and reduced HIV transmission risk among people living with HIV (PLWH) (1–6). However, individuals with socio-structural barriers – such as poverty, unstable housing, and comorbidities like substance use and psychiatric disorders (7–12) – have difficulty adequately engaging in HIV care and achieving viral suppression (1, 13). Individual-level efforts to engage and retain PLWH in HIV care have limited impact on outcomes as they overlook critical fundamental causes (e.g., socio-structural barriers, life course experiences) (14, 15).

Syndemic theory links adverse socio-structural conditions to co-occurring and synergistic health epidemics that systemically and disproportionately burden disadvantaged populations (16). Syndemic theory highlights the socio-structural determinants of health disparities by shifting focus away from individuals and elucidating critical targets for socio-structural intervention. Most applications of syndemic theory to HIV research focus on factors associated with HIV risk (17–25). However, a growing literature points to syndemic co-morbidities (e.g., depression, drug and alcohol use disorders, violence) as important predictors of HIV transmission risk and HIV treatment outcomes among PLWH (12, 26–31). Recent research suggests HIV risk reduction interventions are most effective for individuals with syndemic risk (32).

Despite recent advances in syndemic research, two key methodological issues remain with syndemic theory applications. First is the uniform application of syndemic theory to qualitatively different populations, including HIV-negative men who have sex with men (MSM) (17–20), Hispanic women (21); adolescents (22); transgender women (23); patients receiving clinic-based STI care (24); female sex workers (25); and members of these subgroups who are living with HIV (12, 26–31). Although syndemic mental health, substance use disorders, violence, and sexually transmitted infections (STIs, including HIV) are related and affect the aforementioned populations as a whole, socio-structural theories point to potential differences by both gender and sexuality. For example, the Theory of Gender and Power suggests that heterosexual women experience gendered power inequities compared to heterosexual men (33, 34). The Minority Stress Theory suggests that men who have sex with men (MSM) may experience chronic psychological distress from the fear of stigma based on their sexual identity (35). Furthermore, enhanced socio-structural inequities and resultant chronic stress can affect heterosexual women and MSM differently.

Research suggests that differences in syndemic co-morbidities exist among heterosexual women, MSM, and heterosexual men. Heterosexual women experience significantly higher rates of depression (36, 37), anxiety (38), and intimate partner violence (39), and lower rates of substance use disorders (40, 41), than heterosexual men. MSM experience higher rates of depression (29, 42), substance use disorders (29, 43), and sexual risk behavior (44) than heterosexual men. Likewise, racial and ethnic minority heterosexual men and MSM may

encounter gendered experiences of discrimination and violence (e.g., police, gangs, fights) and can struggle with conforming to masculine ideologies depending on their social context (45–47). However, no study to date has empirically tested for differences in the syndemic among these groups of PLWH to tailor risk reduction or health interventions to sub-group needs.

A second methodological issue is the lack of researcher consistency in measuring syndemics. Whereas some measure syndemics as a summative count of co-morbidity prevalence (26, 48), others fit syndemic co-morbidities to one latent variable (49, 50). Summative approaches are simple and generate easy to interpret outcomes (51), but they have several shortcomings that are addressed by the latent variable approach. Summative approaches: 1) do not account for (potential) synergistic effects of the syndemic, ignoring the covariance structure of syndemic data and throwing away information, 2) assume that each co-morbidity contributes equally to the syndemic, and 3) assume that syndemic co-morbidities are the same and combine equally in all population sub-groups (52). Latent variable approaches capture the full structure of syndemic data, and generate a variable that can be interpreted in light of the syndemic theory (15). Furthermore, research applying the latent variable approach has found that the same co-morbidities comprise the syndemic for both MSM and heterosexual men, but that each co-morbidity had significantly different weight (e.g., greater factor loadings, stronger association) on the syndemic between groups (e.g., risky sex was lower and violent victimization was greater for MSM versus heterosexual men) (50).

This paper empirically explores the similarities and differences in syndemic substance use disorder, violent victimization, and mental health among a sample of substance-using heterosexual men, heterosexual women, and MSM living with HIV in South Florida. Although MSM contact is responsible for the bulk of HIV infections among PLWH in South Florida (68% in Broward County and 69% in Miami Dade County), heterosexual contact comprises a sizeable transmission risk category for men (21% and 19%, respectively) and women (85% to 86%, respectively) (53). This study is among the first to compare the syndemic among three sub-groups of PLWH using structural equation modeling (SEM) confirmatory factor analysis. Figure 1 shows how we operationalize the syndemic construct. We hypothesize that substance use disorder, violent victimization, and poor mental health will adequately measure the syndemic in all groups. In addition, we hypothesize that each co-morbidity among heterosexual women and MSM (compared to heterosexual men) will have greater item loadings (i.e., stronger association) on the underlying syndemic latent variable.

METHODS

Data

Data were collected from a sample of vulnerable, substance-using PLWH in South Florida between 2010 and 2012 (n=503) – details described elsewhere (10, 54). Recruitment was guided by targeted sampling – a method used to recruit hard-to-reach populations – in geographic areas with high HIV prevalence and poverty indices (55). Field staff used direct outreach to recruit indigent HIV-positive substance users through systematic distribution of

study cards and flyers in street venues and HIV service organizations within the target areas. Eligibility requirements were: =>18 years old, recent substance use, confirmed HIV-positive, and prescribed ART. Institutional Review Boards of University of Delaware (predecessor institution) and Nova Southeastern University approved study protocols. An NIH Certificate of Confidentiality was obtained. Trained project staff obtained informed consent and then administered face-to-face structured interviews based on the Global Appraisal of Individual Needs (GAIN) instrument (56), which assessed demographics, substance use and dependence (DSM-IV-R), and mental health status. Standardized instruments assessed violent victimization and stress, among other HIV-related factors (e.g., diagnosis and treatment history, attitudes toward providers, stigma, and ARV knowledge and adherence). Participants received a \$30 stipend upon interview completion (averaging one hour long).

Sample

The analytic sample (n=481) included men and women with information on biological sex, HIV sexual risk behavior, sexual preference (i.e., prefers to have sex with only females, only males, or females and males), and past 90 day sexual activity. We prioritized grouping by sexual risk behavior, beginning with sexual activity and gender of partner in the past 90 days (114 men only had sex with women, 147 MSM, 168 women had sex with men). Among those with missing data (n=58), we grouped individuals according to sexual preference (18 men preferred sex with women, 10 men preferred sex with men, and 8 women preferred sex with men). One man preferred sex with both men and women (MSMW), but was coded as MSM due to a lack of power to conduct sub-analyses with MSMW. Among those still missing a sexual risk group, we grouped individuals according to transmission of HIV via sexual contact (1 man contracted HIV via heterosexual contact, 14 women contracted HIV via heterosexual contact, and 6 remained missing). Our final analytic sub-groups included 133 heterosexual men, 158 MSM, and 190 heterosexual women.

Measures

The Global Appraisal of Individual Needs (GAIN) instrument (57) was used to assess substance use disorder, violent victimization, and mental health variables used to develop the syndemic factor. A General Victimization Scale ($\alpha=0.86$) measured *violent victimization severity* (analyzed as a continuous variable). Questions included physical, sexual, and emotional violence, the recency of such events, and current worry about violent victimization (15-item scale: 0-no victimization to 15-recent, severe victimization) (58).

Mental health was treated as continuous variables in our analysis and measured by scales assessing depression (past year), anxiety (past year), and stress (past 90 days). Depressive symptoms (e.g., feeling sad, lonely, or hopeless; feeling tired or having no energy) were assessed with a 9-item summative scale (0-no/minimal to 9-severe depression; $\alpha=0.87$) (58). Anxiety symptoms (e.g., feeling nervous, anxious, or tense; unable to control worries) were assessed with a 12-item summative scale (0-no anxiety to 12-severe anxiety; $\alpha=0.87$) (58). Stress (e.g., go without food, clothing, housing because no money; get divorced or break up with partner) was measured with an adapted Crisis in Family Systems (CRISYS) scale (0-no stress to 8-severe stress; $\alpha=0.57$) (59). The poor internal consistency of the stress instrument does not pose a serious issue as our analytic approach (e.g., measurement invariance

framework) tests a kind of internal consistency within the latent construct rather than within individual instruments (60).

Substance use disorder (past 12 months) from DSM-IV criteria (eleven items measuring drug problem severity, e.g., using more or longer than intended, withdrawal problems; $\alpha=0.86$) was treated as a continuous variable in our analysis (58).

Data Analysis

Heterosexual men, heterosexual women, and MSM were compared on sociodemographic characteristics and syndemic indicator variables. Chi-square statistic was used for categorical variables and the F-statistic from ANOVA was used for variables treated as continuous to detect statistically significant differences between groups. We then created a scatter plot matrix with correlation coefficients to provide a graphic indication of the covariation between the five co-morbidities across groups (Figure 2). Furthermore, we checked for multivariate outliers as the observed covariance structure and estimated CFA model parameters can be heavily influenced by outliers. Multivariate outliers were accessed using the 15th percentile of the chi-square distribution as the defined threshold for outliers using MATA with the `-bacon-` command in STATA (61, 62). We did not detect any influential outliers in our sample. We then tested hypotheses with confirmatory factor analysis (CFA) using maximum likelihood estimation and the `-sem-` command in STATA 13 (63).

We began by examining the factor structure of the syndemic indicator variables in each group using linear CFA to test for configural invariance. Configural invariance is supported if the model fits the data well in all groups and the pattern of salient and significantly non-zero factor loadings is similar across groups (60). Establishing configural invariance (in this case that the measured variables are all indicators of a latent, or underlying, syndemic variable) allows for comparison between multiple groups in SEM, by demonstrating that the syndemic indicator variables make up a syndemic latent variable with suitable model fit across groups. To test for configural invariance, we fit the syndemic latent variable with free parameters to each group. For identification purposes, the factor loadings for violent victimization severity were fixed to 1 and the intercept was fixed to zero to allow for estimating (identifying) the factor mean (60). It is noteworthy that the syndemic latent variable has an arbitrary (unitless) non-negative scale. Model fit was assessed by evaluating one incremental fit index (Comparative Fit Index–CFI) and two absolute fit indices (Root Mean Squared Error Approximation–RMSEA and the Standardized Root Mean Squared Residual–SRMR). We chose cut off values of CFI 0.95, RMSEA<0.06, and SRMR 0.08, as suggested by Hu and Bentler (64).

Once configural invariance was ascertained, we used multiple group CFA to estimate an unconstrained model (Model 1), and to examine metric invariance (Model 2), strong invariance (Model 3), strict measurement invariance (Model 4), and equal factor means (Model 5) in the syndemic latent variable. By estimating a series of nested models that were increasingly restrictive in their assumptions, we compared: 1) MSM to heterosexual men (Table 3), 2) heterosexual women to heterosexual men (Table 4), and 3) MSM to heterosexual women (Table 5). The fit between models was compared using a chi-square

difference test. A non-significant result would indicate that the fit of the constrained model does not differ significantly from the unconstrained model. Once again, for identification purposes, the factor loadings for violent victimization severity were fixed to 1 and the intercept was fixed to zero to allow for estimating (identifying) the syndemic factor mean (60).

In Model 1, we estimated an *unconstrained model* where factor loadings, item intercepts, and error variances were allowed to vary across groups. In this step, we fitted a single factor model to the covariance matrix of syndemic indicators across groups. In Model 2, we constrained factor loadings for syndemic items to be equal across groups to test for *metric invariance*. A non-significant chi-square difference test between Models 2 & 1 indicates that metric invariance has been satisfied. Metric invariance introduces the concept of equal metrics across groups and, if achieved, differences in item scores can be meaningfully compared across groups (60). In Model 3, we further constrained item intercepts to be equal to test for *strong (or scalar) invariance*, which assumes invariance of factor loadings and item intercepts. Strong invariance implies that any group differences in the mean of the observed items are due to differences in the mean of the underlying syndemic construct (60). A non-significant chi-square difference test between Models 3 & 2 indicates that the intercepts are invariant across groups. In Model 4, we further constrained error variances (residual variances) to be equal to test for *strict invariance*, which assumes invariance of factor loadings, item intercepts, and item variation. As a result, the amount of the variance in the syndemic indicators that was not explained by the common factor was constrained to be equal across groups. A non-significant chi-square difference test between Models 4 & 3 indicates that any observed group differences in the observed syndemic indicators was attributable to a difference with respect to the latent syndemic variable that we measured (65). In Model 5, we further constrained latent means to be equal across groups.

RESULTS

Sample Characteristics

Table 1 provides a description of the overall study sample (n=481) by socio-demographic and syndemic variables. In terms of socio-demographic variables, the overall sample averaged 46.13 years old (range=19–71 years; standard deviation=7.75) and averaged living with HIV for 12.95 years (range=0–30 years; standard deviation=7.31). About half of the sample reported having less than a high school education (44%), the majority were of Black race or African-American (67%), reported a monthly income of at least \$501 (68%), and almost one-third tested positive for an STI in the past year (29%). In terms of syndemic indicator variables, the sample averaged a moderate level (7.06) on the violent victimization scale (range 0–15; standard deviation=4.19). The sample had moderate levels of mental health problems with an average of 5.36 depressive symptoms (range 0–9 standard deviation=3.02), 4.68 anxiety symptoms (range 0–12; standard deviation=3.66), and 2.18 on the stress scale (range 0–8; standard deviation=1.85). The sample averaged a high level (6.87) of past year symptoms of substance use disorder (range 0–11; standard deviation=3.23).

Table 1 also reports significant differences in socio-demographic and syndemic variables by analytic sub-groups (i.e., heterosexual men, MSM, and heterosexual women) with heterosexual men as the comparison group for mean comparisons. On average, MSM were significantly younger (mean age=44.85) than heterosexual men (mean age=47.50; $F=8.55$, $p=0.004$) and lived significantly more years with HIV (mean years=14.32) compared to heterosexual men (mean age=47.50; $F=8.55$, $p=0.004$; mean years with HIV=12.49; $F=4.61$, $p=0.032$). A significantly greater proportion of MSM reported having a high school or more education (75%) than heterosexual men (44%) and heterosexual women (48%, $X^2=43.11$, $p<0.001$). Furthermore, a significantly greater proportion of MSM reported being non-Hispanic White (25%), than heterosexual men (4%) and heterosexual women (13%, $X^2=46.12$, $p<0.001$). For syndemic variables, MSM (mean score=7.05; $F=10.07$, $p=0.002$) and heterosexual women (mean score=8.15) reported significantly more violent victimization on average than heterosexual men (mean score=5.53; $F=32.39$, $p<0.001$).

Figure 2 shows the scatter plot matrices and the correlation coefficients for syndemic variables by analytic sub-group (i.e., heterosexual men, MSM, and heterosexual women). The correlations between syndemic variables are broadly similar in MSM compared to heterosexual men, whereas the correlations between syndemic variables are greater in heterosexual women compared to heterosexual men.

Configural Invariance

Configural invariance for the syndemic we modeled was examined by testing a one-factor solution in each of the three sub-groups (Table 2). In this case, configural invariance examines whether the same items form a syndemic factor in each group. The one factor solution fit well for heterosexual men (CFI=1.00, RMSEA=0.00, SRMR=0.03), MSM (CFI=0.98, RMSEA=0.10, SRMR=0.04), and heterosexual women (CFI=0.97, RMSEA=0.11, SRMR=0.04). Furthermore, the pattern of salient and significantly non-zero factor loadings was similar across groups, indicating support for configural invariance among the three groups.

Metric, Strong, and Strict Invariance

Having established configural invariance for the syndemic variable across all three groups, we next tested invariance with two-group CFAs between: 1) heterosexual men and MSM (Table 3), 2) heterosexual men and heterosexual women (Table 4), and 3) MSM and heterosexual women (Table 5).

Heterosexual men vs. MSM—For men (Table 3), we first examined *metric invariance* (Model 2) in the syndemic in which factor loadings were constrained to be equal. The chi-square difference test (comparing the fit of the constrained model - Model 2 - to that of the unconstrained model - Model 1 - where factor loadings were free to vary), indicated the syndemic factor loadings could be constrained across groups of men without significantly reducing model fit, $\chi^2(4)=2.93$, supporting full metric invariance. Second, we examined *strong invariance* (Model 3) in which item intercepts were constrained to be equal. Strong invariance is a necessary condition for the meaningful comparison of the factor means and, if it holds, it means that differences in the means of the indicators are due to differences in

the level of the factor. The chi-square difference test between Models 3 and 2 was significant, $\chi^2(4)=13.43$, indicating that syndemic item intercepts are not equal across heterosexual men and MSM and could not be constrained across groups without significantly reducing model fit. Heterosexual men had significantly larger item intercepts for substance use disorder (intercept=4.63, 95% CI: 3.0, 6.27), anxiety (intercept=-3.97, 95% CI: -6.69, -1.26), and depression (intercept=-2.23, 95% CI: -4.65, 0.19) than MSM (substance use disorder intercept=2.61, 95% CI: 0.76, 4.45; anxiety intercept=-6.00; 95% CI: -8.83, -3.18; depression intercept=-3.02; 95% CI: -5.34, -0.71) (data not shown). These results suggest that the same syndemic latent factor is being measured across heterosexual men and MSM, but that the item means (expected item values when the value of the factors is equal to zero) of substance use disorder, anxiety, and depression is greater for heterosexual men than MSM. We cannot compare latent means because we did not achieve intercept invariance.

Heterosexual men vs. Heterosexual women—A two-group CFA for the syndemic was then conducted with heterosexual men and heterosexual women (Table 4). We first examined *metric invariance* (Model 2) in the syndemic (factor loadings were constrained to be equal). The chi-square difference test – comparing the fit of the constrained model (Model 2) to that of the unconstrained model (Model 1) – indicated that the syndemic factor loadings could not be constrained to be equal across groups without significantly reducing model fit, $\chi^2(4)=13.91$. This finding does not support metric invariance between heterosexual men and heterosexual women, indicating that heterosexual men and heterosexual women have different factor loadings on syndemic latent variable indicators. Upon further analysis of the syndemic latent variable, the factor loadings for heterosexual women are larger for substance use disorder (loading=0.77, 95% CI: 0.57, 0.99) and smaller for anxiety (loading=1.23, 95% CI: 0.99, 1.48) and depression (loading=0.98, 95% CI: 0.79, 1.18) than for heterosexual men (substance use disorder loading=0.46, 95% CI: 0.18, 0.73; anxiety loading=1.53; 95% CI: 1.07, 2.00; depression loading=1.33; 95% CI: 0.92, 1.75) (data not shown). These results suggest that the same syndemic latent factor is being measured across heterosexual men and heterosexual women, but that substance use disorder may contribute more, and mental health less to the overall syndemic for heterosexual women than for heterosexual men.

MSM vs. Heterosexual women—A two-group CFA for the syndemic was then conducted with MSM and heterosexual women (Table 5). We first examined *metric invariance* (Model 2) in the syndemic in which factor loadings were constrained to be equal. The chi-square difference test – comparing the fit of the constrained model (Model 2) to that of the unconstrained model (Model 1) where factor loadings were free to vary – indicated the syndemic factor loadings could be constrained across groups of men without significantly reducing model fit, $\chi^2(4)=8.16$, supporting full metric invariance. Second, we examined *strong invariance* (Model 3) in which intercepts were constrained to be equal. The chi-square difference test between Models 3 and 2 indicated syndemic factor loadings and intercepts could be constrained across groups without significantly reducing model fit, $\chi^2(4)=8.44$, supporting strong invariance and that differences in the means of the indicators are due to differences in the level of the factor. Third, we examined *strict*

invariance in the syndemic in which residual variances were also constrained to be equal (Model 4). The chi-square difference test between Models 4 and 3 indicated syndemic factor loadings, intercepts, and residual variances could not be constrained across groups without significantly reducing model fit, $\chi^2(4)=13.08$, not supporting strict invariance. MSM had larger residual variances for violent victimization (11.52; 95%CI: 9.13, 14.53) and depressive symptoms (2.37; 95%CI: 1.73, 3.26) than heterosexual women (violent victimization=10.15; 95%CI: 9.13, 14.53; depressive symptoms=1.43; 95%CI: 0.92, 2.23) and heterosexual women had larger residual variances for stress (3.07; 95%CI: 2.49, 3.79) than MSM (2.14; 95%CI: 1.71, 2.69) (data not shown). Furthermore, heterosexual women had a larger factor mean (7.80; 95%CI: 7.29, 8.30) for the syndemic latent variable than MSM (7.53; 95%CI: 7.00, 8.06). These results suggest that the syndemic is comprised of similar indicators, but that there is more variance in violent victimization (MSM), depression (in MSM) and stress (heterosexual women) than captured by the observed items. Furthermore, it appears that the syndemic comprises a similar structure between MSM and heterosexual women, but is larger among heterosexual women than among MSM.

DISCUSSION

This study elucidated a syndemic latent variable in a sample of vulnerable PLWH in South Florida with several key findings. First, the variables used to measure the syndemic appropriately fit each sub-group, which supports research that substance use disorder, violence, and mental health all coincide to create a syndemic in people with HIV. Second, we empirically identify differences in the syndemic between our three sub-groups. Heterosexual men and MSM demonstrated similar syndemic latent variable factor loadings, but significantly different item intercepts. Heterosexual men and heterosexual women demonstrated significantly different similar syndemic latent variable factor loadings, with greater loadings for anxiety and depression in heterosexual men and a greater loading for substance use disorder in heterosexual women. MSM and heterosexual women demonstrated similar syndemic latent variable factor loadings and intercepts, but they had significantly different residual variances and factor means. We situate these findings within the syndemic literature and highlight methodological implications for future syndemic research among PLWH in order to better tailor HIV care and risk reduction interventions to these sub-groups to improve engagement and retention of vulnerable populations in HIV care.

This is one of the first studies to measure the syndemic among PLWH (12, 26–31), whereas most syndemic studies are among individuals at risk for HIV (17–25). Results support the co-occurrence of substance use disorder, violence, and mental health in a vulnerable sample of substance-using PLWH. In a review of the convergence of HIV, substance use disorders, and mental health among PLWH, Walkup et al. (2008) cite numerous studies that support the co-occurrence of these morbidities (66). For example, in a national probability sample of PLWH in medical care in the US (HIV Cost and Services Utilization Study) nearly half (48%) reported recent depressive, anxiety, or panic symptoms and almost 40% used illicit drugs (other than marijuana), with 12% screening as substance dependent (67). Although over half (61%) utilized mental health or substance abuse treatment (prior 6 months) (68), another study found that patient depression was underdiagnosed among those in HIV care (69). It is especially critical to diagnose and treat substance use disorders and mental health

problems among PLWH given their potentially detrimental effects on HIV care outcomes (66). Furthermore, violent victimization was a key co-morbidity of the syndemic in our analysis. A trauma-informed model of care for PLWH is necessary (70) in order to diagnose and address the effects that intimate partner violence and child sexual abuse may have on engagement in HIV care and ART adherence (71, 72).

Despite the syndemic consistently being comprised of substance use disorder, violence, and mental health, this study found that heterosexual men, MSM, and heterosexual women may experience the syndemic differently. For example, our analysis revealed that heterosexual men and MSM demonstrated similar syndemic latent variable factor loadings, but significantly different item intercepts. Heterosexual men had significantly larger item intercepts for substance use disorder (4.63), anxiety (-3.97), and depression (-2.23) compared to MSM (substance use disorder= 2.61, anxiety= -6.00, and depression= -3.02). This finding indicates that heterosexual men have a greater item mean, or expected item value when the factor item value is equal to zero, for substance use disorder, anxiety, and depression compared to MSM. Our study extends upon findings that syndemics undermine health and persist across the lifetime of heterosexual men, as well as MSM (73). It is furthermore critical to approach men's (both heterosexual men and MSM) health holistically by addressing the socio-structural inequities they face that may drive their health behavior and exacerbate their syndemic risk (74).

We also found that the underlying syndemic factor structure for heterosexual women was similar to that of heterosexual men, confirming the syndemic may comprise substance use disorder, violence, and mental health for both groups. However, we find heterosexual men and heterosexual women demonstrated significantly different syndemic latent variable factor loadings. Upon further analysis of the syndemic latent variable, we found that heterosexual men had greater loadings for anxiety (1.53) and depression (1.33) and smaller loadings for substance use disorder (0.46) than heterosexual women (anxiety=1.23; depression=0.98; substance use disorder=0.77). Our findings indicate that systemic integration of screening and care for substance use disorder and mental health co-morbidities for heterosexual men and heterosexual women living with HIV will maximize yields in positive outcomes (66). Moreover, the integration of services can serve to supplement, rather than replace, specialty services in that men and women with severe diagnoses can receive continued long-term care through specialty services (66).

We found that the syndemic had the same underlying factor structure for MSM as for heterosexual women, but our analysis revealed that MSM had significantly greater factor residual variances for violent victimization and depressive symptoms and lower factor residual variances for stress than heterosexual women. This finding suggests that there is more variance in violent victimization and depression for MSM and stress for heterosexual women than what is captured by the observed syndemic indicators. Moreover, heterosexual women had a larger syndemic factor mean than MSM, which indicates that the syndemic burden of experiencing multiple comorbidities is larger among heterosexual women than among MSM.

By identifying differences in the syndemic among PLWH, we can tailor healthcare to sub-group needs. This allows us to: 1) enhance diagnosis of depression, anxiety, and stress and improve patient linkage to mental health care and treatment (66), which may also improve adherence and linkage to care (75, 76), and 2) enhance substance use disorder diagnosis and linkage to treatment, thereby reducing risky behavior associated with substance use disorder and improving adherence and linkage to HIV care (66). Multiple group SEM has been used in other research to examine how socioeconomic status differentially affects mental health by race and ethnicity, age, and gender (77) and how stress and coping measures differ by race in PLWH (78). To our knowledge, measurement invariance modeling has only been applied to syndemics in one other study to identify syndemic differences between MSM, men who have sex with men and women (MSMW), and heterosexual men (50).

Limitations and Strengths

Findings must be considered in light of several important limitations. First, this is a cross-sectional study therefore limiting any assertions of causality. Adverse childhood experiences, including childhood sexual abuse and trauma, are correlated with increased disease and adoption of risky behaviors later in life. However, few studies apply the life course perspective to syndemic HIV research to understand how these traumatic experiences link to syndemic risk later in life. Likewise, understanding the developmental trajectory of the syndemic would ascertain the causal effect of syndemic co-morbidities on the underlying syndemic factor and identify life course targets to address unmet need for prevention and care in vulnerable populations. Second, our study sample is a highly specific group of vulnerable PLWH in South Florida, therefore also limiting the generalizability of findings to other groups of people or regions in the U.S. Third, the study utilized face-to-face in-depth surveys to collect data, which may lend to an interviewer effect or social desirability bias. Nevertheless, research has found that face-to-face interviews are more effective than telephone interviews and equally as effective as computer-based interviews in surveying about drug and alcohol use (79) and psychological distress (80).

Summary Statement

This study measured the syndemic among heterosexual men, heterosexual women, and MSM. We found that substance use disorder, violence, and mental health all coincide to create a syndemic in PLWH. We also identified differences in the syndemic between sub-groups. These findings indicate that substance use disorder, violence, and mental health may co-vary differently among sub-groups to make up distinct syndemics, which require tailored attention by health care providers.

Acknowledgments

This research was funded by PHS Grant Number R01DA023157 (Surratt), T32DA023356 (Tsuyuki), K01 DA036447 (Pitpitan), K01DA036439 (Urada), and K01DA031593 (Stockman) from the National Institute on Drug Abuse (NIDA), R01HD077891-04S1 (Tsuyuki) from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), L60MD003701 (Stockman) from the National Institute of Minority Health and Health Disparities, and K01AA025009 (Tsuyuki) from the National Institute of Alcohol Abuse and Alcoholism (NIAAA). NIDA, NICHD, NIMHD, and NIAAA had no further role in study design; in the collection, analysis and interpretation of data; in the writing of this report; or in the decision to submit the paper for publication.

References

1. Gardner EM, McLees MP, Steiner JF, del Rio C, Burman WJ. The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection. *Clinical infectious diseases*. 2011; 52(6):793–800. [PubMed: 21367734]
2. WHO. Antiretroviral treatment as prevention (TASP) of HIV and TB. 2012.
3. Cohen M, Chen Y, McCauley M, Gamble T, Hosseinipour M, Kumarasamy N, et al. Prevention of HIV-1 infection with early antiretroviral therapy. *New England Journal of Medicine*. 2011; 11(365): 469–505.
4. Baeten JM, Donnell D, Ndase P, Mugo NR, Campbell JD, Wangisi J, et al. Antiretroviral prophylaxis for HIV prevention in heterosexual men and women. *New England Journal of Medicine*. 2012; 367(5):399–410. [PubMed: 22784037]
5. Thigpen MC, Kebaabetswe PM, Paxton LA, Smith DK, Rose CE, Segolodi TM, et al. Antiretroviral preexposure prophylaxis for heterosexual HIV transmission in Botswana. *New England Journal of Medicine*. 2012; 367(5):423–434. [PubMed: 22784038]
6. Das M, Chu PL, Santos G-M, Scheer S, Vittinghoff E, McFarland W, et al. Decreases in community viral load are accompanied by reductions in new HIV infections in San Francisco. *PloS one*. 2010; 5(6):e11068. [PubMed: 20548786]
7. Eberhart MG, Yehia BR, Hillier A, Voytek CD, Fiore DJ, Blank M, et al. Individual and community factors associated with geographic clusters of poor HIV care retention and poor viral suppression. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2015; 69:S37–S43. [PubMed: 25867777]
8. Kamarulzaman A, Altice FL. Challenges in managing HIV in people who use drugs. *Current Opinion in Infectious Diseases*. 2015; 28(1):10–16. [PubMed: 25490106]
9. Metsch L, Philbin MM, Parish C, Shiu K, Frimpong JA. HIV testing, care, and treatment among women who use drugs from a global perspective: progress and challenges. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2015; 69:S162–S168. [PubMed: 25978483]
10. Tsuyuki K, Surratt HL. Antiretroviral Drug Diversion Links Social Vulnerability to Poor Medication Adherence in Substance Abusing Populations. *AIDS and Behavior*. 2015; 19(5):869–881. [PubMed: 25893656]
11. Sprague C, Simon SE. Understanding HIV care delays in the US South and the role of the social-level in HIV care engagement/retention: a qualitative study. *International journal for equity in health*. 2014; 13(1):1–14. [PubMed: 24383432]
12. Sullivan KA, Messer LC, Quinlivan EB. Substance abuse, violence, and HIV/AIDS (SAVA) syndemic effects on viral suppression among HIV positive women of color. *Aids patient care and STDs*. 2015; 29(S1):S42–S48. [PubMed: 25397666]
13. Eldred L, Malitz F. Introduction. *Aids patient care and STDs*. 2007; 21(S1):S-1–S-2.
14. Link BG, Phelan J. Social conditions as fundamental causes of disease. *Journal of Health and Social Behavior*. 1995:80–94. [PubMed: 7560851]
15. Tsai AC, Burns BF. Syndemics of psychosocial problems and HIV risk: A systematic review of empirical tests of the disease interaction concept. *Social Science & Medicine*. 2015; 139:26–35. [PubMed: 26150065]
16. Singer M, Clair S. Syndemics and public health: Reconceptualizing disease in bio-social context. *Medical Anthropology Quarterly*. 2003; 17(4):423–441. [PubMed: 14716917]
17. Kurtz SP, Buttram ME, Surratt HL, Ron S. Resilience, Syndemic Factors, and Serosorting Behaviors among HIV-Positive and HIV-Negative Substance-Using MSM. *AIDS Education and Prevention*. 2012; 24(3):193–205. [PubMed: 22676460]
18. Mustanski B, Garofalo R, Herrick A, Donenberg G. Psychosocial Health Problems Increase Risk for HIV Among Urban Young Men Who Have Sex With Men: Preliminary Evidence of a Syndemic ini Need of Attention. *Annals of Behavioral Medicine*. 2007; 34(1):37–45. [PubMed: 17688395]
19. Bruce D, Harper GW. Operating without a safety net: Gay male adolescents and emerging adults' experiences of marginalization and migrationm and implications for theory of syndemic

- production of health disparities. *Health Education & Behavior*. 2011; 38(4):367–378. [PubMed: 21398621]
20. Dyer TP, Shoptaw S, Guadamuz TE, Plankey M, Kao U, Ostrow D, et al. Application of syndemic theory to black men who have sex with men in the Multicenter AIDS Cohort Study. *Journal of Urban Health*. 2012
 21. Gonzalez-Guarda RM, Vasquez EP, Urrutia MT, Villarruel AM, Peragallo N. Hispanic women's experiences with substance abuse, intimate partner violence, and risk for HIV. *Journal of Transcultural Nursing*. 2011; 22:46–54. [PubMed: 21191036]
 22. Singer MC, Erickson PI, Badiane L, Diaz R, Ortiz D, Abraham T, et al. Syndemics, sex and the city: Understanding sexually transmitted diseases in social and cultural context. *Social Science & Medicine*. 2006; 63:2010–2021. [PubMed: 16782250]
 23. Brennan J, Kuhns LM, Johnson AK, Belzer M, Wilson EC, Garofalo R, et al. Syndemic theory and HIV-related risk among young transgendered women: The role of multiple, co-occurring health problems and social marginalization. *American journal of public health*. 2012; 102(9):1751–1757. [PubMed: 22873480]
 24. Senn TE, Carey MP, Vanable PA. The intersection of violence, substance use, depression, and STDs: Testing of a syndemic pattern among patients attending an urban STD clinic. *Journal of the National Medical Association*. 2010; 102(7):614–620. [PubMed: 20690325]
 25. Buttram ME, Surratt HL, Kurtz SP. Resilience and Syndemic Risk Factors among African American Female Sex Workers. *Psychology, Health and Medicine*. 2013
 26. Blashill AJ, Bedoya CA, Mayer KH, O'Cleirigh C, Pinkston MM, Remmert JE, et al. Psychosocial syndemics are additively associated with worse ART adherence in HIV-Infected individuals. *AIDS and Behavior*. 2014:1–6. [PubMed: 23321946]
 27. Friedman MR, Stall R, Silvestre AJ, Wei C, Shoptaw S, Herrick A, et al. Effects of syndemics on HIV viral load and medication adherence in the multicentre AIDS cohort study. *AIDS*. 2015; 29(9):1087–1096. [PubMed: 25870981]
 28. Mizuno Y, Purcell DW, Knowlton AR, Wilkinson JD, Gourevitch MN, Knight KR. Syndemic vulnerability, sexual and injection risk behaviors, and HIV continuum of care outcomes in HIV-positive injection drug users. *AIDS and Behavior*. 2015; 19(4):684–693. [PubMed: 25249392]
 29. Mimiaga MJ, O'Cleirigh C, Biello KB, Robertson AM, Safren SA, Coates TJ, et al. The effect of psychosocial syndemic production on 4-year HIV incidence and risk behavior in a large cohort of sexually active men who have sex with men. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2015; 68(3):329–336. [PubMed: 25501609]
 30. Mimiaga MJ, Biello KB, Robertson AM, Oldenburg CE, Rosenberger JG, O'Cleirigh C, et al. High prevalence of multiple syndemic conditions associated with sexual risk behavior and hiv infection among a large sample of Spanish-and Portuguese-speaking men who have sex with men in Latin America. *Archives of sexual behavior*. 2015; 44(7):1869–1878. [PubMed: 26159862]
 31. Abdul-Quader AS, Friedman SR, Jarlais DD, Marmor MM, Maslanski R, Bartelme S. Methadone Maintenance and Behavior by Intravenous Drug Users that Can Transmit HIV. *Contemporary Drug Problems*. 1987; 14(3):425–434.
 32. Pitpitan EV, Strathdee SA, Semple SJ, Chavarin CV, Magis-Rodriguez C, Patterson TL. Buffering Syndemic Effects in a Sexual Risk-Reduction Intervention for Male Clients of Female Sex Workers: Results From a Randomized Controlled Trial. *American Journal of Public Health*. 2015; (0):e1–e6.
 33. Connell, RW. *Gender and power*. Stanford, CA: Stanford University Press; 1987.
 34. Wingood GM, DiClemente RJ. Application of the theory of gender and power to examine HIV-related exposures, risk factors, and effective interventions for women. *Health education & behavior*. 2000; 27(5):539–565. [PubMed: 11009126]
 35. Meyer IH. Minority stress and mental health in gay men. *Journal of health and social behaviour*. 1995:38–56.
 36. Nolen-Hoeksema, S., Hilt, LM. *Handbook of depression in adolescents*. Routledge; 2013.
 37. Hatzenbuehler, ML., Hilt, LM., Nolen-Hoeksema, S. *Handbook of gender research in psychology*. Springer; 2010. Gender, sexual orientation, and vulnerability to depression; p. 133-151.

38. McLean CP, Asnaani A, Litz BT, Hofmann SG. Gender differences in anxiety disorders: prevalence, course of illness, comorbidity and burden of illness. *Journal of psychiatric research*. 2011; 45(8):1027–1035. [PubMed: 21439576]
39. Caldwell JE, Swan SC, Woodbrowm VD. Gender differences in intimate partner violence outcomes. *Psychology of Violence*. 2012; 2(1):42.
40. Schulte MT, Ramo D, Brown SA. Gender differences in factors influencing alcohol use and drinking progression among adolescents. *Clinical psychology review*. 2009; 29(6):535–547. [PubMed: 19592147]
41. Substance Abuse and Mental Health Services Administration. Results from the 2009 National Survey on Drug Use and Health: Summary of National Findings, Volume I. Substance Abuse and Mental Health Services Administration; Rockville, MD: 2010. 2010
42. Marshal MP, Dietz LJ, Friedman MS, Stall R, Smith HA, McGinley J, et al. Suicidality and depression disparities between sexual minority and heterosexual youth: A meta-analytic review. *Journal of Adolescent Health*. 2011; 49(2):115–123. [PubMed: 21783042]
43. Marshal MP, Friedman MS, Stall R, King KM, Miles J, Gold MA, et al. Sexual orientation and adolescent substance use: A meta-analysis and methodological review. *Addiction*. 2008; 103:546–556. [PubMed: 18339100]
44. Herrick AL, Marshal MP, Smith HA, Sucato G, Stall RD. Sex while intoxicated: a meta-analysis comparing heterosexual and sexual minority youth. *Journal of Adolescent Health*. 2011; 48(3): 306–309. [PubMed: 21338904]
45. García JJ, Sharif MZ. Black lives matter: a commentary on racism and public health. *American journal of public health*. 2015; 105(8):e27–e30.
46. Martinez, R. *Latino homicide: Immigration, violence, and community*. Routledge; 2014.
47. Pulerwitz J, Barker G. Measuring attitudes toward gender norms among young men in Brazil: development and psychometric evaluation of the GEM scale. *Men and Masculinities*. 2008; 10(3): 322–338.
48. Stall R, Mills TC, Williamson J, Hart T, Greenwood G, Paul J, et al. Association of co-occurring psychosocial health problems and increased vulnerability to HIV/AIDS among urban men who have sex with men. *American journal of public health*. 2003; 93(6):939–942. [PubMed: 12773359]
49. Gonzalez-Guarda RM, McCabe BE, Vermeesch AL, Cianelli R, Florom-Smith AL, Peragallo N. Cultural phenomena and the syndemic factor: substance abuse, violence, HIV, and depression among Hispanic women. *Annals of anthropological practice*. 2012; 36(2):212–231. [PubMed: 24575326]
50. Mustanski B, Andrews R, Herrick A, Stall R, Schnarrs PW. A syndemic of psychosocial health disparities and associations with risk for attempting suicide among young sexual minority men. *American journal of public health*. 2014; 104(2):287–294. [PubMed: 24328641]
51. DiStefano C, Zhu M, Mindrila D. Understanding and using factor scores: Considerations for the applied researcher. *Practical Assessment, Research & Evaluation*. 2009; 14(20):1–11.
52. Starks TJ, Millar BM, Eggleston JJ, Parsons JT. Syndemic factors associated with hiv risk for gay and bisexual men: comparing latent class and latent factor modeling. *AIDS and Behavior*. 2014; 18(11):2075–2079. [PubMed: 25055765]
53. AIDS Vu. Prevalence of HIV by Transmission Category and County. 2013.
54. Surratt HL, Kurtz SP, Cicero TJ, O'Grady C, Levi-Minzi MA. Antiretroviral Medication Diversion Among HIV-Positive Substance Abusers in South Florida. *American journal of public health*. 2013; 103(6):1026–1028. [PubMed: 23597362]
55. Miami Dade County Department of Health. Miami-Dade County Reported HIV (Not AIDS) Cases Through. 2012. 2012
56. Dennis, MLTJ., White, MK., Unsicker, JI., Hodgkins, D. *Global Appraisal of Individual Needs-Initial (GAIN-I)*. Bloomington, IL: Chestnut Health Systems; 2002.
57. Dennis, ML., Titus, JC., White, MK., Unsicker, JI., Hodgkins, D. *Global Appraisal of Individual Needs - Initial (GAIN-I)*. Bloomington, IL: Chestnut Health Systems; 2002.
58. Dennis ML, White M, Titus JC, Unsicker JI. *Global Appraisal of Individual Needs: Administration Guide for the GAIN and Related Measures (Version 5)*. Chestnut Health Systems. 2008

59. Shalowitz MU, Berry CA, Rasinski KA, Dannhausen-Brun CA. A new measure of contemporary life stress: Development, validation, and reliability of the CRISYS. *Health Services Research*. 1998; 33(5):1381–1402. [PubMed: 9865225]
60. Steenkamp J-BE, Baumgartner H. Assessing measurement invariance in cross-national consumer research. *Journal of consumer research*. 1998; 25(1):78–90.
61. Weber S. bacon: An effective way to detect outliers in multivariate data using Stata (and Mata). *Stata Journal*. 2010; 10(3):331.
62. Billor N, Hadi AS, Velleman PF. BACON: blocked adaptive computationally efficient outlier nominators. *Computational Statistics & Data Analysis*. 2000; 34(3):279–298.
63. StataCorp L. Stata version 13.1. College Station, TX: StataCorp LP; 2014.
64. Hu, Lt, Bentler, PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*. 1999; 6(1):1–55.
65. Slof-Op't Landt M, Dolan C, Rebollo-Mesa I, Bartels M, Van Furth E, van Beijsterveldt C, et al. Sex differences in sum scores may be hard to interpret: the importance of measurement invariance. *Assessment*. 2009
66. Walkup J, Blank MB, Gonzalez JS, Safren S, Schwartz R, Brown L, et al. The impact of mental health and substance abuse factors on HIV prevention and treatment. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2008; 47:S15–S19. [PubMed: 18301129]
67. Bing EG, Burnam MA, Longshore D, Fleishman JA, Sherbourne CD, London AS, et al. Psychiatric disorders and drug use among human immunodeficiency virus–infected adults in the United States. *Archives of General Psychiatry*. 2001; 58(8):721–728. [PubMed: 11483137]
68. Burnam MA, Bing EG, Morton SC, Sherbourne C, Fleishman JA, London AS, et al. Use of mental health and substance abuse treatment services among adults with HIV in the United States. *Archives of General Psychiatry*. 2001; 58(8):729–736. [PubMed: 11483138]
69. Asch SM, Kilbourne AM, Gifford AL, Burnam MA, Turner B, Shapiro MF, et al. Underdiagnosis of depression in HIV. *Journal of general internal medicine*. 2003; 18(6):450–460. [PubMed: 12823652]
70. Brezing C, Ferrara M, Freudenreich O. The Syndemic Illness of HIV and Trauma: Implications for a Trauma-Informed Model of Care. *Psychosomatics*. 2015; 56(2):107–118. [PubMed: 25597836]
71. Hatcher AM, Smout EM, Turan JM, Christofides N, Stöckl H. Intimate partner violence and engagement in HIV care and treatment among women: a systematic review and meta-analysis. *AIDS (London, England)*. 2015
72. Meade CS, Hansen NB, Kochman A, Sikkema KJ. Utilization of medical treatments and adherence to antiretroviral therapy among HIV-positive adults with histories of childhood sexual abuse. *Aids patient care and STDs*. 2009; 23(4):259–266. [PubMed: 19260772]
73. Halkitis PN, Kapadia F, Bub KL, Barton S, Moreira AD, Stults CB. A longitudinal investigation of syndemic conditions among young gay, bisexual, and other MSM: the P18 Cohort Study. *AIDS and Behavior*. 2015; 19(6):970–980. [PubMed: 25192900]
74. Halkitis PN, Wolitski RJ, Millett GA. A holistic approach to addressing HIV infection disparities in gay, bisexual, and other men who have sex with men. *American Psychologist*. 2013; 68(4):261. [PubMed: 23688093]
75. Ammassari A, Murri R, Pezzotti P, Trotta MP, Ravasio L, De Longis P, et al. Self-reported symptoms and medication side effects influence adherence to highly active antiretroviral therapy in persons with HIV infection. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2001; 28(5):445–449. [PubMed: 11744832]
76. Yun LW, Maravi M, Kobayashi JS, Barton PL, Davidson AJ. Antidepressant treatment improves adherence to antiretroviral therapy among depressed HIV-infected patients. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2005; 38(4):432–438. [PubMed: 15764960]
77. Meyer OL, Castro-Schilo L, Aguilar-Gaxiola S. Determinants of mental health and self-rated health: A model of socioeconomic status, neighborhood safety, and physical activity. *American journal of public health*. 2014; 104(9):1734–1741. [PubMed: 25033151]

78. Heckman BD, Berlin KS, Heckman TG, Feaster DJ. Psychometric characteristics and race-related measurement invariance of stress and coping measures in adults with HIV/AIDS. *AIDS and Behavior*. 2011; 15(2):441–453. [PubMed: 21153050]
79. Holbrook AL, Green MC, Krosnick JA. Telephone versus face-to-face interviewing of national probability samples with long questionnaires: Comparisons of respondent satisficing and social desirability response bias. *Public opinion quarterly*. 2003; 67(1):79–125.
80. Newman JC, Des Jarlais DC, Turner CF, Gribble J, Cooley P, Paone D. The differential effects of face-to-face and computer interview modes. *American journal of public health*. 2002; 92:294–297. [PubMed: 11818309]




FIGURE 1. Multigroup structural equation model of syndemic health co-morbidities, South Florida, 2013

Structural Model - Configural Invariance:

1. Heterosexual men - AIC = 3,175.75, BIC = 3,219.11, log-likelihood = -1,572.88;
2. MSM - AIC = 3,674.49; BIC = 3,720.43, log-likelihood = -1,822.25;
3. Heterosexual women - AIC = 4,490.03, BIC = 4,538.73, log-likelihood = -2,230.01

Note. AIC=Akaike Information Criterion; BIC=Bayesian Information Criterion.




Figure 2. Scatter plot matrices using jittering with correlation coefficients of syndemic co-morbidities among heterosexual men, MSM, and heterosexual women (n=481)

Table 1
 Characteristics and Syndemic Indicators in Adults Living with HIV in South Florida

	TOTAL (n=481)		Heterosexual Men (n=133)		MSM (n=158)		Women (n=190)	
	M	SD	M	SD	M	SD	M	SD
Age	46.13	7.75	47.50	7.51	44.85**	8.11	46.24	7.47
Years HIV-positive	12.95	7.31	12.49	7.43	14.32*	7.63	12.12	6.82
Violent Victimization Severity	7.06	4.19	5.53	4.00	7.05**	4.05	8.15***	4.12
Anxiety Symptoms (last year)	4.68	3.66	4.51	3.81	4.60	3.50	4.87	3.69
Depressive Symptoms (last year)	5.36	3.02	5.15	3.19	5.26	3.06	5.58	2.87
Substance Use Disorder (last year)	6.87	3.23	7.16	3.10	6.92	3.05	6.64	3.46
Stress (90 days)	2.18	1.85	2.11	1.76	2.13	1.67	2.27	2.04
<i>(F=8.55, p = 0.004)</i>								
<i>(F=4.61, p = 0.032)</i>								
<i>(F=10.07, p = 0.002)</i>								
<i>(F=32.39, p<0.001)</i>								
Categorical variables								
	n	%	n	%	n	%	n	%
<i>(X² =43.11, p<0.001)</i>								
Education***								
High School or more	269	56	59	44	118	75	92	48
Less than High School	212	44	74	56	40	25	98	52
<i>(X² =46.12, p<0.001)</i>								
Race/Ethnicity***								
Black/African-American	324	67	98	74	79	50	147	77
Non-Hispanic White	68	14	5	4	39	25	24	13
Hispanic/Latino	89	19	30	23	40	25	19	10
Income (\$/month)								
501 or more	326	68	84	63	105	66	137	72
0 to 500	155	32	49	37	53	34	53	28
<i>(X² =6.49, p=0.039)</i>								
Positive STI (12 months)*	139	29	29	22	44	28	66	35

Continuous variables	TOTAL (n=481)		Heterosexual Men (n=133)		MSM (n=158)		Women (n=190)	
	M	SD	M	SD	M	SD	M	SD
TOTAL	481	100.0	133	29.9	158	30.1	190	40.0

M=mean; SD=Standard Deviation; F=F-statistic;

* p<0.05,

** p<0.01,

*** p<0.001 difference between heterosexual men, MSM, and women using χ^2 for comparisons among categorical variables & two-way ANOVA for comparison of mean between MSM, women, and heterosexual men (comparison group).

Table 2
Measurement Model of a Syndemic Factor Using Confirmatory Factor Analyses

<i>Indicators</i>	<u>Model 1: Heterosexual Men</u> (n=133)		<u>Model 2: MSM</u> (n=158)		<u>Model 3: Women</u> (n=190)	
	<i>b</i>	SE	<i>b</i>	SE	<i>b</i>	SE
Violent Victimization Severity [‡]	1.00		1.00		1.00	
Substance Use Disorder (last year)	0.46	0.14	0.61	0.13	0.78	0.11
Anxiety Symptoms (last year)	1.53	0.24	1.50	0.20	1.23	0.13
Depressive Symptoms (last year)	1.33	0.21	1.17	0.16	0.98	0.10
Stress (90 days)	0.43	0.09	0.35	0.07	0.40	0.06
Syndemic Mean	5.53	0.35	7.05	0.32	8.15	0.30
<i>Fit Indices</i>						
X ² (d.f.)	4.95(5)	p = 0.422	12.31(5)	p = 0.031	16.26(5)	p = 0.01
CFI	1.00		0.98		0.97	
RMSEA	0.00		0.10		0.11	
SRMR	0.03		0.04		0.04	

b = factor loadings; SE = Standard Error; CFI = Comparative Fit Index; RMSEA = Root Mean Squared Error Approximation; SRMR = Standardized Root Squared Residual;

[‡] marker variable for identification purposes - factor loading fixed to 1 and intercept fixed to 0. Model estimation stopped once model comparisons are significantly different demonstrating worse model fit.

The results of metric invariance tests of syndemic indicators for heterosexual men and MSM (n=291)

Table 3

<i>Model specification</i>	χ^2	df	CFI	RMSEA	SRMR
Model 1: Free all parameters	17.26	10	0.99	0.07	0.04
Model 2: Metric (loadings)	20.19	14	0.99	0.06	0.05
Model 3: Strong (loadings + intercepts)	33.62	18	0.97	0.08	0.05
<i>Model comparisons</i>	χ^2	df	CFI	p-value	
Model 2-Model 1 (Metric invariance)	2.93	4	0.00	0.57	
Model 3-Model 2 (Strong invariance)	13.43	4	-0.02	0.01	

CFI=Comparative Fit Index; RMSEA=Root Mean Squared Error Approximation; SRMR=Standardized Root Mean Squared Residual;

* marker variable for identification purposes - factor loading fixed to 1 and intercept fixed to 0. Model estimation stopped once model comparisons are significantly different demonstrating worse model fit.

Table 4

The results of metric invariance tests of syndemic indicators for heterosexual men and women (n=323)

<i>Model specification</i>	χ^2	df	CFI	RMSEA	SRMR
Model 1: Free all parameters	21.22	10	0.98	0.08	0.03
Model 2: Metric (loadings)	35.13	14	0.97	0.10	0.07
<i>Model comparisons</i>	χ^2	df	CFI	p-value	
Model 2-Model 1 (Metric invariance)	13.91	4	-0.02	0.01	

CFI=Comparative Fit Index; RMSEA=Root Mean Squared Error Approximation; SRMR=Standardized Root Mean Squared Residual;

* marker variable for identification purposes - factor loading fixed to 1 and intercept fixed to 0. Model estimation stopped once model comparisons are significantly different demonstrating worse model fit.

Table 5
The results of metric invariance tests of syndemic indicators for MSM and women (n=348)

<i>Model specification</i>	χ^2	df	CFI	RMSEA	SRMR
Model 1: Free all parameters	28.57	10	0.97	0.10	0.04
Model 2: Metric (loadings)	36.73	14	0.97	0.10	0.06
Model 3: Strong (loadings + intercepts)	45.17	18	0.96	0.09	0.06
Model 4: Strict (loadings + intercepts + residuals)	58.25	23	0.95	0.09	0.05
<i>Model comparisons</i>	χ^2	df	CFI		p-value
Model 2-Model 1 (Metric invariance)	8.16	4	-0.01		0.09
Model 3-Model 2 (Strong invariance)	8.44	4	-0.01		0.08
Model 4-Model 3 (Strict invariance)	13.08	5	-0.01		0.02

CFI=Comparative Fit Index; RMSEA=Root Mean Squared Error Approximation; SRMR=Standardized Root Mean Squared Residual;

[‡]marker variable for identification purposes - factor loading fixed to 1, intercept and residual variance fixed to 0. Model estimation stopped once model comparisons are significantly different demonstrating worse model fit.