



University of Kentucky  
UKnowledge

Sanders-Brown Center on Aging Faculty  
Publications

Aging

3-2017

# Neuroimaging Biomarkers of Caloric Restriction on Brain Metabolic and Vascular Functions

Ai-Ling Lin

University of Kentucky, [ailing.lin@uky.edu](mailto:ailing.lin@uky.edu)

Ishita Parikh

University of Kentucky, [ishita.parikh@uky.edu](mailto:ishita.parikh@uky.edu)

Jared D. Hoffman


University of Kentucky

David Ma

University of Kentucky, [david.ma@uky.edu](mailto:david.ma@uky.edu)

**Right click to open a feedback form in a new tab to let us know how this document benefits you.**

Follow this and additional works at: [https://uknowledge.uky.edu/sbcoa\\_facpub](https://uknowledge.uky.edu/sbcoa_facpub)

 Part of the [Diseases Commons](#), [Neuroscience and Neurobiology Commons](#), and the [Nutrition Commons](#)

## Repository Citation

Lin, Ai-Ling; Parikh, Ishita; Hoffman, Jared D.; and Ma, David, "Neuroimaging Biomarkers of Caloric Restriction on Brain Metabolic and Vascular Functions" (2017). *Sanders-Brown Center on Aging Faculty Publications*. 129.  
[https://uknowledge.uky.edu/sbcoa\\_facpub/129](https://uknowledge.uky.edu/sbcoa_facpub/129)

This Article is brought to you for free and open access by the Aging at UKnowledge. It has been accepted for inclusion in Sanders-Brown Center on Aging Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**Neuroimaging Biomarkers of Caloric Restriction on Brain Metabolic and Vascular Functions**

**Notes/Citation Information**

Published in *Current Nutrition Reports*, v. 6, issue 1, p. 41-48.

© Springer Science+Business Media New York 2017

The copyright holder has granted the permission for posting the article here.

This is a post-peer-review, pre-copyedit version of an article published in *Current Nutrition Reports*. The final authenticated version is available online at: <https://doi.org/10.1007/s13668-017-0187-9>.

**Digital Object Identifier (DOI)**

<https://doi.org/10.1007/s13668-017-0187-9>



Published in final edited form as:

*Curr Nutr Rep.* 2017 March ; 6(1): 41–48. doi:10.1007/s13668-017-0187-9.

## Neuroimaging Biomarkers of Caloric Restriction on Brain Metabolic and Vascular Functions

Ai-Ling Lin, Ph.D.<sup>1,2,3</sup>, Ishita Parikh, Ph.D.<sup>1</sup>, Jared D. Hoffman, M.S.<sup>1,2</sup>, and David Ma<sup>1</sup>

<sup>1</sup>Sanders-Brown Center on Aging, University of Kentucky, 800 South Limestone Street, Lexington, KY, USA 40536

<sup>2</sup>Department of Pharmacology and Nutritional Sciences, University of Kentucky, 800 Roth Street, Lexington, KY, USA 40536

<sup>3</sup>Department of Biomedical Engineering, University of Kentucky, 143 Graham Avenue, Lexington, KY, USA 40506

### Abstract

**Purpose of review**—Non-invasive neuroimaging methods have been developed as powerful tools for identifying *in vivo* brain functions for studies in humans and animals. Here we review the imaging biomarkers that are being used to determine the changes within brain metabolic and vascular functions induced by caloric restriction (CR), and their potential usefulness for future studies with dietary interventions in humans.

**Recent findings**—CR causes an early shift in brain metabolism of glucose to ketone bodies, and enhances ATP production, neuronal activity and cerebral blood flow (CBF). With age, CR preserves mitochondrial activity, neurotransmission, CBF, and spatial memory. CR also reduces anxiety in aging mice. Neuroimaging studies in humans show that CR restores abnormal brain activity in the amygdala of women with obesity and enhances brain connectivity in old adults.

**Summary**—Neuroimaging methods have excellent translational values and can be widely applied in future studies to identify dietary effects on brain functions in humans.

### Keywords

caloric restriction; positron emission tomography (PET); magnetic resonance imaging (MRI); magnetic resonance spectroscopy (MRS); glucose metabolism; ketone bodies; cerebral blood flow; mammalian target of rapamycin (mTOR); brain aging; Alzheimer's disease; memory; anxiety; translational research

---

Correspondence to: Ai-Ling Lin, 307 Sanders-Brown Building, 800 South Limestone Street, Lexington, KY, USA 40536, Telephone: (859) 323-6671, ailing.lin@uky.edu.

#### **Compliance with Ethics Guidelines**

#### **Conflict of Interest**

Ai-Ling Lin, Ishita Parikh, Jared D. Hoffman, and David Ma declare that they have no conflict of interest.

#### **Human and Animal Rights and Informed Consent**

This article does not contain any studies with human or animal subjects performed by any of the authors.

## Introduction

Brain metabolic and vascular integrity plays an important role in determining cognitive capability and mental health. Failure to maintain cerebral metabolic rate of glucose (CMR<sub>glc</sub>) has been shown to lead to cognitive impairment and brain volume atrophy, as observed in normal aging and in patients with diabetes and Alzheimer's disease (AD) [1]. Similarly, studies have shown that neurovascular risk is highly associated with accelerated decline in language ability, verbal memory, attention and visuospatial abilities [2, 3]. Reduced cerebral blood flow (CBF) is linked to anxiety and depression [4–6], and impaired blood-brain barrier is associated with neuroinflammation and synaptic dysfunction [7]. These metabolic and hemodynamic reductions precede brain structural alteration (gray matter and white matter atrophy) and cognitive impairment [8–10]. Therefore, preserving brain metabolism (i.e., glucose oxidative capacity) and hemodynamics are critical for optimizing our lifespan and healthspan [11].

Caloric restriction (CR), without malnutrition, has been repeatedly shown to extend life expectancy as well as enhance brain functions [12–15]. On biochemical and molecular level, CR shows to improve glucose homeostasis and insulin sensitivity [16, 17], up-regulate brain-derived neuro-trophic factor [18–20], and reduce oxidative stress and inflammation [21–24]. CR also shows beneficial effects on vascular system by decreasing blood pressure, atherogenic lipids, inflammatory cytokines and increased cellular stress resistance [25, 26]. In line with this, animals treated with CR had lower incidences of age-related neurodegenerative disorders and diabetes [27, 28]. However, experiments on biochemical and molecular levels may limit CR research to animal models, in conditions of *in vitro* or *ex vivo*, and their findings might not be fully translated to humans.

Noninvasive neuroimaging methods have been developed as powerful tools for identifying *in vivo* metabolic and vascular biomarkers, including positron emission tomography (PET), and magnetic resonance imaging (MRI) and spectroscopy (MRS) [29, 30]. Among these, PET with fluorine-18 (<sup>18</sup>F)-labeled 2-fluoro-2-deoxy-d-glucose (<sup>18</sup>FDG) tracer and proton (<sup>1</sup>H) MRS are the well-established methods to quantify CMR<sub>glc</sub> and neural metabolites, respectively. Other novel techniques, including <sup>1</sup>H[<sup>13</sup>C] proton-observed-carbon-edited (POCE) NMR, have also developed to determine brain bioenergetics in aging brain. MRI-based arterial spin labeling (ASL) uses arterial blood water as an endogenous tracer to determine quantitative cerebral blood flow (CBF), which refers to the rate of delivery of arterial blood to the capillary bed in brain tissue and is quantified in milliliters of blood per 100 g of brain tissue per minute. Table 1 summarizes the imaging techniques. These imaging methods have been widely used in humans, and have been “reversely” translated to be used animal studies. Using these advanced imaging techniques, we are able to identify the CR-induced changes in brain metabolic and vascular functions over time in studies related to aging and neurodegenerative diseases. Here we review the imaging biomarkers being used to determine the CR effects on brain metabolic and vascular functions, and their potential usefulness for future studies with dietary interventions in humans.

## Neuroimaging Evidence of Preserving Brain Metabolic Functions by CR

Mitochondria are the powerhouse of the cells. Mitochondrial oxidative phosphorylation of glucose is the predominate mode of energy generation in mammalian species. CR effects have been extensively studied in isolated mitochondria, showing increased capacity for oxidative phosphorylation resulting from elevated mitochondrial respiration or increased mitochondrial biogenesis. Mammalian brain has the highest energy demand of any organ based on its size; mitochondrial function plays a critical role in supporting neuronal activity and functional processes of the brain.

Using advanced POCE techniques, we are able to trace *in vivo* mitochondrial activity [31]. With the infusion of  $^{13}\text{C}$ -labeled glucose, the mitochondrial glucose oxidation rates in neurons can be measured ( $\text{CMR}_{\text{glc(ox),N}}$ ;  $\mu\text{mol/g/min}$ ). The POCE techniques also provide information for glutamate–glutamine neurotransmitter cycling ( $V_{\text{cycle}}$ ;  $\mu\text{mol/g/min}$ ), which allows us to determine neuronal activity at the same time. We have recently reported the of measure  $\text{CMR}_{\text{glc(ox),N}}$  and  $V_{\text{cycle}}$  in aged rats chronically treated with CR diet (24 months of age). We compared the results from young *ad libitum* (AL; 5 months of age) and age-matched old AL (i.e., healthy aging) rats. We found that, compared with the young AL, neuronal energy production and neurotransmission rates were significantly reduced in healthy aging. However, old CR rats had similar levels of  $\text{CMR}_{\text{glc(ox),N}}$  and  $V_{\text{cycle}}$  compared to those of young AL rats, suggesting preserved mitochondrial functions and neuronal activity (Fig. 1A). We calculated adenosine triphosphate (ATP) production rate for the three groups, and found that old CR rats also had comparable outcomes with the young AL.

$^1\text{H}$  MRS is another *in vivo* imaging technique, which permits visualization of various markers for cellular integrity and function (Fig. 1B). In a recent study, we used  $^1\text{H}$ -MRS to determine energy metabolites in hippocampus in young and old mice fed with either AL or 40% CR diet [32]. We found that young CR mice had significant increases of total creatine (TCr), a high-energy substrate, compared to young AL mice. Although TCr dropped dramatically when the CR mice were getting old, the level was still comparable to the young AL, and higher than that of the old AL. TCr is the sum of creatine and phosphorus creatine (plays a crucial role as an intracellular buffer during the production of ATP), suggesting that CR increases ATP production in young CR mice, and preserves ATP production in old CR mice. We also found significantly elevated Taurine in young CR compared to the young AL mice. As Taurine is associated with neurotransmitter modulation, this may indicate that young CR mice have an early enhancement of neuronal activity compared to the age-matched AL mice. Collectively, data from POCE and  $^1\text{H}$ -MRS indicate that CR enhances brain metabolism and neuronal activity at early stage, and preserves these functions in aging.

As glucose is the predominate energy substrate for sustaining mitochondrial functions and neuronal activity, we further investigated the CR effects on brain glucose uptake using  $^{18}\text{F}$ FDG-PET. Based on the findings with POCE and  $^1\text{H}$ -MRS, we expected that brain glucose uptake would also be enhanced at early stage and preserved with age. To our surprise, CR dramatically reduces brain glucose uptake in young mice, both globally and in the regions related to cognitive functions (e.g., hippocampus and frontal cortex) (Fig. 1C, *left*). In addition, the similar lower level of glucose uptake was maintained with age [32].

Similar results were also found in rats; old CR rats had significantly lower glucose uptake in the various brain regions compared to that of the age-matched AL rats [33]. These CMRglc changes are in line with the evidence of down-regulation mammalian target of rapamycin (mTOR) signaling by CR. mTOR is a nutrient sensor, a serine/threonine protein that is expressed at high levels in the brain. It serves as a potent neuronal survival and division signal, responding to different signals, such as nutrients, growth factors, and stress [34].

The PET imaging findings led us to speculate that brain may use alternative fuel substrates to sustain mitochondrial activity and neuronal functions. In addition to glucose, brain uses ketone bodies as energy source [35]. We thus used mass spectroscopy to measure the levels of ketone bodies, beta-hydroxybutyrate (BHB) from the brain tissue. We found that old rats (24 months of age) with CR diet had significantly increased BHB level, compared with the age-matched and young (5 months of age) AL rats. Similarly, young and old CR mice had significantly higher ketone bodies in the blood compared to their age-matched AL mice [32]. Our findings suggested that CR induces a metabolic shift from glucose to ketone bodies utilization in the brain at very early stage.

Increased ketone bodies metabolism suggest increased oxidative metabolism. Utilization of ketone bodies significantly elevates the oxygen utilization in mitochondria through beta-oxidation of fatty acid [36, 37]. This is supported by evidence from isolated mitochondria, showing CR enhances mitochondrial function and induces bioenergetic efficiency [38]. This is also consistent with our previous imaging findings that old animals with chronic CR diet had preserved oxidative metabolism, mitochondrial functions (TCA cycle flux and ATP production), and neuronal activity (neurotransmission rate) compared to the old AL animals [31]. The increased oxidative metabolism, particularly in brain regions associated with cognition (e.g., frontal cortex and hippocampus), may also play a crucial role for neuronal and cognitive protections.

Previous studies showed that cognition-associated brain regions have non-oxidative glycolysis exceeding the required needs of oxidative phosphorylation, a phenomenon known as aerobic glycolysis (AG) [39]. Excessive AG (or the “Warburg effect”) is a key process that sustains T cell activation and differentiation, and involved in inflammatory-mediated conditions [40]. In line with this, the distribution of AG in normal young adults is spatially correlated with A $\beta$  deposition in AD patients and cognitively normal individuals with elevated A $\beta$  [41, 42]. Animal studies further demonstrated that A $\beta$  plaque formation is an activity dependent process associated with AG [41, 43]. Therefore, increased oxidative metabolism in cognition-related regions may decrease AG and thus reduce the risk for AD, consistent with the findings in CR mice [44, 45]. Reduced AD risk was also found in rhesus monkeys, showing CR impedes age-related iron deposition in the brain, which consequently reduces the potential interaction between metal and A $\beta$ , and thus decelerates the pathogenesis of AD [46, 47].

Taken together, using multi-modal imaging methods we can demonstrate that CR enhances oxidative metabolism at early stage and decelerates the decline with age; the preservation of brain oxidative metabolism may be contributed by the shifted utilization from glucose to ketone bodies.

## Neuroimaging Evidence of Preserving Brain Vascular Functions by CR

CBF is highly correlated with local neuronal activity and metabolism, known as neurovascular coupling. Neurovascular coupling is used as a surrogate marker of brain function [48, 49]. In light of the metabolic preservation by CR, we further investigated CBF in rodents fed with AL and 40% CR diet. We found that CR significantly enhanced CBF when mice were young (Fig. 1C, right) and decelerated the rate of CBF decline in aging [50]. As a result, old CR mice (20–24 months of age) had similar levels of CBF compared to the young AL mice, indicating CBF preservation in aging. Similar results were found in rats, showing that old rats with chronic CR diet had much higher CBF compared to the age-matched animal, and had comparable CBF level compared to the young AL rats [33]. These findings indicate that CR had early enhancement on CBF and is preserved with age, which is consistent with the metabolic results.

We further identified the potential mechanism of the CR-induced changes on CBF. We previously showed that mTOR inhibited activated endothelial nitric oxide synthase (eNOS) and releases vasodilator nitric oxide, which results in increased CBF [51, 52]. To determine whether the enhancement of CBF in young adult mice is also associated with mTOR signaling, we measured the protein levels of mTOR and eNOS in capillaries isolated from young CR and AL mice. We found that, CR mice had significantly lower levels of mTOR compared to AL mice but higher level of eNOS, consistent with our previous findings [51]. In addition, mTOR-induced changes in metabolism (e.g., shift from glucose to ketone bodies metabolism) are also associated with CBF changes. Two studies show that an acute increase in ketone bodies concentration via infusion of BHB, increased CBF without affecting the overall cerebral metabolic activity; suggesting that ketone bodies have direct effect on the cerebral endothelium to increase CBF, independent of metabolic interactions [53, 54]. In addition, ketogenesis is also associated with downregulated mTOR activity [55] and upregulated adenosine monophosphate-activated protein kinase [56]. Both of these changes can activate endothelial nitric oxide synthase signaling and consequently increase CBF [57].

It has been shown that lifelong CR was able to mitigate the age-related endothelial dysfunction in conduit arteries and partially in cerebral resistance arteries. CR has been demonstrated to exert beneficial effects in the cerebrovascular endothelial cells of young and aged rats fed with CR diet compared to the control group, demonstrating that CR could help prevent endothelial dysfunction and alleviate the decrease in CBF that comes with age [58]. This is also consistent with the literature in that CR preserves vascular functions and vascular density in aging [59, 60].

## Correlation of Neuroimaging outcomes with Cognition and Anxiety

CR has been repeatedly shown to improve memory in aging, both for studies in humans and animals [61–64]. We had similar observations in a recent study, showing CR had significantly protective effects on learning and spatial memory for old mice in a Radial Arm Water Maze task [50]. The cognitive outcomes are correlated with CBF in hippocampus and frontal cortex, the brain areas regulating learning and memory. The findings indicate that

level of CBF in cognition-associated brain regions may play a critical role for determining performances on learning and spatial memory.

In the same study, we used elevated plus maze (EPM) to evaluate anxiety of the mice. The EPM consists of two open and two closed arms. Mice with higher anxiety had the tendency to stay in the closed arms because closed arms are perceived as safe zones. We determined the anxiety-related behavior by measuring the time spent in the closed arms over the 5 min test session. For the young mice, we did not find significant differences between the CR and AL groups, though CR mice had a trend of less time in the closed arms. In contrast, old AL mice spent significantly longer time in the closed arms compared to the old CR mice, indicating higher anxiety of the old AL mice. Consequently, when comparing the age-related performances, the AL group showed higher increases in anxiety (31.43% group averaged) compared to the CR mice (12.54% group averaged). Our findings are consistent with those in middle-aged rhesus monkeys, showing CR potentially alters their reaction to stressful situations. The enhanced resilience to stress in the monkeys has been linked to preserved brain volume and density in the emotional-control core and endocrine axis [65].

We further identified that anxiety level in the mice had significant and inverse correlations with CBF in hippocampus and in frontal cortex [50]. These findings indicate that preservation of CBF with age is pivotal for sustaining memory and mental health. More importantly, this positive impact on cognitive functions may also be attributed to early-life changes in neurovascular and neurometabolic functions. A recent study suggested that neuroprotective mechanisms play a major role during early stages and compensatory mechanisms in later stages of neurological diseases [66]. This is consistent with our imaging findings that CR induces early enhancements, and later preservation, on brain metabolic and vascular physiology.

## Neuroimaging Biomarkers for Human Aging and CR Interventions

The non-invasive metabolic and vascular imaging methods have been extensively applied in humans for studies related to normal cerebral physiology [49, 67], aging, and neurodegenerative disorders [29, 30]. Consistent with what we found in the animal models, metabolic and vascular functions decline were also observed in human aging brain.

Cross-sectional <sup>18</sup>F-DG-PET studies have repeatedly shown that brain glucose metabolism decreases with age in frontal cortex, anterior cingulate cortex, ventral and dorsal lateral prefrontal/inferior frontal cortex, medial prefrontal areas and precentral and perisylvian areas [68–70]. These regional changes are paralleled by global CMRglc reductions of approximately 6% per decade [71]. Interestingly, this rate of CMRglc decrease is consistent with decreased brain structural integrity and the estimated rate of decrease in synaptic density measured postmortem in healthy elderly brains [72, 73]. Reduced synaptic density is associated with cognitive impairment [74]. In line with this, longitudinal <sup>18</sup>F-DG-PET studies that monitored changes in cognitive performance in healthy aging subjects showed that CMRglc changes in the parieto-temporal cortex correlate with cognitive decline over time. Studies show that CMRglc in memory and learning-related brain regions among normal aging subjects, including hippocampus and entorhinal cortex, accurately predict future



decline from normal cognition to mild cognitive impairment (MCI) and AD [75, 76]. With a 6–14 years of follow-up scans, the rate of hippocampal CMRglc reduction in healthy elderly individuals was less than 1% per year, whereas patients who decline to MCI and AD had significantly higher rates of CMRglc reductions of 2.4–4.4% per year, respectively.

Using combined infusions of [1-<sup>13</sup>C] glucose and [2-<sup>13</sup>C] acetate with POCE, Boumezbeur *et al.* identified the difference of *in vivo* mitochondrial function between a healthy group of elderly subjects and a group of young adult controls [77]. They found that, compared with young subjects, neuronal mitochondrial metabolism (TCA cycle) and glutamate/glutamine cycle flux was 24–28% lower in elderly subjects. The reduction in individual subjects correlated strongly with reductions in NAA (a neuronal biomarker) and glutamate concentrations consistent with chronic reductions in brain mitochondrial function. In contrast, in the elderly, glial mitochondrial metabolism increased 30% compared with that of young subjects, indicating age-related changes in glial mitochondrial metabolism [78].

The decline of brain glucose metabolism in aging brain needs ketone bodies (KB) as an alternative fuel substrates to sustain brain functions. However, unlike in the condition of CR, the aging brain does not have sufficient KB production or storage to cope with the bioenergetics deficits. Instead, it uses catabolism of myelin lipids to generate KB as an adaptive response to address brain fuel and energy demand [79]. This is consistent with findings that white matter degeneration is as a pathological hallmark in aging and neurodegenerative disorders; and an increase in fatty acids and mitochondrial fatty acid metabolism machinery is coincident with a rise in brain ketone bodies and decline in plasma ketone bodies. This mechanistic pathway links mitochondrial dysfunction early in aging with later age development of white matter degeneration [80].

Neurovascular functions also decline with age. CBF of the cerebral cortex and the basal forebrain correlated significantly with advancing age [81]. Vascular pathological burden as hypoperfusion and neurodegeneration both precede and parallel cognitive decline [82]. In the elderly without dementia but with memory dysfunction, cerebral circulatory and vascular abnormalities have been suggested to contribute to MCI and AD [83]. The neurovascular risk is highly associated with accelerated decline in executive functions and cognition, including language, verbal memory, attention and visuospatial abilities [2, 3]. Mental health, such as anxiety and depression, is also link to reduced CBF [4–6], similar to our observations in animal models [50].

Neuroimaging studies indicate that rodents and humans have similar patterns for vascular and metabolic changes in brain aging. As we have seen neuroprotective effects of CR in rodents, we expect that effects would also be seen in humans. However, quantitative metabolic and vascular measurements induced by CR using neuroimaging have not been reported in human studies. Instead, functional MRI (fMRI), a qualitative determination, has been applied in humans for CR-related studies (Table 1). fMRI is a functional neuroimaging procedure using MRI technology that measures brain activity by detecting changes associated with CBF, with an assumption that CBF and brain activity are tightly coupled [84, 85].

Using fMRI and structural measurements, Witte et al. reported potential effects on memory performance in older adults using resveratrol supplement, a promising CR-mimetic nutrient [86]. Before and after the intervention/control period, they let the participants undergo memory tasks and neuroimaging to assess volume, microstructure, and functional connectivity (FC) of the hippocampus, a key region implicated in memory functions. They also assayed their anthropometry, glucose, lipid metabolism, inflammation, neurotrophic factors, and vascular parameters. They observed a significant effect of resveratrol on word retention for words over 30 min period compared with placebo. In addition, resveratrol led to significant increases in hippocampal FC and leptin levels, and decreases in glycated hemoglobin (HbA1c) and body fat compared with placebo. Increases in FC between the left posterior hippocampus and the medial prefrontal cortex correlated with increases in retention scores and with decreases in HbA1c. This study provides initial evidence that supplementary resveratrol improves memory performance in association with improved glucose metabolism and increased hippocampal FC in older adults.

In another study, using fMRI, Jakobsdottir et al. reported that CR can reverse abnormal brain activity in brain areas (e.g., amygdala) involved in processing visual food related stimuli of postmenopausal women with obesity [87]. They observed that before weight loss, increased activation of the right amygdala was seen in response to food stimuli, and free fatty acids and glucose levels correlated with activity in various areas involved in food reward processing. In the first week of CR, obese postmenopausal women had various metabolic changes occur before clinically relevant weight loss was achieved. After weight loss, fasting ghrelin and sated leptin levels correlated with activity in these areas. Activity in the amygdala region and correlations with metabolic factors differ substantially before and after weight loss.

Collectively, the fMRI studies in humans indicate that CR may have profound impact on brain activity and functional connectivity in the context of brain aging and obesity. However, whether CR has protective effects on brain metabolic and vascular functions as we saw in animal models remain unknown. In the future, it would be important to use quantitative neuroimaging methods (e.g.,  $^{18}\text{F}$ FDG-PET, POCE,  $^1\text{H}$ -MRS, and ASL) to determine the metabolic and vascular changes induced by CR, and monitor effects over time, in humans.

## Conclusion

CR has shown to have various effects in protecting brain functionality, including slowing down brain aging and reducing risks for neurodegenerative disorders. Using multi-modal, non-invasive neuroimaging methods, the beneficial effects of CR on brain metabolic and vascular functions can be detected *in vivo*. In this review, we discuss the imaging evidence of early enhancement and later preservation of metabolic and vascular functions induced by CR, as well as correlation between CBF and improved cognitive functions and reduced anxiety in animal models. As the quantitative neuroimaging methods used in animal models can also be used in human studies, they will have tremendous usefulness in future studies to identify CR and other dietary effects on brain metabolism and vasculature in humans. Understanding nutritional effects on brain function may have profound implications in human aging and age-related neurodegenerative disorders.


## References

1. Everson-Rose SA, Ryan JP. Diabetes, Obesity, and the Brain: New Developments in Biobehavioral Medicine. *Psychosom Med.* 2015; 77(6):612–5. [PubMed: 26163814]
2. Bangen KJ, et al. APOE genotype modifies the relationship between midlife vascular risk factors and later cognitive decline. *J Stroke Cerebrovasc Dis.* 2013; 22(8):1361–9. [PubMed: 23601373]
3. Zlokovic BV. Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders. *Nat Rev Neurosci.* 2011; 12(12):723–38. [PubMed: 22048062]
4. Ebmeier KP, et al. Cerebral perfusion correlates of depressed mood. *Br J Psychiatry.* 1997; 170:77–81. [PubMed: 9068780]
5. Gur RC, et al. The effect of anxiety on cortical cerebral blood flow and metabolism. *J Cereb Blood Flow Metab.* 1987; 7(2):173–7. [PubMed: 3494027]
6. Park J, Moghaddam B. Impact of anxiety on prefrontal cortex encoding of cognitive flexibility. *Neuroscience.* 2016
7. Bell RD, et al. Apolipoprotein E controls cerebrovascular integrity via cyclophilin A. *Nature.* 2012; 485(7399):512–6. [PubMed: 22622580]
8. Reiman EM, et al. Declining brain activity in cognitively normal apolipoprotein E epsilon 4 heterozygotes: A foundation for using positron emission tomography to efficiently test treatments to prevent Alzheimer's disease. *Proc Natl Acad Sci U S A.* 2001; 98(6):3334–9. [PubMed: 11248079]
9. Bookheimer SY, et al. Patterns of brain activation in people at risk for Alzheimer's disease. *N Engl J Med.* 2000; 343(7):450–6. [PubMed: 10944562]
10. Cunnane S, et al. Brain fuel metabolism, aging, and Alzheimer's disease. *Nutrition.* 2011; 27(1):3–20. [PubMed: 21035308]
11. Stranahan AM, Mattson MP. Metabolic reserve as a determinant of cognitive aging. *J Alzheimers Dis.* 2012; 30(Suppl 2):S5–13. [PubMed: 22045480]
12. Redman LM, et al. Effect of caloric restriction in non-obese humans on physiological, psychological and behavioral outcomes. *Physiol Behav.* 2008; 94(5):643–8. [PubMed: 18502453]
13. Colman RJ, et al. Caloric restriction delays disease onset and mortality in rhesus monkeys. *Science.* 2009; 325(5937):201–4. [PubMed: 19590001]
14. Choi JS, Choi KM, Lee CK. Caloric restriction improves efficiency and capacity of the mitochondrial electron transport chain in *Saccharomyces cerevisiae*. *Biochem Biophys Res Commun.* 2011; 409(2):308–14. [PubMed: 21575595]
15. Rahat O, Maoz N, Cohen HY. Multiple pathways regulating the calorie restriction response in yeast. *J Gerontol A Biol Sci Med Sci.* 2011; 66(2):163–9. [PubMed: 21081478]
16. Larson-Meyer DE, et al. Effect of calorie restriction with or without exercise on insulin sensitivity, beta-cell function, fat cell size, and ectopic lipid in overweight subjects. *Diabetes Care.* 2006; 29(6):1337–44. [PubMed: 16732018]
17. Baumeier C, et al. Caloric restriction and intermittent fasting alter hepatic lipid droplet proteome and diacylglycerol species and prevent diabetes in NZO mice. *Biochim Biophys Acta.* 2015; 1851(5):566–76. [PubMed: 25645620]
18. Thrasivoulou C, et al. Reactive oxygen species, dietary restriction and neurotrophic factors in age-related loss of myenteric neurons. *Aging Cell.* 2006; 5(3):247–57. [PubMed: 16842497]
19. Lee J, et al. Dietary restriction increases the number of newly generated neural cells, and induces BDNF expression, in the dentate gyrus of rats. *J Mol Neurosci.* 2000; 15(2):99–108. [PubMed: 11220789]
20. Lee J, Seroogy KB, Mattson MP. Dietary restriction enhances neurotrophin expression and neurogenesis in the hippocampus of adult mice. *J Neurochem.* 2002; 80(3):539–47. [PubMed: 11905999]
21. Agarwal S, et al. Caloric restriction augments ROS defense in *S. cerevisiae*, by a Sir2p independent mechanism. *Free Radic Res.* 2005; 39(1):55–62. [PubMed: 15875812]
22. Gong X, et al. Antioxidant enzyme activities in lens, liver and kidney of calorie restricted Emory mice. *Mech Ageing Dev.* 1997; 99(3):181–92. [PubMed: 9483491]

23. Merry BJ. Oxidative stress and mitochondrial function with aging--the effects of calorie restriction. *Aging Cell*. 2004; 3(1):7–12. [PubMed: 14965349]
24. Sreekumar R, et al. Effects of caloric restriction on mitochondrial function and gene transcripts in rat muscle. *Am J Physiol Endocrinol Metab*. 2002; 283(1):E38–43. [PubMed: 12067840]
25. Mattson MP, Wan R. Beneficial effects of intermittent fasting and caloric restriction on the cardiovascular and cerebrovascular systems. *J Nutr Biochem*. 2005; 16(3):129–37. [PubMed: 15741046]
26. Gomez-Pinilla F. Brain foods: the effects of nutrients on brain function. *Nat Rev Neurosci*. 2008; 9(7):568–78. [PubMed: 18568016]
27. Park SY, et al. Calorie restriction improves whole-body glucose disposal and insulin resistance in association with the increased adipocyte-specific GLUT4 expression in Otsuka Long-Evans Tokushima fatty rats. *Arch Biochem Biophys*. 2005; 436(2):276–84. [PubMed: 15797240]
28. Duan W, Ross CA. Potential therapeutic targets for neurodegenerative diseases: lessons learned from calorie restriction. *Curr Drug Targets*. 2010; 11(10):1281–92. [PubMed: 20840070]
- \*\*29. Lin AL, et al. Multimodal MRI neuroimaging biomarkers for cognitive normal adults, amnesic mild cognitive impairment, and Alzheimer's disease. *Neurol Res Int*. 2012; 2012:907409. A detailed review of using MRI neuroimaging makers to identify different patterns of brain functions between normal aging, mild cognitive impairment, and Alzheimer's Disease. [PubMed: 21949904]
- \*\*30. Lin AL, Rothman DL. What have novel imaging techniques revealed about metabolism in the aging brain? *Future Neurology*. 2014; 9(3):341–354. A comprehensive review of using multimodal neuroimaging biomarkers to identify brain metabolic changes with age. [PubMed: 25214817]
31. Lin AL, et al. Caloric restriction impedes age-related decline of mitochondrial function and neuronal activity. *J Cereb Blood Flow Metab*. 2014; 34(9):1440–3. [PubMed: 24984898]
32. Guo J, Bakshi V, Lin AL. Early Shifts of Brain Metabolism by Caloric Restriction Preserve White Matter Integrity and Long-Term Memory in Aging Mice. *Front Aging Neurosci*. 2015; 7:213. [PubMed: 26617514]
- \*33. Lin AL, et al. Caloric restriction increases ketone bodies metabolism and preserves blood flow in aging brain. *Neurobiol Aging*. 2015; 36(7):2296–303. The original neuroimaging study that shows brain metabolic shift from glucose to ketone bodies utilization in aging rats under caloric restriction. [PubMed: 25896951]
34. Perluigi M, Di Domenico F, Butterfield DA. mTOR signaling in aging and neurodegeneration: At the crossroad between metabolism dysfunction and impairment of autophagy. *Neurobiol Dis*. 2015; 84:39–49. [PubMed: 25796566]
35. Akram M. A focused review of the role of ketone bodies in health and disease. *J Med Food*. 2013; 16(11):965–7. [PubMed: 24138078]
36. Chowdhury GM, et al. The contribution of ketone bodies to basal and activity-dependent neuronal oxidation in vivo. *J Cereb Blood Flow Metab*. 2014; 34(7):1233–42. [PubMed: 24780902]
37. Katewa SD, et al. Intramyocellular fatty-acid metabolism plays a critical role in mediating responses to dietary restriction in *Drosophila melanogaster*. *Cell Metab*. 2012; 16(1):97–103. [PubMed: 22768842]
38. Lanza IR, et al. Chronic caloric restriction preserves mitochondrial function in senescence without increasing mitochondrial biogenesis. *Cell Metab*. 2012; 16(6):777–88. [PubMed: 23217257]
39. Vaishnavi SN, et al. Regional aerobic glycolysis in the human brain. *Proc Natl Acad Sci U S A*. 2010; 107(41):17757–62. [PubMed: 20837536]
40. Palmer CS, et al. Glucose metabolism regulates T cell activation, differentiation, and functions. *Front Immunol*. 2015; 6:1. [PubMed: 25657648]
41. Vlassenko AG, Raichle ME. Brain aerobic glycolysis functions and Alzheimer's disease. *Clin Transl Imaging*. 2015; 3(1):27–37. [PubMed: 26855936]
42. Vlassenko AG, et al. Spatial correlation between brain aerobic glycolysis and amyloid-beta (Abeta) deposition. *Proc Natl Acad Sci U S A*. 2010; 107(41):17763–7. [PubMed: 20837517]
43. Cirrito JR, et al. Synaptic activity regulates interstitial fluid amyloid-beta levels in vivo. *Neuron*. 2005; 48(6):913–22. [PubMed: 16364896]

44. Mouton PR, et al. Caloric restriction attenuates amyloid deposition in middle-aged dtg APP/PS1 mice. *Neurosci Lett*. 2009; 464(3):184–7. [PubMed: 19699265]
45. Schafer MJ, et al. Reduction of beta-amyloid and gamma-secretase by calorie restriction in female Tg2576 mice. *Neurobiol Aging*. 2015; 36(3):1293–302. [PubMed: 25556162]
46. Kastman EK, et al. A calorie-restricted diet decreases brain iron accumulation and preserves motor performance in old rhesus monkeys. *J Neurosci*. 2012; 32(34):11897–904. [PubMed: 23082321]
47. Wang P, Wang ZY. Metal ions influx is a double edged sword for the pathogenesis of Alzheimer's disease. *Ageing Res Rev*. 2016
48. Lin A-L, et al. Functional neuroimaging: a physiological perspective. *Frontiers in Neuroenergetics*. 2010; 2:5. [PubMed: 20725515]
49. Lin AL, et al. Nonlinear coupling between cerebral blood flow, oxygen consumption, and ATP production in human visual cortex. *Proc Natl Acad Sci U S A*. 2010; 107(18):8446–51. [PubMed: 20404151]
- \*50. Parikh I, et al. Caloric restriction preserves memory and reduces anxiety of aging mice with early enhancement of neurovascular functions. *Aging (Albany NY)*. 2016; 8(11):2814–2826. This study provides evidence that we may use neuroimaging as biomarkers to predict cognitive function and mental health in aging. [PubMed: 27829242]
51. Lin AL, et al. Chronic rapamycin restores brain vascular integrity and function through NO synthase activation and improves memory in symptomatic mice modeling Alzheimer's disease. *J Cereb Blood Flow Metab*. 2013; 33(9):1412–21. [PubMed: 23801246]
52. Cheng C, et al. Rapamycin modulates the eNOS vs. shear stress relationship. *Cardiovasc Res*. 2008; 78(1):123–9. [PubMed: 18079107]
53. Hasselbalch SG, et al. Changes in cerebral blood flow and carbohydrate metabolism during acute hyperketonemia. *Am J Physiol*. 1996; 270(5 Pt 1):E746–51. [PubMed: 8967461]
54. Roy M, et al. Long-term calorie restriction has minimal impact on brain metabolite and fatty acid profiles in aged rats on a Western-style diet. *Neurochem Int*. 2013; 63(5):450–7. [PubMed: 23974047]
55. Sengupta S, et al. mTORC1 controls fasting-induced ketogenesis and its modulation by ageing. *Nature*. 2010; 468(7327):1100–4. [PubMed: 21179166]
56. Blazquez C, et al. The AMP-activated protein kinase is involved in the regulation of ketone body production by astrocytes. *J Neurochem*. 1999; 73(4):1674–82. [PubMed: 10501215]
57. Shafique E, et al. Oxidative stress improves coronary endothelial function through activation of the pro-survival kinase AMPK. *Aging (Albany NY)*. 2013; 5(7):515–30. [PubMed: 24018842]
58. Csiszar A, et al. Caloric restriction confers persistent anti-oxidative, pro-angiogenic, and anti-inflammatory effects and promotes anti-aging miRNA expression profile in cerebrovascular endothelial cells of aged rats. *Am J Physiol Heart Circ Physiol*. 2014; 307(3):H292–306. [PubMed: 24906921]
59. Ungvari Z, et al. Mechanisms of vascular aging: new perspectives. *J Gerontol A Biol Sci Med Sci*. 2010; 65(10):1028–41. [PubMed: 20576649]
60. Lynch CD, et al. Effects of moderate caloric restriction on cortical microvascular density and local cerebral blood flow in aged rats. *Neurobiol Aging*. 1999; 20(2):191–200. [PubMed: 10537028]
61. Fontan-Lozano A, et al. Molecular bases of caloric restriction regulation of neuronal synaptic plasticity. *Mol Neurobiol*. 2008; 38(2):167–77. [PubMed: 18759009]
62. Mattson MP. The impact of dietary energy intake on cognitive aging. *Front Aging Neurosci*. 2010; 2:5. [PubMed: 20552045]
63. Valdez G, et al. Attenuation of age-related changes in mouse neuromuscular synapses by caloric restriction and exercise. *Proc Natl Acad Sci U S A*. 2010; 107(33):14863–8. [PubMed: 20679195]
64. Witte AV, et al. Caloric restriction improves memory in elderly humans. *Proc Natl Acad Sci U S A*. 2009; 106(4):1255–60. [PubMed: 19171901]
65. Willette AA, et al. Calorie restriction reduces psychological stress reactivity and its association with brain volume and microstructure in aged rhesus monkeys. *Psychoneuroendocrinology*. 2012; 37(7):903–16. [PubMed: 22119476]

66. Whalley LJ, et al. Cerebral correlates of cognitive reserve. *Psychiatry Res.* 2016; 247:65–70. [PubMed: 26774854]
67. Uh J, et al. Validation of VASO cerebral blood volume measurement with positron emission tomography. *Magn Reson Med.* 2011; 65(3):744–9. [PubMed: 21337407]
68. Herholz K, et al. Discrimination between Alzheimer dementia and controls by automated analysis of multicenter FDG PET. *Neuroimage.* 2002; 17(1):302–16. [PubMed: 12482085]
69. Hsieh TC, et al. Sex- and age-related differences in brain FDG metabolism of healthy adults: an SPM analysis. *J Neuroimaging.* 2012; 22(1):21–7. [PubMed: 21332873]
70. Kalpouzos G, et al. Voxel-based mapping of brain gray matter volume and glucose metabolism profiles in normal aging. *Neurobiol Aging.* 2009; 30(1):112–24. [PubMed: 17630048]
71. Petit-Taboue MC, et al. Effects of healthy aging on the regional cerebral metabolic rate of glucose assessed with statistical parametric mapping. *Neuroimage.* 1998; 7(3):176–84. [PubMed: 9597659]
72. Kochunov P, et al. Loss of cerebral white matter structural integrity tracks the gray matter metabolic decline in normal aging. *NeuroImage.* 2009; 45(1):17–28. [PubMed: 19095067]
73. Peters R. Ageing and the brain. *Postgrad Med J.* 2006; 82(964):84–8. [PubMed: 16461469]
74. Dawson GR, et al. Age-related cognitive deficits, impaired long-term potentiation and reduction in synaptic marker density in mice lacking the beta-amyloid precursor protein. *Neuroscience.* 1999; 90(1):1–13. [PubMed: 10188929]
75. Mosconi L. Brain glucose metabolism in the early and specific diagnosis of Alzheimer's disease. FDG-PET studies in MCI and AD. *Eur J Nucl Med Mol Imaging.* 2005; 32(4):486–510. [PubMed: 15747152]
76. Devanand DP, et al. Hippocampal and entorhinal atrophy in mild cognitive impairment: prediction of Alzheimer disease. *Neurology.* 2007; 68(11):828–36. [PubMed: 17353470]
77. Boumezbeur F, et al. Altered brain mitochondrial metabolism in healthy aging as assessed by in vivo magnetic resonance spectroscopy. *J Cereb Blood Flow Metab.* 2010; 30(1):211–21. [PubMed: 19794401]
78. Lynch AM, et al. The impact of glial activation in the aging brain. *Aging Dis.* 2010; 1(3):262–78. [PubMed: 22396865]
79. Ding F, et al. Early decline in glucose transport and metabolism precedes shift to ketogenic system in female aging and Alzheimer's mouse brain: implication for bioenergetic intervention. *PLoS One.* 2013; 8(11):e79977. [PubMed: 24244584]
80. Klosinski LP, et al. White Matter Lipids as a Ketogenic Fuel Supply in Aging Female Brain: Implications for Alzheimer's Disease. *EBioMedicine.* 2015; 2(12):1888–904. [PubMed: 26844268]
81. Tachibana H, et al. Changing topographic patterns of human cerebral blood flow with age measured by xenon CT. *AJR Am J Roentgenol.* 1984; 142(5):1027–34. [PubMed: 6609553]
82. Koike MA, et al. Long term changes in phospho-APP and tau aggregation in the 3xTg-AD mice following cerebral ischemia. *Neurosci Lett.* 2011; 495(1):55–9. [PubMed: 21419194]
83. Okonkwo OC, et al. Cerebral blood flow is diminished in asymptomatic middle-aged adults with maternal history of Alzheimer's disease. *Cereb Cortex.* 2014; 24(4):978–88. [PubMed: 23236200]
84. Kwong KK, et al. Dynamic magnetic resonance imaging of human brain activity during primary sensory stimulation. *Proc Natl Acad Sci U S A.* 1992; 89(12):5675–9. [PubMed: 1608978]
85. Ogawa S, et al. Functional brain mapping by blood oxygenation level-dependent contrast magnetic resonance imaging. A comparison of signal characteristics with a biophysical model. *Biophys J.* 1993; 64(3):803–12. [PubMed: 8386018]
86. Witte AV, et al. Effects of resveratrol on memory performance, hippocampal functional connectivity, and glucose metabolism in healthy older adults. *J Neurosci.* 2014; 34(23):7862–70. [PubMed: 24899709]
87. Jakobsdottir S, et al. Acute and short-term effects of caloric restriction on metabolic profile and brain activation in obese, postmenopausal women. *Int J Obes (Lond).* 2016; 40(11):1671–1678. [PubMed: 27216819]



**Figure 1.**

(A) The POCE results showing the correlation between mitochondrial activity in neurons ( $CMR_{glc(ox),N}$ ) and the glutamate-glutamine neurotransmission rate ( $V_{cycle}$ ). Old rats treated with CR (Old-CR; 24 months of age) showed levels of  $CMR_{glc(ox),N}$  and  $V_{cycle}$  comparable to those of the Young-AL (6 months of age). In contrast, Old-AL had significantly lower  $CMR_{glc(ox),N}$  and  $V_{cycle}$ . Published data from [31]. (B) A representative  $^1H$ -MRS spectrum, showing choline (Cho), total creatine (TCr), taurine (Tau), glutamate-glutamine complex (Glx), myo-inositol (mI), *N*-acetylaspartate (NAA), in parts per million (ppm). Published data from [32]. (C) (left)  $CMR_{glc}$  visual maps from PET  $^{18}F$ FDG imaging, showing young CR mice had significant lower  $CMR_{glc}$  compared to the young AL mice; (right) CBF visual maps from MRI-ASL imaging, showing young CR mice had significant higher CBF compared to the young AL mice. The color code indicates the  $CMR_{glc}$  and CBF in a linear scale. Published data from [32, 50]. (See description of each imaging method in Table 1).

**Table 1**

List of discussed neuroimaging methods

Imaging methods	Measurements	Quantitative vs. Qualitative
<sup>18</sup> F-DG- PET	Glucose metabolism (CMRglc)	Quantitative
POCE	Mitochondrial function; Neurotransmission rate; neuronal and glial activities	
<sup>1</sup> H-MRS	A variety of essential brain metabolites	
ASL	Cerebral blood flow (CBF)	
fMRI	Neuronal activity in relation to CBF and blood oxygenation	Qualitative

**<sup>18</sup>F-DG**: fluorine-18 (<sup>18</sup>F)-labeled 2-fluoro-2-deoxy-d-glucose; **PET**: positron emission tomography; **POCE**: <sup>1</sup>H[<sup>13</sup>C] proton-observed-carbon-edited NMR; **<sup>1</sup>H-MRS**: proton magnetic resonance spectroscopy; **ASL**: arterial spin labeling; **fMRI**: functional magnetic resonance imaging