

1-10-2019

Allogeneic Hematopoietic Cell Transplantation Provides Effective Salvage Despite Refractory Disease or Failed Prior Autologous Transplant in Angioimmunoblastic T-Cell Lymphoma: A CIBMTR Analysis

Narendranath Epperla
The Ohio State University

Kwang W. Ahn
Medical College of Wisconsin

Carlos Litovich
Medical College of Wisconsin

Sairah Ahmed
M. D. Anderson Cancer Center

Minoo Battiwalla
Sarah Cannon BMT Program


Repository Citation

Epperla, Narendranath; Ahn, Kwang W.; Litovich, Carlos; Ahmed, Sairah; Battiwalla, Minoo; Cohen, Jonathon B.; Dahi, Parastoo; Farhadfar, Noshah; Farooq, Umar; Freytes, Cesar O.; Ghosh, Nilanjan; Haverkos, Bradley; Herrera, Alex; Hertzberg, Mark; Hildebrandt, Gerhard C.; Inwards, David; Kharfan-Dabaja, Mohamed A.; Khimani, Farhad; Lazarus, Hillard; Lazaryan, Aleksandr; Lekakis, Lazaros; Murthy, Hemant; Nathan, Sunita; Nishihori, Taiga; Pawarode, Attaphol; Prestidge, Tim; Ramakrishnan, Praveen; Rezvani, Andrew R.; Romee, Rizwan; and Shah, Nirav N., "Allogeneic Hematopoietic Cell Transplantation Provides Effective Salvage Despite Refractory Disease or Failed Prior Autologous Transplant in Angioimmunoblastic T-Cell Lymphoma: A CIBMTR Analysis" (2019). *Internal Medicine Faculty Publications*. 193.
https://uknowledge.uky.edu/internalmedicine_facpub/193

See next page for additional authors

Right click to open a feedback form in a new tab to let us know how this document benefits you.

Follow this and additional works at: https://uknowledge.uky.edu/internalmedicine_facpub

 Part of the [Diseases Commons](#), [Hematology Commons](#), and the [Oncology Commons](#)

Authors

Narendranath Epperla, Kwang W. Ahn, Carlos Litovich, Sairah Ahmed, Mino Battiwalla, Jonathon B. Cohen, Parastoo Dahi, Nosha Farhadfar, Umar Farooq, Cesar O. Freytes, Nilanjan Ghosh, Bradley Haverkos, Alex Herrera, Mark Hertzberg, Gerhard C. Hildebrandt, David Inwards, Mohamed A. Kharfan-Dabaja, Farhad Khimani, Hillard Lazarus, Aleksandr Lazaryan, Lazaros Lekakis, Hemant Murthy, Sunita Nathan, Taiga Nishihori, Attaphol Pawarode, Tim Prestidge, Praveen Ramakrishnan, Andrew R. Rezvani, Rizwan Romee, and Nirav N. Shah

Allogeneic Hematopoietic Cell Transplantation Provides Effective Salvage Despite Refractory Disease or Failed Prior Autologous Transplant in Angioimmunoblastic T-Cell Lymphoma: A CIBMTR Analysis**Notes/Citation Information**

Published in *Journal of Hematology & Oncology*, v. 12, article no. 6, p. 1-11.

© The Author(s). 2019

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

Due to the large number of authors, only the first 30 and the authors affiliated with the University of Kentucky are listed in the author section above. For the complete list of authors, please download this article or visit: <https://doi.org/10.1186/s13045-018-0696-z>

Digital Object Identifier (DOI)

<https://doi.org/10.1186/s13045-018-0696-z>

RESEARCH

Open Access



Allogeneic hematopoietic cell transplantation provides effective salvage despite refractory disease or failed prior autologous transplant in angioimmunoblastic T-cell lymphoma: a CIBMTR analysis

Narendranath Epperla¹, Kwang W. Ahn², Carlos Litovich², Sairah Ahmed³, Minoo Battiwalla⁴, Jonathon B. Cohen⁵, Parastoo Dahi⁶, Nosha Farhadfar⁷, Umar Farooq⁸, Cesar O. Freytes⁹, Nilanjan Ghosh¹⁰, Bradley Haverkos¹¹, Alex Herrera¹², Mark Hertzberg¹³, Gerhard Hildebrandt¹⁴, David Inwards¹⁵, Mohamed A. Kharfan-Dabaja¹⁶, Farhad Khimani¹⁷, Hillard Lazarus^{18,19}, Aleksandr Lazaryan¹⁷, Lazaros Lekakis^{18,19}, Hemant Murthy²⁰, Sunita Nathan²¹, Taiga Nishihori¹⁷, Attaphol Pawarode²², Tim Prestidge²³, Praveen Ramakrishnan²⁴, Andrew R. Rezvani²⁵, Rizwan Romee²⁶, Nirav N. Shah²⁷, Ana Sureda²⁸, Timothy S. Fenske²⁷ and Mehdi Hamadani^{2,27*}

Abstract

Background: There is a paucity of data on the role of allogeneic hematopoietic cell transplantation (allo-HCT) in patients with angioimmunoblastic T-cell lymphoma (AITL). Using the CIBMTR registry, we report here the outcomes of AITL patients undergoing an allo-HCT.

Methods: We evaluated 249 adult AITL patients who received their first allo-HCT during 2000–2016.

(Continued on next page)

* Correspondence: mhamadani@mcw.edu

²Center for International Blood and Marrow Transplant Research, Department of Medicine, Medical College of Wisconsin, 9200 W. Wisconsin Avenue, Suite C5500, 8701 W. Watertown Plank Rd, Milwaukee, WI 53226, USA

²⁷Division of Hematology and Oncology, Department of Medicine, Medical College of Wisconsin, 8701 Watertown Plank Rd. PO Box 26509, Milwaukee, WI 53226, USA

Full list of author information is available at the end of the article



(Continued from previous page)

Results: The median patient age was 56 years (range = 21–77). Majority of the patients were Caucasians (86%), with a male predominance (60%). Graft-versus-host disease (GVHD) prophylaxis was predominantly calcineurin inhibitor-based approaches while the most common graft source was peripheral blood (97%). Median follow-up of survivors was 49 months (range = 4–170 months). The cumulative incidence of grade 2–4 and grade 3–4 acute GVHD at day 180 were 36% (95% CI = 30–42) and 12 (95% CI = 8–17), respectively. The cumulative incidence of chronic GVHD at 1 year was 49% (95%CI 43–56). The 1-year non-relapse mortality (NRM) was 19% (95% CI = 14–24), while the 4-year relapse/progression, progression-free survival (PFS), and overall survival (OS) were 21% (95% CI = 16–27), 49% (95% CI = 42–56), and 56% (95% CI = 49–63), respectively. On multivariate analysis, chemoresistant status at the time of allo-HCT was associated with a significantly higher risk for therapy failure (inverse of PFS) (RR = 1.73 95% CI = 1.08–2.77), while KPS < 90% was associated with a significantly higher risk of mortality (inverse of OS) (RR = 3.46 95% CI = 1.75–6.87).

Conclusion: Our analysis shows that allo-HCT provides durable disease control even in AITL patients who failed a prior auto-HCT and in those subjects with refractory disease at the time of allografting.

Keywords: Angioimmunoblastic T-cell lymphoma, Allogeneic transplantation, GVL effects,

Background

Angioimmunoblastic T-cell lymphoma (AITL) represents a distinct clinicopathologic entity among the mature T- and NK-cell neoplasms, accounting for approximately 1–2% of all non-Hodgkin lymphomas (NHLs) [1, 2]. AITL patients typically present with advanced stage disease, diffuse lymphadenopathy, hepatosplenomegaly, systemic symptoms, and hypergammaglobulinemia [3]. The clinical course is aggressive and the disease generally carries a poor prognosis even when treated with intensive induction regimens [3]. Standard first-line therapy mostly consists of anthracycline-based regimens with or without etoposide, based on the age [2, 4–6]. With this approach, overall survival (OS) is a little over 30% at 5 years [7]. In an attempt to improve the outcomes, autologous hematopoietic cell transplantation (auto-HCT) consolidation has been applied in this patient population [8–10]. While durable disease control can be observed typically in patients in first complete remission (CR), the outcomes of AITL subjects in partial remission (PR), and in those with refractory disease or treated with ≥ 2 prior therapy lines, following auto-HCT are less encouraging [10].

Allogeneic HCT (allo-HCT) may result in a lower risk of relapse in part due to a *graft-versus-lymphoma* effect mediated by the alloreactive donor cells [11–13]. Several retrospective studies [11, 14–16] have reported excellent disease control with low rates of relapse and a 1-year non-relapse mortality (NRM) ranging from 8 to 25% with allo-HCT in AITL patients. However, these analyses were done mainly in peripheral T-cell lymphoma (PTCL) patients with AITL as a subgroup or reported only a small number of patients with AITL (range $N = 9$ –45 patients; Additional file 1: Table S1). We report here a registry analysis, evaluating the outcomes of patients with AITL undergoing allo-HCT.

Methods

Data sources

The Center for International Blood and Marrow Transplant Research (CIBMTR) is a working group of more than 500 transplantation centers worldwide that contribute detailed data on HCT to a statistical center at the Medical College of Wisconsin (MCW). Participating centers are required to report all transplantations consecutively and compliance is monitored by on-site audits. Computerized checks for discrepancies, physicians' review of submitted data, and on-site audits of participating centers ensure data quality. Observational studies conducted by the CIBMTR are performed in compliance with all applicable federal regulations pertaining to the protection of human research participants. The MCW and National Marrow Donor Program, Institutional Review Boards approved this study.

The CIBMTR collects data at two levels: transplant essential data (TED) and comprehensive report form (CRF) data. TED data includes disease type, age, gender, pre-HCT disease stage and chemotherapy-responsiveness, date of diagnosis, graft type, conditioning regimen, post-transplant disease progression and survival, development of a new malignancy, and cause of death. All CIBMTR centers contribute to TED data. More detailed disease and pre- and post-transplant clinical information is collected on a subset of registered patients selected for CRF data by a weighted randomization scheme. TED- and CRF-level data are collected pre-transplant, 100-days, and 6 months post-HCT and annually thereafter or until death. Data for the current analysis were retrieved from CIBMTR (TED and CRF) report forms.

Patients

Included in this analysis are adult (≥ 18 years) patients with AITL, undergoing their first allo-HCT between 2000 and 2016. Eligible donors included either HLA-identical

Table 1 Baseline patient characteristics of patients with AITL receiving first allo-HCT reported to the CIBMTR from 2000 to 2016

Variable	N = 249 (%)
Median age at HCT, years (range)	56 (21–77)
Male gender	150 (60)
Race	
Caucasian	214 (86)
African American	5 (2)
Others ^a	17 (7)
Missing	13 (5)
Karnofsky performance score ≥ 90	119 (48)
< 90	113 (45)
Missing	17 (7)
HCT-CI	
0	46 (18)
1–2	53 (21)
≥ 3	84 (34)
Not available before 2007	55 (22)
Missing	11 (4)
Interval from diagnosis to HCT, months	
Median (range)	14 (3–118)
Median lines of therapy before HCT (range)	3 (1–5)
Remission status at HCT	
Complete remission	108 (43)
Partial remission	90 (36)
Chemorefractory	38 (15)
Untreated/unknown	13 (5)
History of prior autologous HCT	98 (39)
TBI in conditioning	83 (34)
ATG/alemtuzumab in conditioning ^b	59 (24)
Conditioning intensity ^c	
Myeloablative conditioning	66 (27)
Non-myeloablative/RIC	183 (73)
Graft source	
Bone marrow	8 (3)
Peripheral blood	241 (97)
Donor type	
HLA-identical sibling	140 (56)
Unrelated donor 8/8	109 (44)
Donor/recipient CMV status	
Both negative	72 (29)
Both positive	59 (24)
Either donor/recipient +	69 (28)

Table 1 Baseline patient characteristics of patients with AITL receiving first allo-HCT reported to the CIBMTR from 2000 to 2016 (Continued)

Variable	N = 249 (%)
Missing	49 (19)
Graft-versus-host disease prophylaxis	
Calcineurin inhibitor + MTX ± others ^d (except MMF)	119 (48)
Calcineurin inhibitor + MMF ± others ^d	76 (31)
Calcineurin inhibitor + others (except MMF)	40 (16)
Others ^d	10 (4)
Missing	4 (2)
Year of HCT	
2000–2006	47 (19)
2007–2011	82 (33)
2012–2016	120 (48)
Median follow-up of survivors (range), months	49 (4–170)

ATG antithymocyte globulin, CMV cytomegalovirus, HCT hematopoietic cell transplantation, HCT-CI HCT-Comorbidity index, MMF mycophenolate mofetil, MTX methotrexate, TBI total body irradiation, RIC reduced intensity conditioning

^aOthers: 13 Asian; 3 Hispanic or Latino; 1 race unspecified, non-Hispanic

^bATG/alemtuzumab—49 ATG alone; 10 alemtuzumab alone

^cFor details, refer to Additional file 1: Table S4

^dFor details, refer to Additional file 1: Table S5

sibling donors or unrelated donors (URD) matched at the allele-level at HLA-A, -B, -C, and -DRB1 and graft sources included peripheral blood and bone marrow. Graft-versus-host disease (GVHD) prophylaxis included both calcineurin inhibitor (CNI) and non-CNI-based regimens. Recipients of alternative donor transplantation were excluded due to small numbers (haploidentical allografts, $n = 8$; mismatched unrelated donor, $n = 22$; cord blood grafts, $n = 21$).

Definitions and study endpoints

The intensity of conditioning regimens was defined using consensus criteria [17]. Disease response at the time of HCT was determined using the International Working Group criteria in use during the era of this analysis [18].

The primary endpoint was OS; death from any cause was considered an event and surviving patients were censored at last contact. Secondary endpoints included cumulative incidence of acute GVHD, chronic GVHD, GVHD free, relapse-free survival (GRFS), NRM, progression/relapse, and progression-free survival (PFS). NRM was defined as death without evidence of lymphoma progression/relapse; relapse was considered a competing risk. Progression/relapse was defined as progressive lymphoma after HCT or lymphoma recurrence after a CR; NRM was considered a competing risk. For PFS, a patient was considered treatment failure at the time of progression/relapse or death from any cause. Patients alive without evidence of disease relapse or progression were censored at last follow-up.

Table 2 Univariate Analysis

Outcomes	N Eval	Prob (95% CI)
Neutrophil engraftment	236	
1-year		97 (94–99)%
2-year		97 (94–99)%
Platelet recovery	218	
1-year		91 (87–94)%
2-year		91 (87–95)%
Acute GVHD (II-IV)	239	
180-day		36 (30–42)%
Acute GVHD (III-IV)	229	
180-day		12 (8–17)%
Chronic GVHD	230	
1-year		49 (43–56)%
2-year		58 (51–64)%
Extensive cGVHD	230	
1-year		39 (33–46)%
2-year		46 (39–53)%
GRFS	230	
1-year		35 (29–41)%
2-year		27 (21–33)%
NRM	249	
1-year		19 (14–24)%
2-year		25 (20–31)%
4-year		30 (24–36)%
Progression/relapse	249	
1-year		15 (11–20)%
2-year		19 (15–25)%
4-year		21 (16–27)%
PFS	249	
1-year		66 (60–72)%
2-year		56 (49–62)%
4-year		47 (41–54)%
Overall survival	249	
1-year		73 (68–79)%
2-year		63 (56–69)%
4-year		56 (49–63)%

GVHD graft-versus-host disease, Prob probability, CI confidence interval, N number, NRM non-relapse mortality, PFS progression-free survival, GRFS GVHD free, relapse-free survival

Probabilities of acute GVHD, chronic GVHD, treatment-related mortality and progression/relapse were calculated using the cumulative incidence estimate. Progression-free survival and overall survival was calculated using the Kaplan-Meier product limit estimate

Univariate analysis of alternative donor sources is shown in Additional file 1 Table S6

Acute GVHD [19] and chronic GVHD [20] were graded using standard criteria. Neutrophil recovery was defined as the first of three successive days with absolute neutrophil

Table 3 Multivariate analysis results

	Number	RR	95% CI lower limit	95% CI upper limit	P-value	Overall p value
Chronic GVHD						
ATG/alemtuzumab						
No	174	1				0.02
Yes	55	0.58	0.36	0.93	0.02	
Progression/Relapse						
No significant covariates						
Non-relapse mortality						
No significant covariates						
Progression-free survival						
Disease status						
CR	108	1				0.03
PR	90	1.13	0.76	1.66	0.54	
Chemoresistant	38	1.73	1.08	2.77	0.02	
Missing/Untreated	13	0.43	0.15	1.20	0.11	
Overall survival						
Karnofsky performance score (≤ 6 months) ^a						
$\geq 90\%$	119	1				0.002
< 90%	113	3.46	1.74	6.87	0.0004	
Missing	17	1.95	0.54	6.98	0.31	
Karnofsky performance score (> 6 months) ^a						
$\geq 90\%$	106	1				0.28
< 90%	80	0.66	0.39	1.12	0.12	
Missing	14	0.73	0.29	1.86	0.51	

GVHD graft-versus-host disease, CI confidence interval, ATG anti-thymocyte globulin, CR complete remission, PR partial remission, RR relative risk

Variables tested in the Multivariate analysis are listed in Additional file 1 Table S2


^a6-months was chosen as cut-off based on the maximum likelihood value in the Cox model

p-value < 0.05 is considered significant

count (ANC) $\geq 500/\mu\text{L}$ after post-transplantation nadir. Platelet recovery was defined as achieving platelet counts $\geq 20,000/\mu\text{L}$ for at least 3 days, unsupported by transfusion. For neutrophil and platelet recovery, death without the event was considered a competing risk. The causes of death are reported in accordance to the methodology described previously [21].

Statistical analysis

Probabilities of PFS and OS were calculated using the Kaplan–Meier estimates. Cumulative incidence of NRM, lymphoma progression/relapse, and GVHD were calculated to accommodate for competing risks. Associations among patient-, disease-, and transplantation-related variables and outcomes of interest were evaluated using Cox proportional hazards regression. A stepwise model-building approach



was used to identify covariates that influenced outcomes. Covariates with a $p < 0.05$ were considered statistically significant. The proportional hazards assumption for Cox regression was tested by adding a time-dependent covariate for each risk factor and each outcome. If a variable violated the proportional hazards assumption, it was added as a time-varying covariate. Interactions between the main effect and significant covariates were examined and none were found. Results are expressed as relative risks (RR). The center effect was examined using the random effect score test [22] for OS, PFS, relapse, and NRM. The variables considered in multivariate analysis are shown in Additional file 1: Table S2 of the supplemental appendix. All statistical analyses were performed using SAS version 9.4 (SAS Institute Inc., Cary, NC).

Results

Baseline characteristics

A total of 249 patients met the inclusion criteria and were included in this analysis. The baseline patient-, disease-, and transplantation-related characteristics are shown in Table 1. The median patient age was 56 years (range = 21–77 years). Most of the patients were Caucasians (86%), with

a male (60%) predominance. The majority had a chemosensitive disease at the time of allo-HCT (79%) and received a non-myeloablative/reduced intensity conditioning regimen (73%). Most common type of GVHD prophylaxis included CNI ± methotrexate-based regimens. The graft source used for allo-HCT was predominantly peripheral blood (97%). Pre-transplant (allo-HCT) donor/recipient cytomegalovirus status was available in 200 patients (81%) and the details are provided in Table 1. There was no center effect noted on the outcomes. Median follow-up of survivors was 49 months (range, 4–170 months).

Hematopoietic recovery

On univariate analysis, the cumulative incidence of neutrophil engraftment at 1-year was 97% (95% CI 94–99). The 1-year cumulative incidence of platelet recovery (Table 2) was 91% (95% CI 87–94).

Acute and chronic GVHD

On univariate analysis, the cumulative incidence of grade II–IV acute GVHD was 36% (95% CI 30–42) and grades III–IV acute GVHD was 12% (95% CI 8–17) at day 180 (Table

Table 4 Comparative analysis of AITL patients who received prior auto-HCT vs no prior auto-HCT

Outcomes	No prior auto-HCT (N = 151)		Prior auto-HCT (N = 98)		p value
	N	Prob (95% CI)	N	Prob (95% CI)	
NRM	151		98		0.25
1-year		17 (11–23)%		22 (14–30)%	0.33
2-year		21 (15–28)%		31 (22–41)%	0.08
3-year		22 (16–29)%		33 (23–43)%	0.07
4-year		26 (19–34)%		36 (26–47)%	0.11
Progression/relapse	151		98		0.69
1-year		16 (11–22)%		15 (8–22)%	0.77
2-year		22 (15–29)%		16 (9–24)%	0.23
3-year		23 (16–30)%		17 (10–25)%	0.28
4-year		24 (17–31)%		17 (10–25)%	0.21
PFS	151		98		0.45
1-year		68 (60–75)%		64 (54–73)%	0.56
2-year		57 (49–65)%		53 (43–63)%	0.53
3-year		55 (47–64)%		50 (40–61)%	0.43
4-year		50 (42–59)%		47 (36–57)%	0.60
Overall survival	151		98		0.81
1-year		73 (65–80)%		74 (65–82)%	0.81
2-year		65 (57–72)%		59 (49–69)%	0.43
3-year		61 (53–69)%		58 (47–68)%	0.63
4-year		57 (49–65)%		54 (44–65)%	0.70

Prob probability, CI confidence interval, N number, NRM non-relapse mortality, PFS progression-free survival, HCT hematopoietic cell transplantation

2). None of the tested covariates (Additional file 1: Table S2) affected the risk of the development of acute GVHD.

On univariate analysis, the cumulative incidence of chronic GVHD at 1-year (Table 2) was 49% (95% CI 43–56), while the cumulative incidence of extensive chronic GVHD at 1 year (Table 2) was 39% (95% CI 33–46). Multivariate analysis (Table 3) showed that patients who received anti-thymocyte globulin (ATG) or alemtuzumab had a significantly lower risk of chronic GVHD (RR = 0.58, 95% CI 0.36–0.93, $p = 0.02$) relative to those who did not receive ATG/alemtuzumab.

Transplantation outcomes

On univariate analysis, the cumulative incidence of 1-year GRFS (Table 2) was 35% (95% CI 29–41).

The 1-year NRM rate (Table 2) was 19% (95% CI 14–24) (Fig. 1a). On multivariate analysis, there were no significant covariates affecting the risk of NRM. The cumulative incidence of progression/relapse at 4 years (Table 2) was 21% (95% CI 16–27) (Fig. 1b). On multivariate analysis (Table 3), none of the covariates (Additional file 1: Table S1, including chronic GVHD assessed as a time-dependent variable) significantly affected the relapse risk.

The 4-year PFS and OS (Table 2) were 47% (95% CI 41–54) (Fig. 1c) and 56% (95% CI 49–63) (Fig. 1d), respectively.

On multivariate analysis (Table 3), chemoresistant status at the time of allo-HCT significantly increased the risk for therapy failure (inverse of PFS) (RR = 1.73 95% CI = 1.08–2.77, $p = 0.02$), while KPS < 90% was associated with a significantly higher risk of mortality (inverse of OS) in the first 6-months post allo-HCT (RR = 3.46 95% CI = 1.74–6.87, $p = 0.0004$).

Causes of death

At last follow-up, 45% ($n = 112$) of allo-HCT recipients had died (Additional file 1: Table S3). The most common cause of death was organ failure, 20% ($n = 22$) followed by recurrent/progressive disease, 19% ($n = 21$). GVHD was the cause of death in 17% ($n = 19$) and infectious complications accounted for death in 15% ($n = 17$) of patients. The other causes of death are listed in Additional file 1: Table S3.

Impact of prior autograft and disease status

Among the 249 patients who received first allo-HCT, 98 patients (39%) had received a prior auto-HCT. Univariate analysis looking at the impact of prior auto-HCT (no prior auto-HCT vs prior auto-HCT) on the outcomes showed no significant difference in the 1-year NRM (17% [95% CI 11–23] vs 22% [95% CI 14–30], $p = 0.33$), 4-year progression/relapse (24% [95% CI 17–31] vs 17% [95% CI

10–25], $p = 0.21$), PFS (50% [95% CI 42–59] vs 47% [95% CI 36–57], $p = 0.60$), or OS (57% [95% CI 49–65] vs 54% [95% CI 44–65], $p = 0.70$) (Table 4, Fig. 2).

Among the 198 patients with chemosensitive disease at the time of allo-HCT, 33 patients (17%) were in CR1, while 75 patients (38%) were in CR > 1 and 90 patients (45%) were in PR. Univariate analysis looking at the effect of remission status at allo-HCT, CR1 vs CR > 1 vs PR vs refractory (Table 5), showed a 4-year PFS of 58% vs 45% vs 47% vs 38%, respectively, and a 4-year OS of 70% vs 54% vs 50% vs 52%, respectively. Among patients with chemorefractory AITL, the 1-year NRM was 24%, while the 4-year progression/relapse, PFS, and OS in patients with refractory AITL were 32%, 38%, and 52%, respectively. Figure 3 shows the disease outcomes for AITL patients based on the remission status at allo-HCT (CR vs PR vs chemoresistant).

Discussion

Prospective studies evaluating the outcomes of allo-HCT exclusively in AITL have not been performed given an

overall rarity of this PTCL subtype. Here, we performed a registry analysis of AITL patients receiving first allo-HCT and made several important observations. First, allo-HCT provided durable disease control in patients with AITL as evidenced by 4-year PFS of 47%. Second, the risk of relapse tended to plateau at 2-year post allo-HCT. Lastly, allo-HCT provided durable disease control even in patients with a failed prior auto-HCT and those subjects with refractory disease at the time of allografting.

Auto-HCT has been previously studied as a consolidation modality for patients with AITL in first CR and beyond. While auto-HCT can provide durable disease control in AITL subjects in CR1, the outcomes of patients not in CR, or those with heavily pretreated disease are not optimal [10]. In addition, despite low transplant-related mortality, the risk of relapse following autografting remains high (1- and 2-year relapse risk is 40% and 51%, respectively) [10]. In contrast, allo-HCT provides excellent survival outcomes for patients with AITL with a lower risk of relapse. Additional file 1: Table S1 summarizes the retrospective studies ($n \geq 9$) that have looked at the role of




Table 5 Comparative analysis of AITL patients based on the remission status at the time of allo-HCT

Outcomes	CR1 (N = 33)		CR > 1 (N = 75)		PR (N = 90)		Refractory (N = 38)	
	N	Prob (95% CI)	N	Prob (95% CI)	N	Prob (95% CI)	N	Prob (95% CI)
NRM	33		75		90		38	
1-year		6 (1–17)%		20 (12–30)%		20 (13–29)%		24 (12–38)%
2-year		13 (4–26)%		29 (19–40)%		25 (17–35)%		30 (16–45)%
3-year		17 (6–32)%		31 (21–43)%		25 (17–35)%		30 (16–45)%
4-year		17 (6–32)%		36 (25–49)%		33 (22–44)%		30 (16–45)%
Progression/ relapse	33		75		90		38	
1-year		15 (5–29)%		13 (7–22)%		14 (7–21)%		29 (16–44)%
2-year		25 (12–41)%		16 (9–26)%		19 (11–28)%		29 (16–44)%
3-year		25 (12–41)%		18 (10–28)%		19 (11–28)%		32 (18–48)%
4-year		25 (12–41)%		18 (10–28)%		21 (12–30)%		32 (18–48)%
PFS	33		75		90		38	
1-year		79 (63–91)%		67 (56–77)%		66 (56–76)%		47 (32–63)%
2-year		62 (45–78)%		54 (43–66)%		56 (45–66)%		41 (26–57)%
3-year		58 (41–75)%		50 (38–62)%		56 (45–66)%		38 (23–54)%
4-year		58 (41–75)%		45 (33–58)%		47 (36–58)%		38 (23–54)%
Overall survival	33		75		90		38	
1-year		88 (75–97)%		73 (63–83)%		71 (61–80)%		63 (47–78)%
2-year		78 (62–90)%		62 (51–73)%		59 (48–69)%		52 (36–67)%
3-year		70 (52–85)%		58 (46–70)%		57 (47–68)%		52 (36–67)%
4-year		70 (52–85)%		54 (41–66)%		50 (39–62)%		52 (36–67)%


CR complete response, PR partial response, Prob probability, CI confidence interval, N number, NRM non-relapse mortality, PFS progression-free survival

allo-HCT in AITL [11, 14–16]. The current study is the largest registry validation of these results showing durable responses in patients with AITL following allo-HCT. Though previous studies included patients with prior auto-HCT failure and chemorefractory state, the data are limited by very small patient numbers (for example, the previously published study with a large number of AITL patients [$n = 45$] included 15 patients with prior auto-HCT failure and 18 patients with chemorefractory disease at allo-HCT) [14] limiting the ability to draw meaningful conclusions. Considering the fact that ASBMT Clinical Practice Recommendation Panel [23] endorses the use of auto-HCT in AITL patients in CR1/PR1, and the high rates of disease relapse in patients receiving high-dose therapy, addressing the role of a subsequent allo-HCT is a clinically important question. In the current analysis, we did not observe any statistically significant differences in outcomes for patients who had prior auto-HCT vs no prior auto-HCT. Our results support the curative potential of allo-HCT in high-risk AITL patients who have failed a prior auto-HCT.

Limited data are published on the role of allo-HCT in refractory AITL. Registry data from the European Society for Blood and Marrow Transplantation (EBMT) identified chemorefractory disease as a predictor of

inferior outcomes but included only 18 refractory AITL patients [14]. In the current analysis, the 4-year PFS and OS of chemorefractory patients was 38% and 52% respectively, which supports the use of allografting in this ultra-high-risk subset of patients (who otherwise are fit to undergo allo-HCT). In our study, we did not find a relationship between chronic GVHD and relapse rate in contrast to the previously reported data [14]. The retrospective nature of the registry data does not permit us to analyze the optimal timing of allo-HCT. While the outcomes of CR1 patients in the current study were favorable (4-year PFS and OS 58% and 70%), prior studies have also suggested very encouraging outcomes of AITL patients undergoing auto-HCT in CR1 [10, 24].

AITL is a challenging diagnosis with roughly only 80% concordance even among expert pathologists with access to archival tissue [3, 7]. One of the limitations of the current study is the lack of central pathology review of archival tissue for all patients. The current study included cases as diagnosed by the pathologists at the respective institutions. Of note, disease histology is one of the critical fields CIBMTR examines during its onsite transplant center audits (where diagnosis reported to CIBMTR is audited relative to the pathology records available at the reporting center). In recent CIBMTR



studies involving rare T-cell histologies, >95% concordance was seen between center-reported diagnosis and central review of pathology reports [25, 26]. We acknowledge that this analysis is not a substitute of central review of archival tissue by expert pathologists. At the same time, it is important to note that the majority of prospective clinical trials enrolling AITL subjects accept the patients based on the pathology reports at the participating sites, without a mandatory central review of archival tissue. In addition, the CIBMTR registry does not capture post-relapse salvage therapy, thereby limiting the ability to assess the post-allo-HCT relapse survival.

Conclusions

With a better understanding of the biology and development of prognostic tools, there has been a major effort to study novel drug combinations and immunotherapy agents (including checkpoint inhibitors and chimeric antigen receptor T-cell [CAR-T] therapy) in patients with NHL. Brentuximab vedotin (anti-CD30 antibody-drug conjugate) is being studied in combination with chemotherapy in the

frontline setting in PTCL patients (ECHELON 2 trial, NCT 01777152). The final results are eagerly awaited to assess the impact of CD30-directed therapies in the subset of AITL patients. While the data on CAR-T cell therapy for B-cell NHL (mainly diffuse large B-cell lymphoma) in the relapsed/refractory setting is impressive [27], similar constructs in T-cell NHL have not been translated to the bedside. Our results suggest that allo-HCT offers the potential for cure in AITL patients including those with otherwise chemo-refractory disease. In the foreseeable future, allo-HCT is likely to remain an important therapeutic option for AITL patients.

Additional file

Additional file 1: Table S1. Outcomes of patients with AITL who underwent allogeneic HCT. **Table S2.** Variables tested in Cox proportional hazards regression models. **Table S3.** Causes of Death. **Table S4.** Conditioning Intensity. **Table S5.** Details of GVHD prophylaxis regimens. **Table S6.** Univariate outcomes of AITL patients receiving alternative donor sources. (DOCX 29 kb)

Abbreviations

AITL: Angioimmunoblastic T-cell lymphoma; Allo-HCT: Allogeneic hematopoietic cell transplantation; CNi: Calcineurin inhibitor; CR: Complete remission; CRF: Comprehensive Report Form; GRFS: GVHD free, relapse-free survival; GVHD: Graft-versus-host disease; NRM: Non-relapse mortality; OS: Overall survival; PFS: Progression-free survival; PR: Partial remission; RR: Relative risk; TED: Transplant essential data

Acknowledgments

CIBMTR Support List
The CIBMTR is supported by Public Health Service Grant/Cooperative Agreement U24-CA076518 from the National Cancer Institute (NCI), the National Heart, Lung and Blood Institute (NHLBI), and the National Institute of Allergy and Infectious Diseases (NIAID); a Grant/Cooperative Agreement 5U10HL069294 from NHLBI and NCI; a contract HSH250201200016C with Health Resources and Services Administration (HRSA/DHHS); two Grants N00014-13-1-0039 and N00014-14-1-0028 from the Office of Naval Research; and grants from *Actinium Pharmaceuticals; Allos Therapeutics, Inc.; *Amgen, Inc.; Anonymous donation to the Medical College of Wisconsin; Ariad; Be the Match Foundation; *Blue Cross and Blue Shield Association; *Celgene Corporation; Chimerix, Inc.; Fred Hutchinson Cancer Research Center; Fresenius-Biotech North America, Inc.; *Gamida Cell Teva Joint Venture Ltd.; Genentech, Inc.; *Gentium SpA; Genzyme Corporation; GlaxoSmithKline; Health Research, Inc. Roswell Park Cancer Institute; HistoGenetics, Inc.; Incyte Corporation; Jeff Gordon Children's Foundation; Kiadis Pharma; The Leukemia & Lymphoma Society; Medac GmbH; The Medical College of Wisconsin; Merck & Co, Inc.; Millennium: The Takeda Oncology Co.; *Milliman USA, Inc.; *Miltenyi Biotec, Inc.; National Marrow Donor Program; Onyx Pharmaceuticals; Optum Healthcare Solutions, Inc.; Osiris Therapeutics, Inc.; Otsuka America Pharmaceutical, Inc.; Perkin Elmer, Inc.; *Remedy Informatics; *Sanofi US; Seattle Genetics; Sigma-Tau Pharmaceuticals; Soligenix, Inc.; St. Baldrick's Foundation; StemCyte, A Global Cord Blood Therapeutics Co.; Stemsoft Software, Inc.; Swedish Orphan Biovitrum; *Tarix Pharmaceuticals; *TerumoBCT; *Teva Neuroscience, Inc.; *THERAKOS, Inc.; University of Minnesota; University of Utah; and *Wellpoint, Inc. The views expressed in this article do not reflect the official policy or position of the National Institute of Health, the Department of the Navy, the Department of Defense, Health Resources and Services Administration (HRSA), or any other agency of the U.S. Government.
*Corporate Members.
Morgan Geronimo for administrative Support.

Funding

Not applicable.

Availability of data and materials

Please contact author for data requests.

Authors' contributions

Collection and assembly of data: NE, CL, and MH. Data analysis: KWA, NE, MH, and CL. Data interpretation: All authors. Manuscript writing: First draft prepared by NE and MH. All authors helped revise the manuscript. Final approval of manuscript: All authors.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Division of Hematology, Department of Medicine, The James Cancer Hospital and Solove Research Institute, The Ohio State University, 460 W 10th Ave, Columbus, OH 43210, USA. ²Center for International Blood and Marrow Transplant Research, Department of Medicine, Medical College of Wisconsin, 9200 W. Wisconsin Avenue, Suite C5500, 8701 W. Watertown Plank Rd,

Milwaukee, WI 53226, USA. ³M.D. Anderson Cancer Center, 1515 Holcombe Boulevard, Houston, TX 77030, USA. ⁴Sarah Cannon BMT Program, 2400 Patterson St. Suite 215, Nashville, TN 37206, USA. ⁵Winship Cancer Institute, Emory University School of Medicine, 1365-C Clifton Road NE, Atlanta, GA 30322, USA. ⁶Memorial Sloan Kettering Cancer Center, 1275 York Ave., New York, NY 10065, USA. ⁷Shands Healthcare and University of Florida, PO Box 100278, Gainesville, FL 32610, USA. ⁸University of Iowa Hospitals and Clinics, 200 Hawkins Drive C332 GH, Iowa City, IA 52242, USA. ⁹Texas Transplant Institute, 4410 Medical Drive Suite 410, San Antonio, TX 78229, USA. ¹⁰Levine Cancer Institute, 1021 Morehead Medical Drive Suite 5300, Charlotte, NC 28204, USA. ¹¹University of Colorado Hospital, 1665 Aurora Court F-754, Aurora, CO 80045, USA. ¹²City of Hope National Medical Center, 1500 E Duarte Rd, Duarte, CA 91010, USA. ¹³Prince of Wales Hospital, SEALS Level 4 Campus Building, Barker Street, Randwick, NSW 2031, Australia. ¹⁴University of Kentucky Chandler Medical Center, 800 Rose Street CC 301, Lexington, KY 40536, USA. ¹⁵Mayo Clinic Rochester, 200 First Street SW, Rochester, MN 55902, USA. ¹⁶Mayo Clinic, 4500 San Pablo Rd, Jacksonvile, FL 32224, USA. ¹⁷H. Lee Moffitt Cancer Center and Research Institute, 12902 Magnolia Drive, Tampa, FL 33612, USA. ¹⁸Case Western Reserve University, 11100 Euclid Ave, Cleveland, OH 44106, USA. ¹⁹University of Miami, 1475 NW 12th Ave, Miami, FL 33136, USA. ²⁰Division of Hematology/Oncology, University Florida College of Medicine, 12902 Magnolia Drive, Tampa, FL 33612, USA. ²¹Rush University Medical Center, 849 North Franklin Street Unit 1503, Chicago, IL 60610, USA. ²²The University of Michigan, 322 E Liberty St. Unit 4, Ann Arbor, MI 48104, USA. ²³Starship Children's Health, Level 7 Blood and Cancer Center Park Road, Grafton, Auckland 1142, New Zealand. ²⁴UT Southwestern Medical Center – BMT Program, 7800C Stenton Ave. Apt. 210, Philadelphia, PA 19118, USA. ²⁵Stanford Health Care, 300 Pasteur Drive, Room H0101 MC 5623, Stanford, CA 94305, USA. ²⁶Dana Farber Cancer Institute - Adults, 450 Brookline Avenue, Boston, MA 02215, USA. ²⁷Division of Hematology and Oncology, Department of Medicine, Medical College of Wisconsin, 8701 Watertown Plank Rd. PO Box 26509, Milwaukee, WI 53226, USA. ²⁸Institut Català d'Oncologia - Hospital Duran I Reynals, Avda. Granvía 199-203, 08908 Barcelona, Spain.

Received: 2 November 2018 Accepted: 27 December 2018

Published online: 10 January 2019

References

- Gisselbrecht C, Gaulard P, Lepage E, Coiffier B, Briere J, Haioun C, et al. Prognostic significance of T-cell phenotype in aggressive non-Hodgkin's lymphomas. Groupe d'Etudes des Lymphomes de l'Adulte (GELA). *Blood*. 1998;92(1):76–82.
- Mourad N, Mounier N, Briere J, Raffoux E, Delmer A, Feller A, et al. Clinical, biologic, and pathologic features in 157 patients with angioimmunoblastic T-cell lymphoma treated within the Groupe d'Etude des Lymphomes de l'Adulte (GELA) trials. *Blood*. 2008;111(9):4463–70.
- Federico M, Rudiger T, Bellei M, Nathwani BN, Luminari S, Coiffier B, et al. Clinicopathologic characteristics of angioimmunoblastic T-cell lymphoma: analysis of the international peripheral T-cell lymphoma project. *J Clin Oncol*. 2013;31(2):240–6.
- Siebert W, Agthe A, Griesser H, Schwerdtfeger R, Brittinger G, Engelhard M, et al. Treatment of angioimmunoblastic lymphadenopathy (AILD)-type T-cell lymphoma using prednisone with or without the COPBLAM/IMVP-16 regimen. A multicenter study. *Kiel lymphoma study group. Ann Intern Med*. 1992;117(5):364–70.
- Pautier P, Devidas A, Delmer A, Dombret H, Sutton L, Zini JM, et al. Angioimmunoblastic-like T-cell non Hodgkin's lymphoma: outcome after chemotherapy in 33 patients and review of the literature. *Leuk Lymphoma*. 1999;32(5-6):545–52.
- Schmitz N, Trumper L, Ziepert M, Nickelsen M, Ho AD, Metzner B, et al. Treatment and prognosis of mature T-cell and NK-cell lymphoma: an analysis of patients with T-cell lymphoma treated in studies of the German high-grade non-Hodgkin lymphoma study group. *Blood*. 2010;116(18):3418–25.
- Vose J, Armitage J, Weisenburger D, International TCLP. International peripheral T-cell and natural killer/T-cell lymphoma study: pathology findings and clinical outcomes. *J Clin Oncol*. 2008;26(25):4124–30.
- Schetelig J, Fetscher S, Reichle A, Berdel WE, Beguin Y, Brunet S, et al. Long-term disease-free survival in patients with angioimmunoblastic T-cell lymphoma after high-dose chemotherapy and autologous stem cell transplantation. *Haematologica*. 2003;88(11):1272–8.
- Rodriguez J, Conde E, Gutierrez A, Arranz R, Gandarillas M, Leon A, et al. Prolonged survival of patients with angioimmunoblastic T-cell lymphoma

- after high-dose chemotherapy and autologous stem cell transplantation: the GELTAMO experience. *Eur J Haematol.* 2007;78(4):290–6.
10. Kyriakou C, Canals C, Goldstone A, Caballero D, Metzner B, Kobbe G, et al. High-dose therapy and autologous stem-cell transplantation in angioimmunoblastic lymphoma: complete remission at transplantation is the major determinant of outcome-lymphoma working Party of the European Group for blood and marrow transplantation. *J Clin Oncol.* 2008;26(2):218–24.
 11. Le Gouill S, Milpied N, Buzyn A, De Latour RP, Vernant JP, Mohty M, et al. Graft-versus-lymphoma effect for aggressive T-cell lymphomas in adults: a study by the Societe Francaise de Greffe de Moelle et de Therapie Cellulaire. *J Clin Oncol.* 2008;26(14):2264–71.
 12. Hamadani M, Awan FT, Elder P, Lin TS, Porcu P, Blum KA, et al. Allogeneic hematopoietic stem cell transplantation for peripheral T cell lymphomas; evidence of graft-versus-T cell lymphoma effect. *Biol Blood Marrow Transplant.* 2008;14(4):480–3.
 13. Urbano-Ispizua A, Pavletic SZ, Flowers ME, Klein JP, Zhang MJ, Carreras J, et al. The impact of graft-versus-host disease on the relapse rate in patients with lymphoma depends on the histological subtype and the intensity of the conditioning regimen. *Biol Blood Marrow Transplant.* 2015;21(10):1746–53.
 14. Kyriakou C, Canals C, Finke J, Kobbe G, Harousseau JL, Kolb HJ, et al. Allogeneic stem cell transplantation is able to induce long-term remissions in angioimmunoblastic T-cell lymphoma: a retrospective study from the lymphoma working party of the European group for blood and marrow transplantation. *J Clin Oncol.* 2009;27(24):3951–8.
 15. Doderio A, Spina F, Narni F, Patriarca F, Cavattoni I, Benedetti F, et al. Allogeneic transplantation following a reduced-intensity conditioning regimen in relapsed/refractory peripheral T-cell lymphomas: long-term remissions and response to donor lymphocyte infusions support the role of a graft-versus-lymphoma effect. *Leukemia.* 2012;26(3):520–6.
 16. Smith SM, Burns LJ, van Besien K, Lerademacher J, He W, Fenske TS, et al. Hematopoietic cell transplantation for systemic mature T-cell non-Hodgkin lymphoma. *J Clin Oncol.* 2013;31(25):3100–9.
 17. Bacigalupo A, Ballen K, Rizzo D, Giralto S, Lazarus H, Ho V, et al. Defining the intensity of conditioning regimens: working definitions. *Biol Blood Marrow Transplant.* 2009;15(12):1628–33.
 18. Cheson BD, Pfistner B, Juweid ME, Gascoyne RD, Specht L, Horing SJ, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol.* 2007;25(5):579–86.
 19. Przepiorka D, Weisdorf D, Martin P, Klingemann HG, Beatty P, Hows J, et al. 1994 Consensus Conference on Acute GVHD Grading. *Bone Marrow Transplant.* 1995;15(6):825–8.
 20. Shulman HM, Sullivan KM, Weiden PL, McDonald GB, Striker GE, Sale GE, et al. Chronic graft-versus-host syndrome in man. A long-term clinicopathologic study of 20 Seattle patients. *Am J Med.* 1980;69(2):204–17.
 21. Copelan E, Casper JT, Carter SL, van Burik JA, Hurd D, Mendizabal AM, et al. A scheme for defining cause of death and its application in the T cell depletion trial. *Biol Blood Marrow Transplant.* 2007;13(12):1469–76.
 22. Commenges D, Andersen PK. Score test of homogeneity for survival data. *Lifetime Data Anal.* 1995;1(2):145–56. discussion 57–9
 23. Kharfan-Dabaja MA, Kumar A, Ayala E, Hamadani M, Reimer P, Gisselbrecht C, et al. Clinical practice recommendations on indication and timing of hematopoietic cell transplantation in mature T cell and NK/T cell lymphomas: an international collaborative effort on behalf of the guidelines Committee of the American Society for blood and marrow transplantation. *Biol Blood Marrow Transplant.* 2017;23(11):1826–38.
 24. d'Amore F, Relander T, Lauritzsen GF, Jantunen E, Hagberg H, Anderson H, et al. Up-front autologous stem-cell transplantation in peripheral T-cell lymphoma: NLG-T-01. *J Clin Oncol.* 2012;30(25):3093–9.
 25. Hamadani M, Kanate AS, DiGilio A, Ahn KW, Smith SM, Lee JW, et al. Allogeneic hematopoietic cell transplantation for aggressive NK cell leukemia. A Center for International Blood and Marrow Transplant Research Analysis. *Biol Blood Marrow Transplant.* 2017;23(5):853–6.
 26. Kanate AS, DiGilio A, Ahn KW, Al Malki M, Jacobsen E, Steinberg A, et al. Allogeneic haematopoietic cell transplantation for extranodal natural killer/T-cell lymphoma, nasal type: a CIBMTR analysis. *Br J Haematol.* 2018;182(6):916–20.
 27. Neelapu SS, Locke FL, Bartlett NL, Lekakis LJ, Miklos DB, Jacobson CA, et al. Axicabtagene Ciloleuce CAR T-cell therapy in refractory large B-cell lymphoma. *N Engl J Med.* 2017;377(26):2531–44.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

