

# Journal of Appalachian Health

Volume 1 | Issue 3

Article 3

2019

# Adult food security and the relationship with Adverse Childhood Experiences among residents of Appalachian North Carolina

Manan Roy Appalachian State University, roym1@appstate.edu

Erin Bouldin Appalachian State University, bouldinel@appstate.edu

Maggie Bennett Appalachian State University, bennettme@appstate.edu

Adam Hege Appalachian State University, hegeba@appstate.edu

Follow this and additional works at: https://uknowledge.uky.edu/jah

Part of the Appalachian Studies Commons, Community Health Commons, Food Security Commons, Public Health Commons, Regional Economics Commons, Regional Sociology Commons, and the Rural Sociology Commons

#### **Recommended Citation**

Roy M, Bouldin E, Bennett M, Hege A. Adult food security and the relationship with Adverse Childhood Experiences among residents of Appalachian North Carolina. J Appalach Health 2019;1(3):17–26. DOI: https://doi.org/10.13023/jah.0103.03

Copyright © 2019 Manan Roy, Erin Bouldin, Maggie Bennett, and Adam Hege

This Research Articles is brought to you for free and open access by the College of Public Health at the University of Kentucky.

# Adult food security and the relationship with Adverse Childhood Experiences among residents of Appalachian North Carolina

#### Abstract

**Introduction**: The Appalachian region has worse health outcomes than the remainder of the United States. These disparities are often linked to the underlying social and environmental determinants of health. Adverse childhood experiences (ACEs) are associated with poor health outcomes across the lifespan and have a significant impact on future social determinants as an adult, including food security status.

**Purpose:** To explore the relationships between ACEs and food security among adults in the Appalachian counties of North Carolina and make comparisons with the rest of the state.

**Methods**: Researchers used North Carolina's 2012 Behavioral Risk Factor Surveillance System data; namely, the ACEs optional module which includes 11 items related to experiences respondents had before the age of 18 and a single item from the Social Context optional module to classify food security status. The sample was divided into three age categories (18-44, 45-64, and 65 and older) for statistical comparisons as well as by the indicator for Appalachian county. Using Stata 15, weighted logistic regression was utilized for examining relationships between variables.

**Results:** ACEs were a statistically significant predictor of food insecurity across all respondents; each additional ACE was associated with a 13-21% increase in the odds of food insecurity, depending on age group. However, living in an Appalachian county was only a predictor for those age 45-64.

**Implications:** These findings highlight the long-term effects of childhood experiences on food security generally, and in Appalachia particularly for middle-aged adults. Reducing ACEs could reduce food insecurity and improve health in the region.

#### Keywords

adverse childhood experiences, food insecurity, Appalachia, North Carolina

#### **Creative Commons License**



This work is licensed under a Creative Commons Attribution 4.0 License.

#### **Cover Page Footnote**

The authors would like to thank James Cassell and the NC BRFSS team in the NC Department of Health and Human Services for providing data that enabled us to explore health in Appalachia. No competing financial or editorial interests were reported by the authors of this paper.

# **INTRODUCTION**

Recent reports highlight health disparities across the Appalachian region, which includes 13 states and more than 400 counties. Many of these health disparities are rooted in social determinants of health, including socioeconomic status and opportunity, healthcare access, transportation and the built environment, and food security.<sup>1</sup> In addition, Appalachia is predominantly rural and has a unique social and cultural context that predisposes many areas to face challenges across generations.<sup>2</sup>

Food insecurity, which includes experiences ranging from anxiety about purchasing food of adequate quality or in sufficient quantity to reducing food intake, is linked to numerous adverse health outcomes.<sup>3,4</sup> The growing prevalence of food insecurity in the U.S. is a critical public health issue and is of great concern to policymakers and government officials. National figures showed 12.9% of U.S. households, or approximately 41 million citizens, were food insecure at least some time during 2016.<sup>3</sup>

Adverse childhood experiences (ACEs) include abuse—physical, sexual, and emotional—and household dysfunctions, including substance abuse, separation or divorce, violence between adults, mental illness, and incarceration.<sup>5,6</sup> ACEs are associated with numerous poor health and social outcomes across the lifespan, including poor physical and mental health and mortality.<sup>5,7</sup>

Existing research has established the importance of various social, economic, and behavioral factors that can explain household food insecurity. However, as Temple<sup>8</sup> notes, stressful life events or stressors such as mental illness, not being able to get a job, or death of a family member or close friend, have received scant attention in the empirical literature on food insecurity. Using nationally representative data from Australia, Temple showed that exposure to 18 discrete stressors significantly increased the probability of experiencing food insecurity, while adjusting for the more established correlates of food insecurity, like lower education level, being unmarried, or renting a home. Here in the U.S., Chilton and colleagues<sup>9</sup> found that exposure to ACEs was associated with experiencing food insecurity later in adulthood. Further, using the Midlands Family Study, researchers reported that low-income families with inadequate social support who adjust to adverse life events have a higher probability of child hunger.<sup>10</sup>

# **PURPOSE**

In this light, the current study sought to examine the relationship between ACEs and food insecurity among North Carolina (NC) residents. Four main factors led to studying this relationship. First, the study addresses an important omission in the literature on Appalachian outcomes: population-based long-term effects of ACEs on stress-related food insecurity accounting for residence in an Appalachian county. Second, the study focused on Appalachian NC since NC experienced a food insecurity rate of 14.4% in 2015–2017 while the national average food insecurity rate was 12.3%.<sup>3</sup> Third, the *Appalachian Diseases of Despair* report commissioned by the Appalachian Regional Commission (ARC) finds that in Appalachian NC, the diseases of despair mortality rate was 45% higher than in the non-Appalachian portions in 2015.<sup>11,12</sup> Finally, although not statistically different from the national prevalence rate, the prevalence of 3–8

ACEs among NC residents exceeded the corresponding national prevalence rate in 2016.<sup>11</sup>

#### **METHODS**

#### Data

Data come from the 2012 Behavioral Risk Factor Surveillance System (BRFSS) in NC. BRFSS is a landline and cell phone survey of community-dwelling adults aged 18 and older designed to represent the state's population in terms of age, gender, race/ethnicity, home ownership, and telephone type. The primary independent and dependent variables in this study, ACEs and stress-related food insecurity, were each included in the optional modules of the NC BRFSS only in 2012.

#### Measures

The ACEs optional module includes 11 items that relate to experiences respondents had before age 18. Specifically, respondents were asked whether or not they experienced any of the following:

- (1) lived with someone who was depressed, mentally ill, or suicidal;
- (2) lived with anyone who was a problem drinker or alcoholic;
- (3) lived with anyone who used illegal street drugs or who abused prescription medications;
- (4) lived with anyone who served time or was sentenced to serve time in a prison, jail, or other correctional facility; or
- (5) parents separated or divorced.

In addition, respondents were asked how often (never, once, or more than once) before age 18:

- (6) parents or adults in their home ever slapped, hit, kicked, punched, or beat each other up;
- (7) parent or adult in their home ever hit, beat, kicked, or physically hurt them in any way (except spanking);
- (8) parent or adult in their home ever swore at them, insulted them, or put them down;
- (9) anyone at least 5 years older than them or an adult, ever touched them sexually;
- (10) anyone at least 5 years older than them or an adult tried to make them touch them sexually;
- (11) anyone at least 5 years older than them or an adult forced them to have sex.

Respondents were classified as having experienced an ACE if they had ever experienced it (i.e., "ever" or "once" or "more than once," depending on the item).

Four measures of ACEs were created based on an earlier study by Ford and colleagues<sup>13</sup>:

- a continuous ACE score by summing the number of "yes" and "ever" responses, with a possible range of 0–11 ACEs; and three dichotomous subscales:
- (2) household dysfunction,
- (3) emotional or physical abuse, and
- (4) sexual abuse.

Household dysfunction included items 1–5. Physical or emotional abuse included items 6–8. Sexual abuse included items 9–11.

The BRFSS includes a single question in the Social Context module to classify respondents as experiencing food-related stress (food insecurity): "How often in the past 12 months would you say you were worried or stressed about having enough money to buy nutritious meals?" Respondents who said they were always, usually, or sometimes worried or stressed were classified as experiencing food-related stress (food insecurity) and those who said they rarely or never were worried or stressed were classified as not experiencing food insecurity. Previous studies<sup>14,15</sup> have validated the high correlation between the single measure of food-related stress or food insecurity in the BRFSS with the U.S. Department of Agriculture's Current Population Survey Food Security Supplement. Note the BRFSS measure may reflect marginal food security rather than more severe food insecurity.<sup>16</sup> For consistency, "food insecurity" is used instead of food-related stress.

Respondents were also classified as living within or outside Appalachia based on their county of residence. Counties were considered to be part of Appalachia (n=29) based on the ARC classification. The NC BRFSS coordinator created the Appalachia indicator variable since the public BRFSS files no longer include county.

Respondents self-reported a variety of socioeconomic and demographic characteristics, including age, gender, highest level of education completed, race and ethnicity, current employment status, marital status, the number of children under 18 living in their household, annual household income, home ownership status, and whether they ever served on active duty or in the reserves.

# Statistical Analysis

Existing research<sup>4</sup> finds that younger people and households with children are more likely to be food insecure. Further, mental health problems often emerge in early adulthood such that it is important to study the link between ACEs and food insecurity for that age group.<sup>17</sup> Adults aged ≥45 years are most affected by the ongoing crises of "diseases of despair" in most Appalachian states. Looking at this age group separately may signal a pathway between ACEs and the "diseases of despair" for this age group.<sup>18,19</sup> North Carolina has a higher than average rate of opioid deaths: the rate was 19.8 per 100,000 in the state in 2017 compared to 14.6 per 100,000 nationally.<sup>20</sup> The majority of these deaths in North Carolina occurred among younger and middle-aged adults; in 2017, 30% of opioid-related deaths in the state were among those aged 25–34, 25% were among people aged 35–44, and 21% were among adults aged 45–54.<sup>21</sup> Older adults also experience food insecurity, but in general food insecurity declines with age.<sup>22</sup> Finally, early mortality associated with ACEs may result in differential associations between food insecurity and ACEs by age group.

Accordingly, the sample was divided into three age categories: 18–44, 45–64, and ≥65. T-tests were used to evaluate whether the prevalence of food insecurity, ACE measures, and other demographic characteristics differed significantly across the three age categories. For each age category, logistic regression models were estimated to assess the relationship between each of the ACE measures and food insecurity, controlling for the social, economic, and demographic characteristics listed above as well as an indicator for residing in an Appalachian

county. For each age group, four regression models were estimated, one for each measure of ACE. For each ACE measure, whether living in Appalachia had a differential effect on food insecurity was tested. All analyses were weighted with BRFSS survey weights to represent the state's population. The data were analyzed using Stata 15.

# RESULTS

There were 11,770 respondents included in the study: 3661 aged 18–44; 4424 aged 45–64; and 3685 aged ≥65. The mean ACE score decreased with age (Table 1; all tables are provided under Additional Files). Household dysfunction was the most commonly reported type of ACE across all age groups and sexual abuse was least common; the prevalence of each ACE category generally decreased with age. Respondents aged ≥65 were more likely to be female, white, and veterans than respondents aged 18–44. They also tended to have less education and were less likely to have children in the household and to rent their home than those aged 18–44. Younger respondents; this is not surprising as recent data have shown that rural counties, such as found in Appalachia, have higher rates of residents aged  $\geq 65.^{23}$  Echoing other studies,<sup>4</sup> younger respondents had the highest rate of food insecurity (30%) compared to 25% among those aged 45–64, and only 13.2% for the elderly.

Table 2 (provided under Additional Files) shows the continuous ACE score increased the likelihood of food insecurity for all age groups (p<0.01), after adjusting for covariates. Each additional ACE was associated with a 13%–21% increase in the odds of food insecurity as an adult. No independent effect of living in Appalachia was found in any of the age groups nor were there any differential effect of ACE by Appalachian county. The estimated odds for the differential effect of ACEs by Appalachian county (for Tables 2–5) are available on request.

Table 3 (provided under Additional Files) shows that experiencing household dysfunction as a child increased the odds of food insecurity in adulthood by 44%-54% only in the younger and middle age groups (p<0.01). Similar to the findings in Table 2, there was no independent effect of living in Appalachia in any of the age groups nor were there any differential effect of experiencing household dysfunction in childhood by Appalachian county.

Unlike the household dysfunction subscale of ACE, Table 4 (provided under Additional Files) shows that experiencing any physical and emotional abuse in childhood increased the odds of food insecurity for all age groups: by 56%–90% for all age groups. Moreover, for the 45–64 age group, residing in an Appalachian county increased the odds of food insecurity (p<0.05). However, there was no differential effect of physical and emotional abuse by Appalachian county for any of the age groups.

Finally, as shown in Table 5 (provided under Additional Files), the odds of food insecurity increased by 65%–108% if younger and middle-aged respondents experienced any sexual abuse in childhood (p<0.01). Only for the 45–64 age group, residing in an Appalachian county increased the odds of food insecurity (p<0.05). However, there was no differential effect of experiencing sexual abuse in childhood by Appalachian county.

# **IMPLICATIONS**

The results corroborate earlier findings that ACEs increase the likelihood of being food insecure in adulthood, especially for the groups aged 18–44 and 45–64 years. The significant effect of ACEs on food insecurity in young adulthood could be due to emergence of mental health problems such as depression and substance abuse during this period.<sup>17</sup>

For those aged 45–64, physical and emotional abuse and sexual abuse in childhood had the greatest odds of predicting adult food insecurity. The Opioid Risk Tool, developed by Lynn R Webster, MD, also lists personal pre-adolescent sexual abuse as one of the risk factors for opioid abuse. Although accounting for opioid abuse is beyond the scope of this paper, these independent findings warrant further research about the connections between ACEs, adult food insecurity, and other social determinants of health, and adult opioid use in other areas of the Appalachian region.

The lack of a consistent relationship between ACEs and food insecurity for the elderly could, among others, be explained by potential recall error as well as unwillingness to report or discuss such events. Also, in general, residence in one of the Appalachian counties was not associated with food insecurity after adjusting for other covariates. However, among people in the 45–64 age range, living in an Appalachian county independently increased the odds of food insecurity when controlling for physical/emotional abuse or sexual abuse in childhood. Here too, there were no differential effects by Appalachian county of residence.

The first contribution of the study is assessing the population-based long-term effect of ACEs and adult food insecurity in a mostly rural Appalachian state such as NC. The second contribution is using different definitions of ACEs-the total number of ACEs as well as individual subscales depicting household dysfunction, physical/emotional abuse, and sexual abuse. This distinction throws light on the type of ACE driving the relationship. The other contribution is the focus on different age categories since ACEs have differential effects over the course of a lifetime. Specifically, the study found that physical and emotional abuse in childhood had the greatest odds of predicting adult food insecurity for adults who were at least aged 45. Sexual abuse in childhood was most strongly associated with adult food insecurity for the youngest age group. It is possible that older respondents have had more life experiences and developed coping mechanisms whereby physical and emotional abuse dominate adult life experiences. Interestingly, although household dysfunction was the most commonly reported type of ACE across all age groups, it was not the strongest predictor of adult food insecurity for any age group. These results suggest that individual trauma (physical/emotional abuse or sexual abuse) in childhood may have stronger long-term effects than a dysfunctional environment.

The principal limitation of the study is that a causal relationship between ACEs and food insecurity cannot be estimated, especially in the presence of potential recall error or misreporting ACEs.<sup>24,25</sup> In addition, only a single item was used to assess food insecurity, which may reflect marginal food security rather than more severe food insecurity.

The findings from this study illustrate life course impacts of the social determinants of health, in the form of ACEs and adult food insecurity, and the need for comprehensive and upstream approaches for the challenges facing Appalachia. While the ACE definitions in the study include only household dysfunction, physical and emotional, and sexual abuse, future research should ideally include a broader definition of ACEs such as additional peer- and community-related factors that also count as harmful childhood adversities.<sup>6</sup>

For future generations, it is vital that researchers, program developers, and policymakers translate the growing evidence into action by designing preventative and family-based programs which provide adequate resources and support, such as the Building Community Resilience Model,<sup>26</sup> that systematically address social determinants of health early in life before adverse health effects occur as well as help develop resilience in the face of adversities.<sup>10</sup>

#### SUMMARY BOX

**What is already known about the topic?** Food insecurity and adverse childhood experiences (ACEs) are major public health issues across the United States that are linked to other social determinants of health and poor health outcomes across the lifespan.

**What is added by this report?** This study complements the limited research on the relationship between food insecurity and ACEs in Appalachia. The study uses different definitions of ACEs and focuses on different age categories since ACEs have differential effects over the course of one's life.

What are the implications for public health, practice, policy, and research? Future research should include a broader definition of ACEs such as additional peer- and community-related factors that also count as harmful childhood adversities. Researchers, program developers, and policymakers should focus on designing preventative and family-based programs which provide adequate resources and support to systematically address social determinants of health early in life before adverse health effects occur as well as help develop resilience in the face of adversities.

# REFERENCES

- 1. Goldman L, Benjamin G, Hernandez S, et al. Advancing the Health of Communities and Populations Vital Directions for Health and Health Care 2018.
- 2. Durlauf S. Poverty traps and Appalachia. Washington DC: Brookings Institution Press, 2012.
- 3. Gunderson C, Ziliak, JP. Food insecurity research in the United States: where we have been and where we need to go. Appl Econ Perspect Policy 2018;40(1):119–35.
- 4. Gunderson, C, Ziliak, JP. Food insecurity and health outcomes. Health Affairs 2015;34(11):1830–9.
- 5. Felitti VJ, Anda RF, Nordenberg D, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. Am J Prev Med 1998;14(4):245–58.
- 6. Finkelhor D, Shattuck A, Turner H, Hamby S. Improving the adverse childhood experiences study scale. JAMA Pediatr 2013;167:70–5.
- Hughes K, Bellis M, Hardcastle K, et al. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. Lancet Public Health 2017;2(8):e356–66.
- 8. Temple JB. The association between stressful events and food insecurity: Cross-sectional evidence from Australia. Int J Environ Res Public Health 2018;15(11):2333.
- 9. Chilton M, Knowles M, Rabinowich J, Arnold K. The relationship between childhood adversity and food insecurity: 'It's like a bird nesting in your head'. Public Health Nutr 2015;18(14):2643–53.
- Jones S, Draper C, Bell B, et al. Child hunger from a family resilience perspective. Journal of Hunger & Environmental Nutrition. 2018;13(3):340–61.

https://www.tandfonline.com/doi/full/10.1080/19320248.2017.136418

- Sacks V, Murphey D. The prevalence of adverse childhood experiences, nationally, by state, and by race or ethnicity 2018 [cited 2019 April 11]. Available from: https://www.childtrends.org/publications/prevalenceadverse-childhood-experiences-nationally-state-race-ethnicity.
- Case A, Deaton A. Rising morbidity and mortality in midlife among white non-Hispanic Americans in the 21st century. Proceedings of the National Academy of Sciences, 2015;112(49): 15078–83.
- 13. Ford D, Merrick M, Parks S, et al. Examination of the factorial structure of adverse childhood experiences and recommendations for three subscale scores. Psychol Violence 2014;4(4).
- Irving SM, Njai RS, Siegel PZ. Food insecurity and self-reported hypertension among Hispanic, black and white adults in 12 states, Behavioral Risk Factor Surveillance System, 2009. Prev Chronic Dis 11:140190. DOI: http://dx.doi.org/10.5888/pcd11.140190
- Njai R, Siegel P, Yin S, Liao Y. Prevalence of Perceived Food and Housing Security-15 States, 2013. MMWR Morb Mortal Wkly Rep 2017;66(1):12– 15.
- Gunderson C, Engelhard EE, Crumbaugh AS, Seligman HK. Brief assessment of food insecurity accurately identifies high-risk US adults. Public Health Nutr 2017;20(8):1367–71.
- 17. Mersky JP, Topitzes J, Reynolds AJ. Impacts of adverse childhood experiences on health, mental health, and substance use in early

adulthood: A cohort study of an urban, minority sample in the US. Child Abuse Negl 2013;37(11):817–925.

- Meit M, Hefferman M, Tanenbaum E, Hoffman T. Appalachian Diseases of Despair: Final Report, 2017. https://www.arc.gov/assets/research\_reports/AppalachianDiseasesofDe spairAugust2017.pdf
- 19. Shanahan L, Hill S, Gaydosh L, et al. Does despair really kill? A roadmap for an evidence-based answer. Am J Public Health 2019;109(6):854–8.
- 20. National Institute on Drug Abuse. North Carolina Opioid Summary: Opioid-Involved Overdose Deaths. March 2019. Available at: https://www.drugabuse.gov/opioid-summaries-by-state/north-carolinaopioid-summary
- 21. Henry J. Kaiser Family Foundation. Opioid Overdose Deaths by Age Group (2017). Available at: https://www.kff.org/other/stateindicator/opioid-overdose-deaths-by-agegroup/?dataView=1&currentTimeframe=0&sortModel=%7B%22colId%22: %22Location%22,%22sort%22:%22asc%22%7D.
- Strickhouser S, Wright JD, Donley AM. Food Insecurity Among Older Adults-Full Report, 2015 Update. AARP Foundation, September 2014. Available at: https://www.aarp.org/content/dam/aarp/aarp\_foundation/2015-PDFs/AF-Food-Insecurity-2015Update-Final-Report.pdf.
- USDA. Economic Research Service using data from the U.S. Census Bureau Population Estimates Program. https://www.ers.usda.gov/webdocs/publications/90556/eib-200.pdf?v=5899.2
- Williams LM. Recall of childhood trauma: A prospective study of women's memories of child sexual abuse. J Consult Clin Psychol 1994;62(6):1167– 76.
- 25. Hardt J, Rutter M. Validity of adult retrospective reports of adverse childhood experiences: Review of the evidence. J Child Psychol Psychiatry 2004;45(2): 260–73.
- 26. Ellis WR, Dietz WH. A New Framework for Addressing Adverse Childhood and Community Experiences: The Building Community Resilience Model. Acad Pediatr 2017;17(7):S86–S89.