



University of Kentucky
UKnowledge

Plant and Soil Sciences Faculty Publications

Plant and Soil Sciences

4-19-2019

***trans*-Cinnamic Acid-Induced Leaf Expansion Involves an Auxin-Independent Component**


Jasmina Kurepa

University of Kentucky, jasmina.kurepa@uky.edu

Jan A. Smalle

University of Kentucky, jsmalle@uky.edu

Follow this and additional works at: https://uknowledge.uky.edu/pss_facpub

 Part of the [Integrative Biology Commons](#), [Plant Sciences Commons](#), and the [Soil Science Commons](#)
Right click to open a feedback form in a new tab to let us know how this document benefits you.

Repository Citation

Kurepa, Jasmina and Smalle, Jan A., "*trans*-Cinnamic Acid-Induced Leaf Expansion Involves an Auxin-Independent Component" (2019). *Plant and Soil Sciences Faculty Publications*. 141.

https://uknowledge.uky.edu/pss_facpub/141

This Article is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in Plant and Soil Sciences Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

***trans*-Cinnamic Acid-Induced Leaf Expansion Involves an Auxin-Independent Component**

Abstract

The phenylpropanoid pathway, the source of a large array of compounds with diverse functions, starts with the synthesis of *trans*-cinnamic acid (*t*-CA) that is converted by cinnamate-4-hydroxylase (C4H) into *p*-coumaric acid. We have recently shown that in Arabidopsis, exogenous *t*-CA promotes leaf growth by increasing cell expansion and that this response requires auxin signaling. We have also shown that cell expansion is increased in C4H loss-of-function mutants. Here we provide further evidence that leaf growth is enhanced by either *t*-CA or a *t*-CA derivative that accumulates upstream of C4H. We also show that this growth response pathway has two components: one that requires auxin signaling and another which employs a currently unknown mechanism.

Keywords

Auxin, *trans*-cinnamic acid, phenylalanine ammonia-lyase, cinnamate-4-hydroxylase, leaf expansion

Disciplines

Integrative Biology | Plant Sciences | Soil Science

Notes/Citation Information

Published in *Communicative & Integrative Biology*, v. 12, no. 1, p. 78-81.

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



trans-Cinnamic acid-induced leaf expansion involves an auxin-independent component

Jasmina Kurepa & Jan A. Smalle

To cite this article: Jasmina Kurepa & Jan A. Smalle (2019) *trans*-Cinnamic acid-induced leaf expansion involves an auxin-independent component, Communicative & Integrative Biology, 12:1, 78-81, DOI: [10.1080/19420889.2019.1605814](https://doi.org/10.1080/19420889.2019.1605814)

To link to this article: <https://doi.org/10.1080/19420889.2019.1605814>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 19 Apr 2019.



Submit your article to this journal [↗](#)



Article views: 352



View related articles [↗](#)



View Crossmark data [↗](#)

trans-Cinnamic acid-induced leaf expansion involves an auxin-independent component

Jasmina Kurepa and Jan A. Smalle

Plant Physiology, Biochemistry, Molecular Biology Program, Department of Plant and Soil Sciences, College of Agriculture, Food and Environment, University of Kentucky, Lexington, KY, USA

ABSTRACT

The phenylpropanoid pathway, the source of a large array of compounds with diverse functions, starts with the synthesis of *trans*-cinnamic acid (*t*-CA) that is converted by cinnamate-4-hydroxylase (C4H) into *p*-coumaric acid. We have recently shown that in Arabidopsis, exogenous *t*-CA promotes leaf growth by increasing cell expansion and that this response requires auxin signaling. We have also shown that cell expansion is increased in C4H loss-of-function mutants. Here we provide further evidence that leaf growth is enhanced by either *t*-CA or a *t*-CA derivative that accumulates upstream of C4H. We also show that this growth response pathway has two components: one that requires auxin signaling and another which employs a currently unknown mechanism.

ARTICLE HISTORY

Received 27 January 2019
Revised 1 April 2019
Accepted 8 April 2019



KEYWORDS

Auxin; *trans*-cinnamic acid; phenylalanine ammonia-lyase; cinnamate-4-hydroxylase; leaf expansion

The phenylpropanoid (PP) pathway generates a wide range of structurally and functionally different compounds whose importance for cell growth and development is widely recognized. [1] In the first two steps of this pathway, phenylalanine ammonia-lyase (PAL) deaminates L-phenylalanine to yield *trans*-cinnamic acid (*t*-CA) which is then converted into *p*-coumaric acid by cinnamate-4-hydroxylase (C4H). [1] We have recently shown that feeding with *t*-CA increases the growth rate and final size of Arabidopsis rosette leaves by increasing cell expansion. [2] Two previous studies on plant species other than Arabidopsis have also shown that *t*-CA treatment or increased *t*-CA accumulation leads to increased leaf growth, indicating that the growth promotion by *t*-CA is conserved within higher plants. [3,4] We have also shown that auxin insensitive mutants are resistant to *t*-CA-induced leaf expansion and concluded that *t*-CA-induced leaf growth requires an intact auxin signaling pathway. [2] To test whether *t*-CA itself acts to increase leaf cell expansion or serves as a precursor of a cell-size regulatory PP compound that is synthesized in a downstream branch, we analyzed the cell sizes in lines that have reduced PAL or C4H activity (Figure 1(a)) [2]. In transgenic plants with reduced PAL activity, caused by overexpression of the PAL-targeting ubiquitin ligase subunit KFB20, cells are

smaller compared to the wild type [2] and consequently, the plants are dwarfed. [2,5] On the other hand, in *ref3* mutants, which have reduced C4H activity, [6] cells are larger than in the wild type. Because both *ref3* mutants and KFB20 overexpression plants are predicted to have a reduced carbon flow downstream of C4H, none of the compounds synthesized after the C4H step in the PP pathway can be this promoter of cell expansion and the active PP intermediate(s) is either a PAL product or its derivative. The exact chemical structure of this active compound(s) awaits further investigation.

To further explore the relationship between early PP pathway intermediates and leaf expansion, we analyzed the *t*-CA growth response of *ref3* mutants. We were unable to use the *ref3-1* mutant line for this purpose because when grown on control medium, different seedlings had different rosette growth rates, which separated them into three classes: wild-type-sized, semi-dwarfed and dwarfed, aberrant seedlings (Figure 1(b)). These three different classes of *ref3-1* plants retained their homozygous *ref3-1* genotype as evidenced by the pale seed color of their progeny (Figure 1(b)). [7] In addition, the progeny of each class retained the same pattern of developmental variability (data not shown). Because the *ref3-1* mutant is an auxin hypersensitive response mutant, [2] these developmental

CONTACT Jasmina Kurepa  jsmalle@uky.edu  Plant Physiology, Biochemistry, Molecular Biology Program, Department of Plant and Soil Sciences, College of Agriculture, Food and Environment, University of Kentucky, Lexington, KY 40546, USA

Addendum to: "Modulation of auxin and cytokinin responses by early steps of the phenylpropanoid pathway" by J. Kurepa, T.E. Shull, S.S. Karunadasa, J.A. Smalle BMC plant biology 18 (1), 278

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. *t*-CA-induced rosette growth responses of *ref3* mutants. (a) Summary of previous findings that suggest that *t*-CA or a *t*-CA derivative promotes plant cell expansion. [2] PAL, phenylalanine ammonia-lyase; C4H, cinnamic acid 4-hydrolase; KFB20 OE, Kelch repeat F-box protein 20 overexpressor. (b) Rosettes of 20-day-old plants. The upper panel shows wild type plants (*Ler*) and *ref3-1* plants grown on MS/2 (control) media. The frequencies of *ref3-1* rosette phenotypes are noted as percentages ($n = 89$). Insert shows the seeds of the progeny of each phenotypical class. The lower panel shows rosettes of the *Col-0* wild type and *ref3-3* plants grown on control media. (c) Fresh weight of rosettes of *Col-0*, *ref3-3* and KFB20 OE plants grown on control and *t*-CA-supplemented medium for 20 days. Data are presented as mean \pm SD ($n \geq 12$ pools of 6 plants each). The significance of the difference between the mean fresh weight between lines grown on control media (black arrows) and between controls and the treated samples (dashed, gray arrows) is noted (****, $P < 0.0001$; two-way ANOVA with Tukey's post-test).

abnormalities and their incomplete penetrance are in agreement with the variability observed for some of the previously characterized auxin-related mutant lines. [8–10] The plants of the weaker C4H loss-of-function *ref3-3* mutant did not show any developmental variability. Surprisingly, *ref3-3* rosette leaves were larger than the wild-type leaves as illustrated by the measurements of rosette fresh weight that has been previously shown to be positively correlated with increased leaf size (Figure 1(b,c)). [2] This increase in rosette size stands in contrast to what was reported in the original *ref3* mutant study. [6] In that study, *ref3* plants were sown and grown on soil rather than on a sterile, sucrose-supplemented growth medium. [6] Since many of the products and intermediates of the PP pathway are involved in plant responses to the environment, [11,12] the discrepancy between the results of these two studies is potentially due to differential accumulation of specific PP

compounds in the *ref3* mutants that are less permissive for growth under non-sterile conditions. As expected from our previous study, [2] *t*-CA treatments led to growth increases of $44.4 \pm 4\%$ in the wild-type plants and $420 \pm 33\%$ in the dwarfed KFB20 overexpression plants (Figure 1(c); [2]). *t*-CA feeding also led to an increase of the rosette size in *ref3-3* plants, but this response was attenuated compared to the other lines ($23 \pm 6\%$; Figure 1(c)). Because the rosette size increases caused by C4H loss of function and *t*-CA treatment were not additive (i.e., the *t*-CA-treated wild-type and *ref3-3* plants were of the same size), we concluded that the size increase in *ref3-3* plants is indeed related to the *t*-CA-induced effect on rosette growth. Moreover, the increased rosette size of *ref3-3* plants was in agreement with our earlier observation that treatments with low concentrations of the C4H inhibitor piperonylic acid (PA) lead to an increase in rosette leaf size. [2] Higher doses of PA as

well as strong C4H loss-of- (e.g., *ref3-2*) mutations lead to a reduction in rosette size most likely because of the decreased biosynthesis of compounds downstream of C4H (e.g., lignins) which are needed for growth.

To better understand the role of auxin in *t*-CA-induced leaf expansion, we analyzed the *t*-CA and auxin effects separately and in combination. Treatments with either low concentrations of *t*-CA or IAA promoted leaf expansion. [2] When combined, *t*-CA and IAA led to an increase in leaf size in an additive manner: 1 μ M *t*-CA led to a size increase of $40 \pm 5\%$, 160 nM IAA led to a size increase of $18 \pm 3\%$ and when 1 μ M *t*-CA and 160 nM of IAA were applied together, we observed a growth increase of $67 \pm 4\%$ (Figure 2(a)). Under our experimental conditions, 160 nM IAA was the maximum response dose: 320 nM IAA promoted growth but to a lesser extent than 160 nM IAA (Figure 2(a)). However, even at the highest tested IAA dose, the combined treatment caused a growth increase that was significantly stronger than that observed for the treatment with 160 nM IAA only (Figure 2(a)). This suggests an additive interaction between IAA- and *t*-CA-dependent leaf growth promotion mechanisms.

In conclusion, *t*-CA-induced leaf expansion combines an auxin-dependent and an auxin-independent mechanism. In Figure 2(b), we offer two possible models to explain this dual *t*-CA effect on growth. In common to both models is the auxin-independent positive effect of *t*-CA on cell expansion. The second

component of the first model is a *t*-CA-driven enhancement of auxin signaling that is needed to inhibit an auxin-dependent repressor of cell expansion. The second point of the second model is cell wall loosening, a well-known auxin effect. [13] According to this model, the *t*-CA-dependent cell expansion can commence only if *t*-CA in parallel enhances auxin-dependent cell wall loosening. Regardless of which of these models is true, the fact that the auxin-dependent and *t*-CA-dependent effects are additive, suggests that plants have an intricate and multifaceted *t*-CA-governed mechanism that affects growth.

Materials and methods

The wild-type lines used were Columbia (Col-0) and Landsberg *erecta* (*Ler*) which are the backgrounds for the *ref3-3* and *ref3-1* mutants, respectively. [6] Both *ref3* mutant lines were obtained from the ABRC Seed Stock Center. All plants were grown in sterile conditions on half-strength Murashige and Skoog medium (pH 5.7) containing 1% sucrose and 0.8% PhytoAgar (MS/2 medium). The following chemicals were used for treatments: indole-3-acetic acid (IAA; Sigma) and *trans*-cinnamic acid (*t*-CA; Sigma). All rosette growth response analyses were done as previously described. [2] The descriptive statistics, plotting and hypothesis testing were done using Prism 6 software (GraphPad Software Inc). All data are presented as means \pm SD of at least three independent experiments. When means of more than two




Figure 2. Comparative analysis of the effects of auxin and *t*-CA on rosette growth. (a) Effects of indole-3-acetic acid (IAA) on the fresh weight of rosettes from wild-type plants grown for 20 days on MS/2 media containing the denoted doses of *t*-CA and IAA. Data are presented as mean \pm SD ($n \geq 12$ pools of 8 plants each). All differences between the IAA and IAA/*t*-CA treated samples for each IAA concentration are significant at $P < 0.0001$ (not marked; two-way ANOVA with Sidak's multiple comparisons test). The significance of the difference between the fresh weight of plants grown control media and on different IAA-supplemented media is shown in black and between different *t*-CA control and *t*-CA/IAA co-treated plants treatments is shown in green (****, $P < 0.0001$; two-way ANOVA with Sidak's multiple comparisons test). (b) Hypothetical models for the additive mechanism of action of IAA and *t*-CA in leaf growth promotion.

samples were compared, we used ANOVA followed by post-test to find a significant difference between pairs of means. The significance levels, indicated by asterisks in the figures, illustrate the results of the post-test.

Acknowledgments

The *ref3-1* and *ref3-3* mutant seeds were provided by the Arabidopsis Biological Resource Center at Ohio State University. This work was supported by NIFA HATCH project (1009329) and by the Kentucky Tobacco Research and Development Center.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was supported by the Kentucky Tobacco Research and Development Center [1]; National Institute of Food and Agriculture [HATCH project 1009329].

References

- [1] Vogt T. Phenylpropanoid biosynthesis. *Mol Plant*. 2010;3:2–20.
- [2] Kurepa J, Shull TE, Karunadasa SS, et al. Modulation of auxin and cytokinin responses by early steps of the phenylpropanoid pathway. *BMC Plant Biol*. 2018;18:278.
- [3] Talaat IM, Balbaa LK. Physiological response of sweet basil plants (*Ocimum basilicum* L.) to putrescine and *trans*-cinnamic acid. *Am Eurasian J Agric Environ Sci*. 2010;8:438–445.
- [4] Orlova I, Marshall-Colón A, Schnepf J, et al. Reduction of benzenoid synthesis in petunia flowers reveals multiple pathways to benzoic acid and enhancement in auxin transport. *Plant Cell*. 2006;18:3458–3475.
- [5] Zhang X, Gou M, Liu CJ. Arabidopsis Kelch repeat F-box proteins regulate phenylpropanoid biosynthesis via controlling the turnover of phenylalanine ammonia-lyase. *Plant Cell*. 2013;25:4994–5010.
- [6] Schilmiller AL, Stout J, Weng JK, et al. Mutations in the cinnamate 4-hydroxylase gene impact metabolism, growth and development in Arabidopsis. *Plant J*. 2009;60:771–782.
- [7] Ruegger M, Chapple C. Mutations that reduce sinapoylmalate accumulation in *Arabidopsis thaliana* define loci with diverse roles in phenylpropanoid metabolism. *Genetics*. 2001;159:1741–1749.
- [8] Bennett SRM, Alvarez J, Bossinger G, et al. Morphogenesis in *pinoid* mutants of *Arabidopsis thaliana*. *Plant J*. 1995;8:505–520.
- [9] Lie C, Kelsom C, Wu X. WOX2 and STIMPY-LIKE/WOX8 promote cotyledon boundary formation in Arabidopsis. *Plant J*. 2012;72:674–682.
- [10] Ploense SE, Wu MF, Nagpal P, et al. A gain-of-function mutation in IAA18 alters Arabidopsis embryonic apical patterning. *Development*. 2009;136:1509–1517.
- [11] Dixon RA, Achnine L, Kota P, et al. The phenylpropanoid pathway and plant defence - a genomics perspective. *Mol Plant Pathol*. 2002;3:371–390.
- [12] Lee HI, Leon J, Raskin I. Biosynthesis and metabolism of salicylic acid. *Proc Natl Acad Sci USA*. 1995;92:4076–4079.
- [13] Majda M, Robert S. The role of auxin in cell wall expansion. *Int J Mol Sci*. 2018;19:951.