

## Eastern Kentucky University Encompass

---

EKU Faculty and Staff Scholarship

---

3-2017

# Nutrient contamination from non-point sources: Dissolved nitrate and ammonium in surface and subsurface waters at EKU Meadowbrook Farm, Madison County, Kentucky


Reid E. Buskirk  
*Eastern Kentucky University*

Hunter R. Evans  
*Eastern Kentucky University*

Walter S. Borowski  
*Eastern Kentucky University*

Jonathan M. Malzone  
*Eastern Kentucky University*

Follow this and additional works at: [https://encompass.eku.edu/fs\\_research](https://encompass.eku.edu/fs_research)

 Part of the [Biogeochemistry Commons](#), [Environmental Health and Protection Commons](#), [Environmental Indicators and Impact Assessment Commons](#), and the [Environmental Monitoring Commons](#)

---

### Recommended Citation

Buskirk, R.E., H.R. Evans, W.S. Borowski, J.M. Malzone, 2017. Nutrient contamination from non-point sources: Dissolved nitrate and ammonium in surface and subsurface waters at EKU Meadowbrook Farm, Madison County, Kentucky. *GSA Abstracts with Program*, 49(2), doi: 10.1130/abs/2017NE-290271.

This Conference Presentation is brought to you for free and open access by Encompass. It has been accepted for inclusion in EKU Faculty and Staff Scholarship by an authorized administrator of Encompass. For more information, please contact [Linda.Sizemore@eku.edu](mailto:Linda.Sizemore@eku.edu).

# **Nutrient contamination from non-point sources: Dissolved nitrate and ammonium in surface and subsurface waters at EKV Meadowbrook Farm, Madison County, Kentucky**

Reid E. Buskirk, Hunter R. Evans, Walter S. Borowski, Jonathan M. Malzone  
Department of Geosciences, Eastern Kentucky University, Richmond KY 40475

## **Abstract**

Agricultural activities often contaminate watersheds with excess nutrients leading to poor water quality and eutrophication. We assayed dissolved nutrient levels in surface and subsurface waters of Eastern Kentucky University's Meadowbrook Farm in order to assess levels of dissolved nutrients leaving its farmland and draining into the Muddy Creek watershed. The Farm raises both crops and livestock so that nutrient sources include fertilizer and manure. We sampled springs, runoff, and subsurface pipe drainage as well as Muddy Creek on six days from May to August 2016 under a variety of weather conditions. Using established, standard colorimetric methods, we measured nitrate ( $\text{NO}_3^-$ ; cadmium reduction method) and ammonium ( $\text{NH}_4^+$ ; sodium hypochlorate method) via spectrophotometry with a precision and accuracy of  $\sim 0.1$  mg/L.

Nitrate was the dominant dissolved nitrogen species whereas ammonium was often absent in water samples. Nitrate levels were typically  $< 2$  mg/L N-  $\text{NO}_3$  with the largest values between 7.0 and 14.3 mg/L. Springs and some runoff samples had higher nitrate values. Ammonium generally ranged between 0.0 and 0.5 mg/L N-  $\text{NH}_4$  with concentration spikes between 2.0 and 4.3 mg/L, but from no consistent source.

Dissolved nitrogen concentration values responded to rainfall. Generally, nitrate concentrations increased more than ammonium concentrations during wetter periods. Spring samples maintained higher nitrogen concentrations regardless of different rainfall conditions.

Lastly, nitrate contamination was significantly lower than composite national values from streams draining agricultural lands, whereas ammonium was about equal to the median national average. Median nitrate concentration was  $\sim 1.8$  mg/L N-  $\text{NO}_3$  compared to the national value of  $\sim 2.8$  mg/L, whereas the value for pristine streams is 0.24 mg/L N-  $\text{NO}_3$  (Dubrovsky et al., 2010). Median ammonium values from both data sets are  $\sim 0.1$  mg/L N-  $\text{NH}_4$ ; the national value from pristine streams is  $\sim 0.025$  mg/L N.

Geological Society of America, joint Northeast and North-Central sections, 18-21 March 2017,  
*GSA Abstracts with Program*, 49(2); doi: 10.1130/abs/2017NE-290269.