



Western Washington University  
Western CEDAR

---

Salish Sea Ecosystem Conference

2018 Salish Sea Ecosystem Conference  
(Seattle, Wash.)

---

Apr 5th, 4:00 PM - 4:15 PM

## Effects of environmentally realistic concentrations of neonicotinoid insecticides on an aquatic invertebrate community

Claire Duchet

Washington State Univ., United States, [claire.duchet@wsu.edu](mailto:claire.duchet@wsu.edu)

Alyssa Kraft

Washington State Univ., United States, [akraft3@uw.edu](mailto:akraft3@uw.edu)

John D. Stark

Washington State Univ., United States, [starkj@wsu.edu](mailto:starkj@wsu.edu)

Follow this and additional works at: <https://cedar.wwu.edu/ssec>



Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Natural Resources and Conservation Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

---


Duchet, Claire; Kraft, Alyssa; and Stark, John D., "Effects of environmentally realistic concentrations of neonicotinoid insecticides on an aquatic invertebrate community" (2018). *Salish Sea Ecosystem Conference*. 398.

<https://cedar.wwu.edu/ssec/2018ssec/allsessions/398>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact [westerncedar@wwu.edu](mailto:westerncedar@wwu.edu).

# Effects of environmentally realistic concentrations of neonicotinoid insecticides on an aquatic invertebrate community

Claire Duchet, Alyssa Kraft, Cailin MacKenzie, John D. Stark




30<sup>th</sup> Salish Sea Ecosystem Conference, April 4-6, 2018, Seattle

# Introduction


- ▶ Neonicotinoid insecticides represent 24 % of the global market, and their use is increasing globally.
- ▶ Clothianidin, imidacloprid, thiamethoxam are the most commonly-used neonicotinoids on corn and soybeans.
- ▶ Imidacloprid is also used privately: lawn and garden care, and topical flea medicines.

# Introduction

- ▶ Used against insect pests but toxic to non-target organisms (e.g. crustaceans, bees, flies, birds)



Acetylcholinesterase (AChE) breaks down acetylcholine (ACh), preventing overstimulation and blockage of acetylcholine receptors



Acetylcholinesterase cannot break down neonicotinoids (ACh receptor agonists), leading to paralysis (due to blockage of ACh receptors)

Mode of action of neonicotinoids.

# Introduction

- ▶ Not intended for direct use in water bodies, but they may enter in the aquatic compartment *via* spray drift, runoff or leaching.
- ▶ Soluble in water, persistent in soil, and may be found in surface waters and drinking water (Klarish et al., 2017).

# Introduction

- ▶ In Canada, wetlands close to agricultural fields ([Anderson et al., 2015](#)) :
  - ▶ 3.11 µg/L clothianidin,
  - ▶ 0.256 µg/L imidacloprid,
  - ▶ 1.49 µg/L thiamethoxam.
- ▶ In Western Washington, imidacloprid almost always detected in surface water ([WSDA source](#)):
  - ▶ usually <0.1 µg/L, but >1 µg/L in some cases,
  - ▶ 1.74 µg/L in the Big Ditch slough, a creek providing habitat for salmon in the Skagit wildlife area.


# Objective

- ▶ Very few data about the effects of mixture of neonicotinoids available.
- ▶ Neonicotinoid contamination induce a top-down trophic cascade in a community dominated by invertebrate predators (Miles et al., 2017).
- ▶ Our objective: to test the effect of a mixture of imidacloprid, clothianidin and thiamethoxam on an aquatic invertebrate community.
- ▶ Hypothesis: By affecting predators, indirect positive effect on herbivores.


# Preliminary experiment

- ▶ Test in control conditions on *Ceriodaphnia dubia* (semi-static test): imidacloprid (0.256  $\mu\text{g/L}$ ), clothianidin (3.11  $\mu\text{g/L}$ ), thiamethoxam (1.49  $\mu\text{g/L}$ ), and the mixture

Survival following 8-d exposure



Reproduction following 8-d exposure



■ Control ■ Imidacloprid ■ Clothianidin ■ Thiamethoxam ■ Mixture


\*: Significant differences between the treatment and control (Tukey's HSD test,  $p < 0.05$ ).



# Materials and methods



# Materials and methods




# Materials and methods

Well water +  
pond water +  
leaf litter

Treatment

imidacloprid (0.256  $\mu\text{g/L}$ )  
clothianidin (3.11  $\mu\text{g/L}$ )  
thiamethoxam (1.49  $\mu\text{g/L}$ )  
mixture

Macroinvertebrates  
(predators)



Sampling dates (from 9/13/17 to 10/11/2017)


# Materials and methods

On each sampling date:

- ▶ Environmental parameters (pH, conductivity, dissolved oxygen, temperature)
- ▶ Invertebrate sampling
- ▶ Water samples for chemical analysis
- ▶ Water samples for chlorophyll *a* concentrations
  
- ▶ Dipteran colonization (mosquito oviposition habitat selection): every 2 days

# Results

- Concentrations of the neonicotinoids over time



# Results


- ▶ Community:
  - ▶ Active dispersers (10 taxa):
    - ▶ mosquitoes (*Culex pipiens*, *Culiseta longiareolata*),
    - ▶ non-biting midges (Chironomids),
    - ▶ biting-midges (Ceratopogonidae),
    - ▶ ephidridae larvae,
    - ▶ mayflies (Ephemeroptera),
    - ▶ odonates,
    - ▶ water beetle (Hydrophilidae),
    - ▶ water boatmen (*Anisops sardea*),
    - ▶ hydracarians.

# Results

- ▶ Community:
  - ▶ Passive dispersers (11 taxa):
    - ▶ Copepods (cyclopoids).
    - ▶ Cladocerans:
      - ▶ *Scapholeberis, Chydorus, Pleuroxus, Ceriodaphnia, Daphnia, Simocephalus, Alona, Macrothricidae, Diaphanosoma.*
    - ▶ Ostracods.

# Results

## ► Species evenness




RM ANOVA:  $F_{4,20} = 0.38, p = 0.82$



# Results


## ► Effect on taxa richness



\*: Significant differences between the treatment and control (Fisher's LSD test following RM ANOVA,  $p < 0.05$ ).

# Results

## ► Herbivore biomass



\*: Significant differences between the treatment and control (Fisher's LSD test following RM ANOVA,  $p < 0.05$ ).

# Conclusions - Perspectives

- ▶ Effects on the community:
  - ▶ Decrease of the active dispersers 3 weeks after exposure to the mixture
  - ▶ Increase of the passive dispersers 1 week after exposure to imidacloprid and 3 weeks after exposure to thiamethoxam
  - ▶ Increase of the zooplankton biomass 3 weeks after exposure due to lack of predators?
- ▶ Still have to analyze the chlorophyll *a* data
- ▶ Next study: run the experiment in early stage population development (late spring / early summer)

# Acknowledgments and Funding

- ▶ Funding: Stormwater strategic initiative program



- ▶ *Thank you!*



# Questions?

## References

Anderson, J., Dubetz, C. and Palace, V. 2015 Neonicotinoids in the Canadian aquatic environment: a literature review on current use products with a focus on fate, exposure, and biological effects. *Science of the Total Environment*, 505: 409-422.

Klarich, K.L., Pflug, N.C., DeWald, E.M., Hladik, M.L., Kolpin, D.W., Cwiertny, D.M. and LeFevre, G.H. 2017. Occurrence of neonicotinoid insecticides in finished drinking water and fate during drinking water treatment. *Environmental Science & Technology Letters*, 4(5): 168-173.

WSDA. Ambient Monitoring for Pesticides In Washington State Surface Water. <https://agr.wa.gov/pestfert/natresources/swm/>

Miles, J.C., Hua, J., Sepulveda, M.S., Krupke, C.H. and Hoverman, J.T. 2017. Effects of clothianidin on aquatic communities: Evaluating the impacts of lethal and sublethal exposure to neonicotinoids. *PLoS one*, 12(3), p.e0174171.