



Western Washington University
Western CEDAR

Salish Sea Ecosystem Conference

2018 Salish Sea Ecosystem Conference
(Seattle, Wash.)

Apr 5th, 11:30 AM - 1:30 PM

How did large scale climate anomalies impact 2015 phytoplankton blooms in Puget Sound?

Juhi LaFuente

Washington (State). Department of Ecology, jlaf461@ecy.wa.gov

Christopher Krembs

Washington (State). Department of Ecology, ckre461@ecy.wa.gov

S. L. Albertson

Washington (State). Department of Ecology, salb461@ecy.wa.gov

Allison Brownlee

Washington (State). Department of Ecology, albr461@ecy.wa.gov

Julia Bos

Washington (State). Department of Ecology, jbos461@ecy.wa.gov

See next page for additional authors

Follow this and additional works at: <https://cedar.wwu.edu/ssec>



Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Natural Resources and Conservation Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

LaFuente, Juhi; Krembs, Christopher; Albertson, S. L.; Brownlee, Allison; Bos, Julia; Hermanson, Laura; and Keyzers, Mya, "How did large scale climate anomalies impact 2015 phytoplankton blooms in Puget Sound?" (2018). *Salish Sea Ecosystem Conference*. 183.
<https://cedar.wwu.edu/ssec/2018ssec/allsessions/183>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wwu.edu.

Speaker


Juhi LaFuente, Christopher Krembs, S. L. Albertson, Allison Brownlee, Julia Bos, Laura Hermanson, and Mya Keyzers

Juhi LaFuente^{1,2}, Christopher Krembs¹, Skip Albertson¹, Allison Brownlee¹, Julia Bos¹, Laura Hermanson¹, Mya Keyzers¹
¹Washington State Department of Ecology, ² Washington Conservation Corps

Background

What can we learn from large scale climate anomalies?

The Washington State Department of Ecology's Marine Waters Program has routinely monitored water quality throughout Puget Sound since 1973. Establishing historic baselines at long-term monitoring stations allows us to add context to spatial and temporal trends seen in marine water quality.




Water quality data is collected monthly via floatplane.

In 2015 we observed changes in marine water quality due to the large-scale climate anomaly 'The Blob' – a mass of warm water that entered Puget Sound in the fall of 2014. In conjunction with the Blob, higher than normal air temperatures altered patterns of river discharge in 2015, changing water column stratification and salinity. Changes to hydrological patterns in Puget Sound have the ability to influence nutrient levels and water column stratification, indirectly affecting the timing and amplitude of phytoplankton blooms.

By comparing 2015 marine water quality data to baseline conditions (1999-2008), this study explores how the following played a role in altering the timing and magnitude of phytoplankton blooms in 2015:

- 1) The physical environment
- 2) River discharge
- 3) Nutrient cycling




Water quality samples collected aboard the R/V Skookum.

Why focus on Phytoplankton?

- Ecosystem functioning is reliant on phytoplankton production transferring energy to higher trophic levels.
- Climate impacts modify bloom timing, amplitude and duration resulting in altered energy flow to higher trophic levels.

Methods


- Long-term monitoring stations are visited monthly via floatplane and boat.
- Standard operating procedures are followed for seawater sampling, analysis and data QA/QC.
- Data collected from Central and South Sound in 2015 was compared to an established historic baseline (1999-2008).
- 'Heat' maps were generated to show anomalies in 2015 water quality data.



Central (orange) and South Sound (red) long term monitoring stations. Map by Mya Keyzers

Regional Differences in Phytoplankton Blooms in 2015

Central Sound




- Spring Bloom: Earlier timing, higher amplitude
- Summer Bloom: Expected timing, higher amplitude

Station	Month											
	1	2	3	4	5	6	7	8	9	10	11	12
Admiralty Inlet												
West Point												
Sinclair Inlet												
Elliott Bay												
East Passage												
Commencement Bay												

2015 Central Sound chlorophyll concentrations relative to baseline levels

■ = Higher ■ = Lower □ = Expected ■ = No Data

South Sound



- Spring Bloom: Earlier timing, expected amplitude
- Summer Bloom: Earlier timing, higher amplitude

Station	Month											
	1	2	3	4	5	6	7	8	9	10	11	12
Gordon Point												
Nisqually Reach												
Dana Passage												
Budd Inlet												
Oakland Bay												

2015 South Sound chlorophyll concentrations relative to baseline levels

Factors Influencing Phytoplankton Blooms

1. The Physical Environment

Central Sound

Station	Month											
	1	2	3	4	5	6	7	8	9	10	11	12
Admiralty Inlet												
West Point												
Sinclair Inlet												
Elliott Bay												
East Passage												
Commencement Bay												

Stratification levels were stronger than expected throughout the winter, followed by weaker stratification levels throughout the spring and summer months.

Stratification

South Sound

Station	Month											
	1	2	3	4	5	6	7	8	9	10	11	12
Gordon Point												
Nisqually Reach												
Dana Passage												
Budd Inlet												
Oakland Bay												

Stratification levels varied across all stations throughout 2015.


Salinity

Station	Month											
	1	2	3	4	5	6	7	8	9	10	11	12
Admiralty Inlet												
West Point												
Sinclair Inlet												
Elliott Bay												
East Passage												
Commencement Bay												

Relative to baseline levels salinity was lower in the winter and spring and higher in the summer.

■ = Higher ■ = Lower □ = Expected ■ = No Data

2. River Flow



The Fraser River in Canada is the largest contributor of fresh water to the Salish Sea. Changes in the Fraser River discharge alter the two later exchange of water flowing between Puget Sound and the Pacific Ocean phytoplankton blooms are indirectly affected by river through changes in the the physical and chemical environment.

Winter/Spring


- Premature river discharge
- Stronger stratification levels in Central Sound -> earlier spring bloom
- Lower levels of salinity in both regions

Summer

- Low river flows
- Weaker levels of stratification in Central Sound
- Higher levels of salinity in both regions

South Sound was less affected by stratification likely due to tidal mixing.

3. Nutrient Cycling



In the summer months, higher amounts of reduced nitrogen (low NO₃/DIN ratio) were present in both Central and South Sound. This suggests that more nitrogen was being recycled in the water column compared to previous years.

Conclusions

- Large-scale climate anomalies provide useful information about how warming global and ocean temperatures will impact phytoplankton blooms in Puget Sound.
- Regions in Puget Sound may respond differently to future climate impacts.
- More research on lower trophic level food web dynamics is needed to understand how ecosystem functioning in Puget Sound is affected by changes in the timing and amplitude of phytoplankton blooms.

References

Bos, Julia, et al. "2015 Quality Assurance Monitoring Plan: Long-Term Marine Waters Monitoring, Water Column Program." Washington State Department of Ecology, Olympia, WA. Publication No. 15-03-101 200 pp.
Friedland, Kevin D., et al. "Phenology and time series trends of the dominant seasonal phytoplankton bloom across global scales." Global Ecology and Biogeography, 2018, pp. 1-19
Lewandowska, Aleksandra M., et al. "Effects of Sea Surface warming on marine Plankton." Ecology Letters, 2014
Reum, Jonathan C.P., et al. "Multiscale influence of climate on estuarine populations of forage fish: the role of coastal upwelling, freshwater flow and temperature." Marine Ecology Progress Series, vol. 425, Mar. 14th 2011, pp. 203-215
Scavia, Donald, et al. "Climate Change Impacts on U.S. Coastal and Marine Ecosystems." Estuaries, vol. 25, no. 2, Apr. 2002, pp. 149-164.
Winter, D.F., et al. "The Dynamics of Phytoplankton Blooms in Puget Sound, a Fjord in the Northwestern United States." Marine Biology 1975 vol. 29 pp. 139-176