



3-7-2013

# Lake Whatcom Monitoring Project 2011/2012 Report

Robin A. Matthews

*Western Washington University*, [robin.matthews@wwu.edu](mailto:robin.matthews@wwu.edu)

Michael Hilles

*Western Washington University*, [michael.hilles@wwu.edu](mailto:michael.hilles@wwu.edu)

Joan Vandersypen

*Western Washington University*, [joan.vandersypen@wwu.edu](mailto:joan.vandersypen@wwu.edu)

Robert J. Mitchell

*Western Washington University*, [robert.mitchell@wwu.edu](mailto:robert.mitchell@wwu.edu)

Geoffrey B. Matthews

*Western Washington University*, [geoffrey.matthews@wwu.edu](mailto:geoffrey.matthews@wwu.edu)

Follow this and additional works at: [https://cedar.wwu.edu/lakewhat\\_annualreps](https://cedar.wwu.edu/lakewhat_annualreps)



Part of the [Environmental Monitoring Commons](#)

---

## Recommended Citation

Matthews, Robin A.; Hilles, Michael; Vandersypen, Joan; Mitchell, Robert J.; and Matthews, Geoffrey B., "Lake Whatcom Monitoring Project 2011/2012 Report" (2013). *Lake Whatcom Annual Reports*. 2.

[https://cedar.wwu.edu/lakewhat\\_annualreps/2](https://cedar.wwu.edu/lakewhat_annualreps/2)

This Report is brought to you for free and open access by the Lake Whatcom at Western CEDAR. It has been accepted for inclusion in Lake Whatcom Annual Reports by an authorized administrator of Western CEDAR. For more information, please contact [westerncedar@wwu.edu](mailto:westerncedar@wwu.edu).

# Lake Whatcom Monitoring Project 2011/2012 Report

Dr. Robin A. Matthews  
Michael Hilles  
Joan Vandersypen  
*Institute for Watershed Studies,  
Huxley College of the Environment*

Dr. Robert J. Mitchell  
*Geology Department,  
College of Sciences and Technology*

Dr. Geoffrey B. Matthews  
*Computer Science Department,  
College of Sciences and Technology*

Western Washington University  
Bellingham, Washington 98225

March 7, 2013

---

Funding for this project was provided by the City of Bellingham, as part of their long-term commitment to environmental education and their concern for maintaining the water quality of Lake Whatcom. We thank Marilyn Desmul, James Bucher, Cassie Castrejon, Sean Cooper, Miette Dahlgren, Jonnel Deacon, Neddy Dondup, Olivia Edwards, Chad Elofson, Josh Jones, Laura Junge, Andrew Majeske, Niki Thane, Paul Wanzek-Wild, Chantal Wikstrom, Shaun Weldon, and Jordan Zanmiller for assistance with the project.

(This page blank)

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Lake Whatcom Monitoring</b>	<b>2</b>
2.1	Site Descriptions . . . . .	2
2.2	Field Sampling and Analytical Methods . . . . .	2
2.3	Results and Discussion . . . . .	3
2.3.1	Water temperature . . . . .	4
2.3.2	Dissolved oxygen . . . . .	5
2.3.3	Conductivity and pH . . . . .	7
2.3.4	Alkalinity and turbidity . . . . .	7
2.3.5	Nitrogen and phosphorus . . . . .	8
2.3.6	Chlorophyll, plankton, and Secchi depth . . . . .	11
2.3.7	Coliform bacteria . . . . .	13
2.3.8	Metals . . . . .	14
2.3.9	Total organic carbon and disinfection by-products . . . . .	14
<b>3</b>	<b>Tributary Monitoring</b>	<b>37</b>
3.1	Site Descriptions . . . . .	37
3.2	Field Sampling and Analytical Methods . . . . .	37
3.3	Results and Discussion . . . . .	38
<b>4</b>	<b>Lake Whatcom Hydrology</b>	<b>56</b>
4.1	Hydrograph Data . . . . .	56
4.2	Water Budget . . . . .	56

<b>5</b>	<b>Storm Water Monitoring</b>	<b>69</b>
5.1	Site Descriptions . . . . .	69
5.2	Field Sampling and Analytical Methods . . . . .	69
5.3	Results and Discussion . . . . .	70
<b>6</b>	<b>References and Related Reports</b>	<b>87</b>
6.1	References . . . . .	87
6.2	Related Reports . . . . .	89
<b>A</b>	<b>Site Descriptions</b>	<b>93</b>
A.1	Lake Whatcom Monitoring Sites . . . . .	93
A.2	Tributary Monitoring Sites . . . . .	94
A.3	Storm Water Monitoring Sites . . . . .	95
<b>B</b>	<b>Long-Term Water Quality Figures</b>	<b>101</b>
B.1	Monthly Hydrolab Profiles . . . . .	102
B.2	Long-term Hydrolab Data (1988-present) . . . . .	153
B.3	Long-term Water Quality Data (1988-present) . . . . .	174
B.4	Lake Whatcom Tributary Data (2004-present) . . . . .	235
<b>C</b>	<b>Quality Control</b>	<b>275</b>
C.1	Performance Evaluation Report . . . . .	275
C.2	Laboratory Duplicates, Spikes, and Check Standards . . . . .	277
C.3	Field Duplicate Results . . . . .	308
<b>D</b>	<b>Lake Whatcom Online Data</b>	<b>329</b>

## List of Figures

1	Boxplots showing 1988-2012 surface water temperatures . . . . .	25
2	Relationship between dissolved oxygen and time at Site 1, 12 m .	26
3	Relationship between dissolved oxygen and time at Site 1, 14 m .	27
4	Relationship between dissolved oxygen and time at Site 1, 16 m .	28
5	Relationship between dissolved oxygen and time at Site 1, 18 m .	29
6	Minimum summer, near-surface dissolved inorganic nitrogen concentrations . . . . .	30
7	Median summer, near-surface total phosphorus concentrations . .	31
8	Median summer near-surface chlorophyll concentrations . . . . .	32
9	Log <sub>10</sub> plots of median summer, near-surface algae counts . . . . .	33
10	Log <sub>10</sub> plots of median summer, near-surface Cyanobacteria counts	34
11	Annual total organic carbon concentrations at Sites 1–4. . . . .	35
12	Total trihalomethanes and haloacetic acids concentrations in the Bellingham water distribution system . . . . .	36
13	Austin Creek hydrograph, October 1, 2011–September 30, 2012. .	62
14	Smith Creek hydrograph, October 1, 2011–September 30, 2012. .	63
15	Comparison of Lake Whatcom daily lake volumes for WY2008–WY2012. . . . .	64
16	Summary of 7-day inputs, outputs, and changes in Lake Whatcom storage, October 1, 2011–September 30, 2012. . . . .	65
17	Lake Whatcom watershed direct hydrologic inputs, October 1, 2011–September 30, 2012. . . . .	66
18	Lake Whatcom watershed hydrologic withdrawals, October 1, 2011–September 30, 2012. . . . .	67

19	Summary of 7-day Whatcom Creek flows, water balance runoff estimates, and DHSVM runoff estimates, October 1, 2011–September 30, 2012. . . . .	68
20	Silver Beach Creek storm water monitoring results for Events 17–24: total suspended solids vs. stream flow. . . . .	73
21	Silver Beach Creek storm water monitoring results for Events 17–24: turbidity vs. stream flow. . . . .	74
22	Silver Beach Creek storm water monitoring results for Events 17–24: total phosphorus vs. stream flow. . . . .	75
23	Silver Beach Creek storm water monitoring results for Events 17–24: soluble phosphate vs. stream flow. . . . .	76
24	Silver Beach Creek storm water monitoring results for Events 17–24: total nitrogen vs. stream flow. . . . .	77
25	Silver Beach Creek storm water monitoring results for Events 17–24: nitrate/nitrite vs. stream flow. . . . .	78
26	Correlation between stream flow or stage height and total suspended solids in Silver Beach Creek. . . . .	79
27	Correlation between stream flow or stage height and turbidity in Silver Beach Creek. . . . .	80
28	Correlation between stream flow or stage height and total phosphorus in Silver Beach Creek. . . . .	81
29	Correlation between stream flow or stage height and soluble phosphate in Silver Beach Creek. . . . .	82
30	Correlation between stream flow or stage height and total nitrogen in Silver Beach Creek. . . . .	83
31	Correlation between stream flow or stage height and nitrate in Silver Beach Creek. . . . .	84
32	Correlation between total suspended solids, turbidity, and total phosphorus in Silver Beach Creek. . . . .	85

33	Correlation between stream flow and total phosphorus by storm event in Silver Beach Creek. . . . .	86
A1	Lake Whatcom lake sampling sites. . . . .	97
A2	Lake Whatcom tributary sampling sites. . . . .	98
A3	Silver Beach Creek storm water site. . . . .	99
B1	Lake Whatcom Hydrolab profiles for Site 1, October 4, 2011 . . .	103
B2	Lake Whatcom Hydrolab profiles for Site 2, October 4, 2011 . . .	104
B3	Lake Whatcom Hydrolab profiles for the Intake, October 4, 2011 .	105
B4	Lake Whatcom Hydrolab profiles for Site 3, October 5, 2011 . . .	106
B5	Lake Whatcom Hydrolab profiles for Site 4, October 5, 2011 . . .	107
B6	Lake Whatcom YSI profiles for Site 1, November 1, 2011. . . . .	108
B7	Lake Whatcom YSI profiles for Site 2, November 1, 2011. . . . .	109
B8	Lake Whatcom YSI profiles for the Intake, November 1, 2011. . .	110
B9	Lake Whatcom YSI profiles for Site 3, November 2, 2011. . . . .	111
B10	Lake Whatcom YSI profiles for Site 4, November 2, 2011. . . . .	112
B11	Lake Whatcom YSI profiles for Site 1, December 7, 2011. . . . .	113
B12	Lake Whatcom YSI profiles for Site 2, December 7, 2011. . . . .	114
B13	Lake Whatcom YSI profiles for the Intake, December 7, 2011. . .	115
B14	Lake Whatcom YSI profiles for Site 3, December 6, 2011. The low oxygen value at 80 meters was most likely due to incomplete water column mixing following recent destratification. . . . .	116
B15	Lake Whatcom YSI profiles for Site 4, December 6, 2011. . . . .	117
B16	Lake Whatcom YSI profiles for Site 1, February 3, 2012. . . . .	118
B17	Lake Whatcom YSI profiles for Site 2, February 3, 2012. . . . .	119
B18	Lake Whatcom YSI profiles for the Intake, February 3, 2012. . . .	120
B19	Lake Whatcom YSI profiles for Site 3, February 1, 2012. . . . .	121



B20	Lake Whatcom YSI profiles for Site 4, February 1, 2012. . . . .	122
B21	Lake Whatcom YSI profiles for Site 1, April 14, 2012. . . . .	123
B22	Lake Whatcom YSI profiles for Site 2, April 14, 2012. . . . .	124
B23	Lake Whatcom YSI profiles for the Intake, April 14, 2012. . . . .	125
B24	Lake Whatcom YSI profiles for Site 3, April 10, 2012. . . . .	126
B25	Lake Whatcom YSI profiles for Site 4, April 10, 2012. . . . .	127
B26	Lake Whatcom YSI profiles for Site 1, May 10, 2012. . . . .	128
B27	Lake Whatcom YSI profiles for Site 2, May 10, 2012. . . . .	129
B28	Lake Whatcom YSI profiles for the Intake, May 10, 2012. . . . .	130
B29	Lake Whatcom YSI profiles for Site 3, May 3, 2012. . . . .	131
B30	Lake Whatcom YSI profiles for Site 4, May 3, 2012. . . . .	132
B31	Lake Whatcom YSI profiles for Site 1, June 7, 2012. . . . .	133
B32	Lake Whatcom YSI profiles for Site 2, June 7, 2012. . . . .	134
B33	Lake Whatcom YSI profiles for the Intake, June 7, 2012. . . . .	135
B34	Lake Whatcom YSI profiles for Site 3, June 9, 2012. . . . .	136
B35	Lake Whatcom YSI profiles for Site 4, June 9, 2012. . . . .	137
B36	Lake Whatcom YSI profiles for Site 1, July 7, 2012. . . . .	138
B37	Lake Whatcom YSI profiles for Site 2, July 7, 2012. . . . .	139
B38	Lake Whatcom YSI profiles for the Intake, July 7, 2012. . . . .	140
B39	Lake Whatcom YSI profiles for Site 3, July 5, 2012. . . . .	141
B40	Lake Whatcom YSI profiles for Site 4, July 5, 2012. . . . .	142
B41	Lake Whatcom YSI profiles for Site 1, August 4, 2012. . . . .	143
B42	Lake Whatcom YSI profiles for Site 2, August 4, 2012. . . . .	144
B43	Lake Whatcom YSI profiles for the Intake, August 4, 2012. . . . .	145
B44	Lake Whatcom YSI profiles for Site 3, August 2, 2012. . . . .	146

B45	Lake Whatcom YSI profiles for Site 4, August 2, 2012. . . . .	147
B46	Lake Whatcom YSI profiles for Site 1, September 8, 2012. . . . .	148
B47	Lake Whatcom YSI profiles for Site 2, September 8, 2012. . . . .	149
B48	Lake Whatcom YSI profiles for the Intake, September 8, 2012. . . . .	150
B49	Lake Whatcom YSI profiles for Site 3, September 6, 2012. . . . .	151
B50	Lake Whatcom YSI profiles for Site 4, September 6, 2012. . . . .	152
B51	Lake Whatcom historic temperature data for Site 1. . . . .	154
B52	Lake Whatcom historic temperature data for Site 2. . . . .	155
B53	Lake Whatcom historic temperature data for the Intake. . . . .	156
B54	Lake Whatcom historic temperature data for Site 3. . . . .	157
B55	Lake Whatcom historic temperature data for Site 4. . . . .	158
B56	Lake Whatcom historic dissolved oxygen data for Site 1. . . . .	159
B57	Lake Whatcom historic dissolved oxygen data for Site 2. . . . .	160
B58	Lake Whatcom historic dissolved oxygen data for the Intake. . . . .	161
B59	Lake Whatcom historic dissolved oxygen data for Site 3. . . . .	162
B60	Lake Whatcom historic dissolved oxygen data for Site 4. . . . .	163
B61	Lake Whatcom historic pH data for Site 1. . . . .	164
B62	Lake Whatcom historic pH data for Site 2. . . . .	165
B63	Lake Whatcom historic pH data for the Intake. . . . .	166
B64	Lake Whatcom historic pH data for Site 3. . . . .	167
B65	Lake Whatcom historic pH data for Site 4. . . . .	168
B66	Lake Whatcom historic conductivity data for Site 1. . . . .	169
B67	Lake Whatcom historic conductivity data for Site 2. . . . .	170
B68	Lake Whatcom historic conductivity data for the Intake. . . . .	171
B69	Lake Whatcom historic conductivity data for Site 3. . . . .	172

B70	Lake Whatcom historic conductivity data for Site 4. . . . .	173
B71	Lake Whatcom alkalinity data for Site 1. . . . .	175
B72	Lake Whatcom alkalinity data for Site 2. . . . .	176
B73	Lake Whatcom alkalinity data for the Intake site. . . . .	177
B74	Lake Whatcom alkalinity data for Site 3. . . . .	178
B75	Lake Whatcom alkalinity data for Site 4. . . . .	179
B76	Lake Whatcom turbidity data for Site 1. . . . .	180
B77	Lake Whatcom turbidity data for Site 2. . . . .	181
B78	Lake Whatcom turbidity data for the Intake site. . . . .	182
B79	Lake Whatcom turbidity data for Site 3. . . . .	183
B80	Lake Whatcom turbidity data for Site 4. . . . .	184
B81	Lake Whatcom ammonium data for Site 1. . . . .	185
B82	Lake Whatcom ammonium data for Site 2. . . . .	186
B83	Lake Whatcom ammonium data for the Intake site. . . . .	187
B84	Lake Whatcom ammonium data for Site 3. . . . .	188
B85	Lake Whatcom ammonium data for Site 4. . . . .	189
B86	Lake Whatcom nitrate/nitrite data for Site 1. . . . .	190
B87	Lake Whatcom nitrate/nitrite data for Site 2. . . . .	191
B88	Lake Whatcom nitrate/nitrite data for the Intake site. . . . .	192
B89	Lake Whatcom nitrate/nitrite data for Site 3. . . . .	193
B90	Lake Whatcom nitrate/nitrite data for Site 4. . . . .	194
B91	Lake Whatcom total nitrogen data for Site 1. . . . .	195
B92	Lake Whatcom total nitrogen data for Site 2. . . . .	196
B93	Lake Whatcom total nitrogen data for the Intake site. . . . .	197
B94	Lake Whatcom total nitrogen data for Site 3. . . . .	198

B95	Lake Whatcom total nitrogen data for Site 4. . . . .	199
B96	Lake Whatcom soluble phosphate data for Site 1. . . . .	200
B97	Lake Whatcom soluble phosphate data for Site 2. . . . .	201
B98	Lake Whatcom soluble phosphate data for the Intake site. . . . .	202
B99	Lake Whatcom soluble phosphate data for Site 3. . . . .	203
B100	Lake Whatcom soluble phosphate data for Site 4. . . . .	204
B101	Lake Whatcom total phosphorus data for Site 1. . . . .	205
B102	Lake Whatcom total phosphorus data for Site 2. . . . .	206
B103	Lake Whatcom total phosphorus data for the Intake site. . . . .	207
B104	Lake Whatcom total phosphorus data for Site 3. . . . .	208
B105	Lake Whatcom total phosphorus data for Site 4. . . . .	209
B106	Lake Whatcom chlorophyll data for Site 1. . . . .	210
B107	Lake Whatcom chlorophyll data for Site 2. . . . .	211
B108	Lake Whatcom chlorophyll data for the Intake site. . . . .	212
B109	Lake Whatcom chlorophyll data for Site 3. . . . .	213
B110	Lake Whatcom chlorophyll data for Site 4. . . . .	214
B111	Lake Whatcom Secchi depths for Site 1. . . . .	215
B112	Lake Whatcom Secchi depths for Site 2. . . . .	216
B113	Lake Whatcom Secchi depths for the Intake site. . . . .	217
B114	Lake Whatcom Secchi depths for Site 3. . . . .	218
B115	Lake Whatcom Secchi depths for Site 4. . . . .	219
B116	Lake Whatcom fecal coliform data for Site 1. . . . .	220
B117	Lake Whatcom fecal coliform data for Site 2. . . . .	221
B118	Lake Whatcom fecal coliform data for the Intake site. . . . .	222
B119	Lake Whatcom fecal coliform data for Site 3. . . . .	223

B120	Lake Whatcom fecal coliform data for Site 4. . . . .	224
B121	Lake Whatcom plankton data for Site 1. . . . .	225
B122	Lake Whatcom plankton data for Site 2. . . . .	226
B123	Lake Whatcom plankton data for the Intake Site. . . . .	227
B124	Lake Whatcom plankton data for Site 3. . . . .	228
B125	Lake Whatcom plankton data for Site 4. . . . .	229
B126	Lake Whatcom plankton data for Site 1, with Chrysophyta omitted to show remaining plankton groups. . . . .	230
B127	Lake Whatcom plankton data for Site 2, with Chrysophyta omitted to show remaining plankton groups. . . . .	231
B128	Lake Whatcom plankton data for the Intake Site, with Chrysophyta omitted to show remaining plankton groups. . . . .	232
B129	Lake Whatcom plankton data for Site 3, with Chrysophyta omitted to show remaining plankton groups. . . . .	233
B130	Lake Whatcom plankton data for Site 4, with Chrysophyta omitted to show remaining plankton groups. . . . .	234
B131	Temperature data for Anderson, Austin, Smith, and Whatcom Creeks. . . . .	236
B132	Temperature data for Blue Canyon, Brannian, Carpenter, and Olsen Creek. . . . .	237
B133	Temperature data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	238
B134	Dissolved oxygen data for Anderson, Austin, Smith, and Whatcom Creeks. . . . .	239
B135	Dissolved oxygen data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	240
B136	Dissolved oxygen data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	241

B137 Tributary pH data for Anderson, Austin, Smith, and Whatcom Creeks. . . . .	242
B138 Tributary pH data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	243
B139 Tributary pH data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	244
B140 Conductivity data for Anderson, Austin, Smith, and Whatcom Creeks. . . . .	245
B141 Conductivity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	246
B142 Conductivity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	247
B143 Alkalinity data for Anderson, Austin, Smith, and Whatcom Creeks.	248
B144 Alkalinity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	249
B145 Alkalinity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	250
B146 Total suspended solids data for Anderson, Austin, Smith, and Whatcom Creeks. . . . .	251
B147 Total suspended solids data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	252
B148 Total suspended solids data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	253
B149 Turbidity data for Anderson, Austin, Smith, and Whatcom Creeks.	254
B150 Turbidity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	255
B151 Turbidity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	256

B152 Ammonium data for Anderson, Austin, Smith, and Whatcom Creeks. . . . .	257
B153 Ammonium data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	258
B154 Ammonium data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	259
B155 Nitrate/nitrite data for Anderson, Austin, Smith, and Whatcom Creeks. . . . .	260
B156 Nitrate/nitrite data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	261
B157 Nitrate/nitrite data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	262
B158 Total nitrogen data for Anderson, Austin, Smith, and Whatcom Creeks. . . . .	263
B159 Total nitrogen data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	264
B160 Total nitrogen data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	265
B161 Soluble phosphate data for Anderson, Austin, Smith, and What- com Creeks. . . . .	266
B162 Soluble phosphate data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	267
B163 Soluble phosphate data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	268
B164 Total phosphorus data for Anderson, Austin, Smith, and Whatcom Creeks. . . . .	269
B165 Total phosphorus data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. . . . .	270
B166 Total phosphorus data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. . . . .	271

B167	Fecal coliform data for Anderson, Austin, Smith, and Whatcom Creeks (short scale).	272
B168	Fecal coliform data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks (short scale).	273
B169	Fecal coliform data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain (short scale).	274
C1	Alkalinity laboratory duplicates for the Lake Whatcom monitoring program.	278
C2	Alkalinity high-range check standards for the Lake Whatcom monitoring program.	279
C3	Alkalinity low-range check standards for the Lake Whatcom monitoring program.	280
C4	Chlorophyll laboratory duplicates for the Lake Whatcom monitoring program.	281
C5	Conductivity laboratory duplicates for the Lake Whatcom monitoring program.	282
C6	Dissolved oxygen laboratory duplicates for the Lake Whatcom monitoring program.	283
C7	Ammonium laboratory duplicates for the Lake Whatcom monitoring program.	284
C8	Ammonium matrix spikes for the Lake Whatcom monitoring program.	285
C9	Ammonium high-range check standards for the Lake Whatcom monitoring program.	286
C10	Ammonium low-range check standards for the Lake Whatcom monitoring program.	287
C11	Nitrate/nitrite laboratory duplicates for the Lake Whatcom monitoring program.	288
C12	Nitrate/nitrite matrix spikes for the Lake Whatcom monitoring program.	289



C13	Nitrate/nitrite high-range check standards for the Lake Whatcom monitoring program. . . . .	290
C14	Nitrate/nitrite low-range check standards for the Lake Whatcom monitoring program. . . . .	291
C15	Total nitrogen laboratory duplicates for the Lake Whatcom monitoring program. . . . .	292
C16	Total nitrogen matrix spikes for the Lake Whatcom monitoring program. . . . .	293
C17	Total nitrogen high-range check standards for the Lake Whatcom monitoring program. . . . .	294
C18	Total nitrogen low-range check standards for the Lake Whatcom monitoring program. . . . .	295
C19	Laboratory pH duplicates for the Lake Whatcom monitoring program. . . . .	296
C20	Soluble reactive phosphate laboratory duplicates for the Lake Whatcom monitoring program. . . . .	297
C21	Soluble reactive phosphate matrix spikes for the Lake Whatcom monitoring program. . . . .	298
C22	Soluble reactive phosphate high-range check standards for the Lake Whatcom monitoring program. . . . .	299
C23	Soluble reactive phosphate low-range check standards for the Lake Whatcom monitoring program. . . . .	300
C24	Total phosphorus laboratory duplicates for the Lake Whatcom monitoring program. . . . .	301
C25	Total phosphorus matrix spikes for the Lake Whatcom monitoring program. . . . .	302
C26	Total phosphorus high-range check standards for the Lake Whatcom monitoring program. . . . .	303
C27	Total phosphorus low-range check standards for the Lake Whatcom monitoring program. . . . .	304

C28	Total suspended solids laboratory duplicates for the Lake Whatcom monitoring program. . . . .	305
C29	Total suspended solids check standards for the Lake Whatcom monitoring program. . . . .	306
C30	Turbidity laboratory duplicates for the Lake Whatcom monitoring program. . . . .	307
C31	Alkalinity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . .	309
C32	Alkalinity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). . . . .	310
C33	Chlorophyll field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . .	311
C34	Conductivity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . .	312
C35	Dissolved oxygen field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . .	313
C36	Dissolved oxygen field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). . . . .	314
C37	Ammonium field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . .	315
C38	Ammonium field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). . . . .	316
C39	Nitrate/nitrite field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . .	317
C40	Nitrate/nitrite field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). . . . .	318
C41	Total nitrogen field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . .	319
C42	Total nitrogen field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). . . . .	320

C43 Field duplicates for pH from the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . . 321

C44 Soluble phosphorus field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). . . . . 322

C45 Total phosphorus field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . . 323

C46 Total phosphorus field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). . . . . 324

C47 Total suspended solids field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). . . . . 325

C48 Turbidity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). . . . . 326

C49 Turbidity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). . . . . 327

## List of Tables

1	Summary of IWS, AmTest, Edge Analytical, and City of Bellingham analytical methods and parameter abbreviations. . . . .	16
2	Summary of Site 1 water quality data, Oct. 2011 – Sept. 2012. . .	17
3	Summary of Intake water quality data, Oct. 2011– Sept. 2012. . .	18
4	Summary of Site 2 water quality data, Oct. 2011 – Sept. 2012. . .	19
5	Summary of Site 3 water quality data, Oct. 2011 – Sept. 2012. . .	20
6	Summary of Site 4 water quality data, Oct. 2011 – Sept. 2012. . .	21
7	October hypolimnetic ammonium and hydrogen sulfide concentrations at Sites 1 and 2 . . . . .	22
8	Lake Whatcom 2011/2012 total metals data . . . . .	23
9	Lake Whatcom 2011/2012 total organic carbon data. . . . .	24
10	Comparison of water quality features in Lake Whatcom tributaries.	41
11	Summary of Anderson Creek water quality data, Oct. 2011–Sept. 2012. . . . .	42
12	Summary of Austin Creek water quality data, Oct. 2011–Sept. 2012.	43
13	Summary of Blue Canyon Creek water quality data, Oct. 2011–Sept. 2012. . . . .	44
14	Summary of Brannian Creek water quality data, Oct. 2011–Sept. 2012. . . . .	45
15	Summary of Carpenter Creek water quality data, Oct. 2011–Sept. 2012. . . . .	46
16	Summary of Euclid Creek water quality data, Oct. 2011–Sept. 2012.	47
17	Summary of Millwheel Creek water quality data, Oct. 2011–Sept. 2012. . . . .	48
18	Summary of Olsen Creek water quality data, Oct. 2011–Sept. 2012.	49

19	Summary of Park Place drain water quality data, Oct. 2011–Sept. 2012. . . . .	50
20	Summary of Silver Beach Creek water quality data, Oct. 2011–Sept. 2012. . . . .	51
21	Summary of Smith Creek water quality data, Oct. 2011–Sept. 2012.	52
22	Summary of Whatcom Creek water quality data, Oct. 2011–Sept. 2012. . . . .	53
23	Lake Whatcom tributary data: total metals. . . . .	54
24	Lake Whatcom tributary data: total organic carbon. . . . .	55
25	Annual water balance quantities for the Lake Whatcom watershed, WY2008–WY2012. . . . .	59
26	Monthly input water balance quantities for the Lake Whatcom watershed, October 2011–September 2012. . . . .	60
27	Monthly output water balance quantities for the Lake Whatcom watershed, October 2011–September 2012. . . . .	61
28	Summary of Silver Beach Creek storm events and precipitation at the Bloedel/Donovan precipitation gauge. . . . .	72
A1	Approximate GPS coordinates for Lake Whatcom sampling sites.	96
C1	Single-blind quality control results, WP–183 (06/06/2012). . . . .	276

## Executive Summary

- This report describes the results from the 2011/2012 Lake Whatcom monitoring program. The major objectives were to continue long-term baseline water quality monitoring in Lake Whatcom and selected tributary streams; collect storm runoff water quality data from Silver Beach Creek; continue collection of hydrologic data from Austin and Smith Creeks; and update the hydrologic model for Lake Whatcom.
- This report is part of an on-going series of annual reports and special project reports that provide a complete documentation of the monitoring program over time. A summary of the IWS Lake Whatcom reports, including special project reports, is included in Section 6.2, beginning on page 89.
- During the summer the lake stratified into a warm surface layer (the epilimnion) and a cool bottom layer (the hypolimnion). The water temperatures were near historic median values during most of the year except in June 2012, which was slightly cooler than usual. Despite slightly cooler temperatures, all sites except the Intake were stratified by early June.
- The levels of hypolimnetic oxygen have declined over time at Site 1, causing the lake to be listed by the Department of Ecology on the 1998 303d list of impaired waterbodies in the State of Washington. Following the onset of stratification, the hypolimnetic oxygen concentrations dropped rapidly. By August 8, 2012, the oxygen concentration was  $<1$  mg/L from 12 meters to the bottom.
- Nitrate depletion was evident at all sites in the photosynthetic zone during the summer due to algal uptake of this essential nutrient. Low nitrate in the photosynthetic zone favors the growth of Cyanobacteria. Nitrate depletion also occurred in the hypolimnion at Sites 1 and 2 due to nitrate reduction by bacteria.
- Anaerobic conditions in the hypolimnion at Sites 1 and 2 resulted in elevated concentrations of ammonium by the end of the summer. The concentrations were lower than usual in October 2011, following a cool summer, but were typical for the lake in October 2012.

- The summer near-surface total phosphorus and chlorophyll concentrations have increased significantly over time at most sites. The patterns continue to be somewhat variable, but it appears that the trends may have reached a plateau.
- The concentrations of trihalomethanes in Bellingham’s treated drinking water have been increasing over time, particularly during the late summer/fall (third quarter). The total THMS and HAAS remained below the recommended maximum contaminant levels of 0.080 mg/L and 0.060 mg/L, respectively.
- All of the mid-basin fecal coliforms counts were less than 10 cfu/100 mL. The coliform counts at the Bloedel-Donovan recreational area (collected offshore from the swimming area) were slightly higher than mid-basin counts, but passed the freshwater *Extraordinary Primary Contact Recreational* bacteria standard for Washington State.
- Iron and zinc were often detectable, but were within normal ranges for the lake. Other metals were occasionally detected, but the concentrations were near the limits of detection.
- Beginning in January 2010, 11 lake tributaries and Whatcom Creek were sampled monthly to collect baseline data. Most of the tributaries had relatively low concentrations of total and dissolved solids, low alkalinities and conductivities, and low levels of nitrate and ammonium. Residential streams had higher concentrations of total and dissolved solids, higher alkalinities and conductivities, higher coliform counts, and higher nutrient concentrations.
- A water balance was applied to Lake Whatcom to identify its major water inputs and outputs and to examine runoff and storage. The major inputs into the lake during WY2012<sup>1</sup> included surface and subsurface runoff (74.6%), direct precipitation (19.0%), and water diverted from the Middle Fork of the Nooksack River (6.4%). Outputs included Whatcom Creek (79.9%), the City of Bellingham (9.9%), evaporation (7.0%), the Whatcom Falls Hatchery (2.3%), the Lake Whatcom Water and Sewer District (0.6%)<sup>2</sup>, and the Puget Sound Energy Co-Generation Plant (0.1%)<sup>3</sup>.

---

<sup>1</sup>Water Year 2012 covers the period from October 1, 2011 through September 30, 2012

<sup>2</sup>Formerly Water District #10

<sup>3</sup>This facility currently operates at the former Georgia Pacific site.

- Eight storm events were monitored in Silver Beach Creek using an automated sampler to collect flow-paced, discrete samples. The storm runoff contained elevated levels of total suspended solids, turbidity, and phosphorus that were significantly correlated with flow rates. In addition, total suspended solids, turbidity, and total phosphorus concentrations were highly correlated with each other.



(This page blank)

## 1 Introduction

This report is part of an on-going series of annual reports and special project reports that document the Lake Whatcom monitoring program over time. Many of the reports are available online at <http://www.wvu.edu/iws>. Older reports are available in the IWS library and through the City of Bellingham Public Works Department. A summary of the Lake Whatcom reports, including special project reports, is included in Section 6.2, beginning on page 89.

Lake Whatcom is the primary drinking water source for the City of Bellingham and parts of Whatcom County, including Sudden Valley. Lake Whatcom also serves as a water source for the Puget Sound Energy Co-Generation Plant, which is located at the former Georgia-Pacific Corporation site on Bellingham Bay.<sup>4</sup> The lake and parts of the watershed provide recreational opportunities, as well as providing important habitats for fish and wildlife. The lake is used as a storage reservoir to buffer peak storm water flows in Whatcom Creek. Much of the watershed is zoned for forestry and is managed by state or private timber companies. Because of its aesthetic appeal, much of the watershed is highly valued for residential development.

The City of Bellingham and Western Washington University have collaborated on investigations of the water quality in Lake Whatcom since the early 1960s. Beginning in 1981, a monitoring program was initiated by the City and WWU that was designed to provide long-term data for Lake Whatcom for basic parameters such as temperature, pH, dissolved oxygen, conductivity, turbidity, nutrients (nitrogen and phosphorus), and other representative water quality measurements. The major goal of the long-term monitoring effort is to provide a record of Lake Whatcom's water quality over time.

The major objectives of the 2011/2012 Lake Whatcom monitoring program were to continue long-term baseline water quality monitoring in Lake Whatcom and selected tributary streams; collect storm runoff water quality data from Silver Beach Creek; continue collection of hydrologic data from Austin and Smith Creeks; and update the hydrologic model for Lake Whatcom.

---

<sup>4</sup>The Georgia-Pacific Corporation closed its Bellingham pulp mill operations in 2001, reducing its water requirements from 30–35 MGD to 7–12 MGD. By 2007 the water requirements had been reduced to 0.6–3.88 MGD; the mill closed its operations in December 2007.

Detailed site descriptions can be found in Appendix A. The historic lake data are plotted in Appendix B. The current quality control results can be found in Appendix C. The monitoring data are available online at <http://www.wvu.edu/iws> as described in Appendix D (page 329). Table 1 (page 16) lists abbreviations and units used to describe water quality analyses in this document.

## **2 Lake Whatcom Monitoring**

### **2.1 Site Descriptions**

Water quality samples were collected at five long-term monitoring sites in Lake Whatcom (Figure A1, page 97 in Appendix A.1). Sites 1–2 are located at the deepest points in their respective basins. The Intake site is located adjacent to the underwater intake point where the City of Bellingham withdraws lake water from basin 2. Site 3 is located at the deepest point in the northern sub-basin of basin 3 (north of the Sunnyside sill), and Site 4 is located at the deepest point in the southern sub-basin of basin 3 (south of the Sunnyside sill). Water samples were also collected at the City of Bellingham Water Treatment Plant gatehouse, which is located onshore and west of the intake site.

### **2.2 Field Sampling and Analytical Methods**

The lake was sampled on October 4 & 5, November 1 & 2 and December 6 & 7, 2011; and February 7 & 9, April 10 & 12, May 8 & 10, June 12 & 14, July 10 & 12, August 7 & 9, and September 4 & 6, 2012. Each sampling event is a multi-day task; all samples were collected during daylight hours, typically between 10:00 am and 3:00 pm.

A Hydrolab or a YSI field meter was used to measure temperature, pH, dissolved oxygen, and conductivity.<sup>5</sup> Raw water samples were collected using a VanDorn sampler. All water samples (including bacteriological samples) collected in the

---

<sup>5</sup>The Hydrolab Surveyor 4 field meter was used for field sampling in October 2011, but was replaced to resolve on-going issues with the conductivity and pH probes. Beginning in November 2011, field measurements were collected using a YSI 6600 V2 field meter. No major differences have been observed between the results from the Hydrolab vs. the YSI field meters.

field were stored on ice and in the dark until they reached the laboratory, and were analyzed as described in Table 1 (page 16). Total metals analyses (arsenic, cadmium, chromium, copper, iron, mercury, nickel, lead, and zinc) and total organic carbon analyses were done by AmTest.<sup>6</sup> Plankton samples were placed in a cooler and returned to the laboratory unpreserved. The plankton sample volumes were measured in the laboratory and the samples were preserved with Lugol's solution. The bacteria samples were analyzed by the City of Bellingham.

## 2.3 Results and Discussion

The lake monitoring data include monthly field measurements (conductivity, dissolved oxygen, pH, Secchi depth, and water temperature); laboratory analyses for ambient water quality parameters (ammonium<sup>7</sup>, nitrate/nitrite, total nitrogen, soluble phosphate, total phosphorus, alkalinity, turbidity, chlorophyll); plankton and bacteria counts; and biannual metals and total organic carbon measurements.

Tables 2–6 (pages 17–21) summarize the current field measurements, ambient water quality, and coliform data. The raw data are available online at <http://www.wvu.edu/iws> as described in Appendix D (page 329). The monthly profiles for temperature, dissolved oxygen, conductivity, and pH are plotted in Figures B1–B50 (pages 103–152).

The 2011/2012 lake data are plotted with historic lake data in Figures B51–B130 (pages 154–234). These figures are scaled to plot the full range of Lake Whatcom water quality data including minimum, maximum, and outlier values, and do not provide the best illustration of trends that occur in the lake. Separate tables and figures are provided to show trends and illustrate specific patterns in the data.

---

<sup>6</sup>AmTest, 13600 Northeast 126th Place, Suite C, Kirkland, WA, 98034–8720.

<sup>7</sup>Ammonium ( $\text{NH}_4^+$ ) is ionized ammonia ( $\text{NH}_3$ ). Nearly all ammonia is ionized in surface water. Earlier IWS reports used the term ammonia and ammonium interchangeably to describe ammonium concentrations because it is generally understood that ammonia is usually ionized. To improve clarity, IWS has switched to the term “ammonium” to indicate that we are reporting the concentration of ionized ammonia. This does not represent any change in analytical methods.

### 2.3.1 Water temperature

The mid-winter temperature profiles (e.g., Figures B16–B20, pages 118–122) and the multi-year temperature profiles (Figures B51–B55, pages 154–158) show that the water column mixes during the fall, winter, and early spring. During this time, water temperatures, dissolved oxygen concentrations, pH levels, and conductivities are fairly uniform from the surface to the bottom of the lake, even at Site 4, which is over 300 ft (100 m) deep.

The summer temperature profiles (e.g., Figures B46–B50, pages 148–152) show how the lake stratifies into a warm surface layer (*epilimnion*), and cool bottom layer (*hypolimnion*). The transition zone between the epilimnion and hypolimnion (the *metalimnion*), is a region of rapidly changing water temperature. When stratified, the profiles show distinct differences between surface and bottom temperatures.

Stratification develops gradually, and once stable, persists until fall or winter, depending on location in the lake. Seasonal weather differences alter the timing of lake stratification; if the spring is cool, cloudy, and windy, the lake may stratify later than when it has been hot and sunny.

In Lake Whatcom, all sites except the Intake are usually stratified by late spring or early summer. (The Intake is too shallow to develop a stable stratification.) Stratification may begin as early as April, but is often not stable until May or June. The stability of stratification is determined in part by the temperature differences in the water column, but also by water circulation and local weather patterns. Once the water column temperature differs by at least 5° C ( $\Delta T \geq 5^\circ\text{C}$ ), it is unlikely that the lake will destratify.<sup>8</sup>

The lake cools as the weather becomes colder and days shorten. As the lake cools, the surface and bottom water temperatures become more similar, and eventually the lake will destratify and the water column will mix from the surface to the bottom. Although destratification is relatively abrupt, the process is not instantaneous. In addition, when the lake begins to destratify, water temperatures may be uniform from the surface to the bottom, but the rate of water circulation may not be sufficient to replenish hypolimnetic oxygen concentrations (see November 2006 temperature and oxygen profiles from Sites 1–2: Figures B6 and B7 in Matthews, et al., 2008). Basins 1 and 2 (Sites 1–2) usually destratify by the end

---

<sup>8</sup>The  $\Delta T$  is the difference between the epilimnion and hypolimnion temperatures.

of October but basin 3 (Sites 3–4) is often still stratified in November or early December. Complete destratification of basin 3 usually occurs in December or early January, so by February the temperatures are relatively uniform throughout the water column at all sites.

During the current sampling period, Site 1 was destratified by November 1, 2011 but Site 2 was still slightly stratified (Figures B6–B7, pages 108–109). The oxygen concentrations were still very low near the bottom at both sites, indicating that although the water temperatures were nearly uniform, the water column was not yet completely mixed. Sites 3–4 were still stratified on November 2, 2011, and very weakly stratified December 6, 2011 (Figures B14–B15, pages 116–117).

Historic data reveal that water temperatures in basin 3 are generally cooler than in basins 1 and 2, but the two shallow basins experience more extreme temperature variations. The lowest and highest temperatures measured in the lake since 1988 were at Site 1 (4.2° C on February 1, 1988 and February 26, 1989; 24.1° C on August 4, 2009). The large water volume in basin 3 moderates temperature fluctuations, so water temperatures in basin 3 change slower in response to weather conditions compared to the shallow basins.

The 2012 surface water temperatures were close to the historic median values during most months, but were slightly cooler than usual at Sites 1–2 in June and at Sites 3–4 in July (Figure 1, page 25). The lake was unstratified in April and unstratified or very weakly stratified in May (Figures B21–B30, pages 128–127). Stable stratification was not present until June (Figures B31–B35, pages 133–137).

### **2.3.2 Dissolved oxygen**

Low oxygen conditions are associated with a number of unappealing water quality problems in lakes, including loss of aquatic habitat; release of phosphorus from the sediments; increased rates of algal production due to release of phosphorus; unpleasant odors during lake destratification; fish kills, particularly during lake destratification; release of metals and organics from the sediments; increased mercury methylation; increased drinking water treatment costs; increased taste and odor problems in drinking water; and increased risks associated with disinfection by-products created during the drinking water treatment process.

As in previous years, Sites 1 and 2 developed severe hypolimnetic oxygen deficits by mid-summer (Figures B41–B42 and B56–B57, pages 143–144 and 159–160). Hypolimnetic oxygen depletion only becomes apparent after stratification, when the lower waters of the basin are isolated from the lake’s surface and biological respiration consumes the oxygen dissolved in the water. Biological respiration usually increases when there is an abundant supply of organic matter (e.g., decomposing algae). In basin 3, which has a very large, well-oxygenated hypolimnion, biological respiration has little influence on hypolimnetic oxygen concentrations (Figures B50 and B60, pages 152 and 163). In contrast, there is rapid depletion of the hypolimnetic oxygen concentrations at Sites 1–2 (Figures B46–B47, and B56–B57, pages 148–149 and 159–160). These two sites are in shallow basins that have small hypolimnions compared to their photic zones, so decomposition of algae and other organic matter causes a measurable drop in hypolimnetic oxygen over the summer.<sup>9</sup>

The levels of hypolimnetic oxygen have declined over time at Site 1, causing the lake to be listed by the Department of Ecology as an “impaired” waterbody (Pelletier, 1998).<sup>10</sup> The increasing rate of oxygen loss is most apparent during July and August, after the lake develops a stable thermal stratification but before oxygen levels drops near zero. To illustrate this trend we fitted the July and August data using an exponential function (see discussion by Matthews, et al., 2004). As indicated in Figures 2–5 (pages 26–29), there were significant negative correlations between dissolved oxygen and time for all hypolimnetic samples collected during July and August.<sup>11</sup> Despite slightly cooler temperatures in June, the rate of hypolimnetic oxygen depletion was very rapid. By August 8, 2012, the oxygen concentration was <1 mg/L from 12 meters to the bottom.

A region of supersaturated oxygen was evident in the metalimnion at Site 1 in July and August (Figures B36 and B41, pages 138 and 144). This was caused by the accumulation of phytoplankton along the density gradient between the epilimnion and hypolimnion where light and nutrients are sufficient to support very high lev-

---

<sup>9</sup>The photic zone is the portion of the lake with enough light to support algal photosynthesis. In Lake Whatcom, peak chlorophyll levels are usually at 5–10 meters, so photic zone volumes will be defined as the percent volume  $\leq 10$  meters. Using this definition, the photic zones for basins 1, 2, and 3 occupy 75%, 70%, and 17%, respectively (Mitchell, et al., 2010).

<sup>10</sup><http://www.ecy.wa.gov/programs/wq/303d>.

<sup>11</sup>Correlation analyses examine the relationships between two variables. The test statistic ranges from  $-1$  to  $+1$ ; the closer to  $\pm 1$ , the stronger the correlation. The significance is measured using the p-value; significant correlations have p-values  $< 0.05$ .

els of photosynthesis. Chlorophyll concentrations within the metalimnetic oxygen peak may be 4–5 times higher than those measured near the surface of the lake (Matthews and DeLuna, 2008).

Site 3 developed an oxygen sag near the bottom during late summer and fall (Figures B4–B14, pages 106–116). Sites 3 and 4 developed small oxygen sags near the thermocline (e.g., Figures B4 and B5, pages 106 and 107), which are caused by respiration of heterotrophic bacteria that accumulate along the density gradient between the epilimnion and hypolimnion (Matthews and DeLuna, 2008).

### **2.3.3 Conductivity and pH**

The pH and conductivity data followed trends that were typical for Lake Whatcom (Figures B61–B70, pages 164–173). Surface pH values increased during the summer due to photosynthetic activity. Hypolimnetic pH values decreased and conductivities increased due to decomposition and the release of dissolved compounds from the sediments.

There was a significant long-term trend in the conductivity data. This trend has been attributed to using increasingly sensitive equipment during the past two decades and does not indicate any actual change in the conductivity in the lake (Matthews, et al., 2004). As mentioned in the field sampling description (page 2), the conductivity probe on the Hydrolab Surveyor 4 field meter failed repeatedly, so the conductivity samples collected in October 2011 were measured in the laboratory from water samples collected at 5 meter intervals (Figures B1–B5, pages 103–107). Beginning in November, the new YSI field meter was used to measure conductivity profiles at each site (Figures B6–B50, pages 108–152).

### **2.3.4 Alkalinity and turbidity**

Because Lake Whatcom is a soft water lake, the alkalinity values were fairly low at most sites and depths (Figures B71–B75, pages 175–179). During the summer the alkalinity values at the bottom of Sites 1–2, and occasionally Site 3, increased due to decomposition and the release of dissolved compounds in the lower waters.

Turbidity values in the lake were usually low (1–3 NTU) except during late summer in samples from the bottom of the lake. The high turbidity levels during this



time are an indication of increasing turbulence in the lower hypolimnion as the lake begins to destratify. The highest turbidity peaks were measured at Sites 1–2 (Figures B76–B80, pages 180–184).

Suspended sediments from storm events can also cause elevated turbidity levels in the lake. Major storm events usually occur during winter or early spring when the lake is destratified, so the turbidity levels will be high throughout the water column. Storm-related turbidity peaks are easier to see in samples from the Intake and basin 3 because there are fewer distracting late summer hypolimnetic turbidity peaks (see February 2009 storm-related turbidity peaks in Figures B78 and B79–B80).

Figures B81–B105 (pages 185–209) show the nitrogen and phosphorus data for Lake Whatcom. Nitrogen and phosphorus are important nutrients that influence the amount and type of microbiota (e.g., algae) that grow in the lake. We measured inorganic forms of nitrogen and phosphorus (nitrite, nitrate, ammonium, and soluble phosphate) as well as total nitrogen and total phosphorus, which includes inorganic and organic compounds.<sup>12</sup>

### 2.3.5 Nitrogen and phosphorus

**Nitrogen:** Most algae require inorganic nitrogen in the form of nitrate or ammonium for growth, but some types of algae can use organic nitrogen or even dissolved nitrogen gas.<sup>13</sup> Nitrate depletion was evident at all sites in the photosynthetic zone during the summer (Figures B86–B90, pages 190–194), particularly at Site 1, where the epilimnetic nitrate concentrations often drop below 20  $\mu\text{g-N/L}$  by the end of the summer. Epilimnetic nitrogen depletion is an indirect measure of phytoplankton productivity, and because algal densities have been increasing throughout the lake, epilimnetic dissolved inorganic nitrogen concentrations (DIN)<sup>14</sup> have been declining over time (Figure 6, page 30). Low epilimnetic DIN concentrations favor the growth of Cyanobacteria because many types of Cyanobacteria can use dissolved  $\text{N}_2$  gas as a nitrogen source.

---

<sup>12</sup>Organic nitrogen and phosphorus comes from living or decomposing plants and animals, and may include bacteria, algae, leaf fragments, and other organic particles.

<sup>13</sup>Only Cyanobacteria and a few uncommon species of diatoms can use nitrogen gas.

<sup>14</sup>Dissolved inorganic nitrogen includes ammonium, nitrate, and nitrite. Under most conditions, epilimnetic concentrations of ammonium and nitrite are very low, so epilimnetic DIN is nearly equivalent to nitrate.

Hypolimnetic nitrate concentrations dropped below 20  $\mu\text{g-N/L}$  at Sites 1 and 2. In anaerobic environments, bacteria reduce nitrate ( $\text{NO}_3^-$ ) to nitrite ( $\text{NO}_2^-$ ) and nitrogen gas ( $\text{N}_2$ ). The historic data indicate that nitrate reduction has been common in the hypolimnion at Site 1, but was not common at Site 2 until the summer of 1999. At Site 2 the hypolimnetic nitrate concentrations dropped below 20  $\mu\text{g-N/L}$  from 1999–2006 and 2008–2012, but not in 2007. Matthews, et al. (2008) hypothesized that the higher levels in 2007 were the result of late stratification, which shortened the period of anoxia in the hypolimnion and resulted in less nitrate reduction. The onset of stratification is only one factor involved in hypolimnetic nitrate depletion; the duration of stratification is also important. In 2007, not only did the lake stratify late, Site 2 was nearly destratified by early October and completely mixed by November. The entire period of anoxia was short compared to most years.

Ammonium, along with hydrogen sulfide, is often an indicator of hypolimnetic anoxia.<sup>15</sup> Ammonium is readily taken up by plants as a growth nutrient. In oxygenated environments, ammonium is rarely present in high concentrations because it is rapidly converted to nitrite and nitrate through biological and chemical processes. In low oxygen environments, ammonium accumulates until the lake destratifies. High levels of ammonium (and hydrogen sulfide - see below) are often detected in the hypolimnion at Sites 1 and 2 just before destratification (Table 7, page 22; Figures B81 & B82, pages 185 & 186). Elevated hypolimnetic ammonium concentrations have been common at both sites throughout the monitoring period, but beginning in 1999 the concentrations increased noticeably at Site 2 (Figure B82, page 186).

The hypolimnetic ammonium concentrations in October 2011 were relatively low compared to previous years, which might be related to the cooler water temperatures during the summer of 2011 (Matthews, et al., 2012). As discussed above, Site 2 was still weakly stratified when sampled in November 2011 ( $\Delta T = 2.6^\circ\text{C}$ ), and had an ammonium concentration of 456  $\mu\text{g-N/L}$  at 20 meters. Site 1 was not stratified in November 2011 ( $\Delta T = 0.8^\circ\text{C}$ ), and the ammonium concentrations were low and nearly uniform throughout the water column (21–33  $\mu\text{g-N/L}$ ). The hypolimnetic ammonium concentrations in October 2012 were typical for the lake (275 and 267  $\mu\text{g-N/L}$  at Sites 1 and 2, respectively; Table 7).

---

<sup>15</sup>Ammonium is produced during decomposition of organic matter; hydrogen sulfide is produced by bacteria that use sulfate ( $\text{SO}_4^-$ ) instead of oxygen, creating sulfide ( $\text{S}^{2-}$ ) that reacts with hydrogen ions to form hydrogen sulfide ( $\text{H}_2\text{S}$ ).

Sites 3 and 4 often have slightly elevated ammonium concentrations at 20 m (metolimnion) or near the bottom at 80–90 m (Figures B84–B85, pages 188–189). This is caused by bacterial decomposition of organic matter, but the concentrations never approach the levels found in the hypolimnion at Sites 1–2.

**Site 2 hypolimnetic ammonium and hydrogen sulfide:** The hypolimnion at Site 2 usually has higher concentrations of ammonium and hydrogen sulfide than Site 1 (Table 7, page 22). Although the oxygen concentrations drop to near zero at both sites, basin 2 is slightly shallower than basin 1 (Mitchell, et al., 2010), so a sample from 20 meters is slightly closer to the bottom at Site 2 than Site 1. As a result, the 20 m samples from Site 2 typically contain more of the soluble compounds leaching from the sediments (e.g., ammonium and hydrogen sulfide).

Hydrogen sulfide concentrations are measured in October, which is the latest month that is *consistently* stratified at Sites 1–2. When the lake stratifies late or is unusually cool, the October ammonium and hydrogen sulfide levels will not be as high as in warmer years. The 2012 hydrogen sulfide concentrations were reported as being below the analytical detection limit. This is unlikely, given historic H<sub>2</sub>S concentrations and the strong “rotten egg” smell in both the samples, which indicates hydrogen sulfide. The presence of a rotten egg smell, however, is not a clear indication that the H<sub>2</sub>S levels were above detection. Humans can detect H<sub>2</sub>S at concentrations well below the Edge Analytical H<sub>2</sub>S detection limit of 0.100 mg/L. We have contacted Edge Analytical to request confirmation of the results, and have entered the data as “na” in Table 7 while the issue is being resolved.

**Phosphorus:** Although the Lake Whatcom microbiota require nitrogen, phosphorus is usually what limits microbial growth (Bittner, 1993; Liang, 1994; Matthews, et al., 2002a; McDonald, 1994). The total phosphorus concentration in the water column is a complex mixture of soluble and insoluble phosphorus compounds, only some of which can be used by algae to sustain growth. Soluble forms of phosphorus (e.g., orthophosphate) are easily taken up by algae and other microbiota, and, as a result, are rarely found in high concentrations in the water column. Insoluble phosphorus can be present in the water column bound to the surface of tiny particles or as suspended organic matter (e.g., live or dead algae). Because competition for phosphorus is so intense, microbiota have developed many mechanisms for obtaining phosphorus from the surface of particles or from decomposing organic matter. Liang (1994) and Groce (2011) found that

~50% of the total persulfate phosphorus in soils in the Lake Whatcom watershed was “bioavailable” and could be extracted by algae.

When hypolimnetic oxygen concentrations are low, sediment-bound phosphorus becomes soluble and leaches into the overlying water. Prior to destratification, hypolimnetic phosphorus may be taken up by microbiota in the hypolimnion or metalimnion (see Section 2.3.2 and Matthews and DeLuna, 2008). When the lake mixes in the fall, the hypolimnetic phosphorus will be mixed throughout the water column. As oxygen concentrations increase during mixing, any soluble phosphorus that has not been taken up by biota will usually be converted back into insoluble phosphorus. Because phosphorus moves back and forth between soluble and insoluble forms and between organic and inorganic compounds, it can be difficult to interpret total phosphorus trends. For example, when algal densities increase, their growth usually results in the reduction of soluble and bioavailable fractions of phosphorus in the epilimnion, similar to the epilimnetic DIN reduction that was described for nitrogen. But, since this uptake simply moves the phosphorus into the “live-algae” fraction of organic phosphorus, total phosphorus concentrations may actually increase in the epilimnion.

In Lake Whatcom, total phosphorus and soluble phosphate concentrations were usually low except in the hypolimnion at Sites 1 and 2 just prior to destratification (Figures B96–B100, pages 200–204 and B101–B105, pages 205–209). Epilimnetic total phosphorus concentrations are usually lower than late-summer hypolimnetic peaks. Prior to 2000, the median epilimnetic phosphorus concentrations were  $<5 \mu\text{g-P/L}$  at Sites 2–4 and approximately  $5\text{--}8 \mu\text{g-P/L}$  at Site 1 (Figure 7, page 31). The epilimnetic phosphorus levels have increased significantly at all sites (Figure 7, page 31); however, the pattern is quite erratic, reflecting the complicated nature of phosphorus movement in the water column. It is important to note that low water column phosphorus concentrations do not always predict low algal densities, and may instead indicate rapid and efficient cycling of phosphorus among the lake biota.

### **2.3.6 Chlorophyll, plankton, and Secchi depth**

Site 1 continued to have the highest chlorophyll concentrations of all the sites (Figures B106–B110, pages 210–214). Peak chlorophyll concentrations were usually collected at 0–15 m, while samples from 20 m had relatively low chlorophyll concentrations because light levels are not optimal for algal growth at this depth.

The Lake Whatcom plankton counts were usually dominated by Chrysophyta, consisting primarily *Dinobryon*, *Mallomonas*, and diatoms (Figures B121–B130, pages 225–234). Substantial blooms of bluegreen bacteria (Cyanobacteria) and green algae (Chlorophyta) were also measured at all sites during summer and late fall. Previous analyses of algal biomass in Lake Whatcom indicated that although Chrysophyta dominate the numerical plankton counts, Cyanobacteria and Chlorophyta often dominate the plankton biomass, particularly in late summer and early fall (Ashurst, 2003; Matthews, et al., 2002b). In addition, most of the Cyanobacteria in these samples are counted by colony rather than as individual cells because of the tiny cell size. When the Cyanobacteria density is estimated using settled algae counts (Matthews, et al. 2012), the plankton counts are dominated by tiny Cyanobacteria.

Secchi depths (Figures B111–B115, pages 215–219) showed no clear seasonal pattern because transparency in Lake Whatcom is affected by particulates from storm events and the Nooksack River diversion as well as algal blooms.

**Indications of eutrophication:** Eutrophication is the term used to describe a lake that is becoming more biologically productive. It can apply to an unproductive lake that is becoming slightly more eutrophic, or a productive lake that is becoming extremely eutrophic (see Wetzel, 2001, for more about eutrophication and Matthews, et al., 2005, for a description of the chemical and biological indicators of eutrophication in Lake Whatcom).

The median near-surface summer chlorophyll concentrations were slightly lower in 2012 compared to 2011 (Figure 8, page 32). The chlorophyll concentrations at all sites have increased significantly since 1994, with Site 1 showing the least amount of change and Sites 3–4 showing the greatest change. Although the annual chlorophyll concentrations are quite variable, they seem to have stabilized since 2004, ranging from 3.8–6.7  $\mu\text{g/L}$  at Site 1 and 2.9–4.6  $\mu\text{g/L}$  at Sites 2–4.

Chlorophyll is a direct measure of algal biomass and is best used to evaluate trophic changes in the lake (e.g., is the lake becoming more biologically productive?). We used algal counts rather than chlorophyll to look for trends within the same type of algae (e.g., are the numbers of Cyanobacteria increasing?). The actual relationship between chlorophyll concentration and the algae cell count is complex. The amount of chlorophyll in an algal cell is influenced by the physiological age and condition of the cell, light intensity, nutrient availability, and

many other factors. In addition, while most types of algae are counted by individual cells, a few types must be counted by colonies because the cells are too difficult to see. Even if the amount of chlorophyll was constant in each cell, it would take many tiny cells to equal the chlorophyll biomass in one large colony.

Except for the dinoflagellates<sup>16</sup> the algae counts have also increased significantly since 1994 (Figure 9, page 33). Similarly, there has been a steady increase in the numbers of Cyanobacteria at all sites (Figure 10, page 34). As with the chlorophyll concentrations, the algae and Cyanobacteria counts appear to have stabilized around 2004. The algae count variability looks to be much smaller in Figures 9–10, but that is because the cell counts are plotted using a log<sub>10</sub> scale.

### 2.3.7 Coliform bacteria

The current surface water standards are based on “designated use” categories, which for Lake Whatcom is “Extraordinary Primary Contact Recreation.” The standard for bacteria is described in Chapter 173–201A–200 of the Washington Administrative Code, Water Quality Standards for Surface Waters of the State of Washington:

*Fecal coliform organism levels must not exceed a geometric mean value of 50 colonies/100 mL, with not more than 10 percent of all samples (or any single sample when less than ten sample points exist) obtained for calculating the geometric mean value exceeding 100 colonies/100 mL.*

All of the mid-basin (Sites 1–4) and Intake values for fecal coliforms were less than 10 cfu<sup>17</sup>/100 mL (Figures B116–B120, pages 220–224) and passed the freshwater *Extraordinary Primary Contact Recreation* bacteria standard.

Coliform samples collected offshore from the Bloedel-Donovan swimming area had slightly higher counts than at Site 1 (mid-basin). None of the Bloedel-Donovan counts exceeded 100 cfu/100 mL and the geometric mean was 5 cfu/100 mL, so this site passed both parts of the freshwater *Extraordinary Primary Contact Recreation* bacteria standard.

---

<sup>16</sup>Dinoflagellates are small single-cell algae that are common in Lake Whatcom, but rarely have high densities in the plankton counts.

<sup>17</sup>Colony forming unit/100 mL; cfu/100 mL is sometimes labeled “colonies/100 mL.”

### **2.3.8 Metals**

The metals data for Lake Whatcom are included in Table 8 (page 23). This table includes only the metals listed in our monitoring contract (arsenic, cadmium, chromium, copper, iron, mercury, nickel, lead, and zinc); electronic data files available from IWS contain concentrations for 24 additional metals that are included as part of the analytical procedure used by AmTest.

AmTest has upgraded their equipment, changed analytical procedures, and recalculated detection limits several times since we began collecting metal data from the lake. Because many of the Lake Whatcom metals concentrations are extremely low, changes in equipment or methods can cause the concentration to move from detectable to non-detectable, or vice versa. This type of change does not indicate an actual change in the metals concentration in the lake. Table 1 (page 16) shows the historic and current AmTest detection limits for each metal.

The metals concentrations were within normal concentration ranges for the lake. Iron and zinc concentrations were usually in the detectable range. The highest iron concentration was measured in August at the bottom of Site 1. The elevated iron concentration was the result of sediment-bound iron converting to soluble forms under anaerobic conditions and leaching into the overlying water. Cadmium, copper, and mercury were detected in many of the samples, but at levels close to detection limits, which is typical for Lake Whatcom. Lead was often detected, but the current analytical method has a very low detection limit (0.00005 mg/L). All of the lead concentrations were lower than the historic detection level (<0.001 mg/L, Table 1).

### **2.3.9 Total organic carbon and disinfection by-products**

Total organic carbon concentrations, along with plankton and chlorophyll data, are used to help assess the likelihood of developing potentially harmful disinfection by-products through the reaction of chlorine with organic compounds during the drinking water treatment process. Algae excrete dissolved organic carbon into water, which, along with other decaying organic material, can react with chlorine to form disinfection by-products, predominately chloroform and other trihalomethanes (THMs).

The 2011/2012 total organic carbon concentrations were low at all sites (1.4–2.3 mg/L; Table 9, page 24). The long-term data suggest that total organic carbon concentrations have become more variable. The minimum concentrations measured each year have remained low, usually <1–2 mg/L, but the maximum concentrations have increased (Figure 11, page 35). The data are too variable to determine a specific cause for this pattern.

When algal densities or total organic carbon concentrations increase, we expect to see an increase in THMs. To minimize risk, the Environmental Protection Agency limits the levels of disinfection by-products allowed in treated drinking water through the Safe Drinking Water Act's Disinfection Byproduct Rule. This Rule was adopted in 1979 and has undergone two major revisions (Phase I in 1998; Phase II in 2005). The sampling requirement doubled under Phase II, and beginning with the fourth quarter of 2012 the data will be summarized differently.<sup>18</sup> Figure 12 (page 36) includes data through the end of September 2012 (third quarter). The revised methods will be incorporated into the figures and discussion in future monitoring reports.

The THMs have been increasing in Bellingham's treated drinking water, particularly during the late summer/fall (third quarter; Figure 12, page 36). Haloacetic acids (another disinfection by-product) are not as closely linked to algal concentrations and chlorine dose (Sung, et al., 2000). The Jan-Dec HAAs results were marginally correlated with time (due to the large sample size), but the the third quarter data were not significantly correlated with time. The total THMS and HAAS remained below the recommended maximum contaminant levels of 0.080 mg/L and 0.060 mg/L, respectively, described in Chapter 246–290–310 of Washington Administrative Code, Water Quality Standards for Public Water Supplies of the State of Washington.

---

<sup>18</sup>P. Wendling, pers. comm., City of Bellingham Public Works Dept., December 5, 2012



Abbrev.	Parameter	Method	Historic DL <sup>†</sup>	2011/2012 MDL <sup>†</sup>	Sensitivity or Confidence limit
<b>IWS field measurements:</b>					
cond	Conductivity	Hydrolab (1997) or YSI (2010)	–	–	± 2 µS/cm
do	Dissolved oxygen	Hydrolab (1997) or YSI (2010)	–	–	± 0.1 mg/L
ph	pH	Hydrolab (1997) or YSI (2010)	–	–	± 0.1 pH unit
temp	Temperature	Hydrolab (1997) or YSI (2010)	–	–	± 0.1° C
disch	Discharge	Rantz et al. (1982); SOP-IWS-6	–	–	–
secchi	Secchi depth	Lind (1985)	–	–	± 0.1 m
<b>IWS laboratory analyses:</b>					
alk	Alkalinity	APHA (2012) #2320; SOP-IWS-8	–	–	± 0.4 mg/L
cond	Conductivity	APHA (2012) #2510; SOP-IWS-8	–	–	± 1.4 µS/cm
do	Dissolved oxygen	APHA (2012) #4500-O.C.; SOP-IWS-8	–	–	± 0.1 mg/L
ph	pH-lab	APHA (2012) #4500-H <sup>+</sup> ; SOP-IWS-8	–	–	± 0.03 pH unit
tss	T. suspended solids	APHA (2012) #2540 D; SOP-IWS-13	2 mg/L	0.9 mg/L	± 1.4 mg/L
turb	Turbidity	APHA (2012) #2130; SOP-IWS-8	–	–	± 0.2 NTU
nh4	Ammonium (auto)	APHA (2012) #4500-NH <sub>3</sub> H; SOP-IWS-19	10 µg-N/L	9.9 µg-N/L	± 7.0 µg-N/L
no3	Nitrite/nitrate (auto)	APHA (2012) #4500-NO <sub>3</sub> I; SOP-IWS-19	20 µg-N/L	5.5 µg-N/L	± 3.4 µg-N/L
tn	T. nitrogen (auto)	APHA (2012) #4500-NO <sub>3</sub> I & PJ; SOP-IWS-19	100 µg-N/L	21.8 µg-N/L	± 34.3 µg-N/L
srp	Sol. phosphate (auto)	APHA (2012) #4500-P G; SOP-IWS-19	5 µg-P/L	1.1 µg-P/L	± 1.7 µg-P/L
tp	T. phosphorus (auto)	APHA (2012) #4500-P G & J; SOP-IWS-19	5 µg-P/L	4.8 µg-P/L	± 3.8 µg-P/L
<b>IWS plankton analyses:</b>					
chl	Chlorophyll	APHA (2012) #10200 H; SOP-LW-16	–	–	± 0.1 µg/L
chlo	Chlorophyta	Lind (1985), Schindler trap	–	–	–
cyan	Cyanobacteria	Lind (1985), Schindler trap	–	–	–
chry	Chrysophyta	Lind (1985), Schindler trap	–	–	–
pyrr	Pyrrophyta	Lind (1985), Schindler trap	–	–	–
<b>City coliform analyses:</b>					
fc	Fecal coliform	APHA (2005) #9222 D	–	1 cfu/100 mL	–
<b>Edge Analytical analyses:</b>					
H <sub>2</sub> S	H <sub>2</sub> S	APHA (2012) 4500-S <sup>2</sup>	—	0.100 mg/L	–
<b>AmTest analyses:<sup>‡</sup></b>					
As	T. arsenic	EPA (1994) 200.7	0.01 mg/L	0.02 mg/L	–
Cd	T. cadmium	EPA (1994) 200.7	0.0005 mg/L	0.0015 mg/L	–
Cr	T. chromium	EPA (1994) 200.7	0.001 mg/L	0.0025 mg/L	–
Cu	T. copper	EPA (1994) 200.7	0.001 mg/L	0.005 mg/L	–
Fe	T. iron	EPA (1994) 200.7	0.05 mg/L	0.009 mg/L	–
Pb	T. lead	EPA (1994) 200.8	0.001 mg/L	0.00005 mg/L	–
Hg	T. mercury	EPA (1994) 245.1	0.0001 mg/L	0.00005 mg/L	–
Ni	T. nickel	EPA (1994) 200.7	0.005 mg/L	0.005 mg/L	–
Zn	T. zinc	EPA (1994) 200.7	0.001 mg/L	0.002 mg/L	–
TOC	T. organic carbon	APHA 531 0 B	1.0 mg/L	0.5 mg/L	–

<sup>†</sup> Historic detection limits (DL) are usually higher than current method detection limits (MDL).

<sup>‡</sup> Changes reflect recalculation of detection limits or change in methods.

Table 1: Summary of IWS, AmTest, Edge Analytical, and City of Bellingham analytical methods and parameter abbreviations.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	18.5	19.7	20.6	26.3
Conductivity ( $\mu$ S/cm)	58.0	60.0	61.2	72.9
Dissolved oxygen (mg/L)	0.0	9.8	8.5	12.7
pH	6.0	7.3	7.1	8.4
Temperature (°C)	5.2	9.8	11.0	21.8
Turbidity (NTU)	0.6	1.0	1.4	9.1
Nitrogen - ammonium ( $\mu$ g-N/L)	<10	<10	22.3	180.9
Nitrogen - nitrate/nitrite ( $\mu$ g-N/L)	<20	219.0	194.0	343.0
Nitrogen - total ( $\mu$ g-N/L)	216.3	393.4	364.9	504.1
Phosphorus - soluble ( $\mu$ g-P/L)	<5	<5	<5	8.9
Phosphorus - total ( $\mu$ g-P/L)	<5	10.7	11.8	28.5
Chlorophyll ( $\mu$ g/L)	0.5	3.2	4.3	12.8
Secchi depth (m)	2.8	5.0	4.7	6.4
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	3

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1  $\Rightarrow$  1).

Table 2: Summary of Site 1 water quality data, Oct. 2011 – Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	18.5	19.0	19.2	20.3
Conductivity (μS/cm)	56.0	58.0	58.7	60.0
Dissolved oxygen (mg/L)	8.9	10.7	10.7	12.5
pH	7.1	7.6	7.6	8.1
Temperature (°C)	6.2	13.3	13.3	22.3
Turbidity (NTU)	0.4	0.6	0.6	0.7
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	10.3
Nitrogen - nitrate/nitrite (μg-N/L)	109.8	238.9	228.7	354.5
Nitrogen - total (μg-N/L)	274.9	372.1	374.6	479.1
Phosphorus - soluble (μg-P/L)	<5	<5	<5	<5
Phosphorus - total (μg-P/L)	<5	7.3	7.7	14.5
Chlorophyll (μg/L)	1.1	3.4	3.3	5.5
Secchi depth (m)	4.0	5.5	5.7	8.0
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	2

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 3: Summary of Intake water quality data, Oct. 2011– Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	18.4	18.9	19.4	27.5
Conductivity (μS/cm)	58.0	58.0	59.4	83.0
Dissolved oxygen (mg/L)	0.0	10.3	9.4	12.6
pH	5.9	7.4	7.2	8.0
Temperature (°C)	5.9	10.8	11.7	20.9
Turbidity (NTU)	0.3	0.6	0.9	5.7
Nitrogen - ammonium (μg-N/L)	<10	<10	23.1	456.1
Nitrogen - nitrate/nitrite (μg-N/L)	<20	240.6	242.0	376.4
Nitrogen - total (μg-N/L)	286.8	409.8	410.6	605.8
Phosphorus - soluble (μg-P/L)	<5	<5	<5	<5
Phosphorus - total (μg-P/L)	<5	7.9	8.8	24.8
Chlorophyll (μg/L)	0.7	2.9	2.9	5.7
Secchi depth (m)	3.9	5.5	6.0	9.5
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	3

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 4: Summary of Site 2 water quality data, Oct. 2011 – Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	17.9	18.6	18.7	21.8
Conductivity ( $\mu$ S/cm)	57.0	58.0	58.5	68.0
Dissolved oxygen (mg/L)	0.6 <sup>§</sup>	10.3	10.3	12.8
pH	6.5	7.1	7.2	8.0
Temperature ( $^{\circ}$ C)	6.1	7.0	9.7	21.3
Turbidity (NTU)	0.2	0.4	0.5	3.0
Nitrogen - ammonium ( $\mu$ g-N/L)	<10	<10	<10	13.4
Nitrogen - nitrate/nitrite ( $\mu$ g-N/L)	127.1	355.9	319.3	429.5
Nitrogen - total ( $\mu$ g-N/L)	231.3	468.4	439.6	608.3
Phosphorus - soluble ( $\mu$ g-P/L)	<5	<5	<5	7.0
Phosphorus - total ( $\mu$ g-P/L)	<5	5.5	6.4	25.5
Chlorophyll ( $\mu$ g/L)	0.8	2.9	2.8	5.4
Secchi depth (m)	4.0	5.7	6.0	8.0
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	2

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1  $\Rightarrow$  1).

<sup>§</sup>Atypical value - see discussion in text.

Table 5: Summary of Site 3 water quality data, Oct. 2011 – Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	17.6	18.5	18.6	20.1
Conductivity (μS/cm)	57.0	58.0	58.4	60.5
Dissolved oxygen (mg/L)	8.3	10.3	10.4	12.8
pH	6.4	7.0	7.1	7.8
Temperature (°C)	6.1	6.7	9.4	20.3
Turbidity (NTU)	0.2	0.4	0.4	0.7
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	18.8
Nitrogen - nitrate/nitrite (μg-N/L)	138.5	374.4	338.4	415.3
Nitrogen - total (μg-N/L)	294.8	474.8	452.7	535.1
Phosphorus - soluble (μg-P/L)	<5	<5	<5	5.9
Phosphorus - total (μg-P/L)	<5	6.2	7.0	29.7
Chlorophyll (μg/L)	0.6	3.0	2.6	4.8
Secchi depth (m)	4.2	6.7	6.5	8.3
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	1	1	3

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 6: Summary of Site 4 water quality data, Oct. 2011 – Sept. 2012.

Year	H <sub>2</sub> S (mg/L)		NH <sub>3</sub> (μg-N/L)	
	Site 1	Site 2	Site 1	Site 2
1999 <sup>†</sup>	0.03–0.04	0.40	268.3	424.4
2000 <sup>†</sup>	0.27	0.53	208.8	339.5
2001 <sup>†</sup>	0.42	0.76	168.7	331.9
2002 <sup>†</sup>	0.09	0.32	203.9	383.8
2003 <sup>†</sup>	0.05	0.05	333.8	340.0
2004 <sup>†</sup>	0.25	0.25	300.3	378.3
2005 <sup>‡</sup>	0.13	0.25	257.5	450.4
	0.12	0.42		
2006	0.20	0.42	334.1	354.1
2007	0.40	0.20	324.5	79.3 <sup>§</sup>
2008	0.28	0.38	294.5	404.9
2009	0.15	0.47	271.3	301.2
2010	0.38	0.40	331.3	511.3
2011	0.12	0.16	180.9	209.4
2012	na	na	274.6	267.3

<sup>†</sup>H<sub>2</sub>S samples analyzed by HACH test kit.

<sup>‡</sup>HACH (first value) vs. Edge Analytical (second value)

<sup>§</sup>Atypical result; see discussion by Matthews, et al. (2008)

Table 7: October hypolimnetic ammonium and hydrogen sulfide concentrations at Sites 1 and 2 (20 m). The H<sub>2</sub>S samples have been analyzed by Edge Analytical since 2005. Earlier samples were analyzed using a HACH field test kit.

	Depth (m)	Date	T. As (mg/L)	T. Cd (mg/L)	T. Cr (mg/L)	T. Cu (mg/L)	T. Fe (mg/L)	T. Hg (mg/L)	T. Ni (mg/L)	T. Pb (mg/L)	T. Zn (mg/L)
Site 1	0	Feb 9, 2012	<0.01	0.0008	<0.001	<0.001	0.020	<0.0001	<0.005	<0.00005	0.0060
Site 1	20	Feb 9, 2012	<0.01	0.0010	<0.001	0.002	0.019	<0.0001	<0.005	0.000250	0.0080
Intake	0	Feb 9, 2012	<0.01	0.0011	<0.001	<0.001	0.011	<0.0001	<0.005	<0.00005	0.0060
Intake	10	Feb 9, 2012	<0.01	0.0011	<0.001	<0.001	0.012	0.0001	<0.005	<0.00005	0.0080
Site 2	0	Feb 9, 2012	<0.01	0.0009	<0.001	0.002	0.011	<0.0001	<0.005	<0.00005	0.0120
Site 2	20	Feb 9, 2012	<0.01	0.0010	<0.001	<0.001	0.012	0.0002	<0.005	<0.00005	0.0070
Site 3	0	Feb 7, 2012	<0.01	0.0008	<0.001	<0.001	0.012	<0.0001	<0.005	<0.00005	0.0060
Site 3	80	Feb 7, 2012	<0.01	0.0007	<0.001	0.004	0.016	<0.0001	<0.005	<0.00005	0.0100
Site 4	0	Feb 7, 2012	<0.01	0.0007	<0.001	0.001	0.013	<0.0001	<0.005	<0.00005	0.0070
Site 4	90	Feb 7, 2012	<0.01	<0.0005	<0.001	0.001	0.014	0.0003	<0.005	<0.00005	0.0080
Site 1	0	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.016	0.00012	<0.005	0.000075	0.0037
Site 1	20	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.085	<0.00005	<0.005	<0.00005	0.0028
Intake	0	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.012	0.00020	<0.005	<0.00005	0.0043
Intake	10	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	<0.009	<0.00005	<0.005	<0.00005	0.0033
Site 2	0	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	<0.009	0.00030	<0.005	0.000156	0.0026
Site 2	20	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.024	0.00060	<0.005	<0.00005	0.0058
Site 3	0	Jul 10, 2012	<0.02	<0.0015	<0.0025	<0.005	<0.009	<0.00005	<0.005	0.000241	0.0056
Site 3	80	Jul 10, 2012	<0.02	<0.0015	<0.0025	<0.005	0.015	<0.00005	<0.005	0.000094	0.0262
Site 4	0	Jul 10, 2012	<0.02	<0.0015	<0.0025	<0.005	<0.009	<0.00005	<0.005	0.000092	0.0136
Site 4	90	Jul 10, 2012	<0.02	<0.0015	<0.0025	<0.005	0.011	<0.00005	<0.005	<0.00005	0.0048

Table 8: Lake Whatcom 2011/2012 total metals data. Only the metals specified in the monitoring plan are included in this table; the results for 24 additional metals are available from IWS. AmTest recalculated analytical detection limits between February and July 2012. The February data include original detection limits; July data include new detection limits (see Table 1 for summary of AmTest methods).



Site	Date	Depth	TOC		Date	Depth	TOC	
			(mg/L)				(mg/L)	
Site 1	Feb 9, 2012	0	2.0		Jul 12, 2012	0	2.3	
	Feb 9, 2012	20	2.0		Jul 12, 2012	20	1.8	
Intake	Feb 9, 2012	0	1.9		Jul 12, 2012	0	2.2	
	Feb 9, 2012	10	1.8		Jul 12, 2012	10	2.1	
Site 2	Feb 9, 2012	0	1.6		Jul 12, 2012	0	2.2	
	Feb 9, 2012	20	1.8		Jul 12, 2012	20	1.8	
Site 3	Feb 7, 2012	0	1.9		Jul 10, 2012	0	2.0	
	Feb 7, 2012	80	1.6		Jul 10, 2012	80	1.4	
Site 4	Feb 7, 2012	0	1.8		Jul 10, 2012	0	2.2	
	Feb 7, 2012	90	2.0		Jul 10, 2012	90	1.5	

Table 9: Lake Whatcom 2011/2012 total organic carbon data.




Figure 1: Boxplots showing 1988–2011 surface water temperatures (depth <1 m, all sites and years) with monthly 2012 data (●). Boxplots show medians and upper/lower quartiles; whiskers extend to maximum/minimum values.




Figure 2: Relationship between dissolved oxygen and time at Site 1, 12 m. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 3: Relationship between dissolved oxygen and time at Site 1, 14 m. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 4: Relationship between dissolved oxygen and time at Site 1, 16 m. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 5: Relationship between dissolved oxygen and time at Site 1, 18 m. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 6: Minimum summer, near-surface dissolved inorganic nitrogen concentrations (1994–2012, June–Oct, depths  $\leq 5$  m). Uncensored (raw) data were used to illustrate that minimum values are dropping below analytical detection limits (dashed red line). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; correlations were significant at Sites 1–3.




Figure 7: Median summer, near-surface total phosphorus concentrations (1994–2012, June–Oct, depths  $\leq 5$  m). Uncensored (raw) data were used to illustrate that median values are increasingly above analytical detection limits (dashed red line). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.






Figure 8: Median summer near-surface chlorophyll concentrations (1994–2012, June–October, depths  $\leq 5$  m). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 9:  $\text{Log}_{10}$  plots of median summer, near-surface algae counts (1994-2012, June-October, all sites and depths). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations except Dinoflagellates were significant.




Figure 10: Log<sub>10</sub> plots of median summer, near-surface Cyanobacteria counts (1994–2012, June–October, depths  $\leq 5$  m). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 11: Boxplots of annual total organic carbon concentrations at Sites 1–4. Boxplots show medians and upper/lower quartiles; whiskers show the maximum/minimum values.




Figure 12: Total trihalomethanes (TTHMs) and haloacetic acids (HAAs) concentrations in the Bellingham water distribution system, 1992–2012. Data were provided by the City of Bellingham Public Works Department. Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; correlations for TTHMs (Jan-Dec and Qtr 3) and Jan-Dec HAAs were significant.

### **3 Tributary Monitoring**

The major objective for the tributary monitoring was to provide baseline data for the major tributaries that flow into Lake Whatcom. Whatcom Creek was also sampled to provide baseline data for the lake's outlet. Monthly samples were collected from 2004–2006. The level of effort was reduced from 2007–2009, with samples collected twice each year. Monthly sampling was re-initiated in January 2010 and will continue through December 2012.

#### **3.1 Site Descriptions**

Samples were collected from Anderson, Austin, Blue Canyon, Brannian, Carpenter, Euclid, Mill Wheel, Olsen, Silver Beach, Smith, and Whatcom Creeks and the Park Place drain. The sampling locations for these sites are described in Appendix A.2 and shown on Figure A2, page 98.

#### **3.2 Field Sampling and Analytical Methods**

The tributaries were sampled on October 12, November 9, and December 14, 2011; and on January 10, February 14, March 6, April 5, May 1, June 5, July 17, August 8, and September 12, 2012.

The analytical procedures for sampling the tributaries are summarized in Table 1 (page 16). All water samples (including bacteriological samples) collected in the field were stored on ice and in the dark until they reached the laboratory. Once in the laboratory the handling procedures that were relevant for each analysis were followed (see Table 1). The bacteria samples were analyzed by the City of Bellingham. Total metals analyses (arsenic, cadmium, chromium, copper, iron, mercury, nickel, lead, and zinc) and total organic carbon analyses were done by AmTest.<sup>19</sup> All other analyses were done by WWU.

---

<sup>19</sup>AmTest, 13600 Northeast 126th Place, Suite C, Kirkland, WA, 98034–8720.

### 3.3 Results and Discussion

The monthly tributary data are summarized in Table 10 (page 41), with descriptive statistics for each site listed in Tables 11–22 (pages 42–53). The biannual metals and total organic carbon data are listed in Tables 23–24 (pages 54–55). Historic data from 2004 through the current monitoring period are plotted in Appendix B.4 (Figures B131–B169, pages 236–274). These figures include a dashed (blue) horizontal line that shows the median value for Smith Creek and a solid (red) horizontal line that shows the median value for each creek. Smith Creek was chosen as a reference because it is a major tributary to the lake and has a history of being relatively unpolluted.

Water temperatures and dissolved oxygen concentrations followed predictable seasonal cycles, with most sites having colder temperatures and higher oxygen concentrations during the winter, and warmer temperatures and lower oxygen concentrations during the summer (Figures B131–B136). Whatcom Creek had higher temperatures and lower oxygen concentrations than most other sites, reflecting the influence of Lake Whatcom (Figures B131 and B134). The residential tributaries (Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain) often had slightly elevated temperatures and lower dissolved oxygen concentrations, which is typical for streams in developed watersheds (Figures B133 and B136).

Most of the creeks in the Lake Whatcom watershed had relatively low concentrations of dissolved solids, indicated by conductivities  $\leq 100 \mu\text{S}$  and alkalinities  $\leq 20 \text{ mg/L}$  (Table 10; Figures B137–B145). Sites that did not match this description included the residential tributaries (Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain) and Blue Canyon Creek, which drains an area rich in soluble minerals. Most sites also had low total suspended solids concentrations ( $\leq 5 \text{ mg/L}$ ) and low turbidities ( $\leq 5 \text{ NTU}$ ) except during periods of high precipitation and runoff (Figures B146–B151).

Ammonium concentrations were generally low ( $\leq 10 \mu\text{g-N/L}$ ) except in the residential streams (Table 10; Figures B152–B154). Ammonium does not persist long in oxygenated surface waters. When present in streams, it usually indicates a near-by source such as an upstream wetland with anaerobic soils or a pollution source.

Most of the creeks had lower total nitrogen and nitrate/nitrate concentrations than Smith Creek (Figures B155– B160). The relatively high nitrate and total nitrogen

concentrations in Smith Creek are probably due to the presence of nitrogen-fixing alders (*Alnus rubra*) in the riparian zone upstream from the sampling site. High nitrate and total nitrogen concentrations are not necessarily an indication of water pollution, and low nitrate concentrations actually favor the growth of nuisance Cyanobacteria. The exceptionally low concentrations in Whatcom Creek reflect algal uptake of nitrogen in the lake.

Soluble inorganic phosphate is quickly removed from surface water by biota, so high concentrations of soluble phosphate usually indicate a near-by source such as an anaerobic wetland or a pollution source. In 2011/2012, the median soluble phosphate concentrations were  $\leq 10 \mu\text{g-P/L}$  at all sites except Olsen, Euclid, Mill Wheel, and Silver Beach Creeks and the Park Place drain (Table 10). The historic data indicate that although soluble phosphate concentrations were generally low, nearly all sites have had a few high peaks, and high concentrations were common in residential streams.

Total phosphorus concentrations were higher than soluble phosphate concentrations (Figures B161–B166). The median 2011/2012 concentrations were  $\leq 20 \mu\text{g-P/L}$  at all sites except Carpenter, Euclid, Mill Wheel, and Silver Beach Creeks and the Park Place drain (Table 10). As with soluble phosphate, nearly all sites have had occasional high total phosphorus peaks, and high concentrations were common in samples from residential sites.

High coliform counts are an indicator of residential pollution (Table 10; Figures B167–B169). Although most of the sites had low coliform counts in 2011/2012, five sites exceeded a geometric mean of 50 cfu/100 mL (Carpenter, Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain) and seven sites had more than 10% of the samples with counts  $> 100$  cfu/100 mL (Brannian, Smith, Carpenter, Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain).

The total organic carbon and metals concentrations are included in Tables 23–24. AmTest has upgraded their equipment, changed analytical procedures, and recalculated detection limits several times since we began collecting metal data from the tributaries to Lake Whatcom. Because many of the metals concentrations are extremely low, changes in equipment or methods can cause the concentration to move from detectable to non-detectable, or vice versa. This type of change does not indicate an actual change in the metals concentration in the tributaries. Table 1 (page 16) shows the historic and current AmTest detection limits for each metal.



The metals concentrations were within normal concentration ranges for tributaries to Lake Whatcom. Iron and zinc concentrations were usually in the detectable range. Low concentrations of cadmium, chromium, copper, and mercury were detected in many of the March 2012 samples, but at levels close to detection limits. These elements were all at or below detection in the July samples. Lead was often detected, but the current analytical method has a very low detection limit (0.00005 mg/L). All of the lead concentrations were lower than the historic detection level (<0.001 mg/L, Table 1).

	Typical range	Anderson	Austin	Brannian	Olsen	Smith	Whatcom
Alkalinity	med. $\leq 20$ mg/L	yes	yes	yes	no	yes	yes
Conductivity	med. $\leq 100$ $\mu$ S	yes	yes	yes	yes	yes	yes
pH	6.5–8.0	yes	yes	no	yes	yes	yes
T. susp. solids	med. $\leq 5$ mg/L	yes	yes	yes	yes	yes	yes
Turbidity	med. $\leq 5$ NTU	yes	yes	yes	yes	yes	yes
Ammonium	med. $\leq 10$ $\mu$ g-N/L	yes	yes	yes	yes	yes	yes
Sol. phosphate	med. $\leq 10$ $\mu$ g-P/L	yes	yes	yes	no	yes	yes
T. phosphorus	med. $\leq 20$ $\mu$ g-P/L	yes	yes	yes	yes	yes	yes
F. coliforms	GM $\leq 50$ cfu	yes	yes	yes	yes	yes	yes
	Fewer than 10% exceed 100 cfu	yes	no	yes	no	yes	yes

	Typical range	Blue Canyon	Carpenter	Euclid	Mill Wheel	Park Place	Silver Beach
Alkalinity	med. $\leq 20$ mg/L	no	no	no	no	no	no
Conductivity	med. $\leq 100$ $\mu$ S	no	yes	no	no	no	no
pH	6.5–8.0	no	yes	yes	yes	yes	no
T. susp. solids	med. $\leq 5$ mg/L	yes	yes	yes	no	yes	yes
Turbidity	med. $\leq 5$ NTU	yes	yes	yes	no	yes	yes
Ammonium	med. $\leq 10$ $\mu$ g-N/L	yes	yes	yes	no	no	yes
Sol. phosphate	med. $\leq 10$ $\mu$ g-P/L	yes	yes	no	no	no	no
T. phosphorus	med. $\leq 20$ $\mu$ g-P/L	yes	no	no	no	no	no
F. coliforms	GM $\leq 50$ cfu	yes	no	no	no	no	no
	Fewer than 10% exceed 100 cfu	yes	no	no	no	no	no

Table 10: Comparison of water quality features in Lake Whatcom tributaries.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	12.9	18.2	18.8	26.8
Conductivity ( $\mu$ S/cm)	43.4	58.8	57.6	70.6
Dissolved oxygen (mg/L)	9.2	11.5	11.1	12.6
pH	6.5	7.1	7.0	7.2
Temperature (°C)	4.5	7.7	8.2	13.7
Total suspended solids (mg/L)	<2	2.5	4.8	27.2
Turbidity (NTU)	0.8	2.7	4.5	24.9
Nitrogen - ammonium ( $\mu$ g-N/L)	<10	<10	<10	30
Nitrogen - nitrate/nitrite ( $\mu$ g-N/L)	64.6	443.7	424.9	681.4
Nitrogen - total ( $\mu$ g-N/L)	155.0	653.8	565.3	805.1
Phosphorus - soluble ( $\mu$ g-P/L)	<5	6.9	6.4	11.6
Phosphorus - total ( $\mu$ g-P/L)	8.6	17.4	21.4	51.8
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	10	8	64
(Percent of samples >100 cfu/100 mL = 0)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1  $\Rightarrow$  1).

Table 11: Summary of Anderson Creek water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	10.4	19.9	20.1	35.8
Conductivity (μS/cm)	46.6	72.2	74.6	128.5
Dissolved oxygen (mg/L)	9.8	11.7	11.7	13.5
pH	6.9	7.5	7.4	7.9
Temperature (°C)	2.7	7.7	8.3	15.2
Total suspended solids (mg/L)	<2	<2	2.8	16.5
Turbidity (NTU)	0.6	1.2	2.1	8.2
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	<10
Nitrogen - nitrate/nitrite (μg-N/L)	288.9	661.4	626.7	1038.3
Nitrogen - total (μg-N/L)	362.2	769.8	742.2	1285
Phosphorus - soluble (μg-P/L)	5.6	8.9	9.1	14
Phosphorus - total (μg-P/L)	7.2	17.3	16.3	27.7
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	3	26	25	360
(Percent of samples >100 cfu/100 mL = 25)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 12: Summary of Austin Creek water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	58.1	133.7	125.7	160.2
Conductivity (μS/cm)	153.8	287.1	268.3	319
Dissolved oxygen (mg/L)	10.2	11.4	11.6	13.1
pH	7.7	8.3	8.2	8.4
Temperature (°C)	4.0	8.6	8.7	13.8
Total suspended solids (mg/L)	<2	3.2	3.2	6.5
Turbidity (NTU)	0.5	2.2	2.1	4.4
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	<10
Nitrogen - nitrate/nitrite (μg-N/L)	143.2	419	434	898.4
Nitrogen - total (μg-N/L)	195.0	509.0	527.5	1006.1
Phosphorus - soluble (μg-P/L)	<5	8.6	8.5	14.6
Phosphorus - total (μg-P/L)	5.9	13.9	12.5	17.9
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	2	3	76
(Percent of samples >100 cfu/100 mL = 0)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 13: Summary of Blue Canyon Creek water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	6.1	11	11.9	21.8
Conductivity (μS/cm)	30.8	44.5	44.7	62.7
Dissolved oxygen (mg/L)	7.1	11.1	10.7	12.4
pH	6.4	7	6.9	7.1
Temperature (°C)	4.0	8.0	8.6	14.4
Total suspended solids (mg/L)	<2	<2	2.3	13.7
Turbidity (NTU)	0.4	1.2	1.8	6.1
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	11.5
Nitrogen - nitrate/nitrite (μg-N/L)	136.7	587.6	704.8	1652.5
Nitrogen - total (μg-N/L)	215.4	728.5	818.2	1747.9
Phosphorus - soluble (μg-P/L)	<5	<5	<5	7.1
Phosphorus - total (μg-P/L)	<5	12.3	12.8	23.3
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	8	9	58
(Percent of samples >100 cfu/100 mL = 0)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 14: Summary of Brannian Creek water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	12.1	26.2	28.9	48
Conductivity (μS/cm)	53.4	77.1	82.7	116.7
Dissolved oxygen (mg/L)	9.4	11.5	11.5	13.3
pH	6.8	7.5	7.5	7.9
Temperature (°C)	2.7	7.8	8.6	16.6
Total suspended solids (mg/L)	<2	2.2	4.0	16.8
Turbidity (NTU)	0.4	2.3	2.6	6.5
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	67.2
Nitrogen - nitrate/nitrite (μg-N/L)	165.2	511.5	676.5	1436.3
Nitrogen - total (μg-N/L)	345.7	722.8	889.9	1661.0
Phosphorus - soluble (μg-P/L)	5.3	9.9	12.2	35.1
Phosphorus - total (μg-P/L)	9.8	20.3	23.0	58.3
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	18	140	117	390
(Percent of samples >100 cfu/100 mL = 58)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 15: Summary of Carpenter Creek water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	17.3	45.9	39.9	67.9
Conductivity (μS/cm)	22.3	108.6	100.4	152.1
Dissolved oxygen (mg/L)	8.9	11.4	10.9	12.5
pH	7.0	7.4	7.4	7.6
Temperature (°C)	3.5	8.7	9.0	15.7
Total suspended solids (mg/L)	<2	2.4	2.9	5.5
Turbidity (NTU)	0.5	2.1	2.3	4.7
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	23.2
Nitrogen - nitrate/nitrite (μg-N/L)	179.8	344.2	454.1	898.9
Nitrogen - total (μg-N/L)	358.5	520.2	612.2	1042.0
Phosphorus - soluble (μg-P/L)	8.3	11.1	11.3	16
Phosphorus - total (μg-P/L)	8.3	20.6	21.2	30.6
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	8	72	69	3200
	(Percent of samples >100 cfu/100 mL = 36)			

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 16: Summary of Euclid Creek water quality data, Oct. 2011–Sept. 2012. Euclid Creek had negligible flow on September 12, 2012; no water quality samples were collected under these conditions.



Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	17.8	50.5	46.6	82.9
Conductivity (μS/cm)	66.3	117.7	121.0	184.6
Dissolved oxygen (mg/L)	0.6	10.7	9.0	12.4
pH	6.8	7.3	7.3	7.9
Temperature (°C)	3.4	9.3	11.3	23.6
Total suspended solids (mg/L)	<2	7.8	17.0	93.5
Turbidity (NTU)	2.5	7.5	14.5	55.8
Nitrogen - ammonium (μg-N/L)	<10	28.1	92.9	837.7
Nitrogen - nitrate/nitrite (μg-N/L)	5.6	290.3	517.3	1724.4
Nitrogen - total (μg-N/L)	425.2	1129.2	1426.2	3245.5
Phosphorus - soluble (μg-P/L)	6.5	10.2	13.5	27.2
Phosphorus - total (μg-P/L)	18.1	49.0	128.4	521.8
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	5	345	207	2100
(Percent of samples >100 cfu/100 mL = 58)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 17: Summary of Millwheel Creek water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	9.6	22.1	22.4	46.7
Conductivity (μS/cm)	43.0	67.4	69.3	118.1
Dissolved oxygen (mg/L)	9.7	11.8	11.7	13.4
pH	6.8	7.5	7.4	7.8
Temperature (°C)	2.7	6.9	8.0	16.1
Total suspended solids (mg/L)	<2	4.9	7.8	30.6
Turbidity (NTU)	0.4	3.8	5.6	17.3
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	21.4
Nitrogen - nitrate/nitrite (μg-N/L)	418.1	950.2	916.1	1536.7
Nitrogen - total (μg-N/L)	481.8	1093.8	1038.5	1760.1
Phosphorus - soluble (μg-P/L)	5.9	10.1	10.3	18.5
Phosphorus - total (μg-P/L)	11.3	18.0	20.9	31.8
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	20	14	240
(Percent of samples >100 cfu/100 mL = 25)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 18: Summary of Olsen Creek water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	54.6	82.5	86.6	123.5
Conductivity (μS/cm)	18.4	244	226.6	309.0
Dissolved oxygen (mg/L)	6.2	10.9	10.1	12.4
pH	7.2	7.5	7.5	7.9
Temperature (°C)	5.2	10.1	11.5	22.6
Total suspended solids (mg/L)	<2	<2	<2	4.5
Turbidity (NTU)	0.8	2.5	2.8	5.8
Nitrogen - ammonium (μg-N/L)	<10	23.2	35.3	150.6
Nitrogen - nitrate/nitrite (μg-N/L)	101.7	341.1	493.4	1120.5
Nitrogen - total (μg-N/L)	420.8	573.5	743.8	1353
Phosphorus - soluble (μg-P/L)	14.7	17.5	21.5	38.3
Phosphorus - total (μg-P/L)	21.3	33.7	52.0	153.5
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	7	73	81	490
(Percent of samples >100 cfu/100 mL = 33)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 19: Summary of Park Place drain water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	35.3	89.7	82.2	144
Conductivity (μS/cm)	109.9	210.9	202.3	311.0
Dissolved oxygen (mg/L)	9.1	11.2	11.2	12.9
pH	7.3	8.0	7.9	8.2
Temperature (°C)	3.3	8.8	9.3	18.5
Total suspended solids (mg/L)	<2	2.9	4.4	14.4
Turbidity (NTU)	2.2	4.3	4.5	8.5
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	22.1
Nitrogen - nitrate/nitrite (μg-N/L)	287.5	469.3	580.2	1175.5
Nitrogen - total (μg-N/L)	530.2	733.6	848.9	1499.8
Phosphorus - soluble (μg-P/L)	10.5	15.6	16.6	27.6
Phosphorus - total (μg-P/L)	16.4	35.8	33.9	46.3
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	28	190	172	700
(Percent of samples >100 cfu/100 mL = 75)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 20: Summary of Silver Beach Creek water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	10.0	17.2	18.2	31.6
Conductivity (μS/cm)	43.3	60.2	60.2	86.5
Dissolved oxygen (mg/L)	9.9	12.1	11.9	13.6
pH	7.0	7.4	7.4	7.8
Temperature (°C)	2.8	6.9	8.1	15.6
Total suspended solids (mg/L)	<2	<2	3.1	11.2
Turbidity (NTU)	0.4	1.0	1.9	7.6
Nitrogen - ammonium (μg-N/L)	<10	<10	<10	<10
Nitrogen - nitrate/nitrite (μg-N/L)	476.3	1043.9	1074.1	1908.4
Nitrogen - total (μg-N/L)	550.8	1167.2	1196	2072.9
Phosphorus - soluble (μg-P/L)	<5	7.3	7.8	14
Phosphorus - total (μg-P/L)	9.2	14.3	14.7	21.3
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	4	6	150
(Percent of samples >100 cfu/100 mL = 8)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1 ⇒ 1).

Table 21: Summary of Smith Creek water quality data, Oct. 2011–Sept. 2012.

Variable	Min.	Med.	Mean <sup>†</sup>	Max.
Alkalinity (mg/L CaCO <sub>3</sub> )	19.4	20.8	20.5	21.5
Conductivity ( $\mu$ S/cm)	58.8	60.7	61.0	64.4
Dissolved oxygen (mg/L)	8.6	10.8	10.8	12.6
pH	7.1	7.5	7.5	7.6
Temperature (°C)	5.9	11.5	12.2	22.3
Total suspended solids (mg/L)	<2	<2	<2	3.4
Turbidity (NTU)	0.7	0.9	1.1	2.1
Nitrogen - ammonium ( $\mu$ g-N/L)	<10	<10	<10	31.8
Nitrogen - nitrate/nitrite ( $\mu$ g-N/L)	10.3	159.9	170.0	331.8
Nitrogen - total ( $\mu$ g-N/L)	198.1	317.6	337.9	456.1
Phosphorus - soluble ( $\mu$ g-P/L)	<5	<5	<5	11.8
Phosphorus - total ( $\mu$ g-P/L)	<5	13.3	13.2	28.1
Coliforms - fecal (cfu/100 mL) <sup>‡</sup>	<1	8	8	54
(Percent of samples >100 cfu/100 mL = 0)				

<sup>†</sup>Uncensored arithmetic means except coliforms (geometric mean);

<sup>‡</sup>Censored values replaced with closest integer (i.e., <1  $\Rightarrow$  1).

Table 22: Summary of Whatcom Creek water quality data, Oct. 2011–Sept. 2012.

	Date	T. As (mg/L)	T. Cd (mg/L)	T. Cr (mg/L)	T. Cu (mg/L)	T. Fe (mg/L)	T. Hg (mg/L)	T. Ni (mg/L)	T. Pb (mg/L)	T. Zn (mg/L)
Anderson	Mar 6, 2012	<0.01	0.0013	<0.001	<0.001	0.234	<0.0001	<0.005	0.000160	0.006
Austin (lower)	Mar 6, 2012	<0.01	0.0011	0.001	<0.001	0.528	<0.0001	<0.005	0.000170	0.005
Blue Canyon	Mar 6, 2012	<0.01	0.0017	<0.001	0.001	0.164	<0.0001	<0.005	0.000060	0.006
Brannian	Mar 6, 2012	<0.01	0.0009	<0.001	<0.001	0.104	<0.0001	<0.005	0.000090	0.004
Carpenter	Mar 6, 2012	<0.01	0.0014	<0.001	0.002	0.315	<0.0001	<0.005	0.000150	0.006
Euclid	Mar 6, 2012	<0.01	0.0015	<0.001	<0.001	0.218	<0.0001	<0.005	0.000200	0.007
Millwheel	Mar 6, 2012	<0.01	0.0014	0.002	0.003	1.090	<0.0001	<0.005	0.000710	0.017
Olsen	Mar 6, 2012	<0.01	0.0012	0.001	<0.001	0.364	<0.0001	<0.005	0.000160	0.007
Park Place	Mar 6, 2012	<0.01	0.0021	<0.001	0.003	0.353	0.0003	<0.005	0.000300	0.015
Silver Beach	Mar 6, 2012	<0.01	0.0017	<0.001	0.003	0.476	<0.0001	<0.005	0.000250	0.008
Smith	Mar 6, 2012	<0.01	0.0010	<0.001	<0.001	0.295	<0.0001	<0.005	0.000100	0.006
Whatcom	Mar 6, 2012	<0.01	0.0013	<0.001	0.002	0.039	0.0001	<0.005	0.000070	0.005
Anderson	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.610	<0.00005	<0.005	0.000168	<0.002
Austin (lower)	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.741	<0.00005	<0.005	<0.00005	0.002
Blue Canyon	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.056	<0.00005	<0.005	<0.00005	<0.002
Brannian	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.243	<0.00005	<0.005	0.000064	<0.002
Carpenter	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.124	<0.00005	<0.005	<0.00005	0.004
Euclid	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.216	<0.00005	<0.005	0.000075	<0.002
Millwheel	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.790	<0.00005	<0.005	0.000259	0.002
Olsen	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.099	<0.00005	<0.005	<0.00005	<0.002
Park Place	Jul 12, 2012	<0.02	0.0015	<0.0025	<0.005	0.325	<0.00005	<0.005	<0.00005	0.006
Silver Beach	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.796	<0.00005	<0.005	0.000089	<0.002
Smith	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.023	<0.00005	<0.005	0.000055	<0.002
Whatcom	Jul 12, 2012	<0.02	<0.0015	<0.0025	<0.005	0.071	<0.00005	<0.005	0.000107	<0.002

Table 23: Lake Whatcom tributary data: total metals. Only the metals specified in the monitoring plan are included in this table; the results for 24 additional metals are available from IWS. This parameter is sampled twice each year. AmTest recalculated analytical detection limits between March and July 2012. The March data include the original detection limits; July data include new detection limits (see Table 1 for summary of AmTest methods).

Site	Date	TOC (mg/L)	Date	TOC (mg/L)
Anderson	Mar 6, 2012	2.4	Jul 12, 2012	1.9
Austin (lower)	Mar 6, 2012	2.2	Jul 12, 2012	2.4
Blue Canyon	Mar 6, 2012	3.1	Jul 12, 2012	1.8
Brannian	Mar 6, 2012	2.2	Jul 12, 2012	2.4
Carpenter	Mar 6, 2012	5.2	Jul 12, 2012	3.9
Euclid	Mar 6, 2012	3.9	Jul 12, 2012	3.5
Millwheel	Mar 6, 2012	4.8	Jul 12, 2012	6.9
Olsen	Mar 6, 2012	3.2	Jul 12, 2012	2.8
Park Place	Mar 6, 2012	5.2	Jul 12, 2012	4.5
Silver Beach	Mar 6, 2012	6.1	Jul 12, 2012	5.6
Smith	Mar 6, 2012	3.0	Jul 12, 2012	2.2
Whatcom	Mar 6, 2012	2.1	Jul 12, 2012	2.6

Table 24: Lake Whatcom tributary data: total organic carbon. This parameter is sampled twice each year.



## 4 Lake Whatcom Hydrology

### 4.1 Hydrograph Data

Recording hydrographs are installed in Austin Creek and Smith Creek; the data are plotted in Figures 13–14 (pages 62–63). The location of each hydrograph is described in Appendix A.2. All hydrograph data, including data from previous years, are online at <http://www.wvu.edu/iws>. Detailed field notes and rating curves for each water year are available from the Institute for Watershed Studies. All results are reported as Pacific Standard Time, without Daylight Saving Time adjustment.

### 4.2 Water Budget

A water balance was applied to Lake Whatcom to identify major water inputs and outputs and to examine runoff and storage. The traditional method of estimating a water balance was employed, where  $\text{inputs} - \text{outputs} = \text{change in storage}$  (Table 25, page 59). Inputs into the lake include direct precipitation, runoff (surface runoff + groundwater), and water diverted from the Middle Fork of the Nooksack River. Outputs include evaporation, Whatcom Creek, the Whatcom Falls Fish Hatchery, City of Bellingham, Puget Sound Energy Co-Generation Plant <sup>20</sup>, and the Lake Whatcom Water and Sewer District.<sup>21</sup> The change in storage is estimated from daily lake-level changes. All of these are measured quantities provided by the City of Bellingham except for evaporation, diverted water, and runoff.

Daily direct-precipitation magnitudes on the lake surface were estimated using the precipitation data recorded at the Bloedel Donovan, Geneva gatehouse, North Shore, and Brannian Creek gauges. Due to an equipment malfunction at the North Shore gauge, rainfall data from June 22 to September 30 were replaced with rainfall data from the Geneva gatehouse gauge. Note, however, that only about 9% of the annual rainfall occurred during this time interval. The minimum yearly rainfall (40.0 inches) was recorded at the Bloedel Donovan gauge, the maximum (61.8 inches) was recorded at the Brannian creek gauge. A daily weighted average rainfall average was calculated using a Python script that employed a spatial

---

<sup>20</sup>Located at the Georgia Pacific site

<sup>21</sup>Formerly Water District #10

interpolation technique (inverse distance weighted) in ArcGIS to distribute rainfall from the four gauges over a 10 meter raster of the lake. The average direct-precipitation depth (inches) for a given day was converted to volume in millions of gallons (MG) via a rating curve generated from the lake level-area data (Mitchell et al., 2010). The rating curve accounts for changes in surface area of the lake due to lake level changes. The average annual direct rainfall to the lake for the water year 2011/2012 was 50.3 inches (6778 MG); 69% of which occurred between October 1 and April 1.

Daily diversion volumes were estimated using a hydrograph separation technique based on daily discharge data from the Anderson Creek USGS stream gauge (USGS 12201950), modeled streamflow using the DHSVM, and the outfall valve log-sheet provided by the City of Bellingham. The Distributed Hydrology-Soils-Vegetation Model (DHSVM) is a spatially distributed, physically based numerical model that was calibrated to the Anderson Creek basin (Matthews et al., 2007). The log-sheet documents the dates and times that the diversion was operating and the valve opening percent. These dates and times were located on the hydrograph. The natural streamflow was estimated by the DHSVM and manually removed from the USGS hydrograph. The remaining volume was used to estimate a daily volume discharging to the lake from the diversion. The outfall gate was never open more than 30%, which on average accounted for about 18–20 MG per day during dry periods. As such, if the hydrograph separation technique yielded a value greater than 20 MG during a storm event, it was set to 20 MG. Approximately 2279 MG were diverted into the lake in 2011/2012.

Daily lake evaporation was estimated using a model based on the Penman method (Dingman, 1994). The Penman method is theoretically based model that estimates free-water evaporation using both energy-balance and mass transfer concepts. The method requires daily average incident solar radiation, air temperature, dew point temperature, and wind speed. Hourly data from the North Shore weather station in the watershed were used to estimate daily averages. The daily evaporation depths (inches) predicted by the model were converted to volumes (MG) via a rating curve generated from the lake level-area data developed by Mitchell et al. (2010). The estimated yearly evaporation from the lake is 18.2 inches (2460 MG), 80% of which occurred between April and September.

Daily change in storage was determined by subtracting each day's lake level by the subsequent day's level. This resulted in negative values when the lake level was decreasing and positive values when the lake level was increasing. The minimum

lake level (311.43 ft) was recorded on December 22, 2011; and the maximum lake level (314.65 ft) occurred on July 4, 2012. The change in storage magnitudes are sensitive to the accuracy of the lake level measurements; small lake level changes correspond to large lake volumes. The daily net change in lake level (inches) was converted to a volume (MG) via a rating curve generated from the lake level-volume data developed by Mitchell et al. (2010). The rating curve accounts for changes in volume of the lake due to lake level changes. The median total lake volume in 2011/2012 was 252,758 MG. Figure 15 (page 64) shows daily lake-volume values for the past five years. There was a spike in lake volume when the lake rose from a level of 312.0 feet on January 4, to 315.0 feet on January 9, 2009 due to a 6.3 inch storm event.

Surface runoff and groundwater were combined into a single runoff component that was determined by adding the outputs to the change in storage and subtracting precipitation and diversion volumes. Negative values of runoff estimated from the water budget are likely due to noise in the change in storage estimates or may represent a loss of lake water to deep aquifer systems. The DHSVM was also used to simulate runoff into the lake. Runoff represents 74.6% of the annual input to the lake. About 54% of the total input to the lake occurs as runoff between January 1 and April 30. Predictably, about 50% of the total output of the lake discharges out Whatcom Creek during the same four month interval.

The daily water balance quantities were summed into 7-day totals, which were used to generate Figures 16–19 (pages 65–68). Figure 16 shows 7-day summed totals for inputs, outputs, and change in storage. All the inputs except runoff are shown in Figure 17; all outputs except Whatcom Creek are shown in Figure 18. Due to their much higher magnitude, observed runoff, modeled runoff, and Whatcom Creek data are included on Figure 19.

Yearly water balance totals are listed in Table 25 (page 59) along with data from four previous water years. The total volume of outputs were 13.8% of the median total volume of the lake. Under the assumption that the lake is completely mixed and flow is steady state (inputs = outputs), this would correspond to a 7.2 year residence time.<sup>22</sup> Tables 26 and 27 (pages 60–61) show the 2011/2012 total input and output volumes along with the corresponding monthly percentage of each total.

---

<sup>22</sup>Although the lake is not completely mixed and the flow is not steady state, these assumptions are commonly used to provide a simple estimate of residence time for water in lakes.

	WY2012 (9/30/11–10/1/12)	WY2011 (9/30/10–10/1/11)	WY2010 (9/30/09–10/1/10)	WY2009 (9/30/08–10/1/09)	WY2008 (9/30/07–10/1/08)
<b>Inputs (MG)<sup>†</sup></b>					
Direct Precipitation	6,778 (19.0%)	6,900 (18.0%)	7,350 (23.7%)	5,712 (17.7%)	6,006 (16.7%)
Diversions	2,279 (6.4%)	2,629 (6.9%)	860 (2.8%)	0 (0.0%)	4,902 (13.7%)
Runoff	26,586 (74.6%)	28,709 (75.1%)	22,762 (73.5%)	26,491 (82.3%)	24,989 (69.6%)
Total	35,643 (100%)	38,238 (100%)	30,973 (100%)	32,203 (100%)	35,896 (100%)
<b>Outputs (MG%)</b>					
Whatcom Creek	27,899 (79.9%)	32,351 (81.2%)	22,311 (75.4%)	26,598 (77.5%)	25,793 (76.1%)
Hatchery	807 (2.3%)	851 (2.1%)	875 (3.0%)	856 (2.5%)	931 (2.7%)
Puget Sound Co-Gen	45 (0.1%)	57 (0.1%)	51 (0.2%)	4 (0.01%)	240 (0.7%)
City of Bellingham	3,467 (9.9%)	3,593 (9.0%)	3,522 (11.9%)	3,886 (11.3%)	3,874 (11.4%)
LW Water/Sewer Distr.	225 (0.6%)	226 (0.6%)	239 (0.8%)	250 (0.7%)	237 (0.7%)
Evaporation	2,460 (7.0%)	2,770 (7.0%)	2,592 (8.8%)	2,723 (7.9%)	2,807 (8.3%)
Total	34,903 (%100)	39,847 (100%)	29,589 (100%)	34,317 (100%)	33,883 (100%)
<b>Net change in storage</b>	740	-1,609	1,384	-2,115	2,033
<b>Median lake volume (MG)</b>	252,758	252,637	252,074	252,433	253,003
<b>Outflow percent of volume</b>	13.8%	15.8%	11.7%	13.6	13.4%
<b>Residence time (years)<sup>‡</sup></b>	7.2	6.3	8.5	7.4	7.5

<sup>†</sup>Runoff = surface runoff + groundwater; no diversion inputs in WY2009.

<sup>‡</sup>Based on the assumption that water in the lake is completely mixed and flow is steady state (i. e., inputs = outputs)

Table 25: Annual water balance quantities for the Lake Whatcom watershed, WY2008–WY2012.

Month	Input Percents <sup>†</sup>			Total
	Diversion	Precipitation	Runoff	
Oct	0.00	8.16	-1.36	0.54
Nov	11.31	13.55	4.99	7.02
Dec	3.64	5.52	4.22	4.43
Jan	1.93	12.71	16.84	15.10
Feb	4.43	13.31	18.70	16.76
Mar	7.72	15.12	20.79	18.87
Apr	5.92	11.79	15.95	14.52
May	16.06	5.57	9.55	9.21
Jun	24.54	7.95	6.91	8.23
Jul	23.83	5.94	5.56	6.80
Aug	0.63	0.02	-0.03	0.02
Sep	0.00	0.37	-2.11	-1.51
Input Volume (MG)				
Total	2,279	6,778	26,586	35,643

<sup>†</sup>Runoff = surface runoff + groundwater;

Table 26: Monthly input water balance quantities for the Lake Whatcom watershed, October 2011–September 2012.

Month	Output Percents <sup>†</sup>						Total
	WC	Hatch	PSE	COB	WSD	Evap	
Oct	0.61	5.73	0.19	8.01	8.12	4.63	1.79
Nov	6.51	6.35	1.51	7.32	7.98	2.32	6.29
Dec	4.33	8.10	16.46	7.21	8.53	1.18	4.53
Jan	15.94	8.04	16.35	7.29	9.19	2.41	13.90
Feb	17.69	7.61	11.10	6.70	8.03	2.56	15.23
Mar	16.92	8.89	14.77	7.25	7.33	6.64	14.98
Apr	11.56	9.57	8.16	7.21	7.32	8.23	10.82
May	8.93	8.09	5.35	8.74	8.14	16.33	9.40
Jun	7.77	9.48	2.32	8.20	7.66	13.82	8.27
Jul	6.26	9.83	14.77	9.65	8.78	17.47	7.50
Aug	2.40	9.63	0.00	12.07	9.73	15.52	4.50
Sep	1.09	8.68	9.03	10.36	9.18	8.89	2.80
Output Volume (MG)							
Total	27,899	807	45	3,467	225	2,460	34,903

<sup>†</sup>WC = Whatcom Creek; Hatch = Whatcom Falls Hatchery;  
PSE = Puget Sound Energy Co-Generation Plant;  
COB = City of Bellingham; WSD = Lake Whatcom Water  
Sewer District; Evap = Evaporation

Table 27: Monthly output water balance quantities for the Lake Whatcom watershed, October 2011–September 2012.




Figure 13: Austin Creek hydrograph, October 1, 2011–September 30, 2012. Data were recorded at 15 minute intervals.




Figure 14: Smith Creek hydrograph, October 1, 2011–September 30, 2012. Data were recorded at 15 minute intervals.






Figure 15: Comparison of Lake Whatcom daily lake volumes for WY2008–WY2012. Horizontal line represents median lake volume for the period plotted.




Figure 16: Summary of 7-day inputs, outputs, and changes in Lake Whatcom storage, October 1, 2011–September 30, 2012.




Figure 17: Lake Whatcom watershed direct hydrologic inputs, October 1, 2011–September 30, 2012. Runoff is included on Figure 19 (see Section 4.2 discussion).




Figure 18: Lake Whatcom watershed hydrologic withdrawals, October 1, 2011–September 30, 2012. Whatcom Creek output is included on Figure 19 ( see Section 4.2 discussion).




Figure 19: Summary of 7-day Whatcom Creek flows, water balance runoff estimates, and DHSVM runoff estimates, October 1, 2011–September 30, 2012.

## 5 Storm Water Monitoring

### 5.1 Site Descriptions

The storm water monitoring program was revised in 2009 to focus on collecting baseline data at the Silver Beach Creek outlet and the City of Bellingham's North Shore Drive overlay project. Both sites were monitored in 2009/2010 (see Matthews, et al., 2011). Beginning in 2010, the emphasis focused on collecting additional storm water samples from Silver Beach Creek. For information about other storm water sites that have been monitored by IWS, refer to the annual reports listed in Section 6.2 (page 89).

### 5.2 Field Sampling and Analytical Methods

Flow-paced discrete samples were collected at the USGS gauging site near the mouth of Silver Beach Creek (Figure A3, page 99) using an ISCO sampler provided by the City of Bellingham. A total of eight storm events were sampled between November 2011 and April 2012 (Table 28, page 72). Each storm event was given a unique number (Events 17–24).<sup>23</sup> Six of these storms met the precipitation goal ( $\geq 1$  cm in 24-hr) and included samples from the rising and falling leg of the hydrograph. Two storms (19 and 21) did not meet the precipitation goal, but have been included in this report for general information.

The sampler was calibrated to collect flow proportional samples during each storm event. The samples were analyzed to measure total suspended solids, turbidity, total phosphorus, soluble phosphate, total nitrogen, and nitrate/nitrite following the methods summarized in Table 1 (page 16). Stream elevation (stage height) was recorded at 15 minute intervals during each storm event and when a water sample was collected.<sup>24</sup> Stream flow was estimated from stage height (ft) using the following rating curves.

$$\begin{aligned} \text{Oct - Dec 2010:} \quad \text{Flow (cfs)} &= (2.6402 \times \text{stage height} - 9.1803)^2 \\ \text{Feb 2011 - Apr 2012:} \quad \text{Flow (cfs)} &= (2.7103 \times \text{stage height} - 9.3703)^2 \end{aligned}$$

<sup>23</sup>Events 1–16 were discussed by Matthews, et al. (2011; 2012)

<sup>24</sup>The flow-paced water samples were collected at irregular intervals based on stream flow, so the sampling time rarely coincided with the automatic 15-min stage height measurements.

Sample stage height data (and estimated flow rates) were not collected for a few samples due to instrumentation error. For these samples, the stage height at the time of sampling was estimated using a unweighted average of adjacent 15-min interval stage height data.

### 5.3 Results and Discussion

The amount and intensity of precipitation varied between storm events (Table 28; Figures 20–25, pages 73–78). Four events (17, 18, 20, and 22) had 24-hr maximum precipitation totals of 1.0–2.0 cm during the monitoring event, two events had >2.5 cm (23 and 24), and two events had <1.0 cm (19 and 21). Of the two low-flow events, Event 19 displayed a typical hydrograph and typical water quality patterns in response to the hydrograph. Event 21 was collected during a period of high flow in January 2012. There was no obvious hydrograph peak, and the water quality data collected during this event were atypical for most parameters.

Total suspended solids, turbidity, and total phosphorus increased with stream flow for all events except Event 21 (Figures 20–22). Soluble phosphate and total nitrogen increased with flow during some of the storm events (e.g., Events 18 and 23), but often showed little relationship to the hydrograph (Figures 23–24). Nitrate concentrations were usually diluted by precipitation (Figure 25). Event 21, which occurred during consistently high flow (no hydrograph peak), had nearly constant levels for all of the water quality parameters.

Correlation analysis was used to test the relationship between stream flow, stream elevation (stage height), and water quality (Figures 26–31, pages 79–84). Both stage height and stream flow were included because stream flow is estimated from a rating curve, so it contains uncertainty. Stage height is a direct measurement of the height of the stream when the sample is collected, and produced slightly better correlations with the water quality data. Events 19 and 21 were excluded because they did not meet the precipitation goals and Event 21 did not have a typical hydrograph profile or water quality responses.

All of the water quality parameters were significantly correlated with stream flow and stage height (Figures 26–31); however, the significant positive correlation for nitrate with stream flow and stage height was mostly an artifact of the large sample size. All other parameters (total suspended solids, turbidity, total phosphorus, soluble phosphate, and total nitrogen) had much higher Kendall's  $\tau$  statistics

compared to nitrate. Total suspended solids, turbidity, and total phosphorus were highly correlated with each other (Figure 32). Total phosphorus is often adsorbed to the surface of sediment particles and is transported with sediments in storm runoff.

Part of the scattered “noise” in Figures 26–31 comes from within-storm variation, which can be seen by plotting the storm events separately. For example, Figure 33 shows the correlations between total phosphorus and flow by event. The results varied considerably, with correlation statistics ranging from insignificant (Events 19 and 21) to very highly significant (Event 20  $\tau = 0.95$ ).<sup>25</sup> In theory, the “best” statistical approach would be to evaluate all data separately by storm event. But this is not always feasible, or even desirable, especially if the goal is to develop a simple model of pollutant transport as a function of stream flow.

---

<sup>25</sup>The maximum value for a correlation statistic is  $\pm 1.0$ .



Event	Sampling Period	Event Duration (hr)	Max. 24-hr Precip
17	06:00 Nov 11 to 12:00 Nov 12, 2011	30	0.72 in (1.8 cm)
18	12:00 Nov 16 to 12:00 Nov 18, 2011	48	0.42 in (1.1 cm)
19	14:00 Dec 16 to 09:30 Dec 20, 2011	91	0.20 in (0.51 cm)
20	14:00 Dec 27 to 14:00 Dec 28, 2011	24	0.51 in (1.3 cm)
21	10:00 Jan 25 to 09:45 Jan 26, 2012	23	0.29 in 0.74 cm)
22	20:00 Jan 28 to 08:30 Jan 31, 2012	60	0.64 in (1.6 cm)
23	20:00 Feb 20 to 12:00 Feb 23, 2012	64	1.37 in (3.5 cm)
24	18:00 Apr 19 to 14:00 Apr 20, 2012	20	1.03 in (2.6 cm)

Table 28: Summary of Silver Beach Creek storm events and maximum 24-hr precipitation total at the Bloedel/Donovan precipitation gauge. Precipitation data were provided by the City of Bellingham.




Figure 20: Silver Beach Creek storm water monitoring results for Events 17–24: total suspended solids (●) vs. stream flow (—). Note scale for each event. Results for Events 1–16 were presented in earlier reports (Matthews et al. 2011; 2012).




Figure 21: Silver Beach Creek storm water monitoring results for Events 17–24: turbidity (●) vs. stream flow (—). Note scale for each event. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012).




Figure 22: Silver Beach Creek storm water monitoring results for Events 17–24: total phosphorus (●) vs. stream flow (—). Note scale for each event. Results for Events 1–16 were presented in the earlier reports (Matthews et al., 2011; Matthews et al., 2012).




Figure 23: Silver Beach Creek storm water monitoring results for Events 17–24: soluble phosphate (●) vs. stream flow (—). Note scale for each event. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012).




Figure 24: Silver Beach Creek storm water monitoring results for Events 17–24: total nitrogen (●) vs. stream flow (—). Note scale for each event. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012).




Figure 25: Silver Beach Creek storm water monitoring results for Events 17–24: nitrate/nitrite (●) vs. stream flow (—). Note scale for each event. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012)




Figure 26: Correlation between stream flow or stage height and total suspended solids in Silver Beach Creek (Events 17–18, 20, and 22–24). Events 19 and 21 were excluded because they did not meet precipitation goals. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.






Figure 27: Correlation between stream flow or stage height and turbidity in Silver Beach Creek (Events 17–18, 20, and 22–24). Events 19 and 21 were excluded because they did not meet precipitation goals. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012). Kendall’s  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 28: Correlation between stream flow or stage height and total phosphorus in Silver Beach Creek (Events 17–18, 20, and 22–24). Events 19 and 21 were excluded because they did not meet precipitation goals. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations except Event 19 & 21 were significant.




Figure 29: Correlation between stream flow or stage height and soluble phosphate in Silver Beach Creek (Events 17–18, 20, and 22–24). Events 19 and 21 were excluded because they did not meet precipitation goals. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012). Kendall’s  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 30: Correlation between stream flow or stage height and total nitrogen in Silver Beach Creek (Events 17–18, 20, and 22–24). Events 19 and 21 were excluded because they did not meet precipitation goals. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012). Kendall’s  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 31: Correlation between stream flow or stage height and nitrate in Silver Beach Creek (Events 17–18, 20, and 22–24). Events 19 and 21 were excluded because they did not meet precipitation goals. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 32: Correlation between total suspended solids, turbidity, and total phosphorus in Silver Beach Creek (Events 17–18, 20, and 22–24). Events 19 and 21 were excluded because they did not meet precipitation goals. Results for Events 1–16 were presented in earlier reports (Matthews et al., 2011; Matthews et al., 2012). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.




Figure 33: Correlation between stream flow and total phosphorus by storm event in Silver Beach Creek (Events 17–24). Results for Events 1–16 were presented in earlier reports (Matthews, et al., 2011; Matthews, et al., 2012). Kendall's  $\tau$  correlations were used because the data were not monotonic-linear; all correlations were significant.

## 6 References and Related Reports

### 6.1 References

- APHA. 2012. Standard Methods for the Examination of Water and Wastewater, 22nd Edition. American Public Health Association, American Water Works Association, and Water Environment Federation, Washington, DC.
- Ashurst, S. 2003. Microcosm study of the accumulation of benzo(a)pyrene by Lake Whatcom phytoplankton. M. S. thesis, Huxley College of Environmental Studies, Western Washington University, Bellingham, WA.
- Bittner, C. W. 1993. The response of Lake Whatcom bacterioplankton to nutrient enrichment. M. S. thesis, Huxley College of Environmental Studies, Western Washington University, Bellingham, WA.
- Dingman, S. L. 1994. Physical Hydrology. Macmillan College Publishing Co., New York, NY.
- EPA. 1994. Method 200.7: Determination of Metals and Trace Elements in Water and Wastes by Inductively Coupled Plasma-Atomic Emission Spectrometry, Revision 4.4, U. S. Environmental Protection Agency, Washington DC.
- EPA. 1994. Method 200.8: Determination of Metals and Trace Elements in water and Wastes by Inductively Coupled Plasma-Mass Spectrometry, Revision 5.4, U. S. Environmental Protection Agency, Washington DC.
- Groce, S. 2011. Soils as a Source of Bioavailable Phosphorus in the Lake Whatcom Watershed. M. S. thesis, Huxley College of the Environment, Western Washington University, Bellingham, WA.
- Hydrolab. 1997. Data Sonde 4 Water Quality Multiprobes User Manual, Revision D., August 1997. Hydrolab Corporation, Austin, TX.
- Lind, O. T. 1985. Handbook of Common Methods in Limnology, 2nd Edition. Kendall/Hunt Publishing Co., Dubuque, IA.
- Matthews, R. A. and E. DeLuna, 2008. Metalimnetic oxygen and ammonium maxima in Lake Whatcom, Washington (USA). Northwest Science 82:18–29.



- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. 2012. Lake Whatcom Monitoring Project, 2010/2011 Final Report, February 24, 2012. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. 2011. Lake Whatcom Monitoring Project, 2009/2010 Final Report, March 1, 2011. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. J. Mitchell, and G. B. Matthews. 2008. Lake Whatcom Monitoring Project 2006–2007 Final Report. Final Report prepared for the City of Bellingham Public Works Department, April, 2008, Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. J. Mitchell, and G. B. Matthews. 2007. Lake Whatcom Monitoring Project 2005–2006 Final Report. Final Report prepared for the City of Bellingham Public Works Department, April, 2007, Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. J. Mitchell, and G. B. Matthews. 2005. Lake Whatcom Monitoring Project 2003–2004 Final Report. Final Report prepared for the City of Bellingham Public Works Department, March, 2005, Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. J. Mitchell, and G. B. Matthews. 2004. Lake Whatcom Monitoring Project 2002–2003 Final Report. Final Report prepared for the City of Bellingham Public Works Department, March, 2004, Bellingham, WA.
- Matthews, R., M. Hilles, and G. Pelletier. 2002a. Determining trophic state in Lake Whatcom, Washington (USA), a soft water lake exhibiting seasonal nitrogen limitation. *Hydrobiologia* 468:107–121.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. J. Mitchell, and G. B. Matthews. 2002b. Lake Whatcom Monitoring Project 2000–2001 Final Report. Final Report prepared for the City of Bellingham Public Works Department, March, 2002, Bellingham, WA.
- McDonald, K. R. 1994. Nutrient limitation of phytoplankton in Lake Whatcom. M. S. thesis, Huxley College of Environmental Studies, Western Washington University, Bellingham, WA.

- Mitchell, R., G. Gabrisch, and R. Matthews. 2010. Lake Whatcom Bathymetry and Morphology. Report prepared for the City of Bellingham Public Works Department, December 2, 2010, Bellingham, WA.
- Pelletier, G. 1998. Dissolved oxygen in Lake Whatcom. Trend in the depletion of hypolimnetic oxygen in basin I, 1983–1997. Washington State Department of Ecology Report #98–313, Olympia, WA.
- Rantz, S.E., et al. (1982). Measurement and Computation of Streamflow: Volume 1. Measurement of Stage and Discharge. Geological Survey Water-Supply Paper #2175, U. S. Government Printing Office, Washington, D. C.
- Sung, W., B. Reilly-Matthews, D. K. O’Day, and K. Horrigan. 2000. Modeling DBP Formation. J. Amer. Water Works Assoc. 92:5–53.
- Wetzel, R. G. 2001. Limnology, Third Edition. Academic Press, San Diego, CA.
- YSI. 2010. YSI 6-Series Multiparameter Water Quality Sondes User Manual, Revision G, November 2012. YSI Incorporated, Yellow Springs, OH.

## **6.2 Related Reports**

The following is a list of annual reports and special project reports produced by the Institute for Watershed Studies since 1987 as part of the Lake Whatcom monitoring program sponsored by the City of Bellingham and Western Washington University. Many of the reports are available online at <http://www.wwu.edu/iws> (follow links to the Lake Whatcom project under Lake Studies); older reports are available in the IWS library and through the city of Bellingham Public Works Department. This list does not include research reports, student projects, or publications that were not prepared specifically for the City of Bellingham. Contact IWS for information about additional Lake Whatcom publications.

### **Annual monitoring reports:**

- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. 2012. Lake Whatcom Monitoring Project, 2010/2011 Final Report, February 24, 2012. Report to the City of Bellingham, WA.

- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2009/2010 Final Report, March 1, 2011. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2008/2009 Final Report, March 10, 2010. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2007/2008 Final Report, March 19, 2009. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2006/2007 Final Report, April 2, 2008. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2005/2006 Final Report, April 11, 2007. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2004/2005 Final Report, March 30, 2006. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2003/2004 Final Report, March 15, 2005. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2002/2003 Final Report, April 5, 2004. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2001/2002 Final Report, April 21, 2003. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 2000/2001 Final Report, March 15, 2002. Report to the City of Bellingham, WA.

- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 1999/2000 Final Report, March 23, 2001. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles, J. Vandersypen, R. Mitchell, and G. B. Matthews. Lake Whatcom Monitoring Project, 1998/99 Final Report, March 15, 2000. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles and G. B. Matthews. Lake Whatcom Monitoring Project, 1997/98 Final Report, April 12, 1999. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles and G. B. Matthews. Lake Whatcom Monitoring Project, 1996/97 Final Report, February 10, 1998. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles and G. B. Matthews. Lake Whatcom Monitoring Project, 1995/96 Final Report, March 24, 1997. Report to the City of Bellingham, WA.
- Matthews, R. A., M. Hilles and G. B. Matthews. Lake Whatcom Monitoring Project, 1994/95 Final Report, February 9, 1996. Report to the City of Bellingham, WA.
- Matthews, R. A. and G. B. Matthews. Lake Whatcom Monitoring Project, 1993–1994 Final Report, March 2, 1995. Report to the City of Bellingham, WA.
- Matthews, R. and G. Matthews. Lake Whatcom Monitoring Project, 1992–1993 Final Report, January 31, 1994. Report to the City of Bellingham, WA.
- Matthews, R. and G. Matthews. Lake Whatcom Monitoring Project, 1991–1992 Final Report, March 19, 1993. Report to the City of Bellingham, WA.
- Rector, J. M. and R. A. Matthews. Lake Whatcom Monitoring Program, August 1987 Final Report. Institute for Watershed Studies Report, Western Washington University, Bellingham, WA.

**Other Lake Whatcom reports:**

Matthews, R. A., M. Hilles and J. Vandersypen. Austin Creek and Beaver Creek Sampling Project, October 11, 2005. Report to the City of Bellingham, WA.

Matthews, R. A. Relationship between Drinking Water Treatment Chemical Usage and Lake Whatcom water Quality and Algal Data, October 4, 2004. Report to the City of Bellingham, WA.

Matthews, R. A. Strawberry Sill Water Quality Analysis, March 19, 2004. Report to the City of Bellingham, WA.

Matthews, R. A., M. Saunders, M A. Hilles, and J. Vandersypen. Park Place Wet Pond Monitoring Project, 1994–2000 Summary Report, February 2, 2001. Report to the City of Bellingham, WA.

Carpenter, M. R., C. A. Suczek, and R. A. Matthews. Mirror Lake Sedimentation Study Summary Report, February, 1992. Report to the City of Bellingham, WA.

Walker, S., R. Matthews, and G. Matthews. Lake Whatcom Storm Runoff Project, Final Report, January 13, 1992. Report to the City of Bellingham, WA.

Creahan, K., T. Loranger, B. Gall, D. Brakke, and R. Matthews. Lake Whatcom Watershed Management Plan, December, 1986, revised July, 1987. Institute for Watershed Studies Report, Western Washington University, Bellingham, WA.

## A Site Descriptions

Figures A1–A3 (pages 97–99) show the locations of the current monitoring sites and Table A1 (page 96) lists the approximate GPS coordinates for the lake and creek sites. All site descriptions, including text descriptions and GPS coordinates, are approximate because of variability in satellite coverage, GPS unit sensitivity, boat movement, stream bank or channel alterations, stream flow rates, weather conditions, and other factors that affect sampling location. Text descriptions contain references to local landmarks that may change over time. For detailed information about exact sampling locations, contact IWS.

### A.1 Lake Whatcom Monitoring Sites

**Site 1** is located at 20 m in the north central portion of basin 1 along a straight line from the Bloedel Donovan boat launch to the house located at 171 E. North Shore Rd. The depth at Site 1 should be at least 25 meters.

**Site 2** is located at 18–20 m in the south central portion of basin 2 just west of the intersection of a line joining the boat house at 73 Strawberry Point and the point of Geneva sill.

The **Intake Site** location is omitted from this report at the City's request.

**Site 3** is located in the northern portion of basin 3, mid-basin just north of a line between the old railroad bridge and Lakewood. The depth at Site 3 should be at least 80 m.

**Site 4** is located in the southern portion of basin 3, mid-basin, and just north of South Bay. The depth at Site 4 should be at least 90 m.

## A.2 Tributary Monitoring Sites

**Anderson Creek** samples are collected 15 m upstream from South Bay Rd. Water samples and discharge measurements are collected upstream from the bridge. The Anderson Creek hydrograph<sup>26</sup> is mounted in the stilling well on the east side of Anderson Creek, directly adjacent to the bridge over Anderson Creek (South Bay Rd.), approximately 0.5 km from the mouth of the creek.

The **Austin Creek** hydrograph gauge and sampling site is located approximately 15 m downstream from Lake Whatcom Blvd. From October 2004 through September 2006, three additional sampling sites were sampled in the Austin Creek watershed, so for clarification, the gauged site has been renamed **Lower Austin Creek**.

**Blue Canyon Creek** samples are collected downstream from the culvert under Blue Canyon Rd. in the second of three small streams that cross the road. This site can be difficult to locate and may be dry or have minimal flow during drought conditions; contact IWS for detailed information about the site location.

**Brannian Creek** samples are collected approximately 40 m downstream from South Bay Rd. near the USGS hydrograph gauge. This site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

**Carpenter Creek** samples are collected approximately 7 m upstream from North Shore Dr. near the USGS hydrograph gauge. This site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

**Euclid Ave.** samples are collected from an unnamed tributary located off Decator Rd. near the USGS hydrograph gauge. The site is named for its proximity to Euclid Ave., and was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

**Millwheel Creek** samples are collected approximately 8 m upstream from Flynn St. near the USGS hydrograph gauge. The creek is unnamed on most topographic maps, but has been called “Millwheel Creek” by residents of the watershed due to its proximity to the old mill pond. This site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

---

<sup>26</sup>This hydrograph is no longer maintained by IWS; contact the City of Bellingham for data.

**Olsen Creek** samples are collected just downstream from North Shore Dr. near the USGS hydrograph gauge. This site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

**Park Place** samples are collected from the storm drain that empties into Lake Whatcom at Park Place Ln. Samples from this site include outlet flow from the Park Place storm water treatment facility.

**Silver Beach Creek** samples are collected approximately 15 m upstream from the culvert under North Shore Rd.

The **Smith Creek** hydrograph is mounted on the south wall of a sandstone bluff directly underneath the bridge over Smith Creek (North Shore Rd.) approximately 1 km upstream from the mouth of the creek. Water samples are collected at the gaging station approximately 15 m downstream from North Shore Dr.

**Whatcom Creek** samples are collected approximately 2 m downstream from the foot bridge below the Lake Whatcom outlet spillway. This site was added in October 2004 as part of the monthly 2004–2006 creek monitoring project.

### **A.3 Storm Water Monitoring Sites**

The storm water monitoring program was revised in 2009/2010 to focus on collecting baseline data at the Silver Beach Creek outlet and the North Shore Drive overlay. Both sites were monitored in 2009/2010 (see Matthews, et al., 2011). During the 2010/2011 and 2011/2012 monitoring periods the emphasis was on collecting additional storm water samples from Silver Beach Creek. The 2010/2011 Silver Beach Creek runoff data are described by Matthew, et al. (2012). For information about other storm water sites that have been monitored by IWS, refer to the annual reports listed in Section 6.2 (page 89).


**Silver Beach** storm runoff samples were collected at the USGS gauging site behind the house at 3007 Maynard Place and approximately 150 m upstream from the culvert at North Shore Dr.



Lake Sites	Latitude (°N)	Longitude (°W)
Site 1	48.4536	122.2438
Intake	(GPS omitted)	
Site 2	48.4436	122.2254
Site 3	48.4416	122.2009
Site 4	48.4141	122.1815


Creek Sites	Latitude (°N)	Longitude (°W)
Anderson	48.67335	122.26751
Austin (lower)	48.71312	122.33076
Blue Canyon	48.68532	122.28295
Brannian	48.66910	122.27949
Carpenter	48.75432	122.35449
Euclid	48.74844	122.41005
Millwheel	48.75507	122.41635
Olsen	48.75129	122.35353
Park Place	48.76894	122.40915
Silver Beach	48.76859	122.40700
Smith	48.73191	122.30864
Whatcom	48.75715	122.42229

Table A1: Approximate GPS coordinates for Lake Whatcom sampling sites.




This figure was created using source files provided by Gerald Gabrisch using data obtained from Western Washington University, Skagit County, the Nooksack Tribe, and the City of Bellingham.

Figure A1: Lake Whatcom lake sampling sites.



This figure was created using source files provided by Gerald Gabrisch using data obtained from Western Washington University, Skagit County, the Nooksack Tribe, and the City of Bellingham.

Figure A2: Lake Whatcom tributary sampling sites.



This figure was created using source files provided by Gerald Gabrisch using data obtained from Western Washington University, Skagit County, the Nooksack Tribe, and the City of Bellingham.

Figure A3: Silver Beach Creek storm water site.

(This page blank)

## **B Long-Term Water Quality Figures**

The current and historic Lake Whatcom water quality data are plotted on the following pages. Detection limits and abbreviations for each parameter are listed in Table 1 (page 16).

The historic detection limits for each parameter were estimated based on recommended lower detection ranges (APHA, 1998; Hydrolab, 1997; Lind, 1985), instrument limitations, and analyst judgment on the lowest repeatable concentration for each test. Over time, some analytical techniques have improved so that current detection limits are lower than defined below (see current detection limits in Table 1, page 16). Because the Lake Whatcom data set includes long-term monitoring data that have been collected using a variety of analytical techniques, this report sets conservative historic detection limits to allow comparisons between all years.

In the Lake Whatcom report, unless indicated, no data substitutions are used for below detection values (“bdl” data). Instead, we identify summary statistics that include bdl values, and, if appropriate, discuss the implications of including these values in the analysis.

Because of the length of the data record, many of the figures reflect trends related to improvements in analytical techniques over time, and introduction of increasingly sensitive field equipment (see, for example, Figures B66–B70, pages 169–173, which show the effect of using increasingly sensitive conductivity probes). These changes generally result in a reduction in analytical variability, and sometimes result in lower detection limits.

## **B.1 Monthly Hydrolab Profiles**




Figure B1: Lake Whatcom Hydrolab profiles for Site 1, October 4, 2011. Conductivity profile is not available due to equipment malfunction; discrete results were generated from water samples measured in the laboratory.






Figure B2: Lake Whatcom Hydrolab profiles for Site 2, October 4, 2011. Conductivity profile is not available due to equipment malfunction; discrete results were generated from water samples measured in the laboratory.




Figure B3: Lake Whatcom Hydrolab profiles for the Intake, October 4, 2011. Conductivity profile is not available due to equipment malfunction; discrete results were generated from water samples measured in the laboratory.




Figure B4: Lake Whatcom Hydrolab profiles for Site 3, October 5, 2011. Conductivity profile is not available due to equipment malfunction; discrete results were generated from water samples measured in the laboratory.




Figure B5: Lake Whatcom Hydrolab profiles for Site 4, October 5, 2011. Conductivity profile is not available due to equipment malfunction; discrete results were generated from water samples measured in the laboratory.




Figure B6: Lake Whatcom YSI profiles for Site 1, November 1, 2011.




Figure B7: Lake Whatcom YSI profiles for Site 2, November 1, 2011.




Figure B8: Lake Whatcom YSI profiles for the Intake, November 1, 2011.




Figure B9: Lake Whatcom YSI profiles for Site 3, November 2, 2011.






Figure B10: Lake Whatcom YSI profiles for Site 4, November 2, 2011.




Figure B11: Lake Whatcom YSI profiles for Site 1, December 7, 2011.




Figure B12: Lake Whatcom YSI profiles for Site 2, December 7, 2011.




Figure B13: Lake Whatcom YSI profiles for the Intake, December 7, 2011.




Figure B14: Lake Whatcom YSI profiles for Site 3, December 6, 2011. The low oxygen value at 80 meters was most likely due to incomplete water column mixing following recent destratification.




Figure B15: Lake Whatcom YSI profiles for Site 4, December 6, 2011.




Figure B16: Lake Whatcom YSI profiles for Site 1, February 3, 2012.




Figure B17: Lake Whatcom YSI profiles for Site 2, February 3, 2012.






Figure B18: Lake Whatcom YSI profiles for the Intake, February 3, 2012.




Figure B19: Lake Whatcom YSI profiles for Site 3, February 1, 2012.




Figure B20: Lake Whatcom YSI profiles for Site 4, February 1, 2012.




Figure B21: Lake Whatcom YSI profiles for Site 1, April 14, 2012.




Figure B22: Lake Whatcom YSI profiles for Site 2, April 14, 2012.




Figure B23: Lake Whatcom YSI profiles for the Intake, April 14, 2012.




Figure B24: Lake Whatcom YSI profiles for Site 3, April 10, 2012.




Figure B25: Lake Whatcom YSI profiles for Site 4, April 10, 2012.






Figure B26: Lake Whatcom YSI profiles for Site 1, May 10, 2012.




Figure B27: Lake Whatcom YSI profiles for Site 2, May 10, 2012.




Figure B28: Lake Whatcom YSI profiles for the Intake, May 10, 2012.




Figure B29: Lake Whatcom YSI profiles for Site 3, May 3, 2012.




Figure B30: Lake Whatcom YSI profiles for Site 4, May 3, 2012.




Figure B31: Lake Whatcom YSI profiles for Site 1, June 7, 2012.




Figure B32: Lake Whatcom YSI profiles for Site 2, June 7, 2012.




Figure B33: Lake Whatcom YSI profiles for the Intake, June 7, 2012.






Figure B34: Lake Whatcom YSI profiles for Site 3, June 9, 2012.




Figure B35: Lake Whatcom YSI profiles for Site 4, June 9, 2012.




Figure B36: Lake Whatcom YSI profiles for Site 1, July 7, 2012.




Figure B37: Lake Whatcom YSI profiles for Site 2, July 7, 2012.




Figure B38: Lake Whatcom YSI profiles for the Intake, July 7, 2012.




Figure B39: Lake Whatcom YSI profiles for Site 3, July 5, 2012.




Figure B40: Lake Whatcom YSI profiles for Site 4, July 5, 2012.




Figure B41: Lake Whatcom YSI profiles for Site 1, August 4, 2012.






Figure B42: Lake Whatcom YSI profiles for Site 2, August 4, 2012.




Figure B43: Lake Whatcom YSI profiles for the Intake, August 4, 2012.




Figure B44: Lake Whatcom YSI profiles for Site 3, August 2, 2012.




Figure B45: Lake Whatcom YSI profiles for Site 4, August 2, 2012.




Figure B46: Lake Whatcom YSI profiles for Site 1, September 8, 2012.




Figure B47: Lake Whatcom YSI profiles for Site 2, September 8, 2012.




Figure B48: Lake Whatcom YSI profiles for the Intake, September 8, 2012.




Figure B49: Lake Whatcom YSI profiles for Site 3, September 6, 2012.






Figure B50: Lake Whatcom YSI profiles for Site 4, September 6, 2012.

## **B.2 Long-term Hydrolab Data (1988-present)**

Lake Whatcom temperature data for Site 1, February 1988 through December 2012.




Figure B51: Lake Whatcom historic temperature data for Site 1.

Lake Whatcom temperature data for Site 2, February 1988 through December 2012.




Figure B52: Lake Whatcom historic temperature data for Site 2.

Lake Whatcom temperature data for Intake, February 1988 through December 2012.




Figure B53: Lake Whatcom historic temperature data for the Intake.

Lake Whatcom temperature data for Site 3, February 1988 through December 2012.




Figure B54: Lake Whatcom historic temperature data for Site 3.

Lake Whatcom temperature data for Site 4, February 1988 through December 2012.




Figure B55: Lake Whatcom historic temperature data for Site 4.

Lake Whatcom dissolved oxygen data for Site 1, February 1988 through December 2012.




Figure B56: Lake Whatcom historic dissolved oxygen data for Site 1.



Lake Whatcom dissolved oxygen data for Site 2, February 1988 through December 2012.




Figure B57: Lake Whatcom historic dissolved oxygen data for Site 2.

Lake Whatcom dissolved oxygen data for Intake, February 1988 through December 2012.




Figure B58: Lake Whatcom historic dissolved oxygen data for the Intake.

Lake Whatcom dissolved oxygen data for Site 3, February 1988 through December 2012.




Figure B59: Lake Whatcom historic dissolved oxygen data for Site 3.

Lake Whatcom dissolved oxygen data for Site 4, February 1988 through December 2012.




Figure B60: Lake Whatcom historic dissolved oxygen data for Site 4.

Lake Whatcom pH data for Site 1, February 1988 through December 2012.




Figure B61: Lake Whatcom historic pH data for Site 1.

Lake Whatcom pH data for Site 2, February 1988 through December 2012.




Figure B62: Lake Whatcom historic pH data for Site 2.

Lake Whatcom pH data for Intake, February 1988 through December 2012.




Figure B63: Lake Whatcom historic pH data for the Intake.

Lake Whatcom pH data for Site 3, February 1988 through December 2012.




Figure B64: Lake Whatcom historic pH data for Site 3.



Lake Whatcom pH data for Site 4, February 1988 through December 2012.




Figure B65: Lake Whatcom historic pH data for Site 4.




Figure B66: Lake Whatcom historic conductivity data for Site 1. The decreasing conductivity trend is the result of changing to more sensitive equipment.




Figure B67: Lake Whatcom historic conductivity data for Site 2. The decreasing conductivity trend is the result of changing to more sensitive equipment.




Figure B68: Lake Whatcom historic conductivity data for the Intake. The decreasing conductivity trend is the result of changing to more sensitive equipment.




Figure B69: Lake Whatcom historic conductivity data for Site 3. The decreasing conductivity trend is the result of changing to more sensitive equipment.




Figure B70: Lake Whatcom historic conductivity data for Site 4. The decreasing conductivity trend is the result of changing to more sensitive equipment.

### **B.3 Long-term Water Quality Data (1988-present)**

Lake Whatcom alkalinity data for Site 1, February 1988 through December 2012.




Figure B71: Lake Whatcom alkalinity data for Site 1.



Lake Whatcom alkalinity data for Site 2, February 1988 through December 2012.




Figure B72: Lake Whatcom alkalinity data for Site 2.

Lake Whatcom alkalinity data for Intake, February 1988 through December 2012.




Figure B73: Lake Whatcom alkalinity data for the Intake site.

Lake Whatcom alkalinity data for Site 3, February 1988 through December 2012.




Figure B74: Lake Whatcom alkalinity data for Site 3.

Lake Whatcom alkalinity data for Site 4, February 1988 through December 2012.




Figure B75: Lake Whatcom alkalinity data for Site 4.

Lake Whatcom turbidity data for Site 1, February 1988 through December 2012.




Figure B76: Lake Whatcom turbidity data for Site 1.

Lake Whatcom turbidity data for Site 2, February 1988 through December 2012.




Figure B77: Lake Whatcom turbidity data for Site 2.

Lake Whatcom turbidity data for Intake, February 1988 through December 2012.



Figure B78: Lake Whatcom turbidity data for the Intake site.

Lake Whatcom turbidity data for Site 3, February 1988 through December 2012.




Figure B79: Lake Whatcom turbidity data for Site 3.



Lake Whatcom turbidity data for Site 4, February 1988 through December 2012.




Figure B80: Lake Whatcom turbidity data for Site 4.

Lake Whatcom ammonium data for Site 1, February 1988 through December 2012.




Figure B81: Lake Whatcom ammonium data for Site 1.

Lake Whatcom ammonium data for Site 2, February 1988 through December 2012.




Figure B82: Lake Whatcom ammonium data for Site 2.

Lake Whatcom ammonium data for Intake, February 1988 through December 2012.




Figure B83: Lake Whatcom ammonium data for the Intake site.

Lake Whatcom ammonium data for Site 3, February 1988 through December 2012.




Figure B84: Lake Whatcom ammonium data for Site 3.

Lake Whatcom ammonium data for Site 4, February 1988 through December 2012.




Figure B85: Lake Whatcom ammonium data for Site 4.

Lake Whatcom nitrate/nitrite data for Site 1, February 1988 through December 2012.




Figure B86: Lake Whatcom nitrate/nitrite data for Site 1.

Lake Whatcom nitrate/nitrite data for Site 2, February 1988 through December 2012.




Figure B87: Lake Whatcom nitrate/nitrite data for Site 2.



Lake Whatcom nitrate/nitrite data for Intake, February 1988 through December 2012.




Figure B88: Lake Whatcom nitrate/nitrite data for the Intake site.

Lake Whatcom nitrate/nitrite data for Site 3, February 1988 through December 2012.




Figure B89: Lake Whatcom nitrate/nitrite data for Site 3.

Lake Whatcom nitrate/nitrite data for Site 4, February 1988 through December 2012.




Figure B90: Lake Whatcom nitrate/nitrite data for Site 4.

Lake Whatcom total nitrogen data for Site 1, February 1988 through December 2012.




Figure B91: Lake Whatcom total nitrogen data for Site 1.

Lake Whatcom total nitrogen data for Site 2, February 1988 through December 2012.




Figure B92: Lake Whatcom total nitrogen data for Site 2.

Lake Whatcom total nitrogen data for Intake, February 1988 through December 2012.




Figure B93: Lake Whatcom total nitrogen data for the Intake site.

Lake Whatcom total nitrogen data for Site 3, February 1988 through December 2012.




Figure B94: Lake Whatcom total nitrogen data for Site 3.

Lake Whatcom total nitrogen data for Site 4, February 1988 through December 2012.




Figure B95: Lake Whatcom total nitrogen data for Site 4.



Lake Whatcom soluble reactive phosphate data for Site 1, February 1988 through December 2012.




Figure B96: Lake Whatcom soluble phosphate data for Site 1.

Lake Whatcom soluble reactive phosphate data for Site 2, February 1988 through December 2012.




Figure B97: Lake Whatcom soluble phosphate data for Site 2.

Lake Whatcom soluble reactive phosphate data for Intake, February 1988 through December 2012.




Figure B98: Lake Whatcom soluble phosphate data for the Intake site.

Lake Whatcom soluble reactive phosphate data for Site 3, February 1988 through December 2012.




Figure B99: Lake Whatcom soluble phosphate data for Site 3.

Lake Whatcom soluble reactive phosphate data for Site 4, February 1988 through December 2012.




Figure B100: Lake Whatcom soluble phosphate data for Site 4.

Lake Whatcom total phosphorus data for Site 1, February 1988 through December 2012.




Figure B101: Lake Whatcom total phosphorus data for Site 1.

Lake Whatcom total phosphorus data for Site 2, February 1988 through December 2012.




Figure B102: Lake Whatcom total phosphorus data for Site 2.

Lake Whatcom total phosphorus data for Intake, February 1988 through December 2012.




Figure B103: Lake Whatcom total phosphorus data for the Intake site.



Lake Whatcom total phosphorus data for Site 3, February 1988 through December 2012.




Figure B104: Lake Whatcom total phosphorus data for Site 3.

Lake Whatcom total phosphorus data for Site 4, February 1988 through December 2012.




Figure B105: Lake Whatcom total phosphorus data for Site 4.

Lake Whatcom chlorophyll a data for Site 1, February 1988 through December 2012.




Figure B106: Lake Whatcom chlorophyll data for Site 1.

Lake Whatcom chlorophyll a data for Site 2, February 1988 through December 2012.




Figure B107: Lake Whatcom chlorophyll data for Site 2.

Lake Whatcom chlorophyll a data for Intake, February 1988 through December 2012.




Figure B108: Lake Whatcom chlorophyll data for the Intake site.

Lake Whatcom chlorophyll a data for Site 3, February 1988 through December 2012.




Figure B109: Lake Whatcom chlorophyll data for Site 3.

Lake Whatcom chlorophyll a data for Site 4, February 1988 through December 2012.




Figure B110: Lake Whatcom chlorophyll data for Site 4.

Lake Whatcom Secchi data for Site 1, February 1988 through December 2012.




Figure B111: Lake Whatcom Secchi depths for Site 1.



Lake Whatcom Secchi data for Site 2, February 1988 through December 2012.




Figure B112: Lake Whatcom Secchi depths for Site 2.

Lake Whatcom Secchi data for Intake, February 1988 through December 2012.




Figure B113: Lake Whatcom Secchi depths for the Intake site.

Lake Whatcom Secchi data for Site 3, February 1988 through December 2012.




Figure B114: Lake Whatcom Secchi depths for Site 3.

Lake Whatcom Secchi data for Site 4, February 1988 through December 2012.




Figure B115: Lake Whatcom Secchi depths for Site 4.

Lake Whatcom fecal coliform data for Site 1, February 1988 through December 2012.




Figure B116: Lake Whatcom fecal coliform data for Site 1.

Lake Whatcom fecal coliform data for Site 2, February 1988 through December 2012.




Figure B117: Lake Whatcom fecal coliform data for Site 2.

Lake Whatcom fecal coliform data for Intake, February 1988 through December 2012.




Figure B118: Lake Whatcom fecal coliform data for the Intake site.

Lake Whatcom fecal coliform data for Site 3, February 1988 through December 2012.




Figure B119: Lake Whatcom fecal coliform data for Site 3.



Lake Whatcom fecal coliform data for Site 4, February 1988 through December 2012.




Figure B120: Lake Whatcom fecal coliform data for Site 4.

Lake Whatcom plankton data for Site 1, February 1988 through December 2012.




Figure B121: Lake Whatcom plankton data for Site 1.

Lake Whatcom plankton data for Site 2, February 1988 through December 2012.




Figure B122: Lake Whatcom plankton data for Site 2.

Lake Whatcom plankton data for Intake, February 1988 through December 2012.




Figure B123: Lake Whatcom plankton data for the Intake Site.

Lake Whatcom plankton data for Site 3, February 1988 through December 2012.




Figure B124: Lake Whatcom plankton data for Site 3.

Lake Whatcom plankton data for Site 4, February 1988 through December 2012.




Figure B125: Lake Whatcom plankton data for Site 4.




Figure B126: Lake Whatcom plankton data for Site 1, with Chrysophyta omitted to show remaining plankton groups.




Figure B127: Lake Whatcom plankton data for Site 2, with Chrysochyta omitted to show remaining plankton groups.






Figure B128: Lake Whatcom plankton data for the Intake Site, with Chrysophyta omitted to show remaining plankton groups.




Figure B129: Lake Whatcom plankton data for Site 3, with Chrysochyta omitted to show remaining plankton groups.




Figure B130: Lake Whatcom plankton data for Site 4, with Chrysochyta omitted to show remaining plankton groups.

## **B.4 Lake Whatcom Tributary Data (2004-present)**

The figures in this appendix include the monthly baseline data collected from October 2004 through September 2006, biannual data collected from February 2007 through September 2009, and monthly data collected during the current monitoring period. Each figure includes a dashed (blue) horizontal line that shows the median value for Smith Creek and a solid (red) horizontal line that shows the median value for each creek. Smith Creek was chosen as a reference because it is a major tributary to the lake and has a history of being relatively unpolluted. The figures were scaled to include all but extreme outliers; off-scale outliers are listed in the figure caption. All data are available online at <http://www.wvu.edu/iws>.




Figure B131: Temperature data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B132: Temperature data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B133: Temperature data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B134: Dissolved oxygen data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.






Figure B135: Dissolved oxygen data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B136: Dissolved oxygen data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B137: Tributary pH data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B138: Tributary pH data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B139: Tributary pH data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B140: Conductivity data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B141: Conductivity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B142: Conductivity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.






Figure B143: Alkalinity data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B144: Alkalinity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B145: Alkalinity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B146: Total suspended solids data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek. Two outliers were off-scale (Austin and Anderson Creeks, Jan. 10, 2006).




Figure B147: Total suspended solids data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek. Two outliers were off-scale (Brannian and Olsen Creeks, Jan. 10, 2006).




Figure B148: Total suspended solids data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B149: Turbidity data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek. One outlier was off-scale (Anderson Creek, Jan. 10, 2006).




Figure B150: Turbidity data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.






Figure B151: Turbidity data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B152: Ammonium data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B153: Ammonium data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B154: Ammonium data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek. Three outliers were off-scale (Millwheel Creek, Feb. 8, 2005, July 11, 2011, Sept. 12, 2012).




Figure B155: Nitrate/nitrite data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B156: Nitrate/nitrite data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B157: Nitrate/nitrite data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B158: Total nitrogen data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.






Figure B159: Total nitrogen data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B160: Total nitrogen data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B161: Soluble phosphate data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B162: Soluble phosphate data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B163: Soluble phosphate data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek. One outlier was off-scale (Millwheel Creek, Feb. 8, 2005).




Figure B164: Total phosphorus data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B165: Total phosphorus data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B166: Total phosphorus data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek. Three outliers were off-scale (Millwheel Creek, Sept. 14, 2010, Oct. 12, 2011, Sept. 12, 2012).






Figure B167: Fecal coliform data for Anderson, Austin, Smith, and Whatcom Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek. One outlier was off-scale (Austin Creek, July 17, 2007).




Figure B168: Fecal coliform data for Blue Canyon, Brannian, Carpenter, and Olsen Creeks. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek.




Figure B169: Fecal coliform data for Euclid, Millwheel, and Silver Beach Creeks and the Park Place drain. Dashed (blue) horizontal reference line shows the median value for Smith Creek; solid (red) horizontal reference line shows the median value for each creek. Ten outliers were off-scale (Euclid Creek, Oct. 12, 2011; Millwheel Creek, Sept. 14, 2010, June 5, 2012; Park Place drain, Aug. 1 2006; Silver Beach Creek, Oct. 10, 2005, Sept. 13, 2005, Aug. 1, 2006, July 17, 2007, July 15, 2008, Sept. 13, 2011).

## **C Quality Control**

### **C.1 Performance Evaluation Report**

In order to maintain a high degree of accuracy and confidence in the water quality data all personnel associated with this project were trained according to standard operating procedures for the methods listed in Table 1 (page 16). Single-blind quality control tests were conducted as part of the IWS laboratory certification process (Table C1). All results from the single-blind tests were within acceptance limits.

	Reported Value	True Value	Acceptance Limits	Test Result
Specific conductivity ( $\mu\text{S}/\text{cm}$ at $25^\circ\text{C}$ )	420	418	375–462	accept
Total alkalinity (mg/L as $\text{CaCO}_3$ )	101	103	91.4–113	accept
Ammonium nitrogen, manual (mg-N/L)	7.69	7.74	5.70–9.74	accept
Ammonium nitrogen, autoanalysis (mg-N/L)	7.65	7.74	5.70–9.74	accept
Nitrate/nitrite nitrogen, autoanalysis (mg-N/L)	11.0	10.6	8.64–12.3	accept
Nitrite nitrogen, autoanalysis (mg-N/L)	0.470	0.480	0.349–0.602	accept
Orthophosphate, manual (mg-P/L)	2.37	2.40	1.94–2.88	accept
Orthophosphate, autoanalysis (mg-P/L)	2.45	2.40	1.94–2.88	accept
Total phosphorus, manual (mg-P/L)	1.64	1.66	1.31–2.06	accept
Total phosphorus, autoanalysis (mg-P/L)	1.68	1.66	1.31–2.06	accept
pH	7.78	7.80	7.60–8.00	accept
Solids, non-filterable (mg/L)	84.4	94.0	77.6–104	accept
Turbidity (NTU)	14.8	15.5	12.8–18.2	accept

Table C1: Single-blind quality control results, WP-183 (06/06/2012).

## C.2 Laboratory Duplicates, Spikes, and Check Standards

Ten percent of all lake, storm water, and tributary samples analyzed in the laboratory were duplicated to measure analytical precision. Sample matrix spikes were analyzed during each analytical run to evaluate analyte recovery for the nutrient analyses (ammonium, nitrate/nitrite, total nitrogen, soluble reactive phosphate, and total phosphorus). External check standards were analyzed during each analytical run to evaluate measurement precision and accuracy.<sup>27</sup>

The quality control results for laboratory duplicates, matrix spikes, and check standards are plotted in control charts. Upper and lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper and lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were developed using data from September 2008 through September 2011 (upper examples in Figures C1–C30, pages 278–307), and used to evaluate data from October 2011 through September 2012 (lower examples in Figures C1–C30).

---

<sup>27</sup>External check standards are not available for all analytes.





Figure C1: Alkalinity laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C2: Alkalinity high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.







Figure C3: Alkalinity low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C4: Chlorophyll laboratory duplicates for the Lake Whatcom monitoring program (lake samples). Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C5: Conductivity laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C6: Dissolved oxygen laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C7: Ammonium laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C8: Ammonium matrix spikes for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C9: Ammonium high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C10: Ammonium low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.







Figure C11: Nitrate/nitrite laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C12: Nitrate/nitrite matrix spikes for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C13: Nitrate/nitrite high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C14: Nitrate/nitrite low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C15: Total nitrogen laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C16: Total nitrogen matrix spikes for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C17: Total nitrogen high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C18: Total nitrogen low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.







Figure C19: Laboratory pH duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C20: Soluble reactive phosphate laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C21: Soluble reactive phosphate matrix spikes for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C22: Soluble reactive phosphate high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C23: Soluble reactive phosphate low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C24: Total phosphorus laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data. Slight increase in variability may be due to insufficient persulfate concentration; method revised to increase concentration.





Figure C25: Total phosphorus matrix spikes for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C26: Total phosphorus high-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.







Figure C27: Total phosphorus low-range check standards for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C28: Total suspended solids laboratory duplicates for the Lake Whatcom monitoring program (creek and storm water samples). Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C29: Total suspended solids check standards for the Lake Whatcom monitoring program (creek and storm water samples). Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.





Figure C30: Turbidity laboratory duplicates for the Lake Whatcom monitoring program. Upper/lower acceptance limits ( $\pm 2$  std. dev. from mean pair difference) and upper/lower warning limits ( $\pm 3$  std. dev. from mean pair difference) were calculated based on the preceding two years of lab duplicate data.

### **C.3 Field Duplicate Results**

Field duplicates (Figures C31–C49, pages 309–327) were collected and analyzed for a minimum of 10% of all of the water quality parameters except the Hydrolab or YSI field meter data. To check the field meter measurements, duplicate samples were analyzed for at least 10% of the field meter measurements using water samples collected from the same depth as the field meter measurement.

The absolute mean difference for the field duplicates was calculated using the following equation:

$$\text{Absolute mean difference} = \frac{\sum |\text{Original Sample} - \text{Duplicate Sample}|}{\text{number of duplicate pairs}}$$




Figure C31: Alkalinity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship.




Figure C32: Alkalinity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). Diagonal reference line shows a 1:1 relationship.




Figure C33: Chlorophyll field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship.






Figure C34: Conductivity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship. The high degree of scatter is due to the low concentration of the samples.




Figure C35: Dissolved oxygen field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship. The labeled outliers were collected when the lake was stratified, or recently destratified and incompletely mixed, at depths where extreme gradients were present. Field meter samples were collected at true depth; Winkler samples were collected using a marked line, which is slightly shallower than true depth.




Figure C36: Dissolved oxygen field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). Diagonal reference line shows a 1:1 relationship.




Figure C37: Ammonium field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship; horizontal reference line shows the current detection limits. The high degree of scatter is due to the low concentrations of the samples.




Figure C38: Ammonium field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). Diagonal reference line shows a 1:1 relationship; horizontal reference line shows the current detection limits. The high degree of scatter is due to the low concentrations of the samples.




Figure C39: Nitrate/nitrite field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship; horizontal reference line shows the current detection limits.




Figure C40: Nitrate/nitrite field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). Diagonal reference line shows a 1:1 relationship; horizontal reference line shows the current detection limits.




Figure C41: Total nitrogen field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship. All total nitrogen samples were above the detection limit.






Figure C42: Total nitrogen field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). Diagonal reference line shows a 1:1 relationship. All total nitrogen samples were above the detection limit.




Figure C43: Field duplicates for pH from the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship.




Figure C44: Soluble phosphorus field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). Diagonal reference line shows a 1:1 relationship; horizontal reference line shows the current detection limits. The high degree of scatter is due to the low concentrations of the samples.




Figure C45: Total phosphorus field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship; horizontal reference line shows the current detection limits. The high degree of scatter is due to the low concentrations of the samples.




Figure C46: Total phosphorus field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). Diagonal reference line shows a 1:1 relationship; horizontal reference line shows the current detection limits. The high degree of scatter is due to the low concentrations of the samples.




Figure C47: Total suspended solids field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). Diagonal reference line shows a 1:1 relationship.




Figure C48: Turbidity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (lake samples). Diagonal reference line shows a 1:1 relationship.




Figure C49: Turbidity field duplicates for the 2011/2012 Lake Whatcom Monitoring Project (creek samples). Diagonal reference line shows a 1:1 relationship.



(This page blank)

## D Lake Whatcom Online Data

The following **readme** file describes the electronic data posted at the IWS web site and additional data available from IWS. Please contact the Director of the Institute for Watershed Studies if you have questions or trouble accessing the online data.

```
*****  
* README FILE - LAKE WHATCOM ONLINE DATA  
* THIS FILE WAS UPDATED MARCH 7, 2013  
*****  
Most of the Lake Whatcom water quality data are available in  
electronic format at the IWS website (http://www.wvu.edu/iws) or from  
the IWS Director.
```

The historic and current detection limits and abbreviations for each parameter are listed in the annual reports. The historic detection limits for each parameter were estimated based on recommended lower detection ranges, instrument limitations, and analyst judgment on the lowest repeatable concentration for each test. Over time, some analytical techniques have improved so that current detection limits are usually lower than historic detection limits. Because the Lake Whatcom data set includes long-term monitoring data, which have been collected using a variety of analytical techniques, this report sets conservative detection limits to allow comparisons between years.

All files are comma-separated ascii data files. The code "NA" has been entered into all empty cells in the ascii data files to fill in unsampled dates and depths, missing data, etc. Questions about missing data should be directed to the IWS Director.

Unless otherwise indicated, the electronic data files have NOT been censored to flag or otherwise identify below detection and above detection values. As a result, the ascii files may contain negative values due to linear extrapolation of the standards regression curve for below detection data. It is essential that any statistical or analytical results that are generated using these data be reviewed by someone familiar with statistical uncertainty associated with uncensored data.

\*\*\*\*\*

\* ONLINE LAKE DATA FILES:

\*\*\*\*\*

Hydrolab/YSI data	Water quality data	Plankton data
1988_hl.csv	1988_wq.csv	plankton.csv
1989_hl.csv	1989_wq.csv	
1990_hl.csv	1990_wq.csv	
1991_hl.csv	1991_wq.csv	
1992_hl.csv	1992_wq.csv	
1993_hl.csv	1993_wq.csv	
1994_hl.csv	1994_wq.csv	
1995_hl.csv	1995_wq.csv	
1996_hl.csv	1996_wq.csv	
1997_hl.csv	1997_wq.csv	
1998_hl.csv	1998_wq.csv	
1999_hl.csv	1999_wq.csv	
2000_hl.csv	2000_wq.csv	
2001_hl.csv	2001_wq.csv	
2002_hl.csv	2002_wq.csv	
2003_hl.csv	2003_wq.csv	
2004_hl.csv	2004_wq.csv	
2005_hl.csv	2005_wq.csv	
2006_hl.csv	2006_wq.csv	
2007_hl.csv	2007_wq.csv	
2008_hl.csv	2008_wq.csv	
2009_hl.csv	2009_wq.csv	
2010_hl.csv	2010_wq.csv	
2011_hl.csv	2011_wq.csv	
2012_hl.csv	2012_wq.csv	

The \*\_hl.csv files include: site, depth (m), month, day, year, temp (water temperature, C), pH, cond (specific conductivity, uS/cm), do (dissolved oxygen, mg/L), lcond (lab conductivity quality control data, uS/cm), secchi (secchi depth, m).

The \*\_wq.csv files include: site, depth (m), month, day, year, alk (alkalinity, mg/L as CaCO<sub>3</sub>), turb (turbidity, NTU), nh3 (ammonium, ug-N/L), tn (total persulfate nitrogen, ug-N/L), nos (nitrate/nitrite, ug-N/L), srp (soluble reactive phosphate, ug-P/L), tp (total persulfate phosphorus, ug-P/L), chl (chlorophyll, ug/L).

The plankton.csv file includes: site, depth (m), month, day, year, zoop (zooplankton, #/L), chry (chrysophyta, #/L), cyan (cyanobacteria, #/L), chlo (chlorophyta, #/L), pyrr (pyrrophyta, #/L).

```
*****
* ONLINE HYDROGRAPH DATA FILES:
*****
WY1998.csv
WY1999.csv
WY2000_rev.csv (revised March 8, 2012)
WY2001.csv
WY2002.csv
WY2003.csv
WY2004_rev.csv (revised June 21, 2006)
WY2005.csv
WY2006.csv
WY2007.csv (revised July 31, 2008)
WY2008.csv
WY2009.csv
WY2010.csv
WY2011.csv
WY2012.csv
```

The WY\*.csv files include: month, day, year, hour, min, sec, ander.g (anderson gage height, ft), ander.cfs (anderson discharge, cfs), austin.g (austin gage height, ft), austin.cfs (austin discharge, cfs), smith.g (smith gage height, ft), smith.cfs (smith discharge, cfs). Anderson Creek hydrograph data were deleted in WY2000\_rev.csv due to uncertainty about the gage height; Anderson Creek data are available for WY1998, WY1999, and WY2001-WY2007. Beginning with WY2002, the variable "time" replaced "hour, min, sec," with time reported daily on a 24-hr basis. Data are reported as Pacific Standard Time without Daylight Saving Time adjustment.

```
*****
* STORM WATER DATA FILES
*****
The storm water data include composite and grab samples from numerous sites in the Lake Whatcom watershed (1994--present), representing a variety of study objectives and sampling intensities over time. The electronic data files are not posted online, but may be obtained by contacting the Institute for Watershed Studies.
```

\*\*\*\*\*

\* TRIBUTARY DATA FILES:

\*\*\*\*\*

CURRENT (ONLINE):

The current creek data are listed in creeks.csv. The data file contains the following variables: code (IWS site code), site (descriptive site name), month, day, year, time (24-hr basis), temp (water temperature, C), ph, do (dissolved oxygen, mg/L), cond (specific conductivity, uS/cm), turb (turbidity, NTU), alk (alkalinity, mg/L as CaCO<sub>3</sub>), tp (total phosphorus, ug-P/L), tn (total nitrogen, ug-N/L), nos (nitrite+nitrate, ug-N/L), srp (soluble reactive phosphate, ug-P/L), nh<sub>3</sub> (ammonium, ug-N/L), tss (total suspended solids, mg/L), ts (total solids, mg/L), ecoli (E.coli, cfu/100 mL), fc (fecal coliforms, cfu/100 mL)

HISTORIC STORM WATER MONITORING DATA:

Historic creek data include metals and toc data (creeksmetaltoc.csv); results from an intensive sampling effort in Austin Creek and Creek (creekwalk.csv); a 48-hr creek sampling project; and discharge estimated from ungauged sites. The electronic data are not available online, but may be obtained by contacting the Institute for Watershed Studies.

\*\*\*\*\*

\* SITE CODES

\* ALL FILES - INCLUDES DISCONTINUED SITES AND OFF-LINE DATA

\*\*\*\*\*

The site codes in the data are as follows:

11 = Lake Whatcom Site 1  
 21 = Lake Whatcom Intake site  
 22 = Lake Whatcom Site 2  
 31 = Lake Whatcom Site 3  
 32 = Lake Whatcom Site 4  
 33 = Strawberry Sill site S1  
 34 = Strawberry Sill site S2  
 35 = Strawberry Sill site S3

AlabamaVault inlet = Alabama canister vault inlet  
 AlabamaVault outlet = Alabama canister vault outlet  
 Brentwood inlet = Brentwood wet pond inlet  
 Brentwood outlet = Brentwood wet pond outlet  
 ParkPlace cell1 = Park Place wet pond cell 1  
 ParkPlace cell2 = Park Place wet pond cell 2

ParkPlace cell3 = Park Place wet pond cell 3  
ParkPlace inlet = Park Place wet pond inlet  
ParkPlace outlet = Park Place wet pond outlet  
Parkstone\_swale inlet = Parkstone grass swale inlet  
Parkstone\_swale outlet = Parkstone grass swale outlet  
Parkstone\_pond inlet = Parkstone wet pond inlet  
Parkstone\_pond outlet = Parkstone wet pond outlet  
SouthCampus inlet = South Campus storm water facility inlet  
SouthCampus outletE = South Campus storm water facility east outlet  
SouthCampus outletW = South Campus storm water facility west outlet  
Sylvan inlet = Sylvan storm drain inlet  
Sylvan outlet = Sylvan storm drain outlet  
Wetland outlet = Grace Lane wetland

CW1 = Smith Creek (see alternate code below)  
CW2 = Silver Beach Creek (see alternate code below)  
CW3 = Park Place drain (see alternate code below)  
CW4 = Blue Canyon Creek (see alternate code below)  
CW5 = Anderson Creek (see alternate code below)  
CW6 = Wildwood Creek (discontinued in 2004)  
CW7 = Austin Creek (see alternate code below)

The following tributary site codes were used for the expanded 2004-2006 tributary monitoring project

AND = Anderson Creek (same location as CW5 above)  
BEA1 = Austin.Beaver.confluence  
AUS = Austin.lower (same location as CW7 above)  
BEA2 = Austin.upper  
BEA3 = Beaver.upper  
BLU = BlueCanyon (same location as CW4 above)  
BRA = Brannian  
CAR = Carpenter  
EUC = Euclid  
MIL = Millwheel  
OLS = Olsen  
PAR = ParkPlace (same location as CW3 above)  
SIL = SilverBeach (same location as CW2 above)  
SMI = Smith (same location as CW1 above)  
WHA = Whatcom

\*\*\*\*\*  
\* VERIFICATION PROCESS FOR THE LAKE WHATCOM DATA FILES

\*\*\*\*\*  
During the summer of 1998 the Institute for Watershed Studies began creating an electronic data file that would contain long term data records for Lake Whatcom. These data were to be included with annual Lake Whatcom monitoring reports. This was the first attempt to make a long-term Lake Whatcom data record available to the public. Because these data had been generated using different quality control plans over the years, a comprehensive re-verification process was done.

The re-verification started with printing a copy of the entire data file and checking 5% of all entries against historic laboratory bench sheets and field notebooks. If an error was found, the entire set of values for that analysis were reviewed for the sampling period containing the error. Corrections were noted in the printed copy and entered into the electronic file; all entries were dated and initialed in the archive copy.

Next, all data were plotted and descriptive statistics (e.g., minimum, maximum) were computed to identify outliers and unusual results. All outliers and unusual data were verified against original bench sheets. A summary of decisions pertaining to these data is presented below. All verification actions were entered into the printed copy, dated, and initialed by the IWS director.

The following is a partial list of the changes made to the verified Lake Whatcom data files. For detailed information refer to the data verification archive files in the IWS library.

Specific Deletions: 1) Rows containing only missing values were deleted. 2) All lab conductivity for February 1993 were deleted for cause: meter inadequate for low conductivity readings (borrowed Huxley's student meter). 3) All Hydrolab conductivity from April - December 1993 were deleted for cause: Hydrolab probe slowly lost sensitivity. Probe was replaced and Hydrolab was reconditioned prior to the February 1994 sampling. 4) All 1993 Hydrolab dissolved oxygen data less than or equal to 2.6 mg/L were deleted for cause: Hydrolab probe lost sensitivity at low oxygen concentrations. Probe was replaced and Hydrolab was reconditioned prior to February 1994 sampling. 5) All srp and tp data were deleted (entered as "missing" in 1989) from the July 10, 1989 wq data due to sample contamination in at least three samples. 6) December 2, 1991, Site 3, 0 m conductivity point deleted due to inconsistency with adjacent points. 7) December 15, 1993, Site 4, 80 m lab conductivity point deleted because matching

field conductivity data are absent and point is inconsistent with all other lab conductivity points. 8) November 4, 1991, Site 2, 17-20 m, conductivity points deleted due to evidence of equipment problems related to depth. 9) February 2, 1990, Site 1, 20 m, soluble reactive phosphate and total phosphorus points deleted due to evidence of sample contamination. 10) August 6, 1990, Site 1, 0 m, soluble reactive phosphate and total phosphorus points deleted due to evidence of sample contamination. 11) October 5, 1992, Site 3, 80 m, all data deleted due to evidence of sample contamination in turbidity, ammonium, and total phosphorus results. 12) August 31, 1992, Site 3, 5 m, soluble reactive phosphate and total phosphorus data deleted due to probable coding error. 13) All total Kjeldahl nitrogen data were removed from the historic record. This was not due to errors with the data but rather on-going confusion over which records contained total persulfate nitrogen and which contained total Kjeldahl nitrogen. The current historic record contains only total persulfate nitrogen. Total Kjeldahl nitrogen data were retained in the IWS data base, but not in the long-term Lake Whatcom data files.

\*\*\*\*\*  
\* ROUTINE DATA VERIFICATION PROCESS

\*\*\*\*\*  
1994-present: The Lake Whatcom data are verified using a four step method: 1) The results are reviewed as they are generated. Outliers are checked for possible analytical or computational errors. This step is completed by the Laboratory Analyst and IWS Laboratory Supervisor. 2) The results are reviewed monthly and sent to the City. Unusual results are identified. This step is completed by the IWS Director. 3) The results are reviewed on an annual basis and discussed in the Lake Whatcom Monitoring Program Final Report. Unusual results are identified, and explained, if possible. This step is completed by the IWS Director, IWS Laboratory Supervisor, and Laboratory Analyst. 4) Single-blind quality control samples, laboratory duplicates, and field duplicates are analyzed as specified in the Lake Whatcom Monitoring Program contract and in the IWS Laboratory Certification requirements. Unusual results that suggest instrumentation or analytical problems are reported to the IWS Director and City. The results from these analyses are summarized in the annual report.

1987-1993: The lake data were reviewed as above except that the IWS Director's responsibilities were delegated to the Principle Investigator in charge of the lake monitoring contract (Dr. Robin Matthews).



Prior to 1987: Data were informally reviewed by the Laboratory Analyst and IWS Director. Laboratory and field duplicates were commonly included as part of the analysis process, but no formal (i.e., written) quality control program was in place. Laboratory logs were maintained for most analyses, so it is possible to verify data against original analytical results. It is also possible to review laboratory quality control results for some analyses.