



Summer 1978

Coal Petrology of the Tulameen Coalfield, South Central British Columbia

V. Eileen Williams

Western Washington University

Follow this and additional works at: <https://cedar.wwu.edu/wwuet>



Part of the [Geology Commons](#)

Recommended Citation

Williams, V. Eileen, "Coal Petrology of the Tulameen Coalfield, South Central British Columbia" (1978). *WWU Graduate School Collection*. 810.

<https://cedar.wwu.edu/wwuet/810>

This Masters Thesis is brought to you for free and open access by the WWU Graduate and Undergraduate Scholarship at Western CEDAR. It has been accepted for inclusion in WWU Graduate School Collection by an authorized administrator of Western CEDAR. For more information, please contact westerncedar@wwu.edu.

COAL PETROLOGY
OF THE
TULAMEEN COALFIELD,
SOUTH CENTRAL BRITISH COLUMBIA

A Thesis
Presented to
The Faculty of
Western Washington University

In Partial Fulfillment
Of the Requirements for the Degree
Master of Science

by
V. Eileen Williams
August 1978

COAL PETROLOGY
OF THE
TULAMEEN COALFIELD,
SOUTH CENTRAL BRITISH COLUMBIA

by

V. Eileen Williams

Accepted in Partial Completion
Of the Requirements for the Degree
Master of Science

Dean of Graduate School

Advisory Committee

Chairperson

[Handwritten signature]

[Handwritten signature]

MASTER'S THESIS

In presenting this thesis in partial fulfillment of the requirements for a master's degree at Western Washington University, I grant to Western Washington University the non-exclusive royalty-free right to archive, reproduce, distribute, and display the thesis in any and all forms, including electronic format, via any digital library mechanisms maintained by WWU.

I represent and warrant this is my original work and does not infringe or violate any rights of others. I warrant that I have obtained written permissions from the owner of any third party copyrighted material included in these files.

I acknowledge that I retain ownership rights to the copyright of this work, including but not limited to the right to use all or part of this work in future works, such as articles or books.

Library users are granted permission for individual, research and non-commercial reproduction of this work for educational purposes only. Any further digital posting of this document requires specific permission from the author.

Any copying or publication of this thesis for commercial purposes, or for financial gain, is not allowed without my written permission.

Name: Vera Eileen Williams

Signature: 

Date: 4 June 2018

ABSTRACT

The Tulameen Coalfield lies in a small, southeast plunging syncline in the Intermontane Belt of the Canadian Cordillera in south central British Columbia. The coal is interbedded with fluvial and lacustrine sediments of the Eocene Allenby Formation of the Princeton Group. The abundance of tephra and bentonite indicates volcanic activity was contemporaneous with coal and sediment deposition.

Although coal occurs on both sides of the basin, its economic importance is currently restricted to a 15-21 meter thick coal seam on the southwestern limb of the syncline. The coal is predominantly vitrain and clarain composed of greater than 90 percent vitrinite. Clean coal has 14 percent ash (mostly kaolinite and quartz) and low sulphur. The coal is interbedded with bentonite, mudstone and shale partings which increase in number from south to north. The partings are montmorillonite and/or kaolinite rich.

Coal rank along the southwestern limb of the basin was determined by measuring the percent maximum reflectance of vitrinite. Vitrinite reflectance increases laterally from .62 in the north to .86 in the south but shows no significant vertical variation. According to the A.S.T.M. classification, coal rank ranges from High Volatile C Bituminous to High Volatile B Bituminous. Field relations indicate partial post-deformation coalification.

Petrographic study of the coal suggests the coal-forming peat developed in a predominantly forest moor swamp environment.

TABLE OF CONTENTS

	Page
ABSTRACT	i
TABLE OF CONTENTS	ii
LIST OF FIGURES	iii
LIST OF TABLES	vi
INTRODUCTION	1
ACKNOWLEDGMENTS	3
PREVIOUS WORK	4
GEOLOGICAL SETTING	6
METHODS OF INVESTIGATION	13
STRATIGRAPHY	17
COAL COMPOSITION	
Macerals	27
Microlithotypes	37
Proximate and Ultimate Analyses	37
Mineral Matter	40
COAL RANK	46
DISCUSSION	51
CONCLUSIONS	62
REFERENCES CITED	63
APPENDIX OF MEASURED SECTIONS AND SAMPLE LOCATIONS	68

LIST OF FIGURES

Figure	Page
1. Location of the Tulameen Coalfield and other major coalfields in British Columbia	2
2. Generalized stratigraphic section of the Princeton Group	7
3. Surficial expression of major coal-bearing formations in the Tulameen, Princeton, Merritt and Hat Creek Coalfields	10
4. Geologic Map Tulameen basin	11
5. Map showing location of sample and section sites	14
6. Representative stratigraphic column of the Coal-Bearing Member	18
7. South wall of TR 2 showing bright, bedded appearance of coal and thin bentonite, shale and mudstone partings	19
8. Exposure of main coal seam at Blakeburn open pit (OP-1)	21
9. Stratigraphic section from TR 4 to T-77-4	22
10. North-south stratigraphic columns showing thickness and lithology of the main coal seam	23
11. South wall of TR 4 showing main bentonite bed and other partings in the upper half of the main coal seam	26
12. Undifferentiated vitrinite lacking visible cell structure	29
13. Cell structure in well differentiated vitrinite	29
14. Structured vitrinite divided into telinite and collinite	30
15. Deformed piece of exinite	30

LIST OF FIGURES (CONTINUED)

Figure	Page
16. Elongated resin bodies; thin-walled telinite with resinite fillings; light colored resinite bodies altered to vitrinite; finely disseminated pyrite	32
17. Folded, thick-toothed cutinite	32
18. Thin-walled cutinite in clarite with sporinite, vitrinite and inertinite	33
19. Compressed sporinite in clay-rich clarite	33
20. Sporinite in clarite with cutinite and inertinite	34
21. Isolated multicelled sclerotia with resinite filled cavities	34
22. Concentrated layers of compressed teleutospores in fractured vitrinite	35
23. Deformed semifusinite in dirty vitrinite	35
24. Plastically deformed degradofusinite with poorly preserved cellular structure	36
25. Pyrofusinite with well preserved cellular structure in trimacerite	36
26. Inertinite in clay-rich vitrinite	38
27. Trimacerite with sclerotinite, semifusinite, resinite, cutinite and vitrinite	38
28. Clarite with exinite and vitrinite	39
29. Diffuse clay layers of variable width in vitrinite	39
30. Well defined clay layers and lenses interbedded with vitrinite	42
31. Spherical aggregates of euhedral pyrite crystals; irregular masses of pyrite crystals; pyrite replacing cell walls	42

LIST OF FIGURES (CONTINUED)

Figure	Page
32. Syngenetic pyrite replacing well defined cell walls in vitrinite	44
33. Syngenetic pyrite replacing cell walls and filling cell lumens in vitrinite	44
34. Detrital, sub-angular quartz grain in clay- and pyrite-rich vitrinite	45
35. Carbonate mineral associated with vitrinite and exinite	45
36. Graph showing gradual increase in average vitrinite reflectance from north to south	48
37. Comparison of ASTM system of rank classification with % reflectance of vitrinite, B.T.U. and fixed carbon content of coal	50
38. A. Cordilleran arc-trench system in Paleocene to Late Eocene B. Cordilleran arc-trench system in Late Eocene to Early Miocene	52
39. Principal structural subdivisions of the southern Canadian Cordillera	53
40. Schematic block diagram of the main seam peat formation in the Tulameen forest moor swamp environment	56
41. Diagram showing possible configuration between coalification planes and coal seams in relation to deformation	60

LIST OF TABLES

Table	Page
1. Mineral composition of partings and roof and floor samples	24
2. Results of maceral analyses on selected samples	28
3. Plot of % \bar{R}_0 of vitrinite at designated sites (values arranged in stratigraphic order)	47

INTRODUCTION

The Tulameen Coalfield is one of a number of remnant Tertiary structural basins located in south central British Columbia. It lies just east of the Cascade Mountains approximately 50 km north of the United States-Canadian International Boundary. The location of the Tulameen and other coalfields in British Columbia is shown in Figure 1.

Coal along the southwestern portion of the basin was mined from 1919 to 1940 using underground methods and in 1954 using open cast methods. Although 2,364,561 tons of coal were extracted (Dolmage Campbell & Assoc. Ltd, 1975), potential coal resources are still considerable. Currently an extensive study on the economic feasibility of strip mining the main coal seam along the southwest periphery is in progress. Most of the analytical work to date on the coal has been related to economic use. With the exception of Donaldson (1973) little petrographic work has been done on the coal.

The purpose of the present study is to examine the coal petrology of the main seam along the southwest limb of the Tulameen basin. The stratigraphy, petrographic composition and rank of the coal are examined in order to provide a basis for interpreting the coalification history and the environmental conditions that prevailed during coal formation.




Figure 1. Location of the Tulameen Coalfield and other major coalfields in British Columbia.

ACKNOWLEDGMENTS

I would like to thank and express my appreciation to Dr. Charles A. Ross for suggesting the thesis topic and, along with Drs. Christopher A. Suczek and David A. Pevear, for critically reading the manuscript and offering helpful suggestions. I am grateful to Dr. Dave Pearson, Head of the Coal Division, British Columbia Ministry of Mines and Petroleum Resources, and his assistant, Mr. Dave Grieve, for their patient instruction on the use of equipment and also for their extensive help and aid in discussing and providing information on coal petrology, laboratory procedures and interpretations of data. I thank the British Columbia Ministry of Mines and Petroleum Resources for allowing me to use their equipment for analyzing the coal. I am also thankful to the Cyprus Anvil Mining Corporation and, especially, their exploration geologist, Mr. Tom Adamson, for supplying invaluable information on the Tulameen coal as well as for helping me in the field area each day. Dr. Harold Gluskoter and Research Assistants, Mr. Philip Bowden and Ms. Faith Fiene, of the Illinois State Geological Survey Coal Section kindly low-temperature ashed a number of coal samples for x-ray analyses. I also thank Ms. Sue Kinder and Dr. David Pevear for helping me run and interpret the clay mineralogy of non-coal and ashed coal samples. I thank Mr. George Mustoe for helping me prepare and develop photographs. I would also like to express a special thanks to Dr. June R.P. Ross for initially encouraging and suggesting my going on to graduate school. For their continued support, encouragement and understanding, I am especially grateful to Cindy, Leah, the Sadle and Williams families, Arthur, Jack and Dutchy.

PREVIOUS WORK

The general geology and mineral deposits of the Tulameen area were first described by Charles Camsell (1913) in a series of Canadian government publications from 1906 to 1913. He correlated the coal-bearing sediments of the Tulameen basin with similar Tertiary coal-bearing strata in the Princeton and Hat Creek Coalfields. H.M.A. Rice (1947) published an extensive work which includes a geologic map on the geology and mineral deposits of the Princeton map area.

W.S. Shaw (1952) wrote the first comprehensive report on the Tulameen Coalfield. This publication includes a geologic map of the Tulameen basin as well as a series of stratigraphic columns based on drill core data from the northern limb of the syncline.

The coal petrography of the Blakeburn open pit mine was examined by J. Roger Donaldson (1973). Donaldson's work includes a detailed microlithotype and maceral analysis of a $5\frac{1}{2}$ foot (1.7 meter) column sample as well as some rank determinations based on vitrinite reflectance. In the 1975 publication, Coal Resources of British Columbia, Dolmage Campbell & Associates, Ltd. (1975) describe the coal deposits and mining history of the Tulameen basin. File data from some of the mining companies who have worked in the area is included.

A paper on the Tulameen coal basin by Steve H. Evans (1977) was published in the British Columbia Ministry of Mines and Petroleum Resources report on 1977 geologic fieldwork in British Columbia. Evans' revised field map of the basin reveals the existence of a number of previously unmapped faults.

The most extensive and detailed geologic work to date on the southern limb of the coalfield is found in the 1978 Tulameen Coal Project Report by Tom Adamson of the Cyprus Anvil Mining Corporation. Detailed data contained in his report are included in this thesis.

GEOLOGICAL SETTING

The Tulameen basin rests unconformably on the Upper Triassic Nicola Group. Nicola rocks cover a wide area and are composed of vari-colored lava, argillite, tuff, limestone and chlorite/sericite schist. During the Jurassic and Cretaceous, the Nicola was intruded by a series of intrusions ranging in composition from gabbro to granite (Rice, 1947; Adamson, 1978).

Unconformably overlying the Upper Triassic volcanics and Cretaceous granitic intrusives is the Princeton Group (Figure 2). The two main outcrop areas are in the Tulameen and Princeton basins. The Princeton Group is early Tertiary (Eocene) and consists of an unnamed basal volcanic unit unconformably overlain by a thick sequence of clastic sedimentary rocks (Allenby Formation) which in turn are unconformably overlain by an unnamed upper volcanic unit (Eocene-Oligocene?). The basal volcanics consist of a thick section of massive and banded andesitic and basaltic lavas. The overlying Allenby sediments grade upwards from coarse-grained conglomerates to fine-grained shales. The conglomerates have an abundance of granitic cobbles. Feldspar and volcaniclastic fragments are characteristic constituents in the sandstone. The shales contain mostly clay with some quartz and feldspar (Rice, 1947). Coal horizons (some of mineable thickness) and bentonite beds are found interbedded in the sediments. The coal-bearing strata in the Tulameen basin has the same stratigraphic relations as in the Princeton basin and is most likely correlative with the Allenby Formation in the Princeton basin and, therefore, is here extended into the Tulameen basin.




Figure 2. Generalized stratigraphic section (not to scale) of the Princeton Group in the Tulameen and Princeton basins.

The coal-bearing Allenby Formation forms a sequence more than 800 meters thick in the Tulameen basin. The basal part is 100-150 meters thick and consists of a coarse- to fine-grained sandstone interbedded with siltstone, shale and tephra. Several thin andesitic flows are interbedded with the sediments at the base. The succeeding 130 meter thick section contains shale and minor sandstone interbedded with several coal horizons. The coal is intercalated with numerous shale and bentonite partings. The upper part of the Allenby Formation is over 600 meters thick and is composed of predominantly coarse sandstone and conglomerate. Tuffaceous material is freely dispersed throughout the sediments (Shaw, 1952).

In the southeastern portion of the basin, the Allenby Formation overlaps the basal Princeton Group volcanics and rests unconformably on top of the Nicola Group. The upper volcanic unit of the Princeton Group is absent from the Tulameen basin and, instead, flat-lying Miocene-Pliocene aged volcanics (flood basalts) unconformably cap the sediments. Recent unconsolidated glacial deposits up to 13 meters thick mantle much of the area (Rice, 1947; Shaw, 1952).

In south central British Columbia a number of remnant Tertiary basins are characterized by thick sequences of sedimentary rocks of freshwater origin (Rice, 1947; Cockfield, 1948). The sediments, mostly Eocene, are fluvial and lacustrine and are commonly intercalated with coal and/or carbonaceous shale layers of varying thicknesses. Sediment deposition in the Princeton basin was controlled by block faulting. Paleocurrent data from the sediments indicate a southerly flowing drainage system (Okulitch, et. al., 1977). Major coal deposits

are located in the Tulameen and Princeton Coalfields as well as the Hat Creek and Merritt Coalfields (Figure 3). In the Merritt and Hat Creek Coalfields the coal-bearing sediments are assigned to the Coldwater Formation which consists mainly of conglomerate, sandstone and shale. At Hat Creek the Coldwater lies between two Tertiary volcanic units and at Merritt the Coldwater is part of the Kamloops Group. Though discontinuous, the coal-bearing sediments from the four coalfields probably are related, however, no direct correlation between them has been established. Shaw (1952) suggested that a careful reconstruction of Tertiary drainage patterns in the area would help determine whether or not they are correlative.

The Tulameen Coalfield lies in an oval northwest-southeast trending structural basin 5 km long and 3 km wide. Figure 4 shows a geologic map of the Tulameen basin. The basin is an asymmetric fold, with beds on the northeastern side dipping from 40° to 65° to the southwest and beds on the southwestern side dipping from 25° to 49° to the northeast. The fold axis plunges towards the southeast. Shaw (1952) reported two large fault zones and numerous smaller ones in the southern part of the basin and Evans (1977) mapped two probable faults which parallel the trend and run the length of the fold. Folding took place in the interval of time represented between the top of the Princeton Group and the base of the Miocene-Pliocene "flood" basalts.

Although coal crops out on both sides of the syncline, correlation between seams has not been established (Shaw, 1952). Drilling exploration by mining companies along the northeastern limb disclosed a number of irregularly spaced, steeply dipping coal seams characterized




Figure 3. Surficial expression of major coal-bearing formations in the Tulameen, Princeton, Merritt and Hat Creek Coalfields (modified from Dept. of Mines & Petroleum Resources, British Columbia, 1976).




Figure 4. Geologic Map Tulameen basin (modified from Shaw, 1952 and Adamson, 1978).

by highly crushed coal. On the other side of the basin, the coal has shallower dips and is considerably more coherent. Three underground mines followed the main coal seam down dip for over 300 meters and along strike for 2300 meters.

METHODS OF INVESTIGATION

Field investigations were carried out along the southwestern limb of the Tulameen basin during the summer of 1977. The field area encompasses that portion of the coalfield which may be economically feasible to mine. The coal was followed along strike from the Blakeburn open pit in the south of the basin to the extreme northern edge of the basin. Figure 5 shows the configuration of the field area and the location of section and sample sites.

There are a total of 19 sites: the Blakeburn open pit, 6 trenches spaced approximately 300 meters apart and 12 diamond drill holes. The Blakeburn open pit (OP-1) represents the extreme southern border of the field area. The dip of the beds is 25° to 30° to the east. The bottom of the coal zone is not exposed. The trenches (TR 1 through TR 6) are located in the center of the field area where dips range from 28° to 45°. In each trench the entire thickness of the main coal seam, which shows varying degrees of weathering, is exposed. The 12 diamond drill holes were drilled under the direction of the Cyprus Anvil Mining Corporation and are numbered T-77-1 through T-77-12. T-77-1 through T-77-5 were drilled perpendicular to the dip and T-77-6 through T-77-12 were drilled vertically. T-77-10 represents the extreme northern edge of the field area.

Detailed stratigraphic sections of the main coal seam in OP-1 and TR 1 through TR 6 were measured and described (Appendix). Detailed lithologic logs of all of the drill holes were provided by Cyprus Anvil. Over 200 samples of coal and non-coaly interbeds were collected




Figure 5. Map showing location of sample and section sites.

from OP-1, TR 1 through TR 6 and T-77-8 through T-77-12. Sampling intervals varied between sites. For the location of each sample see the Appendix.

The organic composition of the coal was examined in reflected light under oil immersion. Twenty-three maceral analyses were made. The percentage of vitrinite, exinite and inertinite was measured by counting 1000 points per sample.

In order to help determine the inorganic mineral composition of the coal, 7 samples ground to pass a 60 mesh screen were oxidized at low temperature (<150°C) in an electronically excited oxygen plasma chamber. One sample was ashed at the Coal Division of the British Columbia Ministry of Mines and Petroleum Resources and the others by the Coal Division of the Illinois State Geological Survey. The ashed products were analyzed on a GE XRD-5 x-ray diffractometer using Cu k radiation. Whole rock and oriented specimens of the 2 m fraction of 12 clay and shale partings from TR 4 were also x-rayed. Whole coal analyses on drill cores and bulk samples were provided by Cyprus Anvil.

The rank of 49 coal samples was determined by measuring the maximum percent reflectance (\bar{R}_0) of vitrinite. Vitrinite reflectance is an excellent parameter to use for determining rank because it is not appreciably affected by weathering. Therefore, reliable \bar{R}_0 values were obtained for the weathered coals in the trench and open pit sites as well as for the relatively fresher drill core samples.

Preparation for rank and maceral analyses involved making the samples into pellets. Each coal sample was crushed and passed through a 20 mesh screen. It was then thoroughly mixed with an equal amount of Buehler Transoptic powder and placed into a Buehler, Ltd., PNEUMAT I

mounting press where it was subjected to a pressure of 4500 psi and heated to 140°C. After cooling, the pellet was polished using a five-stage wet polishing technique going from a coarse-grained 240-grit Carbimont paper disc down to a .05 alumina Buehler micropolish. Pellets were allowed to dry for at least 24 hours because moisture influences reflectance values, especially in high volatile coals.

Reflectance was measured using a Leitz 450 photometer at a magnification of x500. Readings were taken on homogeneously appearing vitrinite particles that no longer displayed cellular structure. The vitrinite was viewed in incident light through an oil immersion objective. The stage of the microscope was rotated through 360 and the highest reflectance value was recorded. Fifty readings per sample were taken. Control was maintained by using a glass standard of known reflectance for calibration. The standard was checked after every 25 readings.

STRATIGRAPHY

The coal-bearing middle member of the Allenby Formation in the Tulameen basin is approximately 130 meters thick and consists of interbedded shales, mudstone, tuffs, minor sandstone and coal (Figure 6). Two well defined coal seams are present (Adamson, 1978). The lower seam has an average thickness of 7 to 7.6 meters and can be traced from T-77-11 to T-77-10. It is medium clean to clean banded coal with interbedded bentonite, shale and mudstone. The average percentage of inorganic material intermixed with the coal is high (50-70%).

The upper coal seam has interbeds of bentonite, brown shale and mudstone. The coal is banded, fairly bright bedded coal (Figure 7) with vitrain and clarain predominating. The truly dull coal component consists mainly of durain. Microscopic examination of some of the dull coal samples reveals a large amount of vitrinite and the presence of finely dispersed mineral matter which masks the macroscopic appearance of the coal. This observation also was made by Donaldson (1973). Weathered coal surfaces commonly have a white oxidized coating and iron oxide stain. Clear, amber-colored resin nodules and infrequent, partially coalified wood fragments are scattered throughout the coal. A few small scour channels filled with coal are present. Both calcite and quartz have been identified by x-ray analysis as cleat fillings. In places, the coal exhibits small scale folding and faulting. Finely crushed coal resembling fusain lies along the fault planes. The coal is vitrinite rich throughout the lateral and vertical extent of the coal seam and rank increases slightly from northwest to southeast. The average thickness of the coal in the trench and




Figure 6. Representative stratigraphic column of the Coal-Bearing Member (approx. 130 meters thick).




Figure 7. South wall of TR 2 showing bright, bedded appearance of coal and thin bentonite, shale and mudstone partings. Photograph taken near the base of the coal section.

drill hole area of the field ranges from 15 to 21 meters. In the Blakeburn open pit (Figure 8) the true thickness of the seam is unknown because the floor of the seam is not exposed. Also, determination of the thickness has been complicated by numerous small faults many of which are subparallel to bedding.

Figure 9 shows a cross section drawn down dip from TR 4 to T-77-4 and Figure 10 shows a series of stratigraphic columns that extend laterally from the extreme northern portion of the field area at T-77-10 to the Blakeburn open pit in the southern portion. The measured sections are based upon the author's field measurements of the Blakeburn open pit and surface exposures in Trenches 1 through 6 and upon detailed lithologic logs from T-77-1 through T-77-12.

The percentage and lithologic character of partings change towards the northern portion of the basin. In the Blakeburn open pit, partings are the least numerous and consist mainly of relatively thin layers of bentonite and mudstone. In the interval between T-77-1 and T-77-5, the coal seam changes from 88 to 82% coal and from 12 to 18% partings. The partings are characterized by bentonite, mudstone and brown shale. Several thin scattered interbeds of a dark, chert-like material are almost pure SiO_2 according to x-ray analysis.

Fifteen partings from TR 4 and roof and floor samples were examined by x-ray diffraction. Table 1 lists their location, physical appearance and mineralogy. A montmorillonitic smectite and "b" axis disordered kaolinite are the two predominant minerals and both are distributed throughout the stratigraphic column. The smectite in bentonite partings consists of almost pure, fully expandable montmoril-



Figure 8. Exposure of main coal seam at Blakeburn open pit (OP-1).






Figure 9. Stratigraphic section from TR 4 to 7-77-4.

TABLE 1. MINERAL COMPOSITION OF PARTINGS AND ROOF AND FLOOR SAMPLES

Sample #	Macroscopic Appearance	Mineralogy (in size fractions)	
		< 2 μ m	> 50 μ m
4-70-3	Cream colored B	Almost pure M; minor C	N.D.
4-70-2	Brown coaly Ms	Q and F	N.D.
4-70-1	Cream colored B; Fe stain	Almost pure M	N.D.
4-60-1	Dark brown coaly Ms	Mostly S; K	N.D.
4-45-1	Med. brown colored Ms	S; minor halloysite?	Q,F
4-41-3	Med. buff colored B; Fe stain	Almost pure M; minor C	N.D.
4-34-1	Light brown to buff colored Ms; Fe stained; carbonized plant fragments	K; minor S	N.D.
4-26-1	Med. brown colored Ms; Fe stain; carbonized plant fragments	Almost pure M; minor C; trace Q	N.D.
4-20-1	Brown shaly Ms	S and K	Q
4-18-1	Brown Ms; Fe stain	S	N.D.
4-14-1	Cream to buff colored Ms	K; minor S	Q
4-9-1	Buff colored Ms; Fe stain	S; lesser K	Q
6-63-1	Black cherty rock found interbedded in coal from TR 6	Pure Q	N.D.
OP-1-15-1	Brown fissile Sh; overlies bentonite and main coal seam; from OP-1	Mostly K and Q; minor S	N.D.
1-0-0	Light colored, blocky, laminated Ms; lies at base of coal; from TR 1	Mostly K and Q; expand- able mixed layer clay of smectite group	N.D.

*Samples numbered from 4-9-1 through 4-70-3 are arranged in stratigraphic order and come from TR 4; see Appendix for location of samples.

**B = bentonite; C = cristobalite; F = feldspar; K = kaolinite; M = montmorillonite; Ms = mudstone; Q = quartz; S = smectite; Sh = shale.

lonite show fairly sharp, high intensity peaks indicating good crystallinity (Roberson, 1964). Admixtures of kaolinite and smectite in varying proportions are common in the dark brown mudstones. Some of the partings contain small carbonized plant fragments. All of the kaolinite-bearing partings contain quartz. Kaolinite-rich sediments with quartz and minor amounts of smectite occur both above and below the coal seam.

Between T-77-6 and T-77-10 partings become markedly more numerous. The percentage of coal decreases from south to north from 71% at T-77-6 to 41% at T-77-10. Most of the partings have a volcanic origin and are characterized by tuffs (multi-colored and welded), fine-grained siliceous (sometimes pyritic) rock, very-fine-grained and slightly vesicular flows and bentonitic mudstone. Contacts between the coal and the volcanics are mainly sharp and well-defined (Adamson, 1978).

Occurring near the top of the coal section is a continuous bentonite bed up to 1 meter thick. The bentonite is light gray to cream colored (Figure 11). X-ray diffraction of the $< 2\mu\text{m}$ size fraction shows montmorillonitic smectite to be the predominant clay mineral. Small carbonized plant fragments are dispersed throughout the bed. The bentonite can be traced from the Blakeburn open pit northwards to TR 6. The bed does not appear in the T-77-9, T-77-7 and T-77-8 lithologic logs. However, the T-77-10 lithologic log describes a 76 cm thick bentonite bed which may belong to the key unit. Other partings show little correlation between sites.




Figure 11. South wall of TR 4 showing main bentonite bed (thick, light colored bed in center of photograph) and other partings in the upper half of the main coal seam. The coal is weathered and shows iron staining.

COAL COMPOSITION

Macerals

The microscopic organic constituents of coal are called macerals. There are three maceral groups: vitrinite, exinite and inertinite. Table 2 shows the percentage maceral composition of 23 main seam samples (Appendix) covering the lateral and vertical extent of the field area. The results show a uniformly large percentage of vitrinite and relatively small amounts of exinite and inertinite. Exinite is slightly more abundant than inertinite. Donaldson's (1973) maceral analysis also shows a correspondingly large amount of vitrinite. The abundance of the vitrinite imparts the characteristic macroscopic bright appearance of the coal.

Much of the vitrinite is formed from the woody bark and tissue of trees and shrubs. The vitrinite is mostly undifferentiated and has a uniform "homogeneous" appearance (Figure 12). Although the color of vitrinite in a sample usually varies from medium to light gray, samples from T-77-8 and T-77-10 are generally darker and samples from OP-1 lighter. Some samples contain vitrinite differentiated into telinite and collinite (Figure 13). Most differentiated vitrinite fragments show deformation by folding or compression into microlayers parallel to bedding (Figure 14). The brittleness of vitrinite is shown by numerous microfractures in some samples. Exinite and inertinite macerals and mineral matter are mostly enclosed within vitrinite.

Exinite generally has a black to dark gray appearance. Some of the exinite is difficult to identify (Figure 15). Most samples include minor amounts of resinite, cutinite and sporinite. The resinite often

TABLE 2. RESULTS OF MACERAL ANALYSES ON SELECTED SAMPLES

Sample Site*	Sample No.**	% Vitrinite	Maceral Composition	
			% Exinite	% Inertinite
OP-1	OP-1-43-1	95.0	3.9	1.1
OP-1	OP-1-15-1	95.0	3.2	1.0
OP-1	OP-1-11-1	92.8	6.9	0.8
OP-1	OP-1-1-3	96.1	2.8	1.1
TR 1	1-1-1	99.8	0.2	-
T-77-11	DC 11-Q***	97.4	2.0	0.6
TR 3	3-9-2	93.8	0.7	0.9
T-77-12	DC 12-ss	97.2	2.7	0.1
T-77-12	DC 12-pp	98.8	2.2	1.0
T-77-12	DC 12-hh	98.2	1.4	0.4
T-77-12	DC 12-bb	97.8	1.3	0.9
T-77-12	DC 12-y	97.4	2.0	0.6
T-77-12	DC 12-s	96.1	3.8	0.1
T-77-12	DC 12-j	98.7	0.8	0.6
T-77-12	DC 12-a	98.4	0.7	0.9
TR 5	5-81-1	97.7	1.6	0.7
T-77-9	DC 9-B***	99.2	0.5	0.3
T-77-8	DC 8-E***	94.8	4.9	0.3
T-77-8	DC 8-D***	98.2	1.5	0.3
T-77-8	DC 8-C	98.1	1.3	0.6
T-77-8	DC 8-B	97.6	2.1	0.3
T-77-8	DC 8-A***	99.1	0.5	0.4
T-77-10	DC 10-X	99.4	0.4	0.2

*See Figure 5 for location of sample sites.

**See Appendix for location of samples

***Pyrite-rich samples




Figure 12. Undifferentiated vitrinite (1) lacking visible cell structure. Polished surface, oil immersion, x500.




Figure 13. Cell structure in well differentiated vitrinite; telinite (1) represents cell wall tissue and collinite (2) cell fill material. Polished surface, oil immersion, x500.



Figure 14. Structured vitrinite divided into telinite (1) and collinite (2); cell structure highly compressed in lower half of figure. Polished surface, oil immersion, x500.




Figure 15. Deformed piece of exinite (1)--resinite? (2) filling thin-walled telinite? (3)--surrounded by vitrinite (4). Polished surface, oil immersion, x500.

displays orange internal reflections. Most of the resinite (Figure 16) occurs as isolated elongated or spherical shaped bodies and in some cases as cell lumen fillings. Isolated resinite bodies not altered to vitrinite are seen in a few samples. Thin- and thick-walled cutinite commonly show folded and sometimes toothed structure (Figure 17 and 18). Selective decomposition has formed some concentrated layers of thin-walled untoothed cutinite. Sporinite, sometimes occurring in groups, is present in small quantities in most of the samples. It is formed from the resistant perines and exines of pollens and spores. The sporinite is compressed and in some cases is filled with vitrinite (Figure 19). Microscopic cracks which pass through vitrinite often stop abruptly at sporinite which is very tough and resistant (Figure 20).

Inertinite has a light gray to bright white color and a correspondingly high reflectance. Sclerotinite and semifusinite are the two most abundant inertinite macerals in the coal. Though not abundant, sclerotinite occurs in most samples. The sclerotia (Figure 21) are generally rounded or elliptical and some are multicellular. Cavities are usually empty but some show resinite, vitrinite or pyrite fillings. Concentrated clusters of compressed teleutospores in vitrinite (Figure 22) occur, especially in samples from OP-1. Little semifusinite and rare fusinite are found in trench and drill core samples, but are more common in OP-1 samples. Semifusinite has poorly preserved structure, shows varying degrees of fusinization and is often deformed (Figure 23). Plastically deformed degradofusinite (Figure 24) and pyrofusinite with well preserved cell structure (Figure 25) appear in several samples from OP-1. One piece of macrinite? was identified in




Figure 16. Elongated resin bodies (1) in vitrinite; thin-walled telinite (2) with resinite fillings (3); light colored resinite bodies altered into vitrinite (4); finely disseminated pyrite (5) in vitrinite. Polished surface, oil immersion, x500.




Figure 17. Folded, thick-toothed cutinite (1) in vitrinite; rounded light gray objects represent vitrinitized cell fillings (2). Polished surface, oil immersion, x500.




Figure 18. Thin-walled cutinite (1) in clarite with sporinite (2), vitrinite and inertinite (3). Polished surface, oil immersion, x500.




Figure 19. Compressed sporinite (1) in clay-rich (2) clarite; microspores filled with vitrinite (3). Polished surface, oil immersion, x500.




Figure 20. Sporinite (1) in clarite with cutinite (2) and inertinite (3); toughness of sporinite shown in top of figure where microscopic crack (4) passes through brittle vitrinite and stops at sporinite (5). Polished surface, oil immersion, x500.




Figure 21. Isolated multicelled sclerotia (1) with resinite (2) filled cavities; vitrinite groundmass with vitrinitized cell fillings (3) surrounded by clay (4). Polished surface, oil immersion, x500.




Figure 22. Concentrated layers of compressed teleutospores (1) in fractured vitrinite (2). Polished surface, oil immersion, x500.




Figure 23. Deformed semifusinite (1) in dirty vitrinite; finely disseminated pyrite crystals (2) in vitrinite; aggregate of pyrite crystals (3). Polished surface, oil immersion, x500.




Figure 24. Plastically deformed degradofusinite (1) with poorly preserved cellular structure; thin-walled toothed cutinite (2) and resin bodies (3). Polished surface, oil immersion, x500.



Figure 25. Pyrofusinite (1) with well preserved cellular structure in trimacerite. Polished surface, oil immersion, x500.

in a sample from OP-1 (Figure 26).

Microlithotypes

Characteristic associations of macerals are called microlithotypes. The dominant microlithotype of the main coal seam is vitrinite. Clarite with minor interbands of trimacerite (clarodurite) and durite are also present. Donaldson's (1973) microlithotype analysis shows vitrinite to represent 72.5% and clarite 10.8% of the 5½ foot column that he studied. Figure 27 shows vitrinite, flattened sporinite, sclerotinite and semifusinite in trimacerite (duroclarite). Vitrinite, cutinite and <5% semifusinite are shown in Figure 28 in clarite.

Proximate and Ultimate Analyses

Bulk and drill core analyses (Adamson, 1978) show clean coal has the following specifications:

Proximate Analysis (As Received)

Total Moisture	12.0%
Ash	14.5
Volatile Matter	30.2
Fixed Carbon	<u>43.3</u>
	100.0%
Sulphur	00.6%
Calorific Value	10,000 B.T.U./lb

Ultimate Analysis (As Received)

Ash	13.80%
Carbon	62.73
Hydrogen	4.90
Nitrogen	1.36
Sulphur	0.63
Oxygen	<u>16.57</u>
	99.99%




Figure 26. Inertinite--macrinite? (1)--in clay-rich (2) vitrinite (3). Polished surface, oil immersion, x500.




Figure 27. Trimacerite with sclerotinite (1), semifusinite (2), resinite (3), cutinite (4) and vitrinite (5); the large resilient sclerotinite resists compression. Polished surface, oil immersion, x500.




Figure 28. Clarite with exinite (1) and vitrinite (2); clarite interlayered with vitrinite (3). Polished surface, oil immersion, x500.



Figure 29. Diffuse clay layers (1) of variable width in vitrinite (2). Polished surface, oil immersion, x500.

The ultimate analysis, which determines the major component elements of coal, shows a relatively large amount of hydrogen and small amount of oxygen because of the large vitrinite content.

Mineral Matter

Mineral matter refers to all inorganic material in coal. It includes discrete mineral phases and all inorganic elements other than H, N, O and S which may be present in organic compounds (Ward, 1977). Below are results of the mineral ash analysis of clean coal from bulk and drill core samples (Adamson, 1978):

Mineral Ash Analysis

SiO ₂	69.51%
Al ₂ O ₃	13.54
Fe ₂ O ₃	6.55
CaO	1.17
MgO	0.44
Na ₂ O	0.67
K ₂ O	0.64
TiO ₂	0.04
P ₂ O ₅	0.17
SO ₃	0.51
	<hr/> 93.24%

During mineral ash analysis, coal is heated to high enough temperatures so that all of the inorganic constituents are converted to oxides (Sehgal and Wong, 1974). The silica to iron ratio in the main seam coal is large and the alkali content small. Amounts of sulphur, phosphorous and titanium are significantly small.

Mineral ash differs from low temperature ash (LTA) in that low temperature ashing removes organic matter by slow oxidation at low enough temperatures so that the resultant inorganic residue is essentially unaltered. Seven coal samples were low temperature ashed and then analyzed on the x-ray diffractometer. Three samples are from the lower, middle and upper portions of the exposed coal section in OP-1 and a fourth from the base of the main seam in TR 2 (Appendix). All four samples were weathered. The remaining three samples are relatively fresh and come from the lower, middle and upper portions of the main coal seam in T-77-10 (Appendix). Samples have quartz, kaolinite and minor amounts of an expandable clay (smectite?). Quartz predominates. Results from OP-1 show a larger concentration of quartz in the upper part of the seam and a greater kaolinite to quartz ratio in the lower and middle parts. The absence of identifiable sulphides and carbonates from OP-1 is, in part, the result of weathering. In addition to showing the presence of quartz and minor kaolinite, results from T-77-10 show a concentration of magnesium-rich calcite in the upper part of the seam and the presence of pyrite in the middle and lower parts. The sample from the middle part of the seam has a larger kaolinite to quartz ratio than the other two samples.

Microscopic mineral matter in the main coal seam is readily observable in reflected light, oil immersion. Clay, the most commonly occurring mineral, has a dark gray or black appearance and usually forms thin irregular layers or lenses, or, less commonly, forms pockets or petrifications filling cell lumens. Figures 29 and 30 show some clay layers interbedded with vitrinite.




Figure 30. Well defined clay (1) layers and lenses interbedded with vitrinite (2). Polished surface, oil immersion, x500.




Figure 31. Spherical aggregates of euhedral pyrite crystals (1); irregular masses of pyrite crystals (2); pyrite replacing cell walls (3); vitrinite groundmass. Polished surface, oil immersion, x500.

Pyrite is a minor constituent and is mostly syngenetic. Although it generally appears as small discrete grains, it sometimes occurs in spherical aggregates of euhedral crystals (Figure 31). In some samples it replaces cell walls and/or fills cell lumens (Figures 32 and 33). Drill core samples from the northern part of the study area generally have a greater abundance (up to 11%) of pyrite than drill core samples in the central part.

Some samples have detrital quartz grains. Quartz, because of its low relief and transparent gray color in reflected light under oil, is difficult to distinguish from void spaces. However, when present, compaction structures are used to identify quartz (Figure 34)(Kemežy and Taylor, 1964).

Several coal samples have a carbonate mineral with a mottled gray appearance (Figure 35). Some carbonate mineral crystals appear to replace vitrinite and others are surrounded by compaction structures. Identification of individual carbonate minerals is difficult in reflected light under oil because of their similar appearance.

A rough correlation exists between the chemical analyses of high temperature mineral ash and mineralogical composition of coal derived from low temperature ash and petrographic analyses (Mitchell and Gluskoter, 1975). The high SiO_2 percentage is derived mainly from quartz and to a lesser extent from kaolinite and smectite?. The Al_2O_3 as well as Ca, Mg, Na and K oxides originate in large part from the clays and the Fe_2O_3 and SO_3 primarily come from pyrite. The titanium and phosphorous probably represent inherent mineral matter in the coal forming plants.




Figure 32. Syngenetic pyrite (1) replacing well defined cell walls in vitrinite. Polished surface, oil immersion, x500.




Figure 33. Syngenetic pyrite replacing cell walls and filling cell lumens (2) in vitrinite; dark splotches in vitrinite caused by finely disseminated clay particles. Polished surface, oil immersion, x500.




Figure 34. Detrital, sub-angular quartz grain (1) in clay (2) and pyrite- (3) rich vitrinite; compaction structures enclose the quartz grain; small gray splotches on quartz grain are water marks. Polished surface, oil immersion, x500.




Figure 35. Carbonate mineral (1) associated with vitrinite (2) and exinite (3); exinite-rich band forms compaction structures (4) around the carbonate. Polished surface, oil immersion, x500.

COAL RANK

The average maximum percent reflectance of vitrinite (\bar{R}_O) was measured on 49 coal samples collected from drill cores and from weathered trench and open pit exposures. Forty-five of the samples are from the main seam and four from the lower seam. Table 3 lists the \bar{R}_O per sample at each site in stratigraphic order.

Low \bar{R}_O values from T-77-8 are caused by the abundance of pyrite. In samples with a relatively large quantity of pyrite, vitrinite \bar{R}_O readings are lowered by the high reflectance of pyrite. The larger readings ($\bar{R}_O = .63$ and $.65$) from T-77-8 are from two samples that have little pyrite and are, therefore, more representative of coal rank. Badly scratched polished surfaces also lower readings. Scratching was caused by pellets which disintegrated during the polishing process because of mineral matter intermixed with the coal.

With the possible exception of T-77-11 and TR 2, which are closely associated, no significant vertical variation in reflectance exists. T-77-12, upon which the greatest number of readings were taken, shows little stratigraphic variation between values (.62 to .65).

Figure 36 shows the relationship between the average \bar{R}_O of vitrinite per site and the location of sites with respect to one another. Reflectance gradually increases from .62 in the north to .86 in the south. The average \bar{R}_O for the entire study area is .67.

Coals in North America are usually ranked according to the A.S.T.M. (American Society for Testing Materials) system. The degree of coalification is based upon utilization parameters. High rank coals are classified on a fixed carbon or volatile content calculated on a dry,

TABLE 3. PLOT OF % \bar{R}_O OF VITRINITE AT DESIGNATED SITES (VALUES ARRANGED IN STRATIGRAPHIC ORDER).

	OP-1	TR 1	TR 2	T-77-11	TR 3	T-77-12	TR 5	TR 6	T-77-9	T-77-8	T-77-10
				% \bar{R}_O of Vitrinite at Designated Sites							
U. Section	.90	*	-	-	.62	-	-	-	.54	-	.66
	-	-	-	-	.66	-	-	-	-	-	-
	-	-	.57	.54	.62	.68	.65	.65	*	-	-
	.87	-	.56	-	.66	-	-	-	.55	-	-

Main Coal Seam											
	-	-	-	-	.68	-	-	-	.58	-	.60
	.87	-	-	-	.67	-	-	-	-	-	-
	-	.80	*	.65	*	-	.68	.71	.63	.65	-
	-	-	-	-	.66	-	-	-	-	-	-
	.79	-	-	-	.67	-	-	-	.63	-	-

M. Section											
	.87	-	-	-	.67	-	-	-	-	-	.59
	-	-	.67	-	.67	-	-	-	-	-	-
	.87	.63	.65	.70	.72	.67	*	.71	.64	.52	.64

Avg. % \bar{R}_O =	.86	.71	.60	.63	.67	.66	.67	.69	.64	.58	.62

Lower Seam											
	-	-	-	-	.72	-	-	-	.72	-	.63
	-	-	-	-	-	-	-	-	-	-	.66

*Pellets so badly scratched that reliable \bar{R}_O values were not possible.




Figure 36. Graph showing gradual increase in average vitrinite reflectance from north to south in the study area; shaded area represents the range of values (see Table 3 for values of individual samples).

mineral-matter-free (d.m.m.f.) basis. Lower rank coals having a less than 69% carbon content are classified according to their calorific value calculated on a moist, mineral-matter-free (m.m.m.f.) basis (Williamson, 1967). In the Tulameen Coalfield main seam analyses and washability tests from T-77-1 through T-77-6 and from bulk samples show the carbon content of the coal to lie below 69% (d.m.m.f.). The calorific values range from 11,707 B.T.U./lb to 12,362 B.T.U./lb (m.m.m.f.) (Adamson, 1978). Calorific values for the lower seam are similar to those of the main seam. Therefore, under the A.S.T.M. system the Tulameen coal with the T-77-1 to T-77-6 interval would be classified as High Volatile C Bituminous. According to Figure 37, which correlates maximum reflectance of vitrinite with the A.S.T.M. classification system, vitrinite reflectance values from the northern half of the basin fall within the High Volatile C Bituminous range and coal from OP-1 within High Volatile B to A Bituminous range.




Figure 37. Comparison of ASTM system of rank classification with % reflectance of vitrinite, B.T.U. and fixed carbon content of coal (from Stach, et.al., 1975; Hacquebard and Donaldson, 1970).

DISCUSSION

Widespread late Mesozoic and early Cenozoic sedimentary sections in western North America are associated with a continental margin arc-trench system (Dickinson, 1976). Dickinson (1976) constructed paleotectonic maps illustrating the Cordilleran arc-trench system during the Paleocene to Late Eocene and during the Late Eocene to Early Miocene (Figure 38). The Northern Cascade-Okanogan area during the Late Cretaceous and early Tertiary underwent a period of magmatic cessation characterized by post orogenic sedimentation. Arc magmatism resumed during the Eocene 55 to 50 million years ago and occupied a wide belt lying between 126° to 116° W. Longitude (Davis, 1977; Snyder, et. al, 1976). The Tulameen basin which forms a part of the Intermontane Belt of the Canadian Cordillera (Figure 39) lay within the magmatic arc.

Eocene geology of the Intermontane Belt in south central British Columbia was characterized by concomitant volcanism and sedimentation. Regional north-south block faulting was prevalent and extended across the Canadian/United States border from the Cascade Mountains to Montana (Davis, 1977). Thick successions of volcanic and sedimentary rocks were deposited in lakes and valleys associated with north-south trending block faults (Okulitch, et. al., 1977). Block faulting along with volcanism provided possible mechanisms for the formation of poorly drained areas suitable for coal deposition.

The lithology and thick sequence of clastic sedimentary rocks in the Tulameen basin indicate a fluvial-lacustrine origin (Reineck and Singh, 1975; Hendricks, 1977). The sediments are continental and represent a wide range of particle sizes deposited by fluvial processes.




Figure 38. A. Cordilleran arc-trench system in Paleocene to Late Eocene: SUS, Sustat assemblage; CS, Chuckanut Swauk sequence of North Cascades. B. Cordilleran arc-trench system in Late Eocene to Early Miocene: SA, San Andreas fault. (Dickinson, 1977)




Figure 39. Principal structural subdivisions of the southern Canadian Cordillera (Okulitch, et.al., 1977).

Massive, coarse-grained, feldspathic sandstones and conglomerates were deposited rapidly and were probably derived from a nearby source area. Relatively thick sequences of laminated, fine-grained sandstones, mudstones, fissile shales and intercalated coal horizons represent periods of lower energy environments. The frequent occurrence of tephra and bentonite throughout the sedimentary sequence documents repeated volcanic activity in the area.

Palynomorph assemblages from interior British Columbia imply a warm to subtropical continental climate during Middle and Late Eocene time (Rouse, 1977). Pollens identified from the coal-bearing sediments in the Princeton basin and in four similar locations in interior British Columbia consist of two principle elements: coniferous pollen and deciduous arboreal pollen mainly represented by the Betulaceae and Juglandaceae families (Rouse, 1967). The flora suggests an upland and inland environment. Although interior Eocene assemblages reflect a slightly cooler climate and have different major elements than coastal Eocene assemblages, they have enough similarities to suggest that the present Coastal and Cascade Mountain ranges were much lower during the Eocene than at present (Rouse, 1967).

The mineralogy of the coal determined from LTA and petrographic analyses indicates a fresh water swamp environment. Coals having relatively large quartz and kaolinite contents, as well as little to no carbonate and pyrite, are usually associated with fresh water depositional basins (Millot, 1952; Rao and Gluskoter, 1973). Other Tertiary coal basins in central British Columbia show a similar mineral association (Pearson, personal communication).

Swamp environments are defined on the basis of their predominant type of vegetation. Forest moor environments are dominated by woody trees and shrubs and result in deposits rich in vitrinite (Hacquebard and Donaldson, 1969). The vitrinite concentration and lack of significant petrographic variation in the Tulameen main coal seam indicate that a forest moor environment with a warm, moist climate predominated throughout peat formation. Figure 40 is a schematic block diagram of the Tulameen forest moor swamp as it may have appeared during peat formation.

Poorly drained areas that receive a sufficient inflow of water favor peat development. The type of basin and rate of subsidence help determine the nature of the peat (Hacquebard and Donaldson, 1967; Falini, 1965). Seam thickness and uniform composition (vitrite and exinite-poor clarite) of the Tulameen coal along with numerous thin shale and mudstone partings suggest a relatively high rate of active subsidence (Shibaoka and Smyth, 1972). Alteration of plant material to vitrinite requires anaerobic, reducing conditions with a low pH (Francis, 1961). Such conditions are maintained when the swamp is covered by a moderately high (1 - 1½ meters) stagnating water level (Stach, et. al., 1975). Because forest moor environments promote relatively continuous and rapid accumulation of peat, the groundwater level must steadily rise in order to keep abreast of peat formation. Thin layers of trimacerite and durite, which form subaquatically with increasing oxygenation, record minor periods of a relatively higher water level brought about by an influx of fresh water (Francis, 1961; Stach, et. al., 1975). A too rapid rise in the water level interrupts




Figure 40. Schematic block diagram of the main seam peat formation in the Tulameen forest moor swamp environment (not to scale).

peat deposition and brings about inorganic sedimentation. Thickness of partings is roughly proportional to the duration of the interruption in peat formation (Falini, 1965). Thin partings in the Tulameen coal indicate short interruptions brought about by small scale flooding. The thick interval of sediments between the lower and main coal seams represents a thick clastic incursion which interrupted peat formation for a long period of time. The lower seam may represent a major split in the main coal seam.

Some tectonic control over subsidence is suggested by the presence of active volcanism. The greater abundance of volcanically derived partings in the northern part of the field area suggests a volcanic source lay north of the basin.

Shale and mudstone partings represent an inwashing of detritus during periods of flooding. Kaolinite, which forms a large percent of the sediments, may have been deposited as a detrital sediment that remained unaltered after deposition (Jansa, 1972) or it may have been deposited as a different material that later was altered to kaolinite by interaction with humic acids (Staub and Cohen, 1978). Montmorillonite also occurs in the same coal seam as the kaolinite. Montmorillonite is an alteration product of volcanic ash which requires an alkaline environment for stability (Keller, 1956; Millot, 1970). Because of the nearly monomineralic nature of the bentonites, the presence of montmorillonite could be explained by instantaneous deposition of an ash fall followed by rapid settling which would isolate the particles from the acidic swamp environment and effectually remove them from the chemical system. The glassy tephra would be altered to montmorillonite after deposition

(Pevear, personal communication). The fact that montmorillonite was not altered to another mineral and coexists with kaolinite in many samples suggests that kaolinite was directly deposited into the swamp rather than forming in place. Kaolinite is a common product of intense weathering in warm to subtropical continental environments and its presence seems to be a general characteristic of a large number of western North American Eocene sediments and coal deposits (Murray and Patterson, 1975). Hopkins (1966) reports a thick kaolinite zone from the Upper Eocene Huntingdon Formation on Sumas Mountain which has been interpreted by Horton (1978) as representing detrital kaolinite brought into the basin as a result of intense weathering. Nearly monomineralic bentonite beds are interpreted as ash falls with minimum reworking, whereas shaly partings containing kaolinite and admixtures of kaolinite and montmorillonite are considered to be reworked kaolinite derived by erosion of low-lying intensely weathered areas.

The presence of montmorillonite defines the upper coalification limit of the Tulameen coal. Modifications in clay mineralogy are closely associated with progressive stages of geochemical coalification. Kisch (1968) correlates coal rank and burial-metamorphic mineral facies in terrigenous rocks. Montmorillonite occurs in high volatile bituminous coals and disappears during late diagenesis in lower volatile bituminous coals. The degree of coalification is mainly a function of time and temperature (Kaplan, 1971).



Results from the study of Tulameen coal rank show a gradual increase in rank from High Volatile C Bituminous in the northwestern half of the field area to High Volatile B/A Bituminous in the southeastern

portion. One possible interpretation for the increase in rank places the Blakeburn open pit area nearer to the center of the coal-forming basin based on the apparent greater thickness of the coal and the presence of fewer partings there in comparison to the northern part of the field area. Using such a model, Donaldson (1973) considered the Blakeburn area as the center and interpreted the higher rank as a result of greater depth of burial in conjunction with an abnormally high geothermal gradient caused by volcanic activity in the area.



A second possible interpretation for the increase in rank might be the emplacement of an igneous intrusion to the north or east of OP-1. This intrusion would have to be relatively large because the zone of contact metamorphism is highly dependent upon the size of the intrusion (Chakrabarti, 1969; Stach, et. al., 1975). Mining reports on underground workings in the Blakeburn area mention encountering a dike occupying a large fault zone, however, no details on the size, lithology or source of the dike were included (Shaw, 1952).

A deformation model for explaining the increase in coalification is a third possible interpretation. Figure 41 shows the relationship between major bedding planes and isorank lines in three different coalification situations (Pearson and Grieve, 1977). In coalification that occurs prior to folding, isoranks and major bedding planes are parallel. Rank increases vertically with depth and not down dip. Coalification is related to the original depth of burial that existed after deposition but before folding (Hacquebard and Donaldson, 1970). Post-folding coalification, on the other hand, is related to the maximum depth of burial after deformation. Isorank lines are horizontal

A. Coalification All Pre Folding



B. Coalification All Post Folding



C. Coalification Pre Folding and Post Folding




Figure 4]. Diagram showing possible configurations between coalification planes (dashed lines) and coal seams in relation to deformation (Pearson and Grieve, 1977).

and rank increases down dip with depth. When postfolding is superimposed on prefolding coalification, isorank lines lie obliquely to major bedding planes (Hacquebard and Donaldson, 1974; Stach, 1975; Pearson and Grieve, 1977). The Tulameen coal deposition basin was folded and tilted prior to deposition of the Miocene volcanics. Although the increase in coal rank follows strike along the southwest limb of the Tulameen basin from northwest to southeast, it parallels the down dip direction of the plunging fold axis. The southern part of the basin, therefore, had a greater maximum depth of burial which suggests some degree of post-deformational influence on coalification.

CONCLUSIONS

The Tulameen Coalfield has a fluvial-lacustrine origin. Fluvial sedimentation was probably controlled by regional north-south block faulting. The coal forming basin developed in a poorly drained, fairly low-lying area near an elevated upland. A warm, moist climate favored formation of a forest moor swamp environment. A relatively high water level and continuous subsidence brought about anaerobic reducing conditions favorable for peat growth. Periodic flooding and active volcanism brought inorganic material into the swamp. With time, biological and physical chemical processes altered the peat into coal. Coalification was partially post-deformational.

The coal is predominantly a bright, bedded coal composed primarily of vitrinite and only minor amounts of exinite and inertinite. Kaolinite and quartz are the major inorganic coal constituents. Interbedded within the coal are numerous thin montmorillonitic bentonite and kaolinite- and/or smectite-rich shale and mudstone partings. The partings not only increase in number towards the northern part of the basin but also their lithology becomes increasingly volcanoclastic.

Vitrinite reflectance measurements reveal an increase in coal rank from High Volatile C Bituminous in the north to High Volatile B/A Bituminous in the south of the field area.

REFERENCES CITED

- Adamson, T.J., 1978, Tulameen coal project: report on 1977 field work: Cyprus Anvil Mining Corporation, p. 1-25.
- Camsel, C., 1913, Geology and mineral deposits of Tulameen District, British Columbia: Geol. Surv. Canada Mem. 26, 188 p.
- Cockfield, W.E., 1948, Geology and mineral deposits of Nicola map-area, British Columbia: Geol. Surv. Canada, Dept. of Mines and Res., Mem. 249, 164 p.
- Davis, G.A., 1977, Tectonic evolution of the Pacific Northwest from Precambrian to present: WNP-1/4, PSAR, Amendment 23, Sub-appendix 2RC (report to the Nuclear Regulatory Commission by Washington Public Power Supply Systems), 46 p.
- Dickinson, W.R., 1976, Sedimentary basins developed during evolution of Mesozoic-Cenozoic arc-trench system in western North America: Canadian Jour. Earth Sci., v. 13, p. 1268-1287.
- Dolmage Campbell and Assoc. Ltd., 1975, Similkameen coal, in Coal resources of British Columbia, IV: B.C. Hydro and Power Authority, p. 6-2 - 6-28.
- Donaldson, J.R., 1973, The petrography of the coal from the Blakeburn strip mine in the Tulameen coal area, British Columbia: Geol. Surv. Canada, Dept. of Energy, Mines and Res., Paper 72-39, p. 1-13.
- Evans, S.H., 1977, Tulameen coal basin, in Carter, N.C. and Moir, R.J., eds., Geologic fieldwork 1977: B.C. Min. of Mines and Petrol. Res., p. 83-85.




- Falini, F., 1965, On the formation of coal deposits of lacustrine origin: *Geol. Soc. America Bull.*, v. 76, p. 1317-1346.
- Francis, W., 1961, *Coal, its formation and composition*: London, Edward Arnold Ltd., 806 p.
- Hacquebard, P.A., Birmingham, T.F. and Donaldson, J.R., 1967, Petrography of Canadian coals in relation to environment of deposition, in *Symp. on the science and technology of coal*, Ottawa: Mines Branch, Dept. of Energy, Mines and Res., p. 84-97.
- Hacquebard, P.A. and Donaldson, J.R., 1970, Coal metamorphism and hydrocarbon potential in the Upper Paleozoic of the Atlantic provinces, Canada: *Canadian Jour. Earth Sci.*, v. 7, p. 1139-1163.
- and ———, 1974, Rank studies of coal in the Rocky Mountains and Inner Foothills Belt, Canada: *Geol. Soc. America Spec. Paper* 153, p. 75-94.
- Hendricks, M.L., 1977, Environments of deposition of Coalmont Formation near Coalmont, Jackson County, Colorado (Abs.): *Am. Assoc. Petroleum Geologists Bull.*, v. 61, p. 1381.
- Hopkins, W.S. Jr., 1966, Palynology of Tertiary rocks of the Whatcom Basin, southwestern British Columbia and northwestern Washington: Unpublished Ph.D. thesis, University of British Columbia, 184 p.
- Horton, D.G., 1978, Clay mineralogy and origin of the Huntingdon fire clays on Canadian Sumas Mountain, southwest British Columbia: Unpublished M.S. thesis, Western Washington University, 96 p.
- Jansa, L., 1972, Depositional history of the coal-bearing Upper Jurassic-Lower Cretaceous Kootenay Formation, southern Rocky Mountains, Canada: *Geol. Soc. America Bull.*, v. 83, p. 3199-3222.

- Kaplan, M. Ye., 1971, Criteria for determining zones of catagenesis in terrigenous deposits: *Internat. Geol. Rev.*, v. 13, p. 1365-1376.
- Keller, W.D., 1956, Clay minerals as influenced by environments of their formation: *Amer. Assoc. Petroleum Geologists Bull.*, v. 40, p. 2689-2710.
- Kemežy, M. and Taylor, G.H., 1964, Occurrence and distribution of minerals in some Australian coals: *Jour. Inst. Fuel*, v. 37, p. 389-397.
- Kisch, H.J., 1968, Coal-rank and burial-metamorphic mineral facies, in Schenk, P.A. and Havenaar, I., eds., *Advances in organic geochemistry*: London, Pergamon Press, p. 407-425.
- Millot, G., 1952, The principal sedimentary facies and their characteristic clays: *Clay Minerals Bull.*, v. 1, p. 235-237.
- , 1970, *Geology of clays*: New York, Springer-Verlag, 429 p.
- Mitchell, R.S. and Gluskoter, H.J., 1975, Mineralogy of ash of some American coals: variations with temperature and source: *Fuel*, v. 55, p. 90-96.
- Murray, H.H. and Patterson, S.H., 1975, Kaolin, ball-clay and fire-clay deposits in the United States--their ages and origins, in Bailey, S.W., ed., *Proc. of the Internat. Clay Conf.*, 1975: Wilmette, Illinois, Applied Publ., p. 511-520.
- Okulitch, A.V., Price, R.A. and Richards, T.A., 1977, A guide to the geology of the southern Canadian Cordillera: *Geol. Assoc. Canada--Mineral Assoc. Canada--SEG Field Trip Guidebook 8*, 135 p.
- Pearson, D.E. and Grieve, D.A., 1977, Crowsnest Coalfield, in Carter, N.C. and Moir, R.I., eds., *Geologic fieldwork 1977*: *B.C. Min. of Mines and Petrol. Res.*, p. 47-54.

- Rao, C.P. and Gluskoter, H.J., 1973, Occurrence and distribution of minerals in Illinois coals: Illinois St. Geol. Surv. Circ. 476, p. 1-53.
- Reineck, H.E. and Singh, I.B., 1975, Depositional sedimentary environments: New York, Springer-Verlag, 439 p.
- Rice, H.M.A., 1947, Geology and mineral deposits of the Princeton map-area, British Columbia: Geol. Surv. Canada, Dept. of Mines and Res., Mem. 243, 136 p.
- Rouse, G.E., 1977, Paleogene palynomorph ranges of western and northern Canada, in Elsik, W.C., ed., Contributions to stratigraphic palynology, v.1: Am. Assoc. Strat. Palynologists, Contr. Ser. 5A, p. 48-65.
- Rouse, G.E., Hopkins, W.S. Jr. and Piel, K.M., 1966, Palynology of some Late Cretaceous and Early Tertiary deposits in British Columbia and adjacent Alberta, in Kosanke, R.M., ed., Symposium of palynology of the Late Cretaceous and Early Tertiary: Geol. Soc. America Spec. Paper 127, p. 213-246.
- Sehgal, R.S. and Wong, B., 1974, Significance of calorific value, ultimate analysis, ash fusion and mineral analysis of ash, in Fryer, J.F., Campbell, J.D. and Speight, J.C., eds., Symp. on coal evaluation, Information Series 76: Alberta Res. Council and Coal Assoc. of Canada, p. 21-30.
- Shaw, W.S., 1952, The Tulameen Coalfield, British Columbia: Geol. Surv. Canada, Dept. of Mines and Tech. Surv., Paper 52-19, p. 1-13.

- Shibaoka, M. and Smyth, M., 1975, Coal petrology and the formation of coal seams in some Australian sedimentary basins: *Econ. Geol.*, v. 70, p. 1463-1473.
- Snyder, W.S., Dickinson, W.R. and Silberman, M.L., 1976, Tectonic implications of space-time patterns of Cenozoic magmatism in the western United States: *Earth and Planetary Sci. Letters*, v. 32, p. 32, p. 91-106.
- Stach, E., Taylor, G.H., Mackowsky, M.-Th., Chandra, D., Teichmüller, M. and Teichmüller, R., 1975, *Stach's textbook of coal petrology*: Berlin, Gebruder Borntraeger, 398 p.
- Staub, J.R. and Cohen, A.D., 1978, Kaolinite-enrichment beneath coals; a modern analog, Snuggedy swamp, South Carolina: *Jour. Sed. Petrology*, v. 48, p. 203-210.
- Ward, C.R., 1977, Mineral matter in the Springfield-Harrisburg (No. 5) Coal Member in the Illinois Basin: *Illinois St. Geol. Survey Circ.* 498, 35 p.
- Williamson, I.A., 1967, *Coal mining geology*: London, Oxford Univ. Press, 266 p.


APPENDIX
OF
MEASURED SECTIONS
AND
SAMPLE LOCATIONS




1 meter

OP-1 Measured Section Showing Location of Samples

* ashed coal samples (L.T.A.)




TR 1 Measured Section
Showing Location of
Samples







TR 2 Measured Section Showing Location of samples

* ashed coal sample (L.T.A)




TR 3 Measured Section Showing Location of Samples




TR 4 Measured Section Showing Location of Samples


1 Meter





TR 5 Measured Section Showing Location of Samples




TR6 Measured Section Showing Location of Samples




Location of Samples from T-77-12





Location of Samples from T-77-11



Location of Samples from T-77-9



Location of Samples from T-77-8



Measured Section of T-77-10 from Drill Core Showing Sample Locations

* ashed samples (L.T.A.)