

Marshall University Marshall Digital Scholar

Management Faculty Research

Management and Health Care Administration

10-2018

Telepsychiatry: Access in Rural Areas

David P. Paul III

Brianna Washington
Marshall University

Annie Robinson
Marshall University

Mike Tonnie
Marshall University

Alberto Coustasse
Marshall University, coustassehen@marshall.edu

Follow this and additional works at: https://mds.marshall.edu/mgmt_faculty

 Part of the [Business Commons](#), and the [Psychiatry Commons](#)

Recommended Citation

Paul III, D. P., Washington, B., Robinson, A., Tonnie, M., & Coustasse, A. Telepsychiatry: Access in Rural Areas. Proceedings of the Northeast Business & Economics Association 2018 Conference (pp. 245-252). Galloway, NJ.

This Conference Proceeding is brought to you for free and open access by the Management and Health Care Administration at Marshall Digital Scholar. It has been accepted for inclusion in Management Faculty Research by an authorized administrator of Marshall Digital Scholar. For more information, please contact zhangj@marshall.edu, beachgr@marshall.edu.

Telepsychiatry: Access in Rural Areas

David P. Paul, III
Monmouth University
West Long Branch, NJ 07764
(302) 227-1930
dpaul@monmouth.edu

Brianna Washington
Marshall University
South Charleston, WV 25303
washington43@live.marshall.edu

Annie Robinson
Marshall University
South Charleston, WV 25303
robinsonn1@livemarshall.onmicrosoft.com

Mike Tonnie
Marshall University
South Charleston, WV 25303
mike1@marshall.edu

Alberto Coustasse
Marshall University
South Charleston, WV 25303
coustassehen@marshall.edu

ABSTRACT

Rural areas have experienced higher than average healthcare workforce problems, especially concerning limited access to mental health services. Telepsychiatry may provide at least a partial solution, as it has improved access and quality of care available in rural environments despite implementation problems. As technology continues to advance access, telepsychiatry will also need to strengthen making access more readily available. Additional research is required to identify modalities and diverse methods that can be used to increase access to mental health services further and improve outcomes in rural and underserved areas.

Keywords

Access, benefit, rural communities, telehealth, tele-medicine, telepsychiatry

1 INTRODUCTION

In 2010, 19.3% of the population lived in rural areas (Ratcliffe et al., 2016). According to the United States Census Bureau, a rural area is any population, housing, or territory not defined as an urbanized area or urbanized cluster while an urbanized area contains 50,000 or more individuals and an urban cluster has at least 2,500 but less than 50,000 people (The United States Census Bureau, 2016).

Rural areas experience higher than average healthcare workforce shortages limiting accessibility to healthcare services (RHIhub, 2014). To illustrate, less than 10% of physicians practice in rural areas although 20% of the U.S. population lives in rural communities (Stanford School of Medicine, 2010). Geographic constraints are another challenge (Stanford School of Medicine, 2010). Residents travel further to access different and often limited services (Smalley et al., 2012). Some patients have conveyed substantial burden about time and money (Smalley et al., 2012). Lack of public transportation, distance, hazardous weather conditions, and environmental issues have been among other challenges exacerbated in this environment (Eberhardt and Pamuk, 2004).

For urban and rural areas, the incidence and prevalence of most behavioral disorders have been comparable (Hecke, 2012). A 2014 study found no significant difference in incidence or prevalence of major depression or severe mental illness (SMI) in large metropolitan and rural areas (Breslau et al., 2014). Specifically, a 5.6% prevalence of major depression and SMI was found in large metropolitan areas while a 6% prevalence of major depression and SMI was observed in rural areas (Breslau et al., 2014). Be that as it may, rural areas provide limited availability of mental health services (NRHA, 2017; Douthit et al., 2015). As of September 2017, 59.36% of mental health provider shortage areas were in non-metropolitan areas (HRSA, 2017). Thus, residents were less likely to receive treatment (Eberhardt and Pamuk, 2004). This has led to disparities between rural and urban residents (NRHA, 2017).

Rural areas have higher depression, domestic violence, child abuse, and suicide rates compared to their urban counterparts (Douthit et al. 2015). A lack of providers in addition to other challenges has led to many states developing behavioral health models (Hecke, 2012). Alternatively, telehealth has been utilized. Varying definitions, policies, and regulations across states surround how telehealth has been used (CCHP, 2017). Telehealth, also known as telemedicine, has used technology to deliver healthcare services and clinical information through telecommunications technology (HRSA, 2013; ATA, 2016; AHRQ, 2016).

Telehealth included direct, electronic patient-to-provider interactions in addition to medical devices that transmit and collect health information (ONC, 2017). Medical tools involved have included smartphone applications, activity trackers, automated

reminders, and blood glucose monitors (ONC, 2017). Video conferencing, the internet, store-and-forward imaging, streaming media, and terrestrial and wireless communications technologies have also been utilized to expand access to healthcare services (HRSA, 2015). Tele-behavioral health, also known as telepsychiatry, can be provided in nursing homes, clinics, schools, and other localized community settings (HRSA, 2013). As of 2016, 65% of hospitals in the U.S. have used or implemented telehealth (AHA, 2016).

The purpose of this research was to examine the benefits of telepsychiatry for adults living in rural communities to determine if telepsychiatry improved access and quality. We posit that utilization of telepsychiatry in rural areas increases accessibility to mental health services for adults, and also that the integration of telepsychiatry in primary healthcare settings improved quality of mental health care for rurally located adults.

2 RESULTS

Access to mental health care

Practitioners, psychiatrists, and researchers have long recognized the mental health obstacles exhibited by rural communities and the need for trained mental health professionals (Smalley et al., 2010). In 2015, 46.2 million or 14% of the nation lived in rural communities (U.S. Department of Agriculture, 2016). Rural areas reported a higher percentage of adults with any mental illness, 18.3%, and SMI, 4.7%, compared to large metro areas where 17.3% recorded any mental illness and 3.8% reported SMI (RHIfhub, 2017). Unfortunately, are numerous barriers limit access to mental health services in rural areas of the U.S., including transportation infrastructure, limited numbers of providers, poverty, social stigma, and lack of insurance (Cohn and Hastings, 2013).

Rural patient access to appropriate and adequate mental health services has been limited by various barriers including accessibility, availability, and acceptability. Utilization of telepsychiatry has shown to overcome these limitations by allowing psychiatrists to connect to health personnel staff and patient from a distance while keeping treatment in the patients' community which has proven to help better manage their disease, their symptoms, and lives (Shore, 2013; Trondsen et al., 2012). A range of mental health issues has shown to be efficiently handled using telepsychiatry (Trondsen et al., 2012).

The majority of patients of all ages report high satisfaction with telepsychiatry (Wynchank and Fortuin, 2010). Research has established that increase in patient satisfaction leads to decrease missed appointments, 7%-10% missed appointments compared to non-telepsychiatry rates of 35%-42% of missed visits (Saeed, Diamond and Bloch, 2011). Research has shown that other benefits of telepsychiatry included saved time and reduced need to travel outside the community for services, and that psychiatric and psychological treatment delivered via telepsychiatry have the same clinical outcomes as therapy provided face-to-face (Egede et al., 2013; Fortney et al., 2015; Trondsen et al., 2012). Unfortunately, a review of the literature (Bolton and Dorstyn, 2015, p. 252) characterized most studies as "characterized by small and underpowered samples" such that "the equivalence of telepsychology and face-to-face psychotherapy could not be determined."

Referencing availability of mental health providers, telepsychiatry has been shown not necessarily to create resources, but to reorganize them (Grady, 2012). There has been a continued shortage of mental health providers. A nationally, over 75% of U.S. counties qualify as having a shortage, 85% of deficit stemming from rural areas, and half of all U.S. counties report no mental health providers (RHIfhub, 2017; Smalley et al., 2010; Thomas et al., 2009). Research has shown the shortage of mental healthcare providers in rural communities have been attributed to lower incomes compared to peers in urban/metro settings, few social/educational opportunities, professional isolation, being overburdened by demand, and difficulties adapting to rural life (Cohn and Hastings, 2013; Smalley et al., 2010; Thomas et al., 2009).

Additionally, telepsychiatry has proved to alleviate professional isolation, expand networks that provide telepsychiatry, and increase collaboration between rural mental health clinics and urban/metro mental health clinics (Chung-Do et al., 2012; Grady 2012). Programs such as East Carolina University Telemedicine Center have developed, and expanded services provided by the center by adding sites in 13 eastern North Carolina counties using only three psychiatrists to offer patient services (Saeed, Diamond, and Bloch, 2011). Similar results of expansion have been evident in West Virginia where the Department of Behavioral Medicine and Psychiatry in conjunction with WVU Mountaineer Doctor Television has operated 35 telepsychiatry clinics in 16 counties (West Virginia University, 2015).

The stigma associated with mental health has been shown to act as a barrier to mental health (RHIfhub, 2017; Gamm, Stone and Pittman, 2010). Those living in rural areas often have believed they have less privacy and confidentiality, and they have proven a lack of understanding and knowledge of mental illness, and lack of perceived need for mental health care, which causes hesitation in seeking treatment (Bryant et al., 2013).

Quality issues

When patients have used telepsychiatry in the form of video conferencing, it has been deemed equivalent to face-to-face care received in a physician's office (Fortney, 2015; Morland et al., 2013; Saeed et al., 2017; Whealin et al., 2015). Studies have proven that telepsychiatry has been a method used to provide continuous quality care to several types of adult populations who would have otherwise been left with their mental illnesses untreated (Deslich et al., 2013; Shore, 2015; Saeed et al., 2017).

Rurally located adults usually only have access to a primary care physician for their physical care, as well as their mental care. This has often led to misdiagnosed mental illnesses, lack of appropriate referrals to psychiatrists, and medication mismanagement (Saeed et al., 2017). The integration of mental health care into primary health facilities using a form of telehealth such as telepsychiatry has been a method that has increased the quality of care while treating adults in rural communities (Riding-Malon and Werth, 2014, Lambert et al., 2016). There have been many successful studies developed regarding the integration of telepsychiatry into primary health in rural areas (Fortney et al., 2015; Guerrero et al., 2017; Lu et al., 2014).

In 2012 the U.S. VA was responsible for 179,146 telepsychiatry sessions with rurally located veterans (Fortney et al., 2015). Mental disorders have been diagnosed in more than 30% of Veterans that have used healthcare through the U.S. Department of Veteran's Affairs (VA). Approximately 41% of veterans who are currently using the VA for healthcare live in a rural community (Whealin et al., 2015).

Due to geographical location, access to mental health facilities is scarce, and there are veterans that do not want to leave their homes due to trauma endured while overseas in countries such as Afghanistan and Iraq (Whealin et al., 2015). Telepsychiatry has been a successful treatment method for veterans and other adult patients who felt more comfortable being treated without having to be face-to-face with their psychiatrist (Ganzini et al., 2013; Waugh, Voyles and Thomas, 2015). When telepsychiatry has been used as a treatment with veterans who utilize the VA for healthcare, admissions to hospitals and facilities due to psychiatric issues have decreased by 25 % (Lu et al., 2014).

In a 2017 study, performed by a psychiatric department at Wayne State University in Michigan, Amirsadri et al. (2017) utilized telepsychiatry as a treatment modality after receiving many calls from an elderly patient's daughter asking them to care for her mother who had been diagnosed with schizophrenia in the past. The homebound old patient lived in a rural area and had not followed up on her mental health care in 20 years. Combining in-person care with a virtual type of care such as telepsychiatry was referred to as 'hybrid care' (Hilty and Yellowlees, 2015). The "hybrid care" method used in Amirsadri's et al. (2017) study consisted of a social worker who traveled to the patient's home and used a tablet computer to video conference with a psychiatrist. The case study demonstrated that the patient's mental health was appropriately assessed, and she reported great improvement, but the patient's physical health issues were also assessed and adequately referred by the social worker to other providers.

Another 2017 study took place in Hawaii, in a rural neighborhood island. This rural community did not have any psychiatrists available to care for 1,500 patients who utilized a federally qualified community health center (FQHC). A grant-funded Behavior Health Integration team (BHI) was established as a route to assist in this disparity, and a psychiatrist, who was readily available through video-teleconferencing, was integrated as part of the patients' primary care provided by the FQHC (Guerrero et al., 2017). The integration of psychiatric consultations through video-teleconferencing was very well received by the Hawaiian rurally located adult patients (Fortney et al., 2015; Guerrero et al., 2017). Finally, because the BHI team ensured that psychiatric care was integrated, it has been utilized on five to ten patients a month (Guerrero et al., 2017). Similarly, in a pilot project designed to evaluate a teleconferencing clinical psychology service, the majority of clients reported that they were satisfied with this therapeutic approach, and some expressed a preference for it because they felt it was less confrontational than face-to-face therapy and made them feel more in control of the therapeutic process (Simpson, Deans and Brebner, 2001).

Cost issues

Cost analysis of telepsychiatry is mandatory (Jones, 2001), especially in today's reimbursement-based behavioral healthcare world (Waugh, Voyles and Thomas, 2015). Early studies (see Hyler and Gangure, 2004 for a review) regarding the cost-effectiveness of telepsychiatry yielded mixed results, especially as this approach to mental health care is not suitable for all patients (Norman, 2006). The situation is changing, however, as technology advances and policies shift (Waugh, Voyles and Thomas, 2015), and telepsychiatry is now thought to provide cost-savings to practitioners and healthcare organizations (Brauser, 2012; Deslich et al., 2013; Deslich, Thistlethwaite and Coustasse, 2013). For example, Thomas et al. (2017) determined that telepsychiatry consultations for pediatric psychiatric emergencies, as measured by charges and time, were more cost-efficient from the perspective of the hospital system compared with usual care consisting of in-person consultation at a children's hospital main campus after transportation by ambulance. In a study of the homeless individual in India (a resource-poor country), the majority of whom suffered from long-term mental illness, a telepsychiatry model was found to be 50 times more economical than providing either mental health care at a tertiary care center or via community outreach service (Moirangthem et al., 2017).

The U.S. Government now recognizes telehealth, including mental health care, as an efficient and cost-effective alternative to face-to-face healthcare provision for Medicaid (SecureTelehealth, 2017) and Medicare (CMS, 2016), although there are dramatic variations in claims among states because regulations from state-to-state vary widely (Boyles, 2017). The private sector has adopted telemedicine, including telepsychiatry, even more quickly than the public one, with nearly 60% of employers providing coverage for telemedicine (Jaspin, 2016), and consequently, Griffin (2017) has described behavioral telehealth as “mainstream.”

3 DISCUSSION

The purpose of this research was to examine how having access to telepsychiatry in rural geographic areas can improve the quality of mental health care, reducing the overall symptoms of mental health issues in adults. The results of the literature review have suggested that having access to telepsychiatry can assist the adult rural population in getting mental health care that they may not have received otherwise. Increased access to telepsychiatry to the rural mental adult population has been supported in this literature review. When telepsychiatry has been utilized by rurally located adults, the quality of care that this community receives to reduce mental health issues improves.

It should be a priority for rural areas to have the same access and quality of mental health care as surrounding urban areas (Nelson, 2013, Riding-Malon and Werth, 2014). Access to telepsychiatry has been established as a viable option for adults with mental illnesses living in rural areas. Using telepsychiatry as an option, psychiatric quality of care can be improved, and populations that do not have access can significantly benefit from this healthcare technology (Deslich et al., 2013).

Adults with mental illness who require psychiatric care and live in rural areas could experience personalized, patient-centered care with the utilization of telepsychiatry (Shore, 2015). Incorporating telepsychiatry in some capacity in the primary care setting, rural adults needing mental health treatment would be more likely to be exposed to integrated healthcare which would potentially increase their quality of care.

Telepsychiatry has the potential to impact the quality of care for patients, especially in areas that do not have access to specialists, such as rural communities. However, its more widespread implementation has been limited by problems with reimbursement, issues with licensure, privacy/security/HIPPA concerns, lack of training, cooperation from physicians and technological limitations (Chandra, Pettry and Paul, 2005; Deslich et al., 2013; McGinty et al., 2006; Hyler and Gangure, 2004; Wang and Alexander, 2014).

4 CONCLUSIONS

Telepsychiatry utilization in rural areas has been demonstrated to have significant ability to transform care delivery and clinician productivity. The preponderance of the evidence indicates that telepsychiatry has increased access to mental health services for adults in rural areas, as well as improved mental healthcare delivery, and patient outcomes.

5 REFERENCES

- Agency for Healthcare Research and Quality [AHRQ] (2016), *Telehealth: Mapping the Evidence for Patient Outcomes from Systematic Reviews*. Retrieved from <https://effectivehealthcare.ahrq.gov/topics/telehealth/technical-brief/>
- Amirsadri, Alireza, Jaclynne Burns, Albert Pizzuti and Cynthia L. Arfken (2017), “Home-based Telepsychiatry in US Urban Area,” *Case Reports in Psychiatry*, Volume 2017 (2017), Article ID 6296423. Retrieved from <https://www.hindawi.com/journals/crips/2017/6296423/>
- American Hospital Association [AHA] (2016), *2014, 2015, and 2016 AHA Annual Survey IT Supplement*. Retrieved from <http://www.aha.org/advocacy-issues/factsheets/fs-telehealth.pdf>
- American Telemedicine Association [ATA] (2016), *About Telemedicine*. Retrieved from <http://www.americantelemed.org/about/telehealth-faqs->
- Bolton, A. J., and D. S. Dorstyn (2015), “Telepsychology for Posttraumatic Stress Disorder: A Systematic Review,” *Journal of Telemedicine and Telecare*, 21 (5), 254-267.
- Boyles, Salynn (2017), “Rural Telemedicine for Mental Health Surges - But Medicare Claims Analysis Shows State-By-State Use Varies Widely,” *MedPage Today*, May 2. Retrieved from <https://www.medpagetoday.com/psychiatry/generalpsychiatry/64970>
- Brauser, Deborah (2012), “Pediatric Telepsychiatry Cuts Costs, Improves Symptoms,” *Medscape Medical News*, June 13. Retrieved from <https://www.medscape.com/viewarticle/765618>

- Breslau, Joshua, Grant N. Marshall, Harold A. Pincus and Ryan A. Brown (2014), “Are Mental Disorders More Common in Urban than Rural Areas of the United States?” *Journal of Psychiatric Research*, 56 (September), 50-55.
- Bryant, Keneshia, Nancy Greer-Williams, Nathaniel Willis and Mary Hartwig (2013), “Barriers to Diagnosis and Treatment of Depression: Voices from a Rural African-American Faith Community,” *Journal of the National Black Nurses Association*, 24 (1), 31–38.
- Bureau of Health Workforce Health Resources and Services Administration [HRSA] (2017), *Designated Health Professional Shortage Areas Statistics*. Retrieved from <https://datawarehouse.hrsa.gov/Tools/HDWReports/Reports.aspx>
- The Center for Connected Health Policy [CCHP] (2017), *What is Telehealth?* Retrieved from <http://www.cchpca.org/what-is-telehealth>
- Centers for Medicare & Medicaid Services [CMS] (2016), *Telehealth Services*, Department of Health and Human Services. November. Retrieved from <https://www.cms.gov/Outreach-and-Education/Medicare-Learning-Network-LN/MLNProducts/downloads/TelehealthSrvcsfctsht.pdf>
- Chandra, Ashish, Charles E. Pettry, Jr. and David P. Paul, III (2005), “Telemedicine from a Macromarketing Viewpoint: A Critical Evaluation with Proposed Licensing Strategies,” *Journal of Non-Profit and Public Sector Marketing*, 13 (1/2), 111-135.
- Chung-Do, Jane, Susana Helm, Michael Fukuda, Dan Alicata, Stephanie Nishimura and Iwalani Else (2012), “Rural Mental Health: Implications for Telepsychiatry in Clinical Service, Workforce Development, and Organizational Capacity,” *Telemedicine and e-Health*, 18 (3), 244–246.
- Cohn, Tracey J. and Sarah Hastings (2013), “Building a Practice in Rural Settings: Special Considerations,” *Journal of Mental Health Counseling*, 35 (3), 228-244.
- Deslich, Stacie, Timothy Thistlethwaite and Alberto Coustasse (2013), “Telepsychiatry in Correctional Facilities: Using Technology to Improve Access and Decrease Costs of Mental Health Care in Underserved Populations,” *The Permanente Journal*, 17(3), 80-86.
- Deslich, Stacie, Bruce Stec, Shane Tomblin and Alberto Coustasse (2013), “Telepsychiatry in the 21st Century: Transforming Healthcare with Technology,” *Perspectives in Health Information Management*, 10 (Summer), 1f.
- Douthit, N., S. Kiv, T. Dwolatzky and S. Biswas (2015), “Exposing Some Important Barriers to Health Care Access in the Rural USA,” *Public Health*, 129 (6), 611–620.
- Eberhardt, Mark S. and Elsie R. Pamuk (2004), “The Importance of Place of Residence: Examining Health in Rural and Nonrural Areas,” *American Journal of Public Health*, 94 (10), 1682-1686.
- Egede, Leonard E., Ron Acierno, Rebecca G. Knapp, Carl W. Lejuez, Melba Hernandez-Tejada, Elizabeth H. Payne and B. Christopher Frueh (2015), “Psychotherapy for Depression in Older Veterans via Telemedicine: A Randomised, Open-Label, Non-Inferiority Trial,” *The Lancet – Psychiatry*, 2 (8), 693–701.
- Fortney, John C., James F. Burgess, Jr., Hayden B. Bosworth, Brenda M. Booth and Peter J. Kaboli (2011), “A Re-conceptualization of Access for 21st Century Healthcare,” *Journal of General Internal Medicine*, 26 (Suppl 2), 639–647.
- Fortney, John C., Jeffrey M. Pyne, Eric E. Turner, Kellee M. Farris, Tre M. Normoyle, Marc D. Avery, Donald M. Hilty and Jürgen Unützer (2015), “Telepsychiatry Integration of Mental Health Services into Rural Primary Care Settings,” *International Review of Psychiatry*, 27 (6), 525-539.
- Gamm, Larry D., Sarah Stone and Stephanie Pittman (2010), “Mental Health and Mental Disorders—A Rural Challenge: A Literature Review,” *Rural Healthy People*, 1 (1), 97-114.
- Ganzini, Linda, Lauren M, Denneson, Nancy Press, Matthew J. Bair, Drew A. Helmer, Jennifer Therese Poat and Dteven Dobscha (2013), “Trust is the Basis for Effective Suicide Risk Screening and Assessment in Veterans,” *Journal of General Internal Medicine*, 28 (9), 1215-1221.

- Grady, Brian (2012), "Promises and Limitations of Telepsychiatry in Rural Adult Mental Health Care," *World Psychiatry*, 11 (3), 199-201.
- Griffin, M. (2017), "Behavioral Telemedicine Goes Mainstream," *Wecounsel*, January 11. Retrieved from <https://www.wecounsel.com/behavioral-telemedicine-goes-mainstream/>
- Guerrero, A. P. S., Takesue, C. L., Medeiros, J. H. N., Duran, A. A., Humphry, J. W., Lunsford, R. M., Shaw, D. V., Fukuda, M. H. and Hishinuma, E. S. (2017), "Primary Care Integration of Psychiatric and Behavioral Health Services: A Primer for Providers and Case Report of Local Implementation," *Hawai'i Journal of Medicine & Public Health*, 76 (6), 147-151.
- Health Resources and Services Administration [HRSA] (2013), *Increasing Access to Behavioral Health Care Through Technology*. Retrieved from <https://www.hrsa.gov/sites/default/files/publichealth/guidelines/BehavioralHealth/behavioralhealthcareaccess.pdf>
- Health Resources and Services Administration [HRSA] (2015), *Telehealth Program*. Retrieved from <https://www.hrsa.gov/rural-health/telehealth/index.html>
- Hecke, Saskia V. (2012), "Behavioral Health Aides a Promising Practice for Frontier Communities," *National Center for Frontier Communities*. Retrieved from http://frontierus.org/wp-content/uploads/2012/01/FREP-Behavioral_Health_Aide_Models-2012.pdf
- Hyler, Steven E. and Dinu P. Gangure (2004), "Legal and Ethical Challenges in Telepsychiatry," *Journal of Psychiatric Practice*, 10 (4), 272-276.
- Hilty, Donald M. and Peter M. Yellowees (2015), "Collaborative Mental Health Services Using Multiple Technologies: The New Way to Practice and a New Standard of Practice?" *Journal of the American Academy of Child and Adolescent Psychiatry*, 54 (4), 245-246.
- Jaspin, Bruce (2016), "Most Employers Paying for Doctor Telemedicine Visits," *Forbes [online]*, October 28. Retrieved from <https://www.forbes.com/sites/brucejapsen/2016/10/28/most-employers-now-cover-doctor-telemedicine-visits/#75515d222d0a>
- Jones, Beverly N., III (2001), "Telepsychiatry and Geriatric Care," *Current Psychiatry Reports*, 3 (1), 29-36.
- Lambert, D., Gale, J., Hartley, D., Croll, Z. and Hansen, A. (2016), "Understanding the Business Case for Telemental Health in Rural Communities," *Journal of Behavioral Health Services & Research*, 43 (3), 366-379.
- Lu, M. W., K. I. Woodside, T. L. Chisholm and M. F. Ward (2014), "Making Connections: Suicide Prevention and the Use of Technology with Rural Veterans," *Journal of Rural Mental Health*, 38 (2), 98-108.
- McGinty, Kaye L., Sy Atezaz Saeed, Scott C. Simmons and Yilmaz Yildirim (2006), "Telepsychiatry and e-Mental Health Services: Potential for Improving Access to Mental Health Care," *Psychiatric Quarterly*, 77 (4), 335-342.
- Moirangthem, Sydney, Sabina Rao, Channaveerachari Naveen Kumar, Manjunatha Narayana, Neelaveni Raviprakash and Suresh Bada Math (2017), "Telepsychiatry as an Economically Better Model for Reaching the Unreached: A Retrospective Report from South India," *Indian Journal of Psychological Medicine*, 39 (3), 271-275.
- Morland, Leslie A., Michelle Raab, Margaret-Anne Mackintosh, Craig S. Rosen, Clara E. Dismuke, Carolyn J. Greene and B. Christopher Frueh (2013), "Telemedicine: A Cost-Reducing Means of Delivering Psychotherapy to Rural Combat Veterans with PTSD," *Telemedicine and e-Health*, 19 (10), 754-759.
- National Rural Health Association [NRHA] (2017), *About Rural Health Care*. Retrieved from <https://www.ruralhealthweb.org/about-nrha/about-rural-health-care>
- Nelson, Danielle, Valeria M. Hewell, Larry Roberts, Elizabeth Kersey and Jaedon P. Avey (2013), "Telebehavioral Health Delivery of Clinical Supervision Trainings in Rural Alaska: An Emerging Best Practices Model for Rural Practitioners," *Journal of Rural Mental Health*, 36 (2), 10-15.
- Norman, S. (2006), "The Use of Telemedicine in Psychiatry," *Journal of Psychiatric and Mental Health Nursing*, 13 (6), 771-777.

- Ratcliffe, Michael, Charlynn Burd, Kelly Holder and Alison Fields (2016), *Defining Rural at the U.S. Census Bureau. American Community Survey and Geography Brief*. Retrieved from https://www2.census.gov/geo/pdfs/reference/ua/Defining_Rural.pdf
- Riding-Malon, Ruth and James L. Werth, Jr. (2014), "Psychological Practice in Rural Settings: At the Cutting Edge," *Professional Psychology: Research and Practice*, 45 (2), 85-91.
- Rural Health Information Hub [RHHub] (2014), *Rural Health Disparities*. Retrieved from <https://www.ruralhealthinfo.org/topics/rural-health-disparities>
- Rural Health Information Hub [RHHub] (2017), "Rural Mental Health," in *RHHub*. Retrieved from <https://www.ruralhealthinfo.org/topics/mental-health#workforce-challenges>
- Saeed, Sy Atezaz, John M. Diamond and Richard M. Bloch (2011), "Use of Telepsychiatry to Improve Care for People with Mental Illness in Rural North Carolina," *North Carolina Medical Journal*, 72 (3), 219-222.
- Saeed, Sy Atezaz, Toni Johnson, Mandeep Bagga and Oliver M. Glass (2017), "Training Residents in the Use of Telepsychiatry: Review of the Literature and a Proposed Elective," *Psychiatric Quarterly*, 88 (2), 271-283.
- SecureTelehealth (2017), "Medicare Reimburses for Telehealth," November 2. Retrieved from <http://securetelehealth.com/medicare-reimbursement.html>
- Shore, James (2013), "Telepsychiatry: Videoconferencing in the Delivery of Psychiatric Care," *American Journal of Psychiatry*, 170 (3), 256-262.
- Shore, Jay (2015), "The Evolution and History of Telepsychiatry and Its Impact on Psychiatric Care: Current Implications for Psychiatrists and Psychiatric Organizations," *International Review of Psychiatry*, 27 (6), 469-475.
- Simpson, Susan, George Deans and Eileen Brebner (2001), "The Delivery of a Tele-Psychology Service to Shetland," *Clinical Psychology & Psychotherapy*, 8 (2), 130-135.
- Smalley, K. Bryant, Yancey, C. Thresa, Jacob C. Warren, Karen Z. Naufel, Rebecca Ryan and Kames L. Pugh (2010), "Rural Mental Health and Psychological Treatment: A Review for Practitioners," *Journal of Clinical Psychology*, 66 (5), 479-489.
- Smalley, K. Bryant, Jacob Warren and Jackson Rainer (2012), *Rural Mental Health: Issues, Policies, and Best Practices*. NY: Springer Publishing Company.
- Stanford School of Medicine (2010), "Healthcare Disparities & Barriers to Healthcare," *eCampus Rural Health*, (9). Retrieved from http://ruralhealth.stanford.edu/health-pros/factsheets/downloads/rural_fact_sheet_5.pdf
- The Office of the National Coordinator for Health Information Technology [ONC] (2017), *Health IT Playbook Telehealth*. Retrieved from <https://www.healthit.gov/playbook/patient-engagement/#Telehealth>
- The United States Census Bureau [USCB] (2016), *Urban and Rural*. Retrieved from <https://www.census.gov/geo/reference/urban-rural.html>
- Thomas, J. F., Novins, D. K., Hosokawa, P. W., Olson, C. A., Hunter, D., Brent, A. S., Frunzi, G. and Libby, A. M. (2017), "The Use of Telepsychiatry to Provide Cost-Efficient Care During Pediatric Mental Health Emergencies," *Psychiatric Services*, published ahead of print. Retrieved from <https://ps.psychiatryonline.org/doi/pdf/10.1176/appi.ps.201700140>
- Thomas, Kathleen C., Alan R. Ellis, Thomas R. Konrad, Charles E. Holzer and Joseph P. Morrissey (2009), "County-Level Estimates of Mental Health Professional Shortage in the United States," *Psychiatric Services*, 60 (10), 1323-1328.
- Trondsen, Marianne V., Stein Roald Bolle, Geir Øyvind Stensland and Askel Tjora (2012), "VIDEOCARE: Decentralised Psychiatric Emergency Care through Videoconferencing," *BMC Health Services Research*, 12 (1), 470-473.
- U.S. Department of Agriculture (2016), *Population and Migration. Economic Research Service*. Retrieved from <https://www.ers.usda.gov/topics/rural-economy-population/population-migration.aspx>

Wang, Lidong and Cheryl Ann Alexander (2014), "Telepsychiatry: Technology Progress, Challenges, and Language and Transcultural Issues," *Journal of Translational Medicine and Developmental Disorders*, 1 (1), 1-11.

Waugh, Maryann, Debbie Voyles and Marshall R. Thomas (2015), "Telepsychiatry: Benefits and Costs in a Changing Health-care Environment," *International Review of Psychiatry*, 27 (6), 558-568.

West Virginia University (2015), "Telepsychiatry Program," in *School of Medicine*. Retrieved from https://medicine.hsc.wvu.edu/media/1269/2015-telepsychiatry-flyer_final.pdf

Whealin, Julia M., L. Alana Seibert-Hatalsky, Jennifer Willett Howell and Jack Tsai (2015), "E-mental Health Preferences of Veterans with and without Probable Posttraumatic Stress Disorder," *Journal of Rehabilitation Research & Development*, 52 (6), 725-738.

Wynchank, Sinclair and Jill Fortuin (2010), "Telepsychiatry in South Africa – Present and Future," *South African Journal of Psychiatry*, 16-19.