


Euscorpius

Occasional Publications in Scorpiology



***Mesobuthus caucasicus* (Nordmann, 1840)
(Scorpiones: Buthidae) in Turkey**

Aysegül KARATAŞ

August 2005 – No. 25

Euscorpius

Occasional Publications in Scorpiology

EDITOR: Victor Fet, Marshall University, ‘fet@marshall.edu’

ASSOCIATE EDITOR: Michael E. Soleglad, ‘soleglad@la.znet.com’

Euscorpius is the first research publication completely devoted to scorpions (Arachnida: Scorpiones). *Euscorpius* takes advantage of the rapidly evolving medium of quick online publication, at the same time maintaining high research standards for the burgeoning field of scorpion science (scorpiology). *Euscorpius* is an expedient and viable medium for the publication of serious papers in scorpiology, including (but not limited to): systematics, evolution, ecology, biogeography, and general biology of scorpions. Review papers, descriptions of new taxa, faunistic surveys, lists of museum collections, and book reviews are welcome.

Derivatio Nominis

The name *Euscorpius* Thorell, 1876 refers to the most common genus of scorpions in the Mediterranean region and southern Europe (family Euscorpiidae).

Euscorpius is located on Website ‘<http://www.science.marshall.edu/fet/euscorpius/>’ at Marshall University, Huntington, WV 25755-2510, USA.

The International Code of Zoological Nomenclature (ICZN, 4th Edition, 1999) does not accept online texts as published work (Article 9.8); however, it accepts CD-ROM publications (Article 8). *Euscorpius* is produced in two *identical* versions: online (ISSN 1536-9307) and CD-ROM (ISSN 1536-9293). Only copies distributed on a CD-ROM from *Euscorpius* are considered published work in compliance with the ICZN, i.e. for the purposes of new names and new nomenclatural acts. All *Euscorpius* publications are distributed on a CD-ROM medium to the following museums/libraries:

- **ZR**, Zoological Record, York, UK
- **LC**, Library of Congress, Washington, DC, USA
- **USNM**, United States National Museum of Natural History (Smithsonian Institution), Washington, DC, USA
- **AMNH**, American Museum of Natural History, New York, USA
- **CAS**, California Academy of Sciences, San Francisco, USA
- **FMNH**, Field Museum of Natural History, Chicago, USA
- **MCZ**, Museum of Comparative Zoology, Cambridge, Massachusetts, USA
- **MNHN**, Museum National d'Histoire Naturelle, Paris, France
- **NMW**, Naturhistorisches Museum Wien, Vienna, Austria
- **BMNH**, British Museum of Natural History, London, England, UK
- **MZUC**, Museo Zoologico “La Specola” dell’Universita de Firenze, Florence, Italy
- **ZISP**, Zoological Institute, Russian Academy of Sciences, St. Petersburg, Russia
- **WAM**, Western Australian Museum, Perth, Australia
- **NTNU**, Norwegian University of Science and Technology, Trondheim, Norway

***Mesobuthus caucasicus* (Nordmann, 1840) (Scorpiones: Buthidae) in Turkey**

Aysegül KARATAŞ

Niğde Üniversitesi, Fen-Edebiyat Fakültesi, Biyoloji Bölümü, TR-51200 Niğde, TÜRKİYE
(Division of Natural Sciences, Niğde University, TR-51200 Niğde, TURKEY)
e-mail: karatash@scientist.com

Summary

Although *Mesobuthus caucasicus* was listed from Turkey in the literature, information about this species was confused and needed confirmation. Distribution of *M. caucasicus* is studied and new geographical records are given. As a result of field studies, *M. caucasicus* is confirmed from three certain localities in Turkey. In addition, some bio-ecological information is provided.

Introduction

Turkish scorpiofauna consists of 16 species belonging to the families Buthidae, Euscorpiidae, Iuridae, and Scorpionidae (Pavesi, 1876; Birula, 1898, 1899, 1917; Werner, 1902; Kulczyński, 1903; Schenkel, 1947; Vachon, 1947a, 1947b, 1971; Kinzelbach, 1975, 1980, 1982, 1984; Bonacina, 1980; Francke & Soleglad, 1981; Fet, 1986, 1989; Vachon & Kinzelbach, 1987; Crucitti, 1993, 1999; Lacroix, 1995; Kovařík, 1996; Fet & Braunwalder, 2000; Crucitti & Cicuzza, 2001; Karataş et al., 2004; Fet et al., in press). Buthidae, the richest family in the country, is represented by the genera *Androctonus*, *Buthacus*, *Compsobuthus*, *Hottentotta*, *Leiurus*, and *Mesobuthus*.

The genus *Mesobuthus* Vachon, 1950 currently includes 13 species. It is widespread in the Palaearctic Region from the Balkans to Korea. Four of these species have been recorded in Turkey: *M. caucasicus* (Normann, 1840), *M. eupeus* (C.L. Koch, 1839), *M. gibbosus* (Brullé, 1832), and *M. nigrocinctus* (Ehrenberg, 1828) (Birula, 1917; Kinzelbach, 1975; Vachon & Kinzelbach, 1987; Fet, 1989; Kovařík, 1996, 1998; Fet & Lowe, 2000; Fet et al., 2000; Gantenbein et al., 2000; Karataş, 2003; Karataş & Karataş, 2003; Crucitti & Vignoli, 2002). Of these, the range of *M. gibbosus* (Brullé, 1832) lies in Anatolia (to the west of the Anatolian Diagonal except for the coastal parts of the Black Sea and the Sea of Marmara) and the Balkans. *M. eupeus* is widespread in the Palearctic Region; it ranges from Central Anatolia through Caucasus and Turkestan to China and also found in Iran, Iraq, Afghanistan and Pakistan (Birula, 1917; Vachon, 1951, 1966; Kinzelbach, 1975, 1984; Va-

chon & Kinzelbach, 1987; Fet, 1989, Kovařík, 1996, 1998; Fet & Braunwalder, 2000; Karataş & Karataş, 2001, 2003). *M. gibbosus* has sympatric distribution with *M. eupeus* in Central Anatolia to the west of the Anatolian Diagonal (Kinzelbach, 1975, 1984; Vachon & Kinzelbach, 1987; Fet, 1989, Kovařík, 1996, 1998; Crucitti, 1999; Fet & Braunwalder, 2000; Crucitti & Vignoli, 2002; Karataş & Karataş, 2001, 2003). *M. nigrocinctus* in Turkey was first recorded from Adiyaman by Crucitti & Vignoli (2002); it also occurs in Gaziantep, Malatya, and Hatay Provinces (Karataş, unpubl.). To the east of the Anatolian Diagonal, *M. caucasicus* is syn-topic only with *M. eupeus* in İğdır, Kars, and Van Provinces (Figure 1). A recently described species *M. cyprius*, separated from *M. gibbosus* on the basis of DNA analysis, is found on Cyprus (Gantenbein et al., 2000).

The species *Mesobuthus caucasicus*, found from Turkey to China, was placed in the monotypic genus *Olivierus* by Farzanpay (1987) depending on the number of granules on movable finger of pedipalp. Later, DNA studies by Gantenbein et al. (2003) demonstrated that the genus *Olivierus* was paraphyletic with respect to genus *Mesobuthus*. Therefore, *Olivierus* was synonymized with *Mesobuthus*. Among six subspecies of *M. caucasicus*, Fet & Lowe (2000) listed the nominotypical subspecies as found in Turkey (see also Kovařík, 1998; Crucitti & Vignoli, 2002).

Presence of *M. caucasicus caucasicus* in Turkey was first reported by Birula (1897, 1917) as *Buthus caucasicus* from Aralik. Birula (1897, 1917) also gave morphological characteristics of the species and geographical distribution of the nominotypical subspecies. Vachon (1947a) listed *M. caucasicus* for the scorpion fauna of Turkey, but the study was not based on any specimen.




Figure 1: Map of new (circles) and previous (squares) localities of *M. caucasicus* from Turkey and ranges of *Mesobuthus* spp. in Turkey and surrounding area.

He also identified a female from İğdir Province as *Mesobuthus* sp. Other authors who provided information about *M. caucasicus* in Turkey (Birula, 1898; Vachon, 1947b, 1951; Kinzelbach, 1984; Crucitti, 1999; Fet & Lowe, 2000; Crucitti & Vignoli, 2002) apparently only followed the original information of Birula (1897) and Vachon (1947a). Due to our new data obtained from the field studies, presence of *M. caucasicus* in western Turkey is confirmed with certainty. Below, we provide measurements, drawings and distribution localities for the specimens of *M. caucasicus* from Turkey.

Material and Methods

Scorpions were collected under stones during the day time and placed into 70 % ethanol. Specimens have been deposited in the scorpion collection at Zoology Department of Niğde University (ZDNU). Measurements were made under 0.1 mm accuracy micrometric ocular and drawings were made under the stereomicroscope Olympus SZX9. All measurements are given in millimetres (mm).


Abbreviations

TL: Total Length; Cal: Carapax Length; CaAw: Carapax Anterior Width; CaPw: Carapax Posterior Width; FemL/W: Femur Length/Width; PatL/W: Patella Length/Width; ChL/W/D: Chela Length/Width/Depth; FfL: Fixed Finger Length; MfL: Movable Finger Length; MesL: Mesosoma Length; TailL: Tail Length; Metasomal Segments L/W/H: Length/Width/Height of Metasomal Segments I–V.

Results and Discussion

General features

Morphological characteristics are given here, which are applicable to adult males collected from Van Province (Figs. 2–5). General coloration of body light greenish-yellow (Fig. 6). Metasoma does not increase in width posteriorly. Denticles of the ventrolateral carinae on metasomal segment V somewhat pointed, regularly enlarged posteriorly but not curved outward (Fig. 5). Telson oblong, oval. Lyre-shaped carinal structure be-



Figures 2-5: External morphology of *M. caucasicus* male from Van Province. 2. Carapace carination; 3. Dentition of the fixed finger of pedipalp chela (left) and of the movable finger (right); 4. Right pedipalp chela, lateral (=external) aspect with trichobothrial pattern and scalloping of fingers; 5. Lateral view of metasomal segments III–V and telson. (Scale: 2 mm).

hind the ocular tubercle on carapace not prominent as in *M. eupeus* (Fig. 2). Posterior median carinae are not aligned with central lateral carinae and central median carinae. There are 12 and 13 oblique rows of denticles on pedipalp fixed and movable fingers, respectively (Fig. 3). External accessory denticles are smaller than the inner ones, especially towards fingertips. Basal lobe (scalloping of fingers) is well-developed (Fig. 4). In contrast with *M. eupeus* and *M. gibbosus*, mesosoma of *M. caucasicus* does not have dark stripes. Only the areas between median eyes and around lateral eyes have dark

coloration. Basal two-thirds of metasomal segment V are brownish ventrally and laterally. Legs are yellowish.

Metasomal segment I with 10 complete carinae, segments II and III with eight complete and two lateral accessory carinae. These accessory carinae reach to the middle of segment II but it consist of only 2–3 granules on segment III. Segment IV has eight complete carinae and no accessory carinae.

Metasomal segments in *M. caucasicus* are longer than that of *M. eupeus*. Ventral carinae of metasomal segments I–V are of equal size in *M. caucasicus* but



Figure 6: *M. caucasicus* adult male from Van Prov. Dorsal (above) and ventral (below) views.

enlarged posteriorly, especially on II and III metasomal segments, in *M. eupeus*. *M. caucasicus* has more elongated carapace than that of *M. eupeus* because, with regard to posterior width of carapace, length of carapace and anterior width of carapace are shorter than that of *M. caucasicus*. Lyre-shaped carinal structure on the carapace is more prominent in *M. eupeus* than in *M. caucasicus*. Number of oblique rows of denticles on pedipalp fingers is 10–11 in *M. eupeus*, 12–13 in *M. caucasicus*.

The localities from Turkey clarify the southwestern range boundary of *M. caucasicus*. Birula (1897) reported *Buthus caucasicus* from Aralik, the locality currently situated at the junction of Turkey, Armenia, and Iran (Fet, 1989; Fig. 1). Birula (1917) listed four subspecies within the geographic range of *M. caucasicus*. Of these, the distribution of the nominotypic *M. c. caucasicus* included Eastern Transcaucasia, Daghestan, the Terek Region and Northern Persia. Our measurements agree with the dimensions of adult males giving in Birula (1917). According to the key published by Birula (1917) to distinguish the subspecies of *M. caucasicus*, our specimens correspond to the nominotypic subspecies. Fet (1989), in the catalogue of scorpions of the former USSR, listed the nominotypic subspecies from the modern Armenia, Azerbaijan, Georgia, Russia, and Ukraine. Fet & Lowe (2000) listed the nominotypic subspecies from Armenia, Azerbaijan, Georgia, Turkey and Iraq (?), Russia (Chechnya, Daghestan) and Ukraine (Kherson and Odessa Regions).

Presence of *M. caucasicus* in Turkey is now clearly confirmed by morphological characteristics of specimens as given in this study.

Bio-ecological observations

All specimens were taken from very dry habitats under the stones, remote from human dwellings. İğdir specimens were collected under the stones in the ruins of a house made of concrete.

Kars and Van specimens were observed under the stones in the field covered with steppe vegetation and on the limestone formation. Two adult ♀♂ from Karakoyunlu (İğdir Province) were observed. Each of two females carrying nearly 45 young at first instars on its back in the field on 20.VII.2003 hence they were not taken as field material.

Examined specimens and localities (Fig. 1): 1. *İğdir Province*: Karakoyunlu, Taşburun village ($39^{\circ}58'52''$ N; $44^{\circ}14'40''$ E) (860 m a.s.l.), 20.VII.2003: 2 adult ♀♀; 2. *Kars Province*: Digor, Halimcan village ($40^{\circ}09'24''$ N; $43^{\circ}39'32''$ E) (ca. 1200 m a.s.l.), 21.VII.2003: 1 subadult ♂; 3. *Van Province*: East of Van Lake ($38^{\circ}32'$ N; $43^{\circ}21'$ E) (ca. 1700 m a.s.l.), 15.VI.2001: 2 adult ♂♂.

Literature record: 4. *İğdir Province*: Aralik ($39^{\circ}52'$ N; $44^{\circ}31'$ E) (Birula, 1897).

Morphometric measurements of *M. caucasicus*: Van Province, 2 ♂♂; TL 55.8, 58.9; CaL 5.4, 5.2; CaAw 3, 3.2; CaPw 5.8, 5.5; FemL/W 5/1.6, 4.9/1.6; PatL/W 6.1/2.4, 5.9/2.3; ChL/W/D 10.3/2.6/3.1, 10.3/2.6/3; FfL 5.5, 5.5; MfL 6.6, 6.5; MesL 14.3, 15.3; TailL 34.5, 34.3; Metasomal segments; I, L/W/H 4/3.6/3.1, 3.9/3.4/3; II, 4.1/3.5/3.1, 4.6/3.3/3; III, 4.8/3.4/3.1, 4.9/3.3/3; IV, 5.5/3.2/3, 5.5/3.2/2.9; V, 6/2.9/2.5, 6.3/2.9/2.5, Pectinal tooth counts 25–28 and 27–28. Measurements of the subadult ♂ from Kars and two ♀♀ from İğdir are not given; pectinal tooth counts of the subadult ♂ from Kars were 29–30. İğdir females were not taken from the field because they were carrying newborn litters; they were released under the stones.

Acknowledgements

I am grateful to Dr. Victor Fet and an anonymous reviewer for valuable comments and reviewing the manuscript. Also I wish to thank Dr. Ahmet Karataş for preparing the map of localities and to both of our students and colleagues, Münnir Uçak and Mustafa Kürtüllü, for their help in field studies.

References

- BIRULA, A.A. 1897. Miscellanea scorpionologica. II. Zur Synonymie der russischen Skorpione (Forsetzung). *Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de St.-Pétersbourg*, 2: 377–391.
- BIRULA, A.A. 1898. Ein Beitrag zur Kenntniss der Skorpionenfauna Kleinasiens. *Horae Societatis Entomologicae Rossiae*, 33(1–2): 132–140.
- BIRULA, A.A. 1899. [A new species of scorpions for the Russian fauna]. *Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de St.-Pétersbourg*, 4: XIV–XV [in Russian, Latin diagnosis].
- BIRULA, A.A. [BYALYNITSKII-BIRULYA, A.A.]. 1917. *Chlenistobryukhie paukoobraznye Kavkazskogo Kraja [Arthrogastric Arachnids of Caucasia]. Part I. Scorpiones*. Tiflis: *Mémoires du Musée du Caucase*, Sér. A 5: 1–253. [English translation: Jerusalem: Israel Program for Scientific Translation, 1964].
- BONACINA, A. 1980. Sistematica specifica e sottospecifica del complesso “*Euscorpius germanus*”

- (Scorpiones, Chactidae). *Rivista del Museo Civico di Scienze Naturali "E. Caffi" Bergamo*, 2: 47–78.
- CRUCITTI, P. 1993. Some topics on distribution patterns of the genus *Mesobuthus* in the Near East based on ecological data (Scorpiones: Buthidae). *Biologia Gallo-hellenica*, 20: 69–74.
- CRUCITTI, P. 1999. The scorpions of Anatolia: Biogeographical patterns. *Biogeographica*, 20: 81–94.
- CRUCITTI, P. & D. CICUZZA. 2001. Scorpions of Anatolia: Ecological patterns. Pp. 225–234 in Fet, V. & P.A. Selden (eds.). *Scorpions 2001: In Memoriam Gary A. Polis*. Burnham Beeches, Bucks: British Arachnological Society.
- CRUCITTI, P. & V. VIGNOLI. (2001) 2002. Gli Scorpioni (Scorpiones) dell'Anatolia sud-orientale (Turchia). *Bulletino di Museo regionale di Scienze naturali Torino*, 19 (2): 433–480.
- FARZANPAY, R. 1987 (1366). [Knowing scorpions]. Teheran: Central University Publications, No. 312, Biology 4, 231 pp. [in Farsi, with Latin index].
- FET, V.Ya. 1986. Notes on some *Euscorpius* (Scorpiones: Chactidae) from Greece and Turkey. *Rivista del Museo Civico di Scienze Naturali "E. Caffi" Bergamo*, 9: 3–11.
- FET, V. 1989. A catalogue of scorpions (Chelicerata: Scorpiones) of the USSR. *Rivista del Museo Civico di Scienze Naturali "E. Caffi" Bergamo*, 13 (1988): 73–171.
- FET, V. & M.E. BRAUNWALDER. 2000. The scorpions (Arachnida, Scorpiones) of the Aegean area: current problems in taxonomy and biogeography. *Belgian Journal of Zoology*, 130 (Suppl. 1): 17–22.
- FET, V., B. GANTENBEIN, Ay. KARATAŞ & A. KARATAŞ. (in press). An extremely low genetic divergence across the range of *Euscorpius italicus* (Scorpiones: Euscorpiidae). *Journal of Arachnology*.
- FET, V., B.E. HENDRIXSON, W.D. SISSOM & G. LEVY. 2000. First record for the genus *Mesobuthus* Vachon, 1950 in Israel: *Mesobuthus nigrocinctus* (Ehrenberg, 1828), n. comb. (Scorpiones: Buthidae) from Mt. Hermon. *Israel Journal of Zoology*, 46: 287–295.
- FET, V. & G. LOWE. 2000. Family Buthidae. Pp. 54–286 in FET, V., W.D. SISSOM, G. LOWE & M.E. BRAUNWALDER. 2000. *Catalog of the Scorpions of the World (1758–1998)*. New York, NY: The New York Entomological Society, 690 pp.
- FRANCKE, O.F. & M.E. SOLEGLAD. 1981. The family Iuridae Thorell (Arachnida, Scorpiones). *Journal of Arachnology*, 9: 233–258.
- GANTENBEIN, B., C. KROPF, C.R. LARGIADÈR & A. SCHOLL. 2000. Molecular and morphological evidence for the presence of a new buthid taxon (Scorpiones: Buthidae) on the Island of Cyprus. *Revue suisse de Zoologie*, 107(1): 213–232.
- GANTENBEIN, B., V. FET & A.V. GROMOV. 2003. The first DNA phylogeny of four species of *Mesobuthus* Vachon, 1950 (Scorpiones: Buthidae) from Eurasia. *Journal of Arachnology*, 31(3): 412–420.
- KARATAŞ, A. & Ah. KARATAŞ. 2001. *Mesobuthus eupeus* (C.L. Koch, 1839) from central Anatolia (Scorpiones: Buthidae). Pp. 297–299 in Fet, V. & P.A. Selden (eds.). *Scorpions 2001: In Memoriam Gary A. Polis*. Burnham Beeches, Bucks: British Arachnological Society, UK, 450 pp.
- KARATAŞ, A., F. ÇALIŞKAN, H. ÇALIŞKAN, Ü. ŞİRİN & Ah. KARATAŞ. 2004. The scorpion fauna of Turkey. VII. *Reunion International de Expertos en Envenenamiento por Animales Ponzonosos Cuernavaca*, Morelos, Mexico, Abstracts, p. 119.
- KARATAŞ, A. 2003. New records on the occurrence of *Hottentotta saulcyi* (Simon, 1880) (Scorpiones: Buthidae) in Turkey. *Israel Journal of Zoology*, 49(4): 315–316.
- KARATAŞ, A. & Ah. KARATAŞ. 2003. *Mesobuthus eupeus* (C.L. Koch, 1839) (Scorpiones: Buthidae) in Turkey. *Euscorpius*, 7: 1–6.
- KINZELBACH, R. 1975. Die Skorpione der Ägäis: Beiträge zur Systematik, Phylogenie und Biogeographie. *Zoologische Jahrbücher, Abteilung für Systematik*, 102: 12–50.
- KINZELBACH, R. 1980. Zur Kenntnis des kaukasischen Skorpions *Calchas nordmanni* Birula, 1899 (Scorpionida: Chactidae). *Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg (NF)*, 23: 169–174.
- KINZELBACH, R. 1982. Die Skorpionssammlung des Naturhistorischen Museums der Stadt Mainz. Teil I: Europa und Anatolien. *Mainzer Naturwissenschaftliches Archiv*, 20: 49–66.

- KINZELBACH, R. 1984. Die Skorpionssammlung des Naturhistorischen Museums der Stadt Mainz. Teil II: Vorderasien. *Mainzer Naturwissenschaftliches Archiv*, 22: 97–106.
- KOVAŘÍK, F. 1996. First report of *Compsobuthus matthiesseni* (Scorpionida: Buthidae) from Turkey. *Klapalekiana*, 32: 53–55.
- KOVAŘÍK, F. 1998. *Štíři [Scorpions]*. Jihlava: Mada-gaskar, 175 pp.
- KULCZYŃSKI, W. 1903. Arachnoidea in Asia Minore et ad Constantinopolim a Dre. F. Werner collecta. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften (Wien), Mathematisch-Naturwissenschaftliche Klasse. Abt. I*, 112 (1): 627–680 + i pl.
- LACROIX, J.-B. 1995. *Euscorpius (E.) mingrelicus* Kessler, 1876 en Turquie Anatolienne (Arachnida: Scorpionida). *Arachnides*, 26: 4–6.
- PAVESI, P. 1876. Gli Aracnidi Turchi. *Atti della Società Italiana di Scienze Naturali*, 16: 24–54.
- SCHENKEL, E. 1947. Einige Mitteilungen über Spinnentiere. *Revue suisse de Zoologie*, 54(1): 13–16.
- VACHON, M. 1947a. Remarques préliminaires sur le faune des Scorpions de Turquie. *Bulletin du Muséum National d'Histoire Naturelle, Paris*, 19(2): 161–164.
- VACHON, M. 1947b. Repartition et origine des scorpions de Turquie. *C. R. Société de Biogéographie*, 206(3): 26–29.
- VACHON, M. 1951. A propos de quelques scorpions de Turquie collectés par M. le Professeur Dr. Curt Kosswig. *İstanbul Üniversitesi Fen Fakültesi Mecmuası (Revue de la Faculté des Sciences de l'Université d'Istanbul)*, 16 (B): 341–344.
- VACHON, M. 1971. [Remarques sur le scorpion caucasiens *Calchas nordmanni* Birula (Scorpiones, Chactidae)]. *Entomologicheskoe Obozrenije (Revue d'Entomologie de l'URSS)*, 13: 712–718 [in Russian].
- VACHON, M. & R. KINZELBACH. 1987. On the taxonomy and distribution of the scorpions of the Middle East. Pp. 91–103 in Krupp, F., W. Schneider & R. Kinzelbach (eds.). *Proceedings of the Symposium on the Fauna and Zoogeography of the Middle East*, Mainz, 1985.
- WERNER, F. 1902. Ueber die Faunengebiete Kleinasiens. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien*, 52: 7–9.