



University of Richmond
UR Scholarship Repository

Physics Faculty Publications

Physics

12-11-2010

Sources of the Radio Background Considered

Jack Singal


University of Richmond, jsingal@richmond.edu

L. Stawarz

A. Lawrence

V. Petrosian

Follow this and additional works at: <http://scholarship.richmond.edu/physics-faculty-publications>

 Part of the [External Galaxies Commons](#), and the [Stars, Interstellar Medium and the Galaxy Commons](#)

Recommended Citation

Singal, J., L. Stawarz, A. Lawrence, and V. Petrosian. "Sources of the Radio Background Considered." *Monthly Notices of the Royal Astronomical Society* 409, no. 3 (December 11, 2010): 1172-182. doi:10.1111/j.1365-2966.2010.17382.x.

This Article is brought to you for free and open access by the Physics at UR Scholarship Repository. It has been accepted for inclusion in Physics Faculty Publications by an authorized administrator of UR Scholarship Repository. For more information, please contact scholarshiprepository@richmond.edu.

Sources of the radio background considered

J. Singal,^{1*} Ł. Stawarz,^{1,2,3} A. Lawrence^{4†} and V. Petrosian^{1‡}

¹Kavli Institute for Particle Astrophysics and Cosmology, SLAC National Accelerator Laboratory and Stanford University, 382 Via Pueblo Mall, Stanford, CA 94305-4060, USA

²Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo-ku, Sagami-hara, Kanagawa 252-5510, Japan

³Astronomical Observatory of the Jagiellonian University, ul. Orła 171, 30-244 Kraków, Poland

⁴University of Edinburgh Institute for Astronomy, Scottish Universities Physics Alliance, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ

Accepted 2010 July 17. Received 2010 July 9; in original form 2010 June 7

ABSTRACT

We investigate different scenarios for the origin of the extragalactic radio background. The surface brightness of the background, as reported by the Absolute Radiometer for Cosmology, Astrophysics, and Diffuse Emission 2 (ARCADE 2) collaboration, is several times higher than that which would result from currently observed radio sources. We consider contributions to the background from diffuse synchrotron emission from clusters and the intergalactic medium, previously unrecognized flux from low-surface-brightness regions of radio sources and faint point sources below the flux limit of existing surveys. By examining radio source counts available in the literature, we conclude that most of the radio background is produced by radio point sources that dominate at sub- μ Jy fluxes. We show that a truly diffuse background produced by electrons far from galaxies is ruled out because such energetic electrons would overproduce the observed X-ray/ γ -ray background through inverse Compton scattering of the other photon fields. Unrecognized flux from low-surface-brightness regions of extended radio sources, or moderate flux sources missed entirely by radio source count surveys, cannot explain the bulk of the observed background but may contribute as much as 10 per cent. We consider both radio supernovae and radio-quiet quasars as candidate sources for the background and show that both fail to produce it at the observed level because of an insufficient number of objects and total flux, although radio-quiet quasars contribute at the level of at least a few per cent. We conclude that if the radio background is at the level reported, a majority of the total surface brightness would have to be produced by ordinary star-forming galaxies above redshift 1 characterized by an evolving radio–far-infrared correlation, which changes towards the radio loud with redshift.

Key words: diffuse radiation – radio continuum: general – radio continuum: ISM.

1 INTRODUCTION

Recent results from the Absolute Radiometer for Cosmology, Astrophysics, and Diffuse Emission (ARCADE) 2 project suggest a radio background several times brighter than can be produced by currently observed radio sources (Fixsen et al. 2010; Seiffert et al. 2010; Singal et al. 2010). The extragalactic background, detected from 22 MHz to 8 GHz, has a power-law spectral index of $\alpha \simeq 0.6$ (defined here as $S_\nu \propto \nu^{-\alpha}$, where S_ν is the spectral flux density) and a brightness temperature of 1.17 K at 1 GHz, corresponding to a cosmic radio background (CRB) intensity of

$7 \times 10^{-22} \text{ W Hz}^{-1} \text{ m}^{-2} \text{ sr}^{-1} = 3.6 \times 10^4 \text{ Jy sr}^{-1}$. A significant contribution to the background from free–free emission has been ruled out based on the spectral shape (Seiffert et al. 2010).

It is unlikely that the radio signal reported by the ARCADE 2 collaboration is Galactic or local in origin. As discussed in Kogut et al. (2010), the extragalactic component is separated from the diffuse Galactic foreground by two independent robust indicators, a correlation of radio with C II emission and a cosecant dependence on the Galactic latitude. The inferred residual extragalactic component is several times brighter than the high latitude Galactic level. Comparisons with observed edge-on galaxies indicate that our Galaxy would be quite anomalous to support a radio-emitting halo of the intensity needed to explain the signal.¹ Additionally, assuming a

*E-mail: jsingal@stanford.edu

†Visiting physicist at KIPAC/SLAC National Accelerator Laboratory and Stanford University, USA.

‡Also Departments of Physics and Applied Physics, Stanford University, Stanford, CA 94305, USA.

¹ One may add here that this is true even taking into account the presence of ‘anomalous large-scale radio continuum features’ present in some edge-on spirals (Hummel, van Gorkom & Kotanyi 1983; Elmouttie et al. 1995).

diffuse Galactic origin for the measured radio signal would require a source of radio emission that does not follow the correlation with far-infrared (FIR) emission observed in local galaxies, again making our Galaxy anomalous. Isotropic radio emission from a local region on the scale of the local bubble accounting for the level observed would manifest an all-sky quadrupole polarization pattern at a level visible in *Wilkinson Microwave Anisotropy Probe* 23-GHz data, but such a pattern is not seen. These lines of evidence are summarized in Kogut et al. (2010). Furthermore, the considerations of Section 3 in this paper can be used to place a strong constraint on the amount of the total observed high latitude radio emission that can be Galactic in origin, as in the halo inverse Compton (IC) scattering on the ambient light would produce a much stronger X-ray background than observed, again strongly disfavoring a Galactic origin as an explanation for the signal. In this paper, then, we investigate possible origins for the radio background, assuming that the extragalactic level is that reported by the ARCADE 2 collaboration, although we conclude that producing the background at the level observed is difficult without a mechanism to cause significant redshift evolution in the radio to FIR correlation beyond what has been observed thus far.

Interferometric surveys have achieved a relatively consistent picture of radio source counts above the 10- μ Jy level. At high fluxes, say between 1 mJy and 1 Jy at 1.4 GHz, radio-loud active galactic nuclei (AGN) dominate radio source counts (e.g. Windhorst et al. 1993; Condon 2007), with their differential counts following a power law $dN/dS \propto S^{-\gamma}$ with $\gamma < 2.5$ (the so-called Euclidean value). Below 1 mJy, this trend tends to reverse and the source counts show a more rapid increase with decreasing flux, perhaps indicating emergence of a new population which does not contribute at higher fluxes but dominates in this regime (e.g. Hopkins et al. 2003; Simpson et al. 2006; Condon 2007). Gervasi et al. (2008) have calculated the radio brightness resulting from fitting a two-population model and extrapolating available radio source counts to fainter fluxes at frequencies from 150 MHz to 8.5 GHz. They derive a surface brightness that is three to six times smaller than that reported by the ARCADE 2 collaboration. Clearly, there must be significant contributions to the radio background from sources not considered by Gervasi et al. (2008).

ARCADE 2 and other instruments used to determine the diffuse background intensity have resolutions greater than 1° and cannot distinguish between a background due to discrete sources of angular sizes smaller than 1° , and ones that are truly diffuse. Therefore, we consider several candidates for comprising the background.

The emission mechanism of sources producing the CRB must be synchrotron radiation by relativistic electrons. This puts powerful constraints on potential sources. They must not overproduce the measured diffuse FIR background if related to star-forming activity nor overproduce the measured diffuse X-ray/ γ -ray background through IC scattering of the cosmic microwave background (CMB) and other background photon fields by the same electrons. Additionally, the source spectra must be consistent with the CRB power-law spectrum of index 0.6, mentioned above.

In this paper, we investigate several kinds of possible candidates. In Section 2 we analyse the existing data from radio source counts, and using some approximate fitting procedures estimate the contribution from sources observed in numerous surveys. In Section 3, we use the X-ray/ γ -ray background to limit the contribution from more diffuse emission, clusters and the intergalactic medium (IGM), and in Section 4 we examine whether source count surveys have missed a significant amount of flux. In Sections 5.1 and 5.2 we explore the potential contribution from radio supernovae (RSNe) and radio-

quiet (RQ) quasars, and finally in Section 5.3 we conclude that emission from star-forming galaxies may account for bulk of the measured radio background, but only if the radio/FIR luminosity ratio increases with redshift. A brief summary and discussion is presented in Section 6.

2 RADIO SOURCE COUNTS

Fig. 1 shows a schematic but fairly accurate depiction of radio source counts available in the literature (see Table 1). In the top panel of the figure, we plot the $S^{2.5} dN/dS$ distributions given in different radio surveys (note that the differential count is per steradian per unit flux). The observations were conducted at a range of frequencies (0.151–8.5 GHz). We have converted all fluxes to 1.4 GHz assuming a general power-law spectrum with an average (redshift-independent) spectral index equal to 0.75, the canonical value of the extragalactic radio sources. Note that even though the surveys considered have a wide range of angular resolution (spanning from




Figure 1. Top: schematic description of the observed $S^{2.5} dN/dS$ distribution from different radio surveys. Source counts from 8.4 GHz surveys are in dark blue, 2.7 GHz in light blue, 5 GHz in yellow, 1.4 GHz in green, 610 MHz in pink, 408 MHz in red and 151 MHz in black. Details of the radio surveys used with references are given in Table 1. Data from all frequencies are normalized to flux bins at 1.4 GHz using a 0.75 spectral index. Bottom: the $S^2 dN/dS$ distribution representing the surface brightness per logarithmic flux bin of the same surveys, divided by the surface brightness of CRB at 1.4 GHz reported by the ARCADE collaboration. The source counts tell roughly a consistent story over a wide range of frequencies and resolutions. The peak contribution at ~ 100 mJy comes from bright radio galaxies. The contribution decreases with decreasing flux density, but begins to rise again below ~ 1 mJy as a new population of sources becomes important. Integrating over the entire flux range probed by these surveys gives ~ 26 per cent of the total surface brightness of the CRB at 1.4 GHz.

Table 1. Radio source count data plotted in Fig. 1.

Survey	Frequency (GHz)	Facility ^a	Resolution ^b (arcsec)	S _{limit} ^c (μJy)	γ ^d	k ^d	S _{max} ^e (mJy)
Fomalont et al. (2002) ^f	8.4	VLA	6	7.5	2.11 ± 0.13 ^g	16.8	1
Windhorst et al. (1993)	8.4	VLA	3.5	35.0			
			10	14.5	2.3 ± 0.2 ^g	4.6	1
Ciliegi et al. (2003)	5	VLA	4	50	2.0 ^g	131.7	20
Prandoni et al. (2006)	5	ATCA	11	400	1.93 ^h	289.7	33.5
Gruppioni et al. (1997)	2.7	ATCA	39	400	1.58 ^h	2173.8	21.5
Richards (2000)	1.4	VLA	2	40	2.38 ± 0.13 ^g	8.25	1
Hopkins et al. (2003)	1.4	ATCA	6×12	60	1.83 (≤6 mJy) ⁱ	–	1000
					1.89 (≥6 mJy)	–	
Bondi et al. (2003)	1.4	VLA	31	60	2.28 (≤6 mJy) ^g	27.3	40
					1.79 (≥6 mJy)	1062.3	
Windhorst, va Heerde & Katgert (1984)	1.4	WSRT	12.5	600	1.9 (≤1 mJy) ⁱ	341	10 000
Windhorst et al. (1985)	1.4	VLA	14.7	225	2.1 (≤1 mJy) ^h	175.3	100
					1.9 (≥1 mJy)	867.5	
Fomalont et al. (2006)	1.4	VLA A	1.8	25	2.43 ^g	9.2	10 000
Owen & Morrison (2008)	1.4	VLA A	1.6	15	2.61 ^h	2.0	1500
Simpson et al. (2006)	1.4	VLA C	5×4	100	1.9 ^h	505.2	10 000
Biggs & Ivison (2006) ^f	1.4	VLA A	1.6	15	2.24 ^h	33.1	1153
			1.5	48	2.37 ^h	12.9	1327
			1.5	29	2.44 ^h	7.2	854
Ibar et al. (2009)	1.4	VLA B	4.3×4.2	20	2.08 ^h	38.4	3000
Ibar et al. (2009)	.610	GMRT	7.1×5.6	45	2.09 ^h	284.4	1000
Bondi et al. (2007)	.610	VLA	6	150	1.84 ^h	1346.2	200
Katgert-Merkelijn et al. (1985)	.610	WSRT	57×57	9000	1.94 ^h	1158	900
Wilson & Vallee (1982)	.610	WSRT	60	10000	1.94 ^g	1039	516
Vallée & Roger (1989)	.408	PSRT	204×276	70000	1.9 ± 0.1 ^g	2200	516
Grueff (1988)	.408	NCTRT	156×288	70000	2.28 ^h	1072	10 000
Pearson & Kus (1978)	.408	CMBR	80	10000	1.8 ^h	2284	10 000
Hales, Baldwin & Warner (1988)	.151	CMBR	300	120000	1.9 (≤1 Jy) ^h	3824.3	20 000
					2.3 (≥1 Jy)	3421.2	
McGilchrist et al. (1990)	.151	CMBR	70×100	80000	2.07 (≤1 Jy) ^h	3777.4	10 000
					2.68 (≥1 Jy)	3874.5	

^aVLA = Very Large Array, ATCA = Australia Telescope Compact Array, WSRT = Westerbork Synthesis Radio Telescope, PSRT = Penticon Synthesis Radio Telescope, NCTRT = Northern Cross Transit Radio Telescope, CMBR = Cambridge Radio Telescope Survey.

^bThe resolution quoted here is the full width at half-maximum of the composite image. If the beam is not round, values for two axes are given.

^cThe low-flux limit of the survey is a factor determined by the survey authors (usually five) times the rms noise.

^dFrom fits to dN/dS of the form $dN/dS = k \times S^{-\gamma}$, where the dN/dS distribution is expressed in $\text{sr}^{-1} \text{Jy}^{-1}$, and the flux S in Jy.

^eThe highest flux object observed or the highest flux listed in a dN/dS table or fit to dN/dS .

^fSeparate results for different fields reported.

^gSource counts power law directly given.

^h dN/dS given in table and power law determined with fit.

ⁱSource counts given by equation in S .

1 to 300 arcsec) there is not only a clear agreement among the counts at $S > 1$ mJy, but also in regard to the presence of the – relatively poorly constrained – low-flux population emerging at $S < 1$ mJy.

In the bottom panel of Fig. 1, we plot a proxy for 1.4-GHz total surface brightness due to radio sources, $S^2 (dN/dS)$, divided by the observed CRB brightness as reported by the ARCADE collaboration in Fixsen et al. (2010), i.e.

$$\frac{S^2 (dN/dS)}{B_{\text{CRB}}(\nu)} \Big|_{1.4\text{GHz}} \equiv \frac{S^2 (dN/dS)}{2k\nu^2 c^{-2} T_{\text{CRB}}(\nu)} \Big|_{1.4\text{GHz}}, \quad (1)$$

where

$$T_{\text{CRB}}(\nu) = 1.17 \times \left(\frac{\nu}{\text{GHz}} \right)^{-2.6} \text{K}. \quad (2)$$

The true surface brightness

$$B_{\text{counts}}(S) = \int_S^\infty S \frac{dN}{dS} dS = \frac{1}{(\gamma - 2)} S^2 \frac{dN}{dS} \quad (3)$$

for a power law $dN/dS \equiv S^{-\gamma}$ is equal to the area under the $S^2 (dN/dS)$ curve, and thus the bottom panel of Fig. 1 provides directly a minimum fractional contribution to the CRB from the resolved objects. We estimate this fraction to be $\simeq 26$ per cent. We note that the high-flux population is dominated by bright radio-loud AGN while the low-flux one is generally thought to be dominated by star-forming galaxies (see Ballantyne 2009, and references therein), with an additional possible contribution from RQ and low-radio-power AGN (e.g. Ibar et al. 2009; Padovani et al. 2009). Surveys probing fluxes as low as a few μJy are consistent with the two-population model (e.g. Fomalont et al. 2002; Biggs & Ivison 2006).

2.1 High-flux population

The integrated (fractional) contribution of the high-flux population ($S > S_0 \simeq 1$ mJy) to the measured CRB, $\int_{S_0}^\infty dS S (dN/dS) / B_{\text{CRB}}$, is about $\simeq 16$ per cent at 1.4 GHz, and the corresponding surface

number density $N(> S_0) = \int_{S_0}^{\infty} dS (dN/dS) \simeq 4.7 \times 10^5 \text{ sr}^{-1}$. One could ask if a significant portion of the flux from the high-flux population has been missed by radio surveys. If this were the case, then the contribution of the high-flux population to the measured background would have been underestimated. One possible source of the missing flux could be extended low-surface-brightness sources. However, as mentioned above, the surveys included in Fig. 1, in spite of the fact that they span a wide range of resolutions and are obtained from different interferometer arrays, show very good agreement on the integrated contribution to the background. We will return to the contribution of low-surface-brightness sources and quantify the possible missed flux further in Section 4.

2.2 Low-flux population

Unlike the high-flux population, the total contribution of the low-flux population is not fixed by existing surveys, since these do not constrain the low flux peak on the $S^2(dN/dS)$ – S plot. Even though at low fluxes ($S < S_0$) source counts indicate a power-law distribution $dN/dS \propto S^{-\gamma}$, the measured faint-end index γ varies substantially between different surveys, ranging from $\gamma = 2.11$ claimed by Fomalont et al. (2002) to $\gamma = 2.61$ claimed by Owen & Morrison (2008). The value of $\gamma = 2.5$ is what would be seen in a static, isotropic and homogeneous Euclidean universe. In extrapolating to lower fluxes, as long as the faint-end index is above 2, the integrated contribution from lower flux sources to the background will continue to increase. Obviously, at some eventual lower flux, the faint-end index must drop below 2, to avoid Olbers' paradox.

Assuming

$$\frac{dN}{dS} = k_0 \left(\frac{S}{S_0} \right)^{-\gamma} \quad \text{for } S_{\min} < S < S_0, \quad (4)$$

one can calculate the required value of S_{\min} as a function of γ such that the sources in this range provide the rest of the CRB. It is easy to show that for $\gamma > 2$

$$S_{\min} = S_0 \left[\frac{(1-H)(\gamma-2)B_{\text{CRB}}}{k_0 S_0^2} + 1 \right]^{1/(2-\gamma)}, \quad (5)$$

where the factor H is the estimated fractional contribution from sources above $S_0 \simeq 1 \text{ mJy}$, which we determine to be 0.16. The top panel of Fig. 2 shows the variation of S_{\min} with γ at $\nu = 1.4 \text{ GHz}$. We can also estimate the corresponding minimum number of required sources to be

$$N(> S_{\min}) - N(> S_0) = \frac{k_0 S_0^{1-\gamma}}{\gamma-1} \left[\left(\frac{S_{\min}}{S_0} \right)^{1-\gamma} - 1 \right], \quad (6)$$

whose dependence on γ is shown in the bottom panel of Fig. 2. Thus, if the background is to be made primarily from low-flux sources, their faint-end index below 1 mJy should be close to $\gamma = 2.5$ in order to not exceed reasonable estimates for the total number of non-dwarf galaxies in the observable Universe. In other words, in order for low-flux density sources to account for the observed CRB, they must reach low-flux values of less than 1 μJy and have a surface density of $\gtrsim 10^{10} \text{ sr}^{-1}$, but not much higher than this value.

A modelling of the dN/dS distribution of the low-flux ($< \text{mJy}$) population in terms of a single power law, although customary, is more than likely an over-simplification. A more realistic description would involve a smoothly curved function with the integrated contribution per log flux bin (Fig. 1, bottom panel) peaking around S_{\min} and then falling off, but including some contribution from objects




Figure 2. Top: the lower flux density of sources S_{\min} at 1.4 GHz required so that the extrapolated source counts from $S_0 \simeq 1 \text{ mJy}$, with a faint-end index γ , can account for observed CRB as reported by the ARCADE 2 collaboration, including a 16 per cent (solid lines) or 25 per cent (dashed line; see Section 4) contribution to the background from $S \geq S_0$ sources. The upper and lower solid curves are for the two extreme values of the 1-mJy normalizations k_0 of the surveys we consider. For $\gamma \sim 2.5$ an extrapolation to $S_{\min} \sim 0.1 \mu\text{Jy}$ is needed, while the required lower flux density falls rapidly with a decreasing faint-end index. Bottom: same as the top panel, but for the minimum number of sources below S_0 needed to produce the observed CRB.

with fluxes lower than S_{\min} . The large population of dwarf galaxies may be the sources that increasingly become relevant at fluxes below S_{\min} .

3 DIFFUSE SOURCES

In this section, we consider the possibility that the CRB may result from truly diffuse emission associated with the large-scale structure of the Universe, such as the intracluster medium (ICM), IGM or filaments connecting the clusters containing warm-hot gas. First, we consider very general constraints and then look at specific possibilities.

3.1 General constraints on diffuse emission

The simplest constraint on any population of electrons producing the CRB is that it must have a relatively flat energy spectrum, in order to produce the observed spectral index of $\alpha \sim 0.6$. Less obviously, we show below that it must be associated with a magnetic field of at least 1 μG . This is because otherwise the magnetic field energy density in such systems will be much less than the energy density of the CMB and other background radiations. As a result, the

relativistic electrons responsible for the radio emission would lose most of their energy producing hard X-ray and gamma radiation via IC scattering of the other background fields. We now derive this limit more carefully.

Given the observed power-law spectrum, the energy density of the radio background per frequency dex at ν_r is given by

$$[\nu_r U_{\nu_r}] = \frac{4\pi}{c} [\nu_r B_{\text{CRB}}(\nu_r)] = 1.17 \frac{8\pi k_B \nu_*^3}{c^3} \left(\frac{\nu_r}{\nu_*} \right)^{0.4}, \quad (7)$$

where $\nu_* = 1$ GHz. We are considering the background spanning frequencies from $\nu_1 \sim 1$ MHz to $\nu_2 \sim 10$ GHz.

For a density of ultrarelativistic electrons with a power-law energy spectrum

$$n_e(\gamma_e) = k_e \gamma_e^{-s} \quad \text{for } \gamma_{e1} \leq \gamma_e \leq \gamma_{e2}, \quad (8)$$

where k_e is a normalization constant in units of cm^{-3} and γ_{e1} and γ_{e2} correspond to the Lorentz factors of the electrons producing radiation primarily around ν_1 and ν_2 , respectively, the synchrotron emissivity may be approximated as (see e.g. Rybicki & Lightman 1979)

$$[\nu_r j_{\nu_r}] \simeq \frac{c\sigma_T U_B}{6\pi} k_e \left(\frac{\nu_r}{\nu_{\text{cr}}} \right)^{\frac{3-s}{2}}, \quad (9)$$

where $U_B \equiv B^2/8\pi$ is the energy density of the magnetic field and $\nu_{\text{cr}} \equiv (3eB/4\pi mc) \gamma_e^2 \simeq 4.2(B/\mu\text{G}) \gamma_e^2$ Hz is the critical (radio) synchrotron frequency for a given γ_e .

For production of the observed radio background, we need $s = 2.2$, $\gamma_{e1} = 5 \times 10^2 (B/\mu\text{G})^{-1/2}$ and $\gamma_{e2} = 5 \times 10^4 (B/\mu\text{G})^{-1/2}$, with the value of k_e being determined from the following relation between the emissivity and the observed energy density.

We relate the (radio) synchrotron energy density to the emissivity with

$$\begin{aligned} [\nu_r U_{\nu_r}] &= \frac{4\pi}{c} \int \frac{dV}{dz} \frac{dz}{4\pi d_L^2(z)} [\tilde{\nu}_r j_{\tilde{\nu}_r}] \\ &= [\nu_r j_{\nu_r}] \frac{4\pi}{H_0} \int \frac{F_{\text{syn}}(z) dz}{(1+z)^{(s+1)/2} E(z)}, \end{aligned} \quad (10)$$

where $\tilde{\nu}_r = \nu_r(1+z)$, $H_0 = 70 \text{ km}^{-1} \text{ Mpc}^{-1}$ is the Hubble constant and $F_{\text{syn}}(z)$ describes the evolution of the product $U_B \times k_e$. Here $E(z) \equiv \sqrt{\Omega_M(1+z)^3 + \Omega_\Lambda}$ for the assumed flat cosmology, and the comoving volume element is

$$\frac{dV}{dz} \equiv \frac{c^3 4\pi}{H_0^3 E(z)} \left[\int_0^z \frac{dz'}{E(z')} \right]^2 = \frac{4\pi c d_L^2}{H_0 (1+z)^2 E(z)}. \quad (11)$$

Similarly, we can get the IC emissivity resulting from the same population of electrons as²

$$[\nu_{\text{IC}} j_{\nu_{\text{IC}}}] = \frac{3ch\sigma_T}{16\pi} \int d\nu_0 \int d\gamma_e n(\nu_0) n_e(\gamma_e) \frac{\nu_{\text{IC}}^2 f_T}{\gamma_e^2 \nu_0} \quad (12)$$

(Blumenthal & Gould 1970), where $n(\nu_0) \equiv [\nu_0 U_{\nu_0}]_{\text{tot}}/h\nu_0^2$ is the total spectral number density of the extragalactic background photon field (including radio, CMB, IR, optical/UV, X-ray, etc.):

$$f_T = 2q_T \ln q_T + q_T + 1 - 2q_T^2,$$

$$q_T \equiv \frac{\nu}{4\nu_0 \gamma_e^2} \quad \text{and} \quad \frac{1}{4\gamma_e^2} \leq q_T \leq 1. \quad (13)$$

² The following expressions are valid in the Thompson regime and are a good approximation for scattering of photons with $h\nu_0 \sim 1$ eV by the highest energy electrons γ_2 . For scattering of photons above this energy, one must use the Klein–Nishina cross-section. Few relevant photons lie above this range, so in what follows we approximate the Klein–Nishina suppression by a sharp cut-off.

The IC emissivity can be related to the IC energy density by an equation similar to equation (10) with $F_{\text{syn}}(z)$ being replaced by $F_{\text{IC}}(z)$ which now describes the evolution of the product $n(\nu_0) \times k_e$. Ignoring the differences between these two evolutions [i.e. assuming that the ratio of the integrals over redshift involving $F_{\text{syn}}(z)$ and $F_{\text{IC}}(z)$ is of order unity] and eliminating k_e , we can express the IC energy density in terms of the synchrotron energy density as

$$\begin{aligned} [\nu_{\text{IC}} U_{\nu_{\text{IC}}}] &\simeq \frac{[\nu_r U_{\nu_r}]}{B^2/8\pi} \left(\frac{4.2(B/\mu\text{G})}{\nu_*} \right)^{0.4} \nu^2 \\ &\times \int_{\gamma_{e1}(B)}^{\gamma_{e2}(B)} d\gamma_e \int_{\nu_{\text{min}}}^{\nu_{\text{max}}} d\nu_0 \nu_0^{-3} \gamma_e^{-4.2} [\nu_0 U_{\nu_0}]_{\text{tot}} f_T, \end{aligned} \quad (14)$$

with $\nu_{\text{max}} = \min[\nu_2, \nu, m_e c^2/4h\gamma_e]$ and $\nu_{\text{min}} = \max[\nu_1, \nu/4\gamma_e^2]$. We note that because the IC emissivity is dominated by upscattering of the CMB photon field, for which the energy density increases with redshift, the presented evaluation of $[\nu_{\text{IC}} U_{\nu_{\text{IC}}}]$ with the cosmological evolution neglected corresponds strictly to a lower limit.

Fig. 3 shows the energy density of the observed extragalactic background light (thick curve), and the expected IC energy density resulting from upscattering of these background photons by electrons producing the radio background as given by equation (14), for different magnetic fields $B = 0.001, 0.01, 0.1$ and $1 \mu\text{G}$ (dotted, dashed, dot–dashed and solid curves, respectively). Spectral energy densities of the IR/optical, X-ray and γ -ray cosmic photon fields were constructed to be in agreement with the background levels provided by Franceschini, Rodighiero & Vaccari (2008), Gilli, Comastri & Hasinger (2007) and Sreekumar et al. (1998), respectively. Clearly, any magnetic field weaker than $1 \mu\text{G}$ would result in X-ray/ γ -ray emission exceeding the observed background, and regions with such magnetic fields may be excluded as significant sources of the CRB.

Importantly, this consideration excludes our own Galactic halo as the origin of the bulk of the isotropic radio signal. Taylor, Stil & Sunstrum (2009) use the measurements of the rotation measure of 37 000 polarized extragalactic radio sources to determine the




Figure 3. The thick black curve shows the measured energy density of the radio, microwave, IR, optical, UV, X-ray and γ -ray extragalactic backgrounds. The other curves show the energy density produced by IC scattering of the photon backgrounds by electrons necessary to produce the radio background reported by the ARCADE 2 collaboration via synchrotron emission. The dotted, dashed, dot–dashed and solid curves are for a 1-nG, 10-nG, 100-nG and 1- μG level average magnetic field. Because the intergalactic magnetic field is known to be $\leq 1 \mu\text{G}$, the observed level of the X-ray background rules out a significant portion of the radio background being produced by electrons far from galaxies. Spectral energy densities of the IR/optical, X-ray and γ -ray cosmic photon fields were constructed to be in agreement with the background levels provided by Franceschini et al. (2008), Gilli et al. (2007) and Sreekumar et al. (1998), respectively.

intensity of the magnetic field in the Galactic halo, concluding with a value of approximately $1 \mu\text{G}$. In our Galactic halo, the level of the ambient optical and IR photon fields will be even higher than that considered in the calculation here, by an amount depending on the distance from the Galactic plane, predicting an X-ray background many times larger than that observed. There are Galactic and Solar system components to the observed diffuse X-ray background, but these are significant only below 1 keV (Hickox & Markevitch 2006). The observed level of the X-ray background therefore strongly disfavours a Galactic origin for the observed isotropic radio signal.

3.2 Diffuse emission from the IGM

The CRB could in principle result from a population of relativistic electrons pervading the IGM as a whole, perhaps resulting from many generations of AGN. Large-scale radio sources can easily expand to sizes of many Mpc on gigayear time-scales, and so overlap. Almost certainly, however, adiabatic expansion losses would result in a very low energy density and synchrotron losses would lead to a steep energy spectrum. Furthermore, a variety of arguments suggest that the magnetic field in the IGM is likely to be very weak, $B \leq 0.2 \mu\text{G}$ (see Vallée 2004 and references therein). Diffuse emission from the IGM therefore seems unlikely to be the solution.

3.3 Diffuse emission from clusters

There is evidence for diffuse radio emission in some but not all clusters of galaxies (e.g. Feretti & Giovannini 2008). Ever since the discovery of this emission from the Coma cluster (Large, Mathewson & Haslam 1959), there have been improved observations of this system (Schlickeiser, Sievers & Thiemann 1987; Wilson 1970) as well as more extensive surveys by Giovannini & Feretti (2000) who have identified more than 40 clusters with diffuse radio emission. These are classified as haloes or relics and are often associated with dynamically disturbed (merging) clusters. The origin of the relativistic electrons is controversial. They may be injected by AGN jets (McNamara & Nulsen 2007, and references therein) or produced as secondary pairs from ultrarelativistic cosmic ray hadrons' scattering of ICM protons. Finally, non-thermal electrons within IGM may also be picked up directly from the thermal pool and accelerated to ultrarelativistic energies by ICM shocks or turbulence ('primary' electrons; see e.g. Petrosian 2001; Brunetti & Lazarian 2006). We note that the most recent observations of a few clusters at γ -ray energies seem to exclude the 'secondary' nature of the radio halo electrons (e.g. Aleksić et al. 2010).

The fraction of clusters with radio emission increases rapidly with increasing soft X-ray luminosity, indicating that *strong* diffuse radio emission is not a common property of all clusters. However, deeper radio observations, especially at low frequencies, suggest that the presence of extended, low-surface-brightness radio structures in galaxy clusters is relatively common (e.g. Rudnick & Lemmerman 2009) but at lower flux levels. Possibly then, the integrated effect of a large number of such weakly emitting clusters could be a significant contribution to the CRB. We are not able to predict the level of this integrated emission, but we can check the cluster hypothesis against our two general constraints – magnetic field and spectral index.

There are estimates of the magnetic fields in some clusters via Faraday rotation measurements, which indicate (line-of-sight-averaged) magnetic fields from 1 to $10 \mu\text{G}$ (e.g. Kim & Kronberg

1990; Taylor, Fabian & Allen 2002). In particular, in Coma, Kronberg et al. (2007) measure a line-of-sight-averaged intergalactic magnetic field in the range of $0.2\text{--}0.4 \mu\text{G}$. However, the actual magnetic field will be larger by a factor of $(R/H_B)^{1/2}$ if it is tangled or chaotic on a distance scale H_B that is smaller than the cluster size R . For $H_B \sim 10 \text{kpc}$, one can get fields as high as few μG . Overall, it is quite plausible that some cluster emission passes the magnetic field test.

However, the observed radio spectra are relatively steep (index $\alpha > 1$; see Liang et al. 2000) which does not match the index of the CRB. It therefore seems that diffuse ICM/IGM emission, while possibly widespread, is not a dominant contributor to the CRB.

4 MISSING LOW-SURFACE-BRIGHTNESS EMISSION

It is possible that the calculated strength of the CRB based on source counts is underestimated in two ways, due to the surface brightness limits of surveys. First, the fluxes of some extended objects may have been undercharacterized, if their sizes are large, or they have extended low-surface-brightness components. Secondly, if there is a wide range of source surface brightnesses extending to low values, there may exist sources which will not be detected. In order to discuss this issue in more detail, it is useful to consider high- and low-flux sources separately.

4.1 High-flux sources

At high fluxes, source counts are dominated by radio galaxies and radio-loud quasars that very often have large-scale morphology. It is useful to distinguish the two well-known Fanaroff–Riley (FR) morphological types (Fanaroff & Riley 1974). The low-power FR type Is (FR Is) have surface brightness profiles which fade outwards with no clear end, and therefore part of their total flux will be undetected in interferometric radio surveys. However, these outer components typically have steep spectra (e.g. Hardcastle 1999; Liang et al. 2006, 2008), which is inconsistent with being the dominant component of the CRB. Moreover, they have only a modest cosmological evolution (Willott et al. 2001; Jamrozy 2004; Rigby, Best & Snellen 2008; Smolčić et al. 2009) and so although significant at sub-mJy fluxes (see Rigby et al. 2008; Padovani et al. 2009), they are unlikely to be the dominant low-flux population.³

'Classical doubles' (FR IIs), on the other hand, have clearly defined boundaries of their extended radio lobes, but are expected to decrease in surface brightness as they age and grow in size (Kaiser, Dennett-Thorpe & Alexander 1997). Such extended structures, which can reach Mpc-scale sizes, may fall below typical survey surface brightness detection limits, especially at high redshifts. Even the cores may remain undetected, as they often have very low levels of central (AGN-like) activity (Machalski, Jamrozy & Zola 2001; Machalski et al. 2006; Dwarakanath & Kale 2009). However, the known low-surface-brightness giant radio galaxies tend to have steep spectra (see Jamrozy et al. 2008; Konar et al. 2008), which is not consistent with that of the CRB.

A new survey, the Australia Telescope Low Brightness Survey (Subrahmanyam et al. 2010; Subrahmanyam et al. in preparation), has addressed the question of the number of potentially missed

³ This conclusion is in agreement with the modest estimated contribution of FRI sources to the extragalactic X-ray and γ -ray backgrounds (Celotti & Fabian 2004; Stawarz, Kneiske & Kataoka 2006, respectively)

low-surface-brightness radio galaxies or their extended components, using an 8.4-deg² survey down to mJy fluxes, with a surface brightness threshold being a factor of 5 lower than previous comparable surveys. Subrahmanyan et al. (2010) state that 30 per cent of their sources have at least half of their flux between 5 and 30 arcsec and that 10 per cent of their sources have a size at least 1.5 times their beamsize of 50 arcsec, indicating that there is a significant, but not dominant, component of extended flux. This survey discovered a few new giant radio galaxies. The authors have not yet published source counts, but their initial analysis finds 500 sources in 8.4 deg² to a completeness limit of 1 mJy at 1.4 GHz, corresponding to an integrated source density of $1.95 \times 10^5 \text{ sr}^{-1}$. This is not very different from our estimate based on other surveys discussed in Section 2.1, suggesting that any missing population of high-flux but low-surface-brightness sources is not large enough to explain the CRB.

In summary, while there are indications that some high-flux density, low-surface-brightness sources may have been missed in earlier source count surveys, it seems unlikely that this is a very large effect, at most a 50 per cent correction, which can increase the maximum integrated contribution of the high-flux population to the CRB to ~ 25 per cent.

4.2 Low-flux sources

Below 1 mJy, optical spectra of identified sources indicate that source counts become dominated by star-forming galaxies (e.g. Benn et al. 1993) and the characteristic radio source size changes from ~ 10 to ~ 2 arcsec (Coleman & Condon 1985), consistent with a change in population from double-lobed, elliptical-hosted radio galaxies to star-forming ‘regular’ spiral galaxies and Seyferts. Sizes of a few arcsecond are large enough that measured peak fluxes will often be an underestimate of the total flux, but not by a large factor. The usual procedure is to apply a statistical correction to the fluxes, assuming a given source size distribution according to the formula of Windhorst, Mathis & Neuschaefer (1990).

For example, Hopkins et al. (2003), with a beam size of 6 arcsec, apply the Windhorst et al. (1990) formula for the angular flux distribution to estimate a correction factor of $f \sim 1.3$ for an assumed median source size of 2 arcsec. A median size of 4 arcsec (considerably bigger than expected for spiral galaxies at the redshift $z = 1$) would instead require $f \sim 1.7$. This effect is therefore unlikely to be very large for the star-forming galaxy population.

One could imagine a population of large- and low-surface-brightness radio-emitting galaxies whose fluxes are badly underestimated. Then for median source sizes of 10 and 50 arcsec, the correction factors would be $f \sim 2.5$ and $f \sim 5.9$, respectively, the later being nearly sufficient to explain the observed CRB. However, a completely missing population at $S \sim 0.1$ mJy seems unlikely. In compiling the source counts shown in Fig. 1, we examined normalization against survey resolution and found no significant consistent effect. There is also some clear evidence in a specific field. With a beam size of 6 arcsec, Coleman & Condon (1985) surveyed a Very Large Array field to μJy levels which had previously been surveyed at the same frequency with a beam size of 19 arcsec by Condon & Mitchell (1984). Of 159 sources found in the first (large beam) survey, only eight were not seen in the later (small beam) survey.

Summarizing the situation for low-flux sources, it seems unlikely that significant extended flux has been missed (i.e. not already corrected for).

5 POINT SOURCES

The above results indicate that a class of radio sources not previously considered is needed to produce the bulk of the radio background. In particular, the background cannot be formed from diffuse emission or relatively few high luminosity sources, but rather many lower luminosity sources. As discussed in Section 2.2, these sources should be numerous enough to dominate source counts from below the ~ 10 - μJy limit of current radio surveys down to the $\sim 10^{-2}$ - μJy level. In addition, the sources should have high magnetic fields, $B > 1 \mu\text{G}$, and should be produced by a relatively flat spectrum of relativistic electrons. In this section, we consider some possible candidates.

5.1 Radio supernovae

RSNe produce significant radio flux and could be a contributor to the background. RSNe associated with Type II supernovae have a mean spectral index roughly comparable to the observed CRB (Weiler et al. 1986). Colina et al. (2001) discovered an RSN in a radio monitoring campaign of the galaxy NGC 7469 with an 8.4-GHz luminosity of $\simeq 1.1 \times 10^{21} \text{ W Hz}^{-1}$, which is more than a thousand times more luminous than Cassiopeia A, the brightest RSN remnant in the Milky Way, and which would correspond to flux densities in the range $\mu\text{Jy} < S < \text{mJy}$ between $z = 0.1$ and $z = 2$. This object would not have been found in conventional optical searches.

Are RSNe of this luminosity numerous enough to account for the background? As discussed in Weiler et al. (2004) significant radio emission is only seen in core collapse supernovae, and approximately 10 per cent of core collapse supernovae exhibit radio loudness. Madau, Della Valle & Panagia (1998) estimate the number of core collapse supernovae integrating across all redshifts to be $3 \text{ arcmin}^{-2} \text{ yr}^{-1}$. Mannucci, Della Valle & Panagia (2007) update this with the number of supernovae obscured optically by starbursts, limiting the correction to a factor of 2 at most, giving $\leq 1 \times 10^9 \text{ yr}^{-1}$ core collapse supernovae over the whole sky. With the period of radio loudness lasting on the order of a year (Weiler et al. 2004), we have only $\leq 1 \times 10^8$ RSNe on the sky at any time. Noting that $z = 2$ is the peak era of supernova activity and assuming a mean RSN 8.4-GHz luminosity of $1 \times 10^{21} \text{ W Hz}^{-1}$, this would give a contribution to the background intensity at the level of $\lesssim 10 \text{ Jy sr}^{-1}$, falling short of the observed CRB by a factor of ≥ 1000 . For a broad luminosity function (LF) extending below $1 \times 10^{21} \text{ W Hz}^{-1}$, the contribution will be even less. Also note that a fraction of the RSNe population at this luminosity located at significantly lower redshifts would have sufficiently high flux to be included in the radio surveys considered in Section 2 and Fig. 1, and therefore cannot explain the ‘missing’ portion of the background.

A higher fraction of core collapse supernovae could be more luminous in the radio if they exploded in denser environments. At very high redshifts, nearly all population III stars could be very massive and explode in so-called pair-instability supernovae. However, the supernova rate from population III stars is estimated by Wise & Abel (2005) to be only $10^{-4} \text{ arcmin}^{-2} \text{ yr}^{-1}$, which is much less than regular core collapse supernovae.

Finally, if RSNe were significant contributors to the radio fluxes of spiral galaxies, they would be seen as compact sources in high-resolution images of nearby spiral galaxies. However, they are not. Young RSNe can sometimes be seen as individual sources, but they contribute only a tiny fraction of the total flux. We conclude that RSNe are not a significant component of the radio background.

5.2 Radio-quiet quasars

RQ quasars, as is well known, are not radio silent and thus could potentially account for a missing population of CRB sources. They are typically unresolved on sub-arcsecond scales, with 1.4-GHz luminosities of the order of 10^{38} – 10^{41} erg s⁻¹ (Elvis et al. 1994; Blundell & Kuncic 2007; White et al. 2007). The radio emission of higher luminosity RQ quasars is most often produced by mildly/non-relativistic electrons in nuclear outflows originating in the inner parts of accretion discs, though large-scale relativistic jets are sometimes (surprisingly) detected in such systems (Blundell, Beasley & Bicknell 2003). The expected strong magnetization of disc winds ($B \gg 1 \mu\text{G}$) requires low Lorentz factors and low energies of radio-emitting electrons which remove the IC problem discussed in Section 3. We note that the radio emission from most low-luminosity ($< 10^{22}$ W Hz⁻¹) RQ quasars may not originate in nuclear outflows but rather from star-forming host galaxies (Kellerman et al. 1994), but their space density is much lower than that of comparably luminous sources in normal star-forming galaxies. The observed radio continua of RQ quasars are flat, possibly in agreement with the CRB spectrum (Blundell, private communication). RQ quasars are also numerous, though under-represented in existing radio surveys due to the flux limits of the surveys. In addition, White et al. (2007) have demonstrated that stacking seemingly empty radio images of optically identified quasars which are individually below the noise level of the radio survey leads to a composite image of significant radio flux. Finally, Simpson et al. (2006), Ibar et al. (2009) and Padovani et al. (2009) have recently claimed that RQ AGN constitute a significant fraction of the sub-mJy radio source population.

On the other hand, the idea that RQ quasars produce the bulk of the CRB presents several problems. From Fig. 2, even with a faint-end index of 2.5 below 100 μJy , the number of sources needed to make the background is perhaps only a factor of 10 lower than the reasonable estimates of the total number of non-dwarf galaxies in the observable Universe and as such is much higher than the expected number of quasars, i.e. high-accretion rate objects. Also, optical quasar number counts (e.g. Richards et al. 2006) point to a peak in the contribution to $S^2 dN/dS$ occurring at $\sim 50 \mu\text{Jy}$ in the i band. Converting this to a radio flux using a radio loudness parameter (the ratio of the 1.4-GHz radio to 2500-Å optical luminosity) on the order of 1 for RQ quasars, and adjusting the optical flux between the i band and 2500 Å according to standard spectral models, indicates that the peak of $S^2 dN/d \log(S)$ at 1.4 GHz would fall near 10 μJy . Below this peak flux, the faint-end index would drop to a sub-Euclidean value. This turnover would occur at least one order of magnitude higher than that necessitated by Fig. 2.

Let us now estimate the expected contribution of RQ quasars to the measured CRB. Knowing the *radio* LF of RQ quasars, $\psi(L_R, z)$, where $L_R \equiv [v_r L_{v_r}]$ is the radio luminosity at frequency v_r (in what follows, we will use $v_r = 5$ GHz), their contribution to the CRB energy density can be evaluated in a manner equivalent to that of equation (10) with the emissivity replaced by the LF multiplied by L or $L_R \psi(L_R, z)$, namely

$$\begin{aligned} [v_r U_{v_r}] &= \frac{1}{c} \int dz \int dL_R \frac{dV}{dz} \frac{\psi(L_R, z) L_R}{(1+z)^{\alpha-1} 4\pi d_L^2} \\ &= \int dz \int dL_R \frac{\psi(L_R, z) L_R}{H_0 (1+z)^{1+\alpha} E(z)}, \end{aligned} \quad (15)$$

where $\alpha = d \ln L_{v_r} / d \ln v_r$ is the radio spectral index.

We do not have direct knowledge of the radio LF, but we can estimate as described below by relating it to the known optical

bolometric LF of quasars $\psi(L_{\text{bol}}, z)$ which may evolve as a pure luminosity evolution, pure density evolution or luminosity-dependent density evolution as constrained by Hopkins, Richards & Hernquist (2007). Assuming similar evolution for the radio LF, by definition,

$$\psi(L_R, z) dL_R = \psi(L_{\text{bol}}, z) dL_{\text{bol}}. \quad (16)$$

To evaluate the above integral, we need the relation between the bolometric and radio luminosities. White et al. (2007) give a relation between the radio luminosity at 5 GHz and the absolute 2500-Å magnitude M_{UV} , which can be written as $\log[L_{5 \text{ GHz}} / (\text{erg s}^{-1} \text{ Hz}^{-1})] \simeq 22.07 - 0.34 M_{\text{UV}}$ or $\log[L_R / (\text{erg s}^{-1})] = 31.76 - 0.34 M_{\text{UV}}$. We convert the 2500-Å magnitude to the bolometric quasar luminosity using the approximate relations $L_{\text{bol}} \simeq 3 \times v_{\text{UV}} L_{v_{\text{UV}}}$, $m_{\text{UV}} - M_{\text{UV}} = 5 \log[d_L / \text{pc}] - 5 + K$, where K stands for the K -correction factor, and $m_{\text{UV}} = -2.5 \log[S_{\text{UV}} / 3500 \text{ Jy}]$, to obtain

$$\left(\frac{L_R}{10^{36} \text{ erg s}^{-1}} \right) \simeq 0.9 \times \left(\frac{L_{\text{bol}}}{10^{41} \text{ erg s}^{-1}} \right)^{0.85}, \quad (17)$$

where the luminosities L_{bol} , $v_{\text{UV}} L_{v_{\text{UV}}}$ and L_R are expressed in the units of erg s⁻¹.

After integrating equation (15) over the redshift range⁴ $z = 0-6$ and the radio luminosity range $L_R = 10^{36}$ – 10^{42} erg s⁻¹ (corresponding to the bolometric luminosity range $L \simeq 10^{41}$ – 10^{48} erg s⁻¹), for the fit parameters regarding the quasar LF as provided in Hopkins et al. (2007), we find the expected contribution of RQ quasars to the extragalactic radio background at 5 GHz to be at the level of $\simeq 1.4$ – 1.7 per cent for pure luminosity or luminosity-dependent density evolution models. In the case of a pure density evolution model, the expected contribution rises to 4 per cent. This contribution is therefore smaller by at least a factor of 10 than needed. One has to keep in mind, however, that the presented calculations correspond strictly to the lower limits, because in constructing the radio LF, we considered only type 1 quasars. However, including the population of type 2 sources, which cannot outnumber the population of unobscured quasars by a large factor, is not expected to account for the order-of-magnitude difference.

In order for emission from RQ quasars to comprise the bulk of the background, therefore, their radio loudness must evolve with redshift and they must be significantly more numerous at redshifts between 1 and 3 than previously thought. Although Blundell (2003) has claimed that significant flux from RQ quasars may be missed by interferometric observations, we conclude that emission from RQ quasars, while significant (at the few per cent level at least), does not likely explain the bulk of the missing contribution to the radio background, unless there is some as of yet undetected but significant evolution in the quasar radio LF.

5.3 Star-forming galaxies

Given the large ($\gtrsim 10^{10}$ sr⁻¹) number of sources needed to produce the CRB, the constraints on extended and non-galactic emission discussed in Sections 4 and 3, the required strong magnetic fields ($B > 0.2 \mu\text{G}$) and flat spectra, and the shortcomings of RSNe and RQ quasars (Sections 5.1 and 5.2), it seems that ordinary galaxies may

⁴ While some high accretion rate objects may be present before $z = 6$, the number must necessarily fall far short of the number of objects needed to make the background outlined in Section 2.2, as the number of galaxies present at these redshifts is a small fraction of the total in the observable universe.

be more reasonable candidates for production of most of the CRB. Ordinary star-forming galaxies are thought to dominate the low-flux population of the resolved radio sources (e.g. Richards 2000). The radio continua produced in star-forming regions are relatively flat, being characterized by the spectral indices close to those of the CRB $\alpha = 0.6$ (see Ibar et al. 2009). Finally, the expected magnetic fields $B \gtrsim 1 \mu\text{G}$ in star-forming regions circumvent the IC problem discussed in Section 3.

5.3.1 Constraints from the far-infrared background

The radio emission originating in local star-forming regions is observed to be correlated with the FIR flux (e.g. Dwek & Barker 2002; Ibar et al. 2008, and references therein), which, combined with the observed level of the IR background, constrains the contribution of these sources to the CRB. Dwek & Barker (2002) give a relation between the 1.4-GHz radio power, $P_{1.4\text{GHz}}$, and the FIR luminosity, L_{FIR} , in the wavelength range of 10–1000 μm , which can be written as $L_{\text{r}} \equiv [\nu P(\nu)]_{1.4\text{GHz}} \sim \times 10^{-6} L_{\text{FIR}}$. Therefore, if the radio background is produced in the star-forming regions of Seyferts, spirals and ultraluminous IR galaxies, then from a naive application of this ratio one should expect the observed FIR background at the level of

$$[\nu U_{\nu}]_{\text{FIR}} \simeq 10^6 [\nu U_{\nu}]_{1.4\text{GHz}} \simeq 2 \times 10^{-13} \text{ erg cm}^{-3},$$

which is higher than that observed (Marsden et al. 2010) by a factor of >10 . This is in agreement with the Dwek & Barker (2002) result showing that the expected surface brightness of the sky at 178 MHz for different models of the cosmologically evolving star formation rate is between 3 and 30 K, about 3–30 per cent of the ARCADE result $T_{\text{CRB}}(178 \text{ MHz}) \simeq 104 \text{ K}$ (Fixsen et al. 2010).

An alternative way to formulate this problem is as follows. Appleton et al. (2004) find that individual star-forming galaxies locally show a ratio of FIR to radio fluxes given by the value $q_{70} \equiv \log(S_{70\mu\text{m}}/S_{1.4\text{GHz}}) = 2.15$. If we take the observed CRB and assume that it is made of star-forming galaxies following this ratio, then, ignoring for now any K -correction issues, we predict an FIR surface brightness at 70 μm of $178 \text{ nW m}^{-2} \text{ sr}^{-1}$, which is about 25 times the observed level reported in Dole et al. (2006). Thus, we can conclude that the contribution of systems that obey the local radio–FIR correlation to the radio background is on the level of approximately $\lesssim 5$ per cent or, alternately, that the radio background must be made in large part from sources that appear more radio loud than the local radio–FIR correlation by a factor of ~ 20 .

5.3.2 An evolving radio–far-infrared correlation

Since star formation in the Universe has evolved and the rate was much higher at redshifts of 1 and above (e.g. Madau et al. 1998), the bulk of both the radio and IR backgrounds is produced at these redshifts. Therefore, an evolution in the observed FIR to radio flux ratio towards greater radio loudness with redshift is necessary to have the radio and FIR backgrounds produced at the observed levels by star forming galaxies.

It is important to note that the ratio of the relative contribution to the FIR background and radio background surface brightnesses from a galaxy (or a class of galaxies) located at a particular redshift is given by the non- K -corrected FIR to radio flux ratio rather than the K -corrected ratio that is often reported for higher redshifts. Relating an observed flux ratio or ratio evolution to a K -corrected one, or vice versa, is dependent on the source spectral energy distribution

(SED) assumed and the wavelengths in question. For observations at 70 μm and 1.4 GHz and a typical starburst galaxy SED, a fixed intrinsic q_{70} (where q_x is the log of the flux ratio at IR frequency x to 1.4-GHz radio) from a galaxy at $z \sim 2$ will show an observed value reduced by $\Delta q \sim 0.6$ (see example calculations in references below). To explain the observed discrepancy between radio and 70- μm backgrounds shown in Section 5.3.1, we would need to observe a high redshift evolution of $\Delta q(\text{observed}) \sim -1.3$ or $\Delta q(\text{intrinsic}) \sim -0.7$.

A number of papers have explored the evolution of q with redshift at either mid-infrared, FIR or sub-mm wavelengths. These determinations are complicated by selection effects, and in the case of reports of the K -corrected correlation, by template spectra. Although some papers conclude that there is no evidence for a change in intrinsic q (e.g. Ibar et al. 2008; Sargent et al. 2010) others do find possible evolution in intrinsic q (e.g. Vlahakis, Eales & Dunne 2007; Beswick et al. 2008; Seymour et al. 2009; Bourne et al. 2010), although it is not clear that this is large enough to explain the radio background. Recent data from the Balbon-Borne Large-Aperture Submillimeter Telescope (BLAST) instrument (Ivison et al. 2010a) also show the intrinsic correlation evolving, and new results from the Herschel instrument (Ivison et al. 2010b) are consistent with this as well. The most suitable reported results for comparison are those of Sargent et al. (2010) and Bourne et al. (2010) who use *Spitzer* 70- μm data and present the evolution in observed q_{70} versus redshift (fig. 12 in Sargent et al. and fig. 9 in Bourne et al.). These figures agree in showing that by $z \sim 2$ a change of $\Delta q(\text{observed}) \sim -0.7$ is observed, i.e. a factor of 5 decline in the FIR to radio flux ratio. While significant, this is not enough to explain the FIR/radio background discrepancy. However, these figures show considerable scatter, and Sargent et al. show that a radio-selected, as opposed to IR-selected, sample shows a systematically lower value, with $\Delta q(\text{observed}) \sim -0.4$. We again note that determinations of the value of the correlation at higher redshifts are complicated by selection biases and contamination by AGN emission.

An evolving radio to FIR luminosity ratio towards the radio loud with increasing redshift would suggest that one or more of the following is true at higher redshifts: (1) a larger portion of the energy of star formation goes into relativistic particles, for some reason possibly related to a different structure of the interstellar medium shaping the cosmic ray acceleration efficiency at shocks driven by supernova, (2) a larger portion of the stars are of high mass, resulting in more supernovae as well as an enhanced cosmic ray acceleration at shocks driven by winds from massive stars, (3) synchrotron emissivity is enhanced by higher interstellar magnetic fields or (4) AGN activity in ordinary galaxies is proportionally more important.

There is no concrete information on point 1, while point 2 requires evolution of the initial mass function, which if true would alter the interpretation of the data on the star formation rate. The third possibility can be neither rejected nor supported by current data.

Here, let us only comment on the possibility that the cosmological evolution of supermassive black holes (SMBHs) in the Galactic Centres results in enhanced AGN activity in late-type galaxies at high redshifts and thus in their enhanced radio emission relative to IR. Note that in this context, it is now established that a large fraction of nearby spiral galaxies show weak AGN activity (Ho, Filippenko & Sargent 1997; Ho 2008). In addition, several authors have argued that SMBHs were spinning more rapidly at early epochs (Wang et al. 2009). For spiral-hosted SMBHs the effect may be even stronger than for the elliptical hosted SMBHs that give rise to quasars, due to the different character of the dominant

accretion events which determine the black hole spin evolution (see the discussion in Sikora, Stawarz & Lasota 2007; Volonteri, Sikora & Lasota 2007). In the framework of the spin paradigm for jet production, critically re-examined recently by Sikora et al. (2007), one could therefore expect more powerful jets and therefore more AGN-related radio emission (for the same accretion luminosity) at higher redshifts in these systems.

If the radio background is indeed formed from $\sim 0.1\text{-}\mu\text{Jy}$ sources, then this corresponds to about $\sim 10^{21}\text{ W Hz}^{-1}$ radio (1.4-GHz) luminosity at $z = 2$, which would be between 10^8 and $10^9 L/L_{\odot}$ at $70\text{ }\mu\text{m}$ with an evolving (K -corrected) q_{70} . This IR luminosity is below the ‘knee’ of the IR LF, i.e. the radio background would be made by relatively normal spiral galaxies, in contrast to the IR background, which appears to be dominated by luminous and ultra-luminous IR galaxies (Magnelli et al. 2009). We note the difficulty that locally the peak contribution to radio emission is from relatively luminous galaxies with 1.4-GHz radio luminosities of around 10^{22} W Hz^{-1} (Condon, Cotton & Broderick 2002), while the peak contribution to the radio background, from galaxies beyond redshift 1, is at 1.4-GHz luminosities of around 10^{21} W Hz^{-1} .

6 DISCUSSION

We have considered several mechanisms as the origin of the ‘missing’ radio flux necessary to account for the radio background, assuming that it is at the level reported by the ARCADE 2 collaboration. As shown in Section 3, diffuse, low-surface-brightness synchrotron emission from large-scale structures (IGM, ICM and WHIM), and from our own Galactic halo, is limited by the observed level of the X-ray/ γ -ray background. As discussed in Section 2.1, radio sources detected in many surveys above 1 mJy at 1.4 GHz contribute a total of 16 per cent of the background, and considerations of possible missed flux from extended sources discussed in Section 4 may increase this number by at most a factor of 1.5. The resolved sources with fluxes from below 1 mJy to the current lower limit of interferometric surveys at $\sim 10\text{ }\mu\text{Jy}$ contribute another ~ 10 per cent, leaving between 60 and 75 per cent of the CRB unaccounted for.

We find it difficult to explain the level of the radio background reported by ARCADE 2 without a new population of low-flux radio sources. These sources should be numerous and faint enough to dominate source counts below the $\sim 10\text{-}\mu\text{Jy}$ limit of current radio surveys and must extend to the $\sim 10^{-2}\text{-}\mu\text{Jy}$ (at 1.4-GHz) level. Moreover, they should have an observed ratio of radio to IR output a factor of 5 above what is observed in local galaxies. As discussed in Sections 5.1 and 5.2, RSNe and RQ quasars do not seem to produce adequate radio emission to account for the bulk of the background, although the latter class of objects is expected to contribute to the CRB at the level of at least a few per cent. We conclude that for the radio background to be at the level reported by ARCADE 2, it must be largely comprised of emission from ordinary galaxies at $z > 1$ in which the radio to FIR observed flux ratio increases significantly with redshift.

ACKNOWLEDGMENTS

We thank Philip Best, Katherine Blundell, Chi C. Chueng, Sebastian Jester, Dan Marrone, Matthew Turk and Tom Abel for their input. JS thanks the ARCADE team, S. Kahn and R. Schindler for their encouragement. LS was supported by the Polish Ministry of Science and Higher Education through the project N N203 380336, and

also by the Scandinavian NORDITA programme on ‘Physics of Relativistic Flows’.

REFERENCES

- Aleksić J. et al., 2010, *ApJ*, 710, 634
 Appleton P. et al., 2004, *ApJS*, 154, 147
 Ballantyne D. R., 2009, preprint (arXiv:0904.0996)
 Benn C., Rowan-Robinson M., McMahon R., Broadhurst T., Lawrence A., 1993, *MNRAS*, 263, 98
 Beswick R., Muxlow T., Thrall H., Richards A., Garrington S., 2008, *MNRAS*, 385, 1143
 Bigs A., Ivison R., 2006, *MNRAS*, 371, 963
 Blumenthal G. R., Gould R. J., 1970, *Rev. Modern Phys.*, 42, 237
 Blundell K. M., 2003, *New Astron. Rev.*, 47, 593
 Blundell K., Kuncic Z., 2007, *ApJ*, 668, 103
 Blundell K. M., Beasley A. J., Bicknell G. V., 2003, *ApJ*, 591, L103
 Bondi M. et al., 2003, *A&A*, 403, 857
 Bondi M. et al., 2007, *A&A*, 463, 519
 Bourne L., Dunne L., Ivison R., Maddox S., Dickinson M., Frayer D. 2010, *MNRAS*, in press (arXiv:1005.3155)
 Brunetti G., Lazarian A., 2006, in Van der Hucht, ed., *Proc. 26th General Assembly IAU*, 13
 Celotti A., Fabian A. C., 2004, *MNRAS*, 353, 523
 Ciliegi P., Zamorani G., Hasinger G., Legmann I S. G., Wilson G., 2003, *A&A*, 398, 801
 Coleman P., Condon J., 1985, *AJ*, 90, 1431
 Colina L., Alberdi A., Torrelles J., Panagia N., Wilson S., 2001, *ApJ*, 553, L19
 Condon J., 2007, in Afonso J., Ferguson H. C., Mobasher B., Norris R., eds, *ASP Conf. Ser. Vol. 380, At the Edge of the Universe: Latest Results from the Deepest Astronomical Surveys*. Astron. Soc. Pac., San Francisco, p. 189
 Condon J., Mitchell K., 1984, *AJ*, 87, 1429
 Condon J., Cotton W., Broderick J., 2002, *AJ*, 124, 675
 Dole H. et al., 2006, *A&A*, 451, 417
 Dwarakanath K. S., Kale R., 2009, *ApJ*, 698, 163
 Dwek E., Barker M., 2002, *ApJ*, 575, 7
 Elmouttie E., Haynes R. F., Jones K. L., Ehle M., Beck R., Wielebinski R., 1995, *MNRAS*, 275, L53
 Elvis M. et al., 1994, *ApJS*, 95, 1
 Fanaroff B. L., Riley J. M., 1974, *MNRAS*, 167, 31P
 Feretti L., Giovannini G., 2008, in Plionis M., Lopez-Cruz O., Hughes D., eds, *Lect. Notes Phys. Vol. 740, A Pan-Chromatic View of Clusters of Galaxies and the Large-Scale Structure*. Springer-Verlag, Berlin, p. 143
 Fixsen D. et al. 2010, *ApJ*, submitted (arXiv:0901.0559)
 Fomalont E., Kellerman K., Partridge R., Windhorst R., Richards E., 2002, *ApJ*, 123, 2402
 Fomalont E. et al., 2006, *ApJS*, 167, 103
 Franceschini A., Rodighiero G., Vaccari M., 2008, *A&A*, 487, 837
 Gervasi M., Tartari A., Zannoni M., Boella G., Sironi G., 2008, *ApJ*, 682, 223
 Gilli R., Comastri A., Hasinger G., 2007, *A&A*, 463, 79
 Giovannini G., Feretti L., 2000, *New Astron. Rev.*, 5, 335
 Grueff G., 1988, *A&A*, 193, 40
 Grupponi C., Zamorani G., de Ruiter H., Parma P., Mignoli M L. C., 1997, *MNRAS*, 286, 470
 Hales S., Baldwin J., Warner P., 1988, *MNRAS*, 234, 919
 Hardcastle M., 1999, *A&A*, 349, 381
 Hickox R., Markevitch M., 2006, *ApJ*, 645, 95
 Ho L., 2008, *ARA&A*, 46, 475
 Ho L., Filippenko A., Sargent W., 1997, in Peterson B. M., Cheng F.-Z., Wilson A. S., eds, *Proc. IAU Vol. 159, Emission Lines in Active Galaxies: New Methods and Techniques*. Astron. Soc. Pac., San Francisco, p. 429
 Hopkins A., Alfonso J., Chan B., Cram L., Georgakakis A., Mobasher R., 2003, *AJ*, 125, 465

- Hopkins P. F., Richards G. T., Hernquist L., 2007, *ApJ*, 654, 731
- Hummel E., van Gorkom J. H., Kotanyi C. G., 1983, *ApJ*, 267, L5
- Ibar E. et al., 2008, *MNRAS*, 386, 953
- Ibar E., Ivison R., Biggs A., Lal D., Best P., Green D., 2009, *MNRAS*, 397, 281
- Ivison R. et al., 2010a, *MNRAS*, 402, 245
- Ivison B. et al., 2010b, *A&A*, 518, 31
- Jamrozy M., 2004, *A&A*, 419, 63
- Jamrozy M., Konar C., Machalski J., Saikia D. J., 2008, *MNRAS*, 385, 1286
- Kaiser C., Dennett-Thorpe J., Alexander P., 1997, *MNRAS*, 292, 723
- Katgert-Merkelijn J., Robertson J., Windhorst R., Katgert P., 1985, *A&AS*, 61, 517
- Kellerman K., Sramek R., Schmidt M., Green R., Shaffer D., 1994, *AJ*, 108, 1163
- Kim K., Kromberg P., 1990, *ApJ*, 355, 29
- Kogut A. et al. 2010, *ApJ*, submitted, preprint (arXiv:0901.0562)
- Konar C., Jamrozy M., Saikia D. J., Machalski J., 2008, *MNRAS*, 383, 525
- Kronberg P., Kothes R., Salter C., Perillat P., 2007, *ApJ*, 659, 257
- Large M., Mathewson D., Haslam C., 1959, *Nat*, 183, 1663L
- Liang H., Hunstead R., Birkinshaw M., Andreani P., 2000, *ApJ*, 544, 686
- Liang R. A., Canvin J. R., Cotton W. D., Bridle A. H., 2006, *MNRAS*, 368, 48
- Liang R. A., Bridle A. H., Parma P., Feretti L., Giovannini G., Murgia M., Perley R. A., 2008, *MNRAS*, 386, 657
- McGilchrist M., Baldwin J., Riley J., Titterton D., Waldram E., Warner P., 1990, *MNRAS*, 246, 110
- Machalski J., Jamrozy M., Zola S., 2001, *A&A*, 371, 445
- Machalski J., Jamrozy M., Zola S., Koziel D., 2006, *A&A*, 454, 85
- McNamara B. R., Nulsen P. E. J., 2007, *ARA&A*, 45, 117
- Madau P., Della Valle M., Panagia N., 1998, *MNRAS*, 297, L17
- Magnelli B., Elbaz D., Chary R., Dickinson M., LeBorgne D., Frayer D., Willmer C., 2009, *A&A*, 496, 57
- Mannucci F., Della Valle M., Panagia N., 2007, *MNRAS*, 377, 1229
- Marsden G. et al., 2010, *ApJ*, 707, 1729
- Owen F., Morrison G., 2008, *ApJ*, 136, 1889
- Padovani P., Mainieri V., Tozzi P., Kellermann K. I., Fomalont E. B., Miller N., Rosati P., Shaver P., 2009, *ApJ*, 694, 235
- Pearson T., Kus A., 1978, *MNRAS*, 182, 273
- Petrosian V., 2001, *ApJ*, 557, 560
- Prandoni L., Parma P., Wieringa M., de Ruiter H., Gregorini L., Mignano A., Vettolani G., Ekers R., 2006, *A&A*, 457, 517
- Richards E., 2000, *ApJ*, 533, 611
- Richards E. et al., 2006, *AJ*, 131, 2766
- Rigby E. E., Best P. N., Snellen I. A. G., 2008, *MNRAS*, 385, 310
- Rudnick L., Lemmerman J. A., 2009, *ApJ*, 697, 1341
- Rybicki G. B., Lightman A. P., 1979, *Radiative Processes in Astrophysics*. Wiley, New York
- Sargent M. et al., 2010, *ApJ*, 186, 341
- Schlickeiser R., Sievers A., Thiemann H., 1987, *A&A*, 182, 21
- Seiffert M. et al., 2010, submitted (arXiv:0901.0559)
- Seymour N. et al., 2009, *MNRAS*, 398, 1573
- Sikora M., Stawarz Ł., Lasota J. P., 2007, *ApJ*, 658, 815
- Simpson C. et al., 2006, *MNRAS*, 372, 741
- Singal J. et al., 2010, *ApJ*, accepted (arXiv:0901.0546)
- Smolčić V. et al., 2009, *ApJ*, 696, 24
- Sreekumar P. et al., 1998, *ApJ*, 494, 523
- Stawarz Ł., Kneiske T. M., Kataoka J., 2006, *ApJ*, 637, 693
- Subrahmanyan R., Ekers R., Saripalli L., Sadler M., 2010, *MNRAS*, 402, 2792
- Taylor G., Fabian A., Allen S., 2002, *MNRAS*, 334, 769
- Taylor A., Stil J., Sunstrum C., 2009, *ApJ*, 702, 1230
- Vallée J. P., 2004, *New Astron. Rev.*, 48, 763
- Vallee J., Roger R., 1989, *A&AS*, 77, 31
- Vlahakis C., Eales S., Dunne L., 2007, *MNRAS*, 379, 1042
- Volonteri M., Sikora M., Lasota J., 2007, *ApJ*, 667, 704
- Wang J. et al., 2009, *ApJ*, 697, 704
- Weiler K., Sramek R., Panagia N., van der Hulst J., Salvati M., 1986, *ApJ*, 301, 790
- Weiler K., van Dyk S., Sramek R., Panagia N., 2004, *New Astron. Rev.*, 9, 17
- White R., Helfand D., Becker R., Glickman E., De Vries W., 2007, *ApJ*, 654, 99
- Willott C. J., Rawlings S., Blundell K. M., Lacy M., Eales S. A., 2001, *MNRAS*, 322, 536
- Wilson M., 1970, *MNRAS*, 151, 1
- Wilson A., Vallee J., 1982, *A&AS*, 47, 601
- Windhorst R., va Heerde G., Katgert P., 1984, *A&A*, 58, 1
- Windhorst R., Miley G., Owen F., Kron R., Koo D., 1985, *ApJ*, 289, 494
- Windhorst R., Mathis D., Neuschaefer L., 1990, *ASP Conf. Ser. Vol. 10*, Edwin Hubble Centennial Symposium. Astron. Soc. Pac., San Francisco, p. 389
- Windhorst R., Fomalont E., Partridge R., Lowenthal J., 1993, *ApJ*, 405, 9498
- Wise J., Abel T., 2005, *ApJ*, 629, 615

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.

Copyright of Monthly Notices of the Royal Astronomical Society is the property of Wiley-Blackwell and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.