

1969

# The role of endogenous hyperlipemia in experimental atherosclerosis

Brian David Altman  
*Yale University*

Follow this and additional works at: <http://elischolar.library.yale.edu/ymtdl>

---

## Recommended Citation

Altman, Brian David, "The role of endogenous hyperlipemia in experimental atherosclerosis" (1969). *Yale Medicine Thesis Digital Library*. 2336.  
<http://elischolar.library.yale.edu/ymtdl/2336>

This Open Access Thesis is brought to you for free and open access by the School of Medicine at EliScholar – A Digital Platform for Scholarly Publishing at Yale. It has been accepted for inclusion in Yale Medicine Thesis Digital Library by an authorized administrator of EliScholar – A Digital Platform for Scholarly Publishing at Yale. For more information, please contact [elischolar@yale.edu](mailto:elischolar@yale.edu).



T113

+Y12

2044

YALE UNIVERSITY LIBRARY



3 9002 06584 7189

THE ROLE OF ENDOGENOUS HYPERLIPEMIA  
IN EXPERIMENTAL ATHEROSCLEROSIS

Brian David Altman

1969

MUDD  
LIBRARY  
Medical



YALE



MEDICAL LIBRARY

YALE




MEDICAL LIBRARY









Digitized by the Internet Archive  
in 2017 with funding from  
The National Endowment for the Humanities and the Arcadia Fund

<https://archive.org/details/roleofendogenous00altm>

THE ROLE OF ENDOGENOUS HYPERLIPEMIA  
IN EXPERIMENTAL ATHEROSCLEROSIS

A Thesis Submitted  
in Partial Fulfillment  
of the Requirements for the Degree  
of Doctor of Medicine

by

Brian David Altman  
Yale University School of Medicine  
New Haven, Connecticut


1969



THE UNIVERSITY OF CHICAGO  
LIBRARY

PHYSICS DEPARTMENT  
5712 S. UNIVERSITY AVE.

CHICAGO, ILL. 60637



PHYSICS DEPARTMENT  
5712 S. UNIVERSITY AVE.  
CHICAGO, ILL. 60637

To my Father, Wife, and Unborn Child



## ACKNOWLEDGEMENT

I am deeply indebted to Dr. Levin L. Waters for his constant encouragement and superb teaching. He has always been available for constructive criticism, objective evaluation and genuine support.

I am also grateful to Edward Iannucci, Peter Integlia, Helen Cavallero, Sharon Williams and my wife Ann for many hours of technical assistance and manuscript preparation. This thesis was in part supported by U.S.P.H.S. grants.





## CONTENTS

	<u>Page</u>
Introduction	1
Lipoprotein Classification of Fredrickson, et al.	2
Presentation of a patient with Type III Pattern	5
Experimental - Materials and Methods	6
Results	11
Discussion	
Spontaneous Arterial Lesions in Rabbits	17
Cholesterol Feeding Experiments in Brief	18
Endogenous Lipemias and Triton	19
Experimental Vascular Injury	21
Summary and Conclusions	24
Illustrations	26
Charts	40
Bibliography	57



## INTRODUCTION

More lives are claimed in our country by atherosclerosis and its complications than by any other disease process. In order to prevent atherosclerosis, investigators have been endeavoring to understand its pathogenesis. Various experimental models have been developed for this purpose. Saltykow<sup>120a</sup>, in 1908, and Ignatowski<sup>62</sup>, in 1909, first discovered that milk, meat and egg diets could produce atherosclerotic lesions in rabbits. Cholesterol feeding in one or another form has since been the primary method of induction of the characteristic arterial changes.

More recently the role of endogenous lipids rather than dietary cholesterol in the process of atherogenesis has been considered. Certain questions could then be asked: where do atherogenic lipids originate in the body? Which types circulate in the bloodstream? Why do they localize in arterial walls at certain regions? Are endogenous lipids involved in the atherogenetic process?

During the past decade workers have raised these and similar questions. They have become increasingly concerned with carrier states of lipids in the bloodstream (lipoproteins)<sup>45</sup>. Distinct combinations of plasma lipoproteins have been documented in human population groups chiefly by Fredrickson and his associates at the National Heart Institute<sup>38,39,81-3</sup>.

The following investigation is a study of the role in the experimental atherosclerotic process of induced endogenous lipidemia. Included are anatomic studies of resultant lesions and a consideration of the chemical nature of lipoprotein types involved.



### LIPOPROTEIN CLASSIFICATION

Fredrickson and his associates at the National Heart Institute have been studying human serum lipoprotein disorders for several years. In 1965<sup>38</sup> and again in 1967<sup>39</sup> and 1968<sup>83</sup> they published extensive reviews of current findings. They have described five types of "essential" human hyperlipemias, readily identifiable by their electrophoretic patterns. Many of the characteristic patterns also appear secondary to other disorders such as biliary obstruction, hypothyroidism and diabetes.

The first pattern, Type I, is known as hyperchylomicronemia or severe exogenous hyperlipemia. Of less than fifty known patients reported, all have had an early onset. Attacks of abdominal pain are frequent complaints, and no diabetes or severe atherosclerosis has been observed in this group. Deficiency of "lipoprotein lipase" leads to low post-heparin lipolytic activity, and numerous chylomicrons in fasting overnight plasma samples, even while these individuals are on a low-fat diet.

"Essential familial hypercholesterolemia", Type II hyperlipemia, is the most common. Normal types of beta lipoprotein concentrations are increased, and a moderate increase in pre-beta lipoproteins is noted. Homozygous individuals may have fatal atheromatosis in early childhood. Coronary artery disease is common. Patients with hypothyroidism often display this pattern of lipoproteins.

The Type III pattern individuals discussed by Gofman in 1954<sup>45a</sup> may be recognized clinically by their palmar xanthomas. These patients often have a family history of diabetes and have a greatly enhanced incidence of occlusive peripheral vascular and coronary artery disease. Their





turbid plasma contains greatly increased quantities of cholesterol and glycerides. A broad beta band including pre-beta components distinguishes the electrophoretic patterns in this group.

Greatly elevated pre-beta lipoproteins are observed in Type IV patients. Too many glycerides are produced by their livers, and obesity and diabetes are the rule. Similar patterns are seen in patients with hypothyroidism, glycogen storage disease, diabetes mellitus, nephrotic syndrome, and in some individuals after myocardial infarcts.


The final group of hyperlipemic patients are classified as Type V, a combination of "endogenous and exogenous" hyperlipemia patterns. These persons experience bouts of abdominal pain often in their third decade but they usually have diabetes mellitus and/or a strong family history.



# CLASSIFICATION

## BLOOD CHEMICAL AND LIPOPROTEIN PATTERNS\*

### Primary or Familial Hyperlipidemia








EXAMPLE OF A PATIENT WITH A SECONDARY HYPERLIPEMIA

W. S. is a 52-year-old white German male who has a fifteen year history of diabetes mellitus requiring about 75 units of N.P.H. insulin per day. He has had two right sided C.V.A.'s with regaining of most function, has moderate diabetic retinopathy with some blurred vision, no xanthelasma, only mild hypertension (B.P. 150/95), rather massive obesity (wt. 230 lbs., ht. 65 in.), calcified abdominal aorta, intermittent claudication, no angina pectoris, warm extremities with palpable pulses, and one flight dyspnea without jugular venous distention or peripheral edema. In October 1968 his fasting morning blood showed: gross lipemia-milky appearance

total cholesterol 249 mg.%  
free cholesterol 36%  
fatty acids 18.3 meq./l  
tri-glycerides 680 mg.%

His serum lipoprotein electrophoresis (courtesy Dr. R. Scheig) showed a prominent beta and markedly increased pre-beta band consistent with Fredrickson's Type III pattern.



The patient is now on a four-month trial of Atromid-S (Clofibrate), and repeated dietary counseling has been given. Prior control with diet alone was unsatisfactory.



EXPERIMENTAL - MATERIALS AND METHODS

As indicated in the Introduction, the objective of this study was to test the role of endogenous lipidemia in the pathogenesis of experimental atherosclerosis. It is well known that local, acute lesions of animals' arteries, however produced, may be readily modified to changes of the arteriosclerotic type by a cholesterol-rich diet. Therefore the plan of the present experiments was to produce such lesions in the arteries of rabbits and to test the ability of induced endogenous lipidemia to modify the basic lesions in a similar fashion. The goal was to obtain a combined experimental procedure which yielded frequent arterial lesions and a significant lipemia in the majority of animals.

Seventy-four adult New Zealand white rabbits, randomly sexed, weighing 2-5 kg. were maintained on a low-fat (2 percent) diet (Purina rabbit chow), with drinking water ad libitum, in individual wire mesh cages in the Yale animal care unit. Blood samples were drawn with sterile disposable syringes from either the femoral vessels or via cardiac puncture. Drugs were administered (except where noted) via the marginal ear veins. Drugs employed were as follows:

Epinephrine 4 ml. vials 1:1000 (Parke-Davis)  
Levophed<sup>(R)</sup> Bitartrate 0.2% 4 ml. vials (Winthrop)  
Tyramine Hydrochloride Powder (Nutritional Biochemicals)  
Horse Serum, Pooled Sterile (Baltimore Biological)  
Pituitary Powder, Beef Defatted and Whole Beef  
(Nutritional Biochemical)  
Triton WR 1339 (Ruger).

Initial experiments on vascular injury were conducted with epinephrine. Five rabbits were given different doses intravenously. Two animals survived the first series of injections. Only minor histological changes, mainly edema of many tissues, were seen up to one week after the experiments.



Levophed<sup>(R)</sup> (Nor-epinephrine) was employed intravenously for two additional rabbits, but aside from the interesting clinically observable hemodynamic effects (noted later), no significant histologic lesions were seen.

Two more animals were subjected to tyramine injections following the method of Duff<sup>32</sup>. Again, significant arterial lesions could not be demonstrated. (Our early experiments are summarized on Chart 1).

It was then decided to use the experimental hypersensitivity method of Rich and Gregory<sup>111-113</sup> to induce vascular injury. Rabbits were given 10 cc./kg. body weight of sterile pooled frozen horse serum on three successive weekly occasions. Many of these rabbits also received whole dried beef pituitary extract to induce an endogenous lipemia according to procedures outlined by both Rudman<sup>120</sup> and by Kellner<sup>70</sup> for raising serum lipid concentrations. Unfortunately these methods did not yield arterial lesions consistently although sporadic lesions and lipemias were noted. (See Chart 2 for a summary of the experimental findings).

Finally Triton WR 1339, a non-ionic detergent was utilized to induce readily repeatable endogenous lipemias. A single 10 cc./kg. intravenous injection of 10% sterile Triton solution was administered under sterile conditions (according to the technique of Courtice and Schmidt-Diedrichs<sup>24,25</sup> and of Kellner<sup>69</sup>).

A modification of the technique of Kelly<sup>71,142</sup> was used to injure hyperlipemic rabbits' aortas in vivo by freezing. Instead of ethyl chloride spray aortic segments were frozen by the application of dry ice directly at laparotomy.





Anesthesia for aortic freezing experiments was 15-20 cc. of Nembutal in a single dose supplemented when necessary by open drop ether inhalation. Surgical operations were conducted under aseptic conditions. Aortic freezing was performed as follows: the rabbit was immobilized in a supine position by tying all four extremities and administering a single dose of intravenous Nembutal. After shaving from midchest through abdomen, the ventral skin was cleansed with standard Zephiran solution. After checking for regular unobstructed respirations and keeping an ether cone ready, a generous midline abdominal incision was made. The peri-aortic connective tissue was dissected away and the aorta was mobilized for 2-4 cm. well below the origin of the renal arteries. A kelly clamp was inserted behind the aorta, and the vessel was gently lifted, spreading the clamp's jaws 1-2 cm. apart, until the pulsations ceased. At this point the flat edge of a suitably-sized piece of dry ice was applied uniformly along the raised aortic segment for 30 seconds. The aorta was then allowed to fall back into place and the kelly clamp removed. Five to ten cc. of sterile saline at room temperature was then poured over the frozen vessel. Often a primary aneurysmoid bulging was observed before the segment again began to pulsate and regain its tone. A fine silk ligature was inserted transversely in the left psoas muscle mass to mark the region of injury. The abdomen was closed with silk in three layers. The skin was closed with a continuous inverted mattress silk suture. Rabbits were closely observed until they recovered, usually in about 2-4 hours. They were then replaced in their original cages in the animal room, with drinking water ad libitum, and, as before, Purina rabbit chow.



Rabbits were sacrificed at appropriate times using 4-5 cc. intravenous Nembutal. Organs were fixed in 10% formalin for at least 24-48 hours. Tissues removed and examined included: brain and pituitary (occasionally), heart, lungs, aorta (intact, when possible), a portion of liver, adrenals, kidneys, spleen, abdominal fat and musculature. Animals dying overnight were not included in the study unless found within 1-2 hours of death.

Histological preparations were made both by frozen section followed by staining with Sudan IV for fat, and by routine hematoxylin and eosin staining for permanent sections.

Black and white photomicrographs were taken with an American Optical binocular microscope fitted with a Kodak Color Snap 35 mm. camera. Kodak Plux-X film (ASA 125) was used with a Wratten 58 (green) filter.

Serum lipid determinations were conducted as follows: fatty acids according to Man and Gildea's modification of Stoddard and Drury<sup>87</sup> followed by the method of Stern and Shapiro<sup>138</sup> when Triton was found to interfere with fatty acid precipitation in the former, phospholipids according to Hawk and Oser<sup>51</sup>, total and free cholesterol following Schoenheimer and Sperry<sup>127</sup>; triglycerides were obtained by the following calculations:

1-convert cholesterol and phospholipids to meq./liter.

a)  $\frac{\text{total-free cholesterol}}{38.6}$  or if no free cholesterol

total cholesterol x 0.0186

b) phospholipids x 0.58

2-triglycerides (meq./l.)=fatty acids (meq./l.) -

$[\text{chol. (total-free) (meq./l.)} + \text{phospholipids (meq./l.)}]$

Blood samples were immediately centrifuged and sera were refrigerated if not analyzed promptly.



Electrophoretic patterns were obtained by a cellulose acetate method (Beckman) courtesy of John Kelly<sup>72</sup>. The serum lipoprotein electrophoresis pattern of W.S., the patient reported herein, was obtained through the kindness of Dr. R. Scheig, Department of Medicine, Yale-New Haven Hospital.

Finally, hyperlipemic serum (removed from rabbits both twenty-two hours and four days after an injection of Triton) was injected aseptically through a 30 guage hypodermic needle into the superficial central portions of the corneas of anesthetized rabbits: these animals were then sacrificed at appropriate times and their corneas removed for both gross and microscopic observations.





RESULTS

The five rabbits given intravenous epinephrine showed no changes indicative of recent gross arterial injury. Such acute findings as petechial abdominal wall hemorrhages, pulmonary edema, generalized congestion and isolated regions of foam cells in peri-aortic abdominal arterioles were noted. The latter change was considered to be spontaneous. Only rabbit WA-4 showed a significant finding: A left ventricular apical aneurysm with subendocardial hemorrhages.

The two animals receiving Levophed<sup>(R)</sup> demonstrated no experimentally induced lesions, but one was soon able to tolerate huge doses of Levophed<sup>(R)</sup> with comparatively slight clinical effect. Bradycardia lasting about three minutes was noted after each dose.

One percent tyramine hydrochloride was given to two other rabbits: the first tolerated daily 7.5 cc. doses for six days and had no remarkable post-mortem changes. WA-9, however, succumbed with cardiac arrest following grand mal seizures minutes after a single dose of tyramine. No remarkable experimentally induced post-mortem findings were noted. (See Chart 1 for a summary of these preliminary experiments.)

Rabbits 10,12-42 received beef pituitary extract in doses of 100-200 mg. per animal subcutaneously in one of several forms (see Chart 2) in attempts to induce significant endogenous hyperlipemia according to procedures outlined by both Rudman<sup>120</sup> and by Kellner<sup>70</sup>. In addition, numbers WA-10-12, 21-24, 34-37, and 39-42 received one to three doses of sterile pooled horse serum intravenously to produce concomitant vascular injury following the technique of Rich and Gregory<sup>113</sup>. Doses of pituitary above 500 mg. occasionally yielded gross lipemia but more than 800 mg.



was toxic within twenty-four hours. Fatty acids, only, were sometimes elevated to 100-400 meq./l. No consistent experimentally induced vascular lesion was noted, and the narrow margin of safety with pituitary extract made the above methods unsatisfactory.

Intravenously administered non-ionic surface-active detergents were found by Kellner<sup>68</sup> and again by Cornforth<sup>23</sup> in 1951 to yield sustained endogenous hyperlipemias. It was decided to combine the use of detergents and direct aortic freezing, according to Taylor (1950)<sup>142</sup> and Kelley et. al. (1952)<sup>71</sup>, to obtain regularly reproducible hyperlipemia concomitant with known vascular injury. With one intravenous dose of ten percent sterile Triton solution grossly evident lipemias were consistently obtained. The chemical effects on the blood of a single 7-11 cc. dose were evident within a few hours. Sera remained lactescent for approximately six days and chemical hyperlipemias were evident for about eight to ten days (see Charts 5 through 8). Total and free cholesterol rose from average baseline values of 69.3 mg.% (range 33-161 mg.%) and 21.3 mg.% (range 9.1-48 mg.%) to peaks of 470-740 mg.% and 335-490 mg.% respectively between three and seven days. Slightly elevated values were still evident two weeks after the single dose of Triton.

Fatty acids abruptly increased from average baseline amounts of 11.0 meq./l. (range 4.0-18.0 meq./l.) to values of 125-290 meq./l. within 48-72 hours followed by a steady decline to control after nine to thirteen days. Triglycerides followed in like fashion. Triton was found to interfere with precipitation of fatty acids in Stoddard and Drury's modified procedure, consequently, Stern and Shapiro's method as indicated was employed with success. A comparison of the two procedures performed to



verify accuracy on sera not containing Triton may be found on Chart 4. Close agreement of determinations was noted.

Phospholipids peaked in two to five days in the range of 22-55 mg.%. Pre-Triton levels of 5.2 mg.% (range 2.5-8.6 mg.%) were approximated after two weeks.

It was remarkable that every rabbit given a single dose of about 250 mg. Triton responded with a predictable rise in fatty acids (and, therefore, triglycerides), followed by steady elevations in both phospholipids and total and free cholesterol. The 22 hour pooled centrifuged sera with moderate total cholesterol elevations (from rabbits WA-57 and WA-58) were found to invoke only slight corneal foam cellular reactions in rabbits WA-61 and WA-62. (see Figures 1 and 2).

By contrast four day post-Triton serum was noted to induce more corneal reaction with moderate numbers of foam cells and lipid observable between and apparently within collagen fibrils. (WA-65 and 66). This serum contained high amounts of cholesterol and less lipid in the form of triglycerides. Corneal foam cellular reactions obtained by Grauer<sup>48</sup> with sera containing greatly increased cholesterol in the range of 100 mg.% were significantly more intense (Figure 4). It would appear, then, that the high cholesterol lipid fractions exert a greater atherogenetic effect in the rabbit corneal models than do other lipid types.

Frozen sections from the aortas, coronary, and pulmonary arterioles of rabbits sacrificed within 24-48 hours after a single Triton injection, however, demonstrated significant sudanophilic material. Increased uptake of lipid was noted in selectively injured aortic segments. Grossly fatty livers and peri-aortic postoperative inflammation and hemorrhage were seen.





After 4-5 days the livers were still markedly fatty and blood cholesterol was elevated substantially. Segments of aortas formerly subjected to freezing in the first 24 hours of lipemia showed extensive periadventitial hemorrhage and granulation tissue with polymorphonuclear leucocytes and mononuclear cells, medial hyalinization and necrosis, and subintimal separation and proliferation (Figures 5-8,19). Higher power views revealed the subendothelial space accumulating monocytes and small foam cells (Figures 8,9,12,13). So called "tadpole" cells were observed in the aortic lumen (Figures 10,11) and foam cells were seen migrating through the internal elastic lamina (Figure 15). Lipid was also present between intimal elastic laminae in intracellular droplets (Figures 8,9,13-18).

Three types of controls were examined. Unfrozen, unmanipulated portions of the aorta, and other arteries in animals subjected to either Triton or to aortic freezing were scrutinized for signs of the effects of Triton. No lipid infiltration was noted in uninjured aortic segments. In addition, aortas from rabbits WA 68-72 (animals subjected solely to Triton and periodic blood collection) showed no regions of lipid deposition. Pulmonary arterioles, however, frequently contained sudanophilic intimal and medial deposits sometimes with foam cellular responses.

Two rabbits (WA-58 and 59) were given intravenous sterile normal saline instead of Triton and subsequently underwent aortic freezing. Injured aortic segments after eleven days contained medial calcification, subintimal proliferation and only trace amounts of lipid (Figures 20,21). Pulmonary arterioles showed subintimal swelling without any stainable lipid. Lungs were essentially normal with some regions of isolated congestion. Heart, liver and kidney were considered normal. Rabbit WA-53,





that was sacrificed at the same time, but after both Triton and freezing, proved to have similar aortic findings but with the addition of slight liquefaction and lipid in the inner media, and small amounts of remaining hepatic lipid.

WA-55 and 56 received Triton but their abdominal aortas were touched with aneurysm hooks instead of dry ice for thirty seconds. The remainder of their treatment was identical to that of rabbits whose aortas were frozen. After one week, the aortas had thickened folded internal elastic lamellae but no subintimal proliferation or stainable lipid. Peri-aortic arteriolar medial lipid was noted however. No stainable fat was seen in the livers. WA-55 had phlebitis of the inferior vena cava, and renal arterial intimal edema was noted in conjunction with focal round cellular renal infiltrates in WA-56.

Twenty days elapsed after the aorta of WA-49 was frozen. Virtually no hepatic lipid was noted, nor lipid in the aorta. Slight subintimal proliferation and extensive medial calcification and necrosis was observed (Figure 22). The lungs were of interest; they contained intimal proliferation of the pulmonary arterioles, and lipoid pneumonia was present (Figure 24) with hemorrhage, foam cells (Figure 25), mononuclear cells, and giant cells (Figure 26). (For comparison, Figure 23 illustrates a region of lung from WA-60, an animal receiving no Triton.) Finally, phlebitis of the inferior vena cava was evident. By comparison, lipid was noted in the aorta of WA-50 (Figure 27) twenty-eight days after freezing, and subintimal proliferation (Figure 27) along with widespread medial necrosis were seen (Figure 28). The liver still contained some lipid (Figure 29).



The final animals (WA-45 and 51) were sacrificed three months after a single injection of Triton followed by aortic freezing. One (WA-45) showed medial liquefaction and calcification with trace amounts of stainable lipid. Both rabbits had pulmonary arteriolar intimal proliferation. WA-45 had a myocardial infarction located in the interventricular septum.

An incidental finding was that rabbits given Triton suffered weight losses of 1-2 pounds in the first two weeks. They ate and drank during this period (although precise amounts were not measured) and they occasionally had diarrhea. Abdominal fat reserves were slightly reduced.

No glomerular foam cell lesions were noted (reported by Vidone and Lowman<sup>148</sup> in dogs). Our rabbits, however, received only single doses of Triton.



## DISCUSSION

### A. Spontaneous Arterial Lesions in the Rabbit.

Three types of spontaneous aortic lesions have been described in the rabbit. Israel, in 1881<sup>65</sup> named "chronic endo-aortitis deformans", a spontaneous medial degeneration and calcification. Quadri (1907)<sup>109</sup>, Miles and Johnstone (1907)<sup>92</sup>, Ophuls (1907)<sup>97</sup>, Pearce (1908)<sup>102</sup>, Lucie and Parisot (1908)<sup>85</sup>, Hill (1910)<sup>54</sup>, and Nazum et al. (1930)<sup>95</sup> found similar lesions but reported different incidences. Kesten<sup>73</sup> found aortic and iliac artery medial calcifications in 87% of eight month old rabbits. Atheromatous yellow plaques were seen in the adult aortic arch by Ophuls in 1907 and by Nazum et al. in 1930. Finally, a raised polypoid region consisting of variously-directed layers of smooth muscle cells in a mucopolysaccharide stroma was described by De Faria in 1955<sup>28</sup> and confirmed by Haust and More ten<sup>50</sup> years later.

In addition, spontaneous muscular pulmonary artery thickening with splitting of the internal elastic lamina, increased number of elastic fibers, and a narrowed vessel lumen were presented by Prior et al. in 1961<sup>108</sup>.

As Haust and More indicate, however, the frequency of these lesions varies greatly according to observers, and all factors, such as strain background, living conditions (temperature, humidity, barometric pressure, availability of food and water, exposure to other species and each other and parasitic and infectious diseases), and age. In general, however, spontaneous atherosclerosis is rare in rabbits.<sup>26</sup>





B. Cholesterol Feeding Experiments in Brief.

In 1908, milk, eggs and meat were noted to be atherogenic in rabbits<sup>120a</sup>. Ignatowski<sup>62</sup> corroborated these findings in 1909. Many investigators have administered cholesterol and cholesterol-modified diets since that time. Reviews of progress in experimental atherogenesis using cholesterol feeding models appear periodically<sup>26,160,22</sup>. At this time prolonged cholesterol feeding in animals is the basic experimental model for human atherosclerosis.

Many difficulties are encountered in cholesterol feeding. Somatic afflictions of experimental animals<sup>22</sup>, and long duration are two examples. Vascular lesions produced do not always occur in regions comparable to natural lesions in humans.





### C. Endogenous Lipemias and Triton.

Perhaps the first experimental induction of a lipemia not related to feeding of exogenous lipids was reported in 1909 by Boggs and Morris<sup>7</sup>. They found rabbits subjected to daily bleedings of 25 cc. developed hyperlipemia after eight to sixteen days. Friedland's experience with desiccated thyroid will be mentioned later<sup>40</sup>. Reticulo-endothelial blockade with substances such as India ink has also led to lipemia.

Increased blood lipids are noted during exposure to cold, therapy with corticoids<sup>114</sup> and in individuals afflicted with the nephrotic syndrome<sup>46,118,131</sup>. Zarafonitis mentioned these as well as femoral fracture, protamine and diisopropylfluorophosphate for inducing lipemia. Cobaltous chloride led to similar blood lactescence in experiments reported by Caplan and Block<sup>14</sup> and by Brody<sup>9</sup>. Additional hyperlipemic conditions were reviewed by Scanu in 1965. Cushing's syndrome, essential xanthomatosis, excess growth hormone, increased catecholamines, obstructive jaundice and androgen influence were also mentioned.

Triton, a non-ionic detergent, was first associated with hyperlipemia by Hueper in 1944<sup>61</sup>. Triton in a 0.1 percent solution was administered both orally and intravenously to rabbits and was associated with severe toxic manifestations. Foam cellular lesions were noted in pulmonary arteries and aorta. The toxicity of Triton apparently deterred other investigators until, in 1951, Cornforth<sup>23</sup> found elevation of blood lipids after testing Triton for experimental tuberculosis therapy. Later that year, Kellner, Correll and Ladd<sup>68,69</sup> reemphasized the hyperlipemic effects of Triton.



From this point on, investigators wondered about the mechanism of action of Triton. Frantz<sup>37</sup>, Friedman<sup>41,42</sup>, Hirsch and Kellner<sup>55,56,57</sup>, Kellner<sup>70</sup>, Otway<sup>98</sup>, Pawar<sup>101</sup>, Pethica<sup>103</sup>, Radding<sup>110</sup>, Scanu<sup>121-4</sup>, Schotz<sup>128</sup>, Vidone<sup>148</sup>, Zarafonitis<sup>165,166</sup> advanced theories about Triton's action in the plasma, reticuloendothelial system, liver and posterior pituitary. The present author believes its actions holding lipids in the plasma and inhibiting removal by the liver are particularly significant. Triton may interfere with specific enzyme systems and block reactions at membrane surfaces.

Triton-induced lipemia has a characteristic pattern. During the first few hours the serum contains increasing amounts of fatty acids until lactescence is noted. The next day cholesterol (both total and free) build up to values several times normal, even in animals maintained on low-fat diets. Triglycerides and fatty acids reach peak amounts on the second day and steadily decline, while phospholipids and cholesterol remain markedly elevated for about one week. Electrophoretic patterns of this lipemia demonstrate a pattern similar to Fredrickson's Type III; both beta and pre-beta fractions are increased.



D. Experimental Vascular Injury.

Interest in experimental production of vascular injury has steadily increased since the turn of the century. In 1903, Josué reported experimental aortic lesions after repeated injections of adrenalin<sup>(R)</sup><sup>66</sup>. Waterman<sup>149</sup> noted medial necrosis and calcification of small muscular arterioles in 1908. Anitchkow<sup>2</sup> employed adrenalin<sup>(R)</sup> for vascular injury in 1913. Schmidtman obtained selective focal arterial damage in 1929 using vitamin D overdosage<sup>126</sup>. Desiccated thyroid was found by Friedland to produce experimental arterial injury (cited by Anitchkow in E.V. Cowdry, 1933)<sup>40</sup>.

Goldblatt<sup>46</sup> reported arteriolonecrosis in dogs with malignant hypertension secondary to induced renal ischaemia in 1938. Duff and Rich in 1936 used subcutaneous trypsin injections, Duff, again, in 1939<sup>32</sup> employed tyramine in rabbits and Waters, nine years<sup>150</sup> later obtained lesions using allylamine in dogs.

More recent chemical procedures for producing vascular injury have included: seven percent saline diets given to rats to induce nephrotic syndromes with edema, hypertension, anemia, lipemia, hypoproteinemia and severe arteriolar disease (1958), aminopropionitriles, according to Schwartz<sup>130</sup> (1959), and diethylstilbestrol injections that produced dissecting aneurysms and aortic ruptures in turkeys (1966)<sup>134</sup>. Norepinephrine<sup>24,140,141</sup> was noted to induce arterial and myocardial damage in patients.

Serum sickness has served as an elegant manner of inducing experimental vascular injury since studies by Fleischer et. al<sup>36</sup> (1931), Clark and Kaplan<sup>17</sup> (1937) and Rich and Gregory's well-known reports in 1942 and 1943<sup>111-113</sup>.





Direct physical trauma to arteries has consistently resulted in vascular injury. Ssolowjew<sup>136,137</sup>, in 1929, cauterized blood vessels. Juvenile rabbits' aortas were frozen by Taylor et. al. in 1950<sup>142</sup>. Kelly, Taylor and Hass<sup>71</sup> directly froze cholesterol-fed rabbits' abdominal aortas with ethyl chloride spray two years later. Prior and Hartman<sup>106</sup> scraped aortic intimas of hyperlipemic animals with a hooked needle and found increased atherogenesis in 1956. Williams, five years later<sup>159</sup>, had similar results with only surgical mobilization of hyperlipemic animals' aortas, and Baumgartner et. al.<sup>4</sup> and Gutstein et. al.<sup>49</sup> stretched aortas to obtain severe vascular disease. In 1968, Hoff<sup>58</sup> merely subjected aortas to brief ligation to observe changes.

A controversy exists in the literature regarding the ability of Triton to induce morphologic counterparts of atherosclerosis in experimental animals. Courtice and Schmidt-Diedrichs<sup>25</sup> reported no retained lipid or foamy lipophages in carotid arteries of rabbits injured by local infusions of heated saline. At the times of injury these animals had a Triton-induced hyperlipidemia. In contrast, Vidone and Lowman<sup>148</sup> found extensive free and foam-cellular intimal lipids in aortas and coronary and pulmonary arteries of dogs subjected to repeated injections of Triton alone. Grossly these arterial lesions appeared as fatty streaks. These dogs received no other form of experimental vascular injury.

The results of the present experiments would indicate that an experimental endogenous lipemia, similar to Fredrickson's Type III in humans, does indeed alter acute arterial lesions in rabbits to morphological sequences like those of early atherosclerosis in man. This evidence broadens our concepts of the role of lipids in atherogenesis and indicates



that the emphasis on study of exogenous dietary cholesterol in the process may have dealt with only part of the problem.



SUMMARY AND CONCLUSIONS

Seventy-four rabbits on low-fat diets were employed in a study of experimental vascular disease with concomitant endogenous lipemia. The resultant lesions were considered in relation to human atherosclerosis and to the lipoprotein patterns present in the animals' blood. Preliminary experiments with epinephrine, norepinephrine, tyramine and sensitization with horse serum proved unsatisfactory methods for producing vascular damage. Injection of pituitary extract produced only occasional mild hyperlipemias insufficient in magnitude for our purposes. Also this procedure was very toxic to the experimental animals.

Direct physical trauma to intact abdominal aortic segments in vivo provided predictable vascular injury with medial necrosis and calcification and subintimal proliferation. Spontaneous subintimal reaction in pulmonary arterioles was often seen.

Triton WR 1339, a non-ionic detergent, when administered as a single intravenous dose of 250 mg./kg. body weight led to significant elevation in plasma lipids; first triglycerides, then cholesterol (total and free) accompanied by phospholipids rose to amounts sufficient to produce grossly milky serum. Lipemias remained at peak levels for three to seven days and were approaching control levels in nine to fourteen days. Electrophoresis revealed serum lipoproteins in a pattern similar to Fredrickson's Type III described in patients. An example of a patient with a secondary Type III pattern was presented for comparison.

Histological preparations of aortas from animals with concomitant vascular injury and Triton hyperlipemias revealed aortic lesions with many hallmarks of human atherosclerosis including subintimal proliferation,



foam cellular accumulation with lipid deposits in the intima and media, medial necrosis and liquefaction with calcification, and periadventitial reaction. "Tadpole cells" (young foam cells) were seen in the aortic lumen and penetrating the internal elastic membrane. The membrane, itself, was seen accumulating lipid droplets within its cells, and one cell was photographed in mitosis. Pulmonary arterioles were noted to accumulate stainable lipid occasionally. Livers remained fatty for between one and two weeks after Triton. Lipoid pneumonia with giant cells and foamy alveolar macrophages was noted in some instances.

Rabbit corneas were used as sites for ancillary models of atherosclerosis following concepts and procedures outlined by Waters<sup>154,155,157</sup> and by Grauer<sup>48</sup>. Twenty-two hour, post-Triton serum produced little corneal stromal lipophagic reaction whereas four day, post-Triton serum containing high cholesterol values and moderately increased fatty acids yielded mild foam cellular responses. Even these reactions were less than the known marked responses in corneas injected with serum having very high cholesterol levels<sup>48</sup>. It would appear, then, that cholesterol fractions are more atherogenic, at least in the corneal models.

Aortic freezing and Triton injection provide a reliable and readily obtainable model with many characteristics of human atherosclerosis.








Figure 1. WA-61 rabbit cornea eight days after instillation of twenty-two hour post-Triton serum. Scarce foam cells.  
(H & E, 450x original magnification)




Figure 2. WA-61 cornea. Foam cells.  
(H & E, 1000x original magnification)





Figure 3. WA-61 cornea. Droplets of lipid within collagen fibrils and extracellularly.  
(Sudan IV. Frozen section. 450x original magnification)




Figure 4. WLG-994-RC cornea six days after instillation of hypercholesterolemic rabbit serum. Abundant intra- and extracellular lipid.  
(Sudan IV. Frozen section. 100x original magnification. Courtesy L. Graver, 1968)








Figure 5. WA-44 aorta five days after single injection of Triton and four days after aortic freezing. Intimal proliferation, medial necrosis, periadventitial hemorrhage, granulation and fat. (H & E, 100x original magnification).




Figure 6. WA-44 aorta. Subintimal foam cellular accumulation, medial necrosis, adventitial hemorrhage. (H & E, 100x original magnification).








Figure 7. WA-44 aorta. Subintimal proliferation, medial necrosis, adventitial reaction with polys and mononuclear cells. (H & E, 100x original magnification).




Figure 8. WA-44 aorta. Subendothelial space with many foam cells. (H & E, 450x original magnification).






Figure 9. WA-44 aorta. Monocytes and young foam cells in subendothelial space.  
(H & E, 450x original magnification).




Figure 10. WA-44 aorta. "Tadpole" cell (young foam cell) in aortic lumen.  
(H & E, 1000x original magnification).






Figure 11. WA-44 aorta. Intimal foam cells, "tadpole" cell in lumen. (H & E, 450x original magnification).




Figure 12. WA-44 aorta. Foam cell within subendothelial space. (H & E, 1000x original magnification).








Figure 13. WA-44 aorta. Subendothelial foam cells, multinucleated endothelial cell.  
(H & E, 1000x original magnification).




Figure 14. WA-44 aorta. Same as Figure 13 but different exposure.  
(H & E, 1000x original magnification).








Figure 15. WA-44 aorta. Lipid-laden macrophage entering subendothelial space.  
(H & E, 1000x original magnification).




Figure 16. WA-44 aorta. Lipid droplets within foam cell beneath internal elastic lamina.  
(Sudan IV, frozen section. 150x original magnification).






Figure 17. WA-44 aorta. Foam cells with lipid droplets within intima.  
(Sudan IV, frozen section. 450x original magnification).




Figure 18. WA-44 aorta. Foam cellular and "free" lipid deep within  
intima.  
(Sudan IV, frozen section, 450x original magnification).





Figure 19. WA-44 aorta. Endothelial cell in mitosis and young sub-endothelial foam cells.  
(H & E, 1000x original magnification).




Figure 20. WA-60 aorta thirteen days after intravenous saline and eleven days after freezing. Marked subintimal cellularity and proliferation.  
(H & E, 100x original magnification).







Figure 21. WA-59 or 60 aorta. Intimal proliferation with little visible lipid in aortic wall.  
(Sudan IV, frozen section, 100x original magnification).




Figure 22. WA-49 aorta twenty-four days after Triton and twenty days after freezing. Proliferated intimal region, medial necrosis with calcification.  
(H & E, 100x original magnification).






Figure 23. WA-60 lung demonstrates no pneumonia.  
(H & E, 40 x original magnification).




Figure 24. WA-49 lung. Lipoid pneumonia with mononuclear cells, giant cells, hemorrhage and foam cells.  
(H & E, 100x original magnification).








Figure 25. WA-49 lung. Lipoid pneumonia with foamy lipophages.  
(H & E, 450x original magnification).




Figure 26. WA-49 lung. Lipoid pneumonia with giant cell.  
(H & E, 450x original magnification).





Figure 27. WA-50 aorta thirty-two days after Triton and twenty-eight days after aortic freezing. Subintimal proliferation and slight stainable lipid. (Sudan IV, frozen section, 100x original magnification).




Figure 28. WA-50 aorta. Extensive medial necrosis. (H & E, 40x original magnification).







Figure 29. WA-50 liver. Small amount of lipid still present thirty-two days after single Triton injection. (Sudan IV, frozen section, 40x original magnification).



## Chart 1

## Summary of Preliminary Experiments with Pressors

<u>Rabbit</u>	<u>Pressor Used</u>	<u>Wt. (lb.)</u>	<u>Sacrifice</u>	<u>Results</u>
WA-1	epinephrine 1 cc. 1:10000 10-1 cc. injections/15 min.	~ 6	*20 min.	(mydriasis, cyanotic, weak. died 5 min. after final inject.) Pulmonary congestion and edema. abdominal wall musc. petechiae.
WA-2	epinephrine 1:10000 4-1 cc. injections/15 min.	~6	*2-4 hrs.	died 2-4 hrs. after epinephrine.
WA-3	epinephrine 1:10000 3-1 cc. injections/90 min.	6	*12 hrs.	renal tubular edema.
WA-4	epinephrine 1:10000 2-1 cc., then 1-2 cc. injection q. 24 hrs.	~ 6	5th. day	fibrin clot right pleural cavity. left ventricular aneurysm. pulmonary congestion.
WA 5	epinephrine 1:10000 1-2 cc. injection.	4.7	54 hrs.	petechial pulmonary hemorrhages
WA-6	Levophed <sup>(R)</sup> (norepinephrine) 1 cc. 1/5 strength	7.4	45 hrs.	double left renal artery. lungs clear pulmonary effusion.
WA-7	Levophed 2 cc., then 5 cc. 1/5 strength at 48, 96 hrs.	6.5	1 week	rapid induction tolerance. congested liver, lungs, kidneys. sacrificed week after <u>first</u> Levophed.
WA-8	Tyramine-HCl 7.5 cc. of 1% over 2 min./ repeat q. 24 hrs. for 6 days.	5	8th. day	perivascular edema in lungs, heart. sacrificed 8 days after <u>first</u> Levophed.
WA-9	Tyramine-HCl 7.5 cc. of 1% over 1 min.	8.2	*3 min.	5x7 mm. accessory spleen. tachycardia → cardiac arrest, seizures → death.

\* Did not survive length of experiment.



## Chart 2

## Summary of Experiments Using Pituitary Extract and/or Horse Serum

Key to types of pituitary extract used: 1-whole dried beef pituitary extracts/10 cc. sterile saline.  
 2-AS#1, but filtered with ether.  
 3-AS#1, but boiled in ethanol/1 hour.  
 4-Defatted beef pituitary extract in saline.

<u>Rabbit</u>	<u>Pituitary Extract</u> <u>X type (mg)</u>	<u>Days</u> <u>Given</u>	<u>Horse Serum</u> <u>(cc. and # of</u> <u>times given)</u>	<u>Days</u> <u>Given</u>	<u>Findings</u>
WA-10	130 mg. <sup>1</sup>	22	25 cc. x 2	1,20	No lipemia.
WA-11			25 cc.	1	Found dead overnight.
WA-12	130 mg. <sup>1</sup>	22	25 cc. x 2	1,20	No lipemia.
WA-13	100 mg. <sup>4</sup>	1			Found dead overnight.
WA-14	100 mg. <sup>1</sup> , 420 mg. <sup>3</sup>	1,20			Serum samples at 24, 52, 100 hours.
WA-15	200 mg. <sup>4</sup> , 600 mg. <sup>2</sup>	1,20			Serum samples milky at 24, 52, 100 hours.
WA-16	200 mg. <sup>1</sup>	1			Serum sample at 24 hours, then died.
WA-17	2000 mg. <sup>1</sup>	1			Found dead next day, autolysed.
WA-18	1000 mg. <sup>1</sup>	1			Found dead next day, autolysed.
WA-19	1000 mg. <sup>1</sup>	1			Died next day after blood drawn; lipemia, fatty liver.
WA-20	1000 mg. <sup>1</sup>	1			Serum at 24 hours lipemia; died one hour later; pulmonary edema.





<u>Rabbit</u>	<u>Pituitary Extract X type (mg)</u>	<u>Days Given</u>	<u>Horse Serum (cc. and # of times given)</u>	<u>Days Given</u>	<u>Findings</u>
WA-21			20 cc. x 3	1,7,14	Died 4 minutes after third serum; anaphylaxis.
WA-22	600 mg. <sup>2</sup> x 2	17,21	20 cc. x 3	1,7,14	Serum not ipemic.
WA-23	600 mg. <sup>2</sup> x 2	17,21	20 cc. x 3	1,17, 14	
WA-24	600 mg. <sup>2</sup> x 2	17,21	20 cc. x 2 then 12 cc.	1,17, 14	Died with broken back.
WA-25	750 mg. <sup>1</sup>	1			Diarrhea 45 min. after pituitary; found dead next AM.
WA-26	500 mg. <sup>1</sup>	1			Died in 48 hours; autolysed; left renal cortical scar.
WA-27	750 mg. <sup>1</sup>	1			Found dead next AM with abdominal hemorrhage.
WA-28	1000 mg. <sup>3</sup>	1			Died day 2 while being bled.
WA-29	375 mg. <sup>1</sup>				No gross lipemia.
WA-30	200 mg. <sup>2</sup> , 400 mg. <sup>2</sup> , 600 mg. <sup>2</sup> , 500 mg. <sup>2</sup>	1,3, 9,10			No gross lipemia.
WA-31	200 mg. <sup>2</sup> , 400 mg. <sup>2</sup> , 600 mg. <sup>2</sup> , 500 mg. <sup>2</sup> , 1000 mg. <sup>2</sup>	1,3 4,5 18			



<u>Rabbit</u>	<u>Pituitary Extract</u> X type (mg)	<u>Days</u> <u>Given</u>	<u>Horse Serum</u> (cc. and # of times given)	<u>Days</u> <u>Given</u>	<u>Findings</u>
WA-32	2 200 mg., 400 mg., 600 mg., 650 mg.	1,3,9, 10			
WA-33	2 350 mg., 400 mg.	1,3			Sacrificed day 4
WA-34	2 700 mg.	14	20 cc. x 3	1,7, 13	Serum slightly lipemic on day 15.
WA-35			20 cc. x 3	1,7, 13	Diarrhea day 8; died day 13; anaphylaxis and congested liver.
WA-36	2 700 mg.	15	20 cc. x 3	1,8, 14	
WA-37	2 700 mg.	15	20 cc. x 3	1,8, 14	Serum milky on day 16.
WA-38	2 600 mg., 500 mg.	1,2			Diarrhea AM day 2
WA-39	2 650 mg.	25	33 cc. x 3	1,14, 21	One pulmonary arteriole with fibrinoid necrosis; pericarditis; liver not fatty; diffuse plural effusion; normal aortas and kidneys.
WA-40	2 650 mg.	25	20 cc. x 1	1*	
WA-41	2 650 mg.	25	33 cc. x 3	1,14, 21	
WA-42	2 650 mg.	25	33 cc. x 3	1,14 21	

\* Stolen from animal room day 2.



Chart 3

Baseline Serum Lipid Values of Rabbits

<u>Rabbit</u>	<u>Weight (lbs.)</u>	<u>TTL Chol. mg.%</u>	<u>Free Chol. mg.%</u>	<u>Fatty Acids meq./.</u>	<u>Phos-L. mg.%</u>	<u>Triglyc. meq./1</u>
WA-13	5.4	60.6	33.0	15.0	6.9	11.3
14	4.8	81.0	48.0	15.0	6.0	10.6
15	5.25	60.6	19.0	15.0	6.1	10.4
16	5.0	72.0	33.0	17.0	7.0	12.9
17	-----Died-----					
18	5.0	74.5	21.0	12.0	5.5	7.4
19	7.0	46.0	20.0	12.6	4.1	9.5
20	7.0	72.5	43.0	12.0	4.8	8.5
21	4.75	-----Died-----				
22	4.75	65.1	13.0	13.6	5.1	9.3
23	4.75	92.1	13.0	10.4	4.3	6.8
24	5.0	67.1	9.1	8.0	3.8	4.3
25	5.0	-----Died-----				
26	5.0	128.5	---	14.4	5.8	8.6
27	7.0	-----Died-----				
28	5.0	74.3	---	11.2	5.6	6.6
29	5.0	39.4	9.1	11.2	4.0	8.1
30	7.0	47.0	13.9	12.0	4.3	8.7
31	7.0	44.0	11.7	8.0	3.8	5.0
32	7.0	77.5	17.0	11.2	5.6	6.4
33	7.0	39.0	27.0	7.2	2.8	5.3
38	4.5	65.5	10.2	15.7	6.7	9.9
43	8.9	36.4	---	6.8	3.0	4.4
44	8.25	47.6	---	8.0	5.2	4.1
45	8.4	70.2	---	18.0	6.0	13.2
46	10.4	23.0	---	6.0	2.5	4.1
47	7.9	47.6	---	8.0	3.3	5.2
48	9.4	24.4	---	18.0	4.4	15.0
49	8.5	54.0	---	8.0	5.8	3.6
50	9.3	29.8	---	4.0	2.5	1.9
51	8.5	74.4	---	12.0	6.4	6.9
53	8.5	106.8	---	10.7	8.6	3.7
54	8.5	72.4	---	9.7	6.0	4.9
55	7.9	60.6	---	8.3	5.6	3.9
56	8.4	108.2	---	8.3	5.5	3.1
59	6.1	148.4	---	18.0	9.0	10.0
60	9.25	161.0	---	14.5	8.5	1.2
67	---	67.0	---	9.5	5.5	5.1
68	---	83.0	---	9.5	5.0	3.1
69	---	95.0	---	9.5	6.0	4.2
70	---	59.6	---	10.2	3.1	6.3
71	---	43.0	---	5.0	4.0	1.9
72	---	85.4	---	7.5	6.0	2.4



Chart 4

A Comparison of Fatty Acid Determinations on Baseline Samples

<u>Rabbit</u>	<u>Fatty Acids meq./1 Stoddard &amp; Drury</u>	<u>Fatty Acids meq./1 Stern and Shapiro</u>
53B	10.0	11.3
54B	9.3	10.0
43-9 days	13.0	9.5
55B	8.6	8.0
56B	8.6	8.0
49#3	7.3	8.7
53#3	40.0	40.5
55#3	18.0	15.5
56#3	52.6	54.5





Chart 5

Triton-Chemistries

<u>Rabbit</u>	<u>Time</u>	<u>Cholesterol</u> mg.%		<u>Fatty</u> <u>Acids</u> meq./l.	<u>Phospho-</u> <u>lipids</u> mg.%	<u>Trigly-</u> <u>cerides</u> meq./l.
		<u>Total</u>	<u>Free</u>			
WA-43	Base	36.4		6.8	3.0	4.4
	5½ hr	118.1	34.3	32.5	8.5	25.4
	23 hr	330.0	182.0	90.0	25.0	71.7
	9 days	145.0	41.5	11.8*	7.8	
WA-44	Base	47.6		8.0	5.2	4.1
	23½ hr	366.6	149.0	150.0	37.5	122.6
	120 hr	507.9	350.0	125.0	45.0	95.1
WA-45	Base	70.2		18.0	6.0	13.2
	23½ hr	461.6	270.0	180.0	41.3	151.2
WA-46	Base	23.0		6.0	2.5	4.1
	17 hr	238.0	68.0	110.0	22.5	92.6
WA-47	Base	47.6		8.0	3.3	5.2
	17 hr	226.6	103.0	75.0	20.0	60.2
	48 hr	482.2	322.0	250.0	43.5	221.4
WA-48	Base	24.4		18.0	4.4	15.0
	17 hr	225.0	113.8	142.5	23.8	125.8
	48 hr	456.5	351.0	210.0	42.5	182.4
WA-49	Base	54.0		8.0	5.8	3.6
	70 hr	594.3	491.0	205.0	47.5	174.4
	19 days					
WA-50	Base	29.8		4.0	2.5	1.9
	70 hr	556.0	299.0		53.2	
	27 days					
WA-51	Base	74.8		12.0	6.4	6.9
	70 hr	443.0	385.0	165.0	43.5	138.1
WA-53	Base	106.8		10.7*	8.6	
	18 hr	280.0	170.0	107.0	28.0	
	12 days			40.2*		?#3
WA-54	Base	72.4		9.7	6.0	
	18 hr	220.0	102.5	95.5	24.0	
WA-55	Base	60.6		8.3	5.6	
	20 hr	150.0	106.0	55.0	17.5	?#3
	7 days			16.8		



<u>Rabbit</u>	<u>Time</u>	<u>Cholesterol</u> mg.%		<u>Fatty</u> <u>Acids</u> meq./l.	<u>Phospho-</u> <u>lipids</u> mg.%	<u>Trigly-</u> <u>cerides</u> meq./l.
		<u>Total</u>	<u>Free</u>			
WA-56	Base	108.2		8.3	5.5	
	20 hr	247.5	167.4	84.0	21.5	?#3
	7 days			53.5		

FOR CORNEAL INJECTION

WA-57		224.1	120.0		8.3	5.5
WA-58		269.8	175.5	10.0	2.5	6.1
Supernate x 0.1		70.0	34.1	15.0	8.8	8.9
Subnate		168.0	127.5	35.0	20.0	22.3
WA-59	Base	148.4		18.0	9.0	10.0
(Saline Control)	42 hr	160.7	34.1	12.6	11.0	3.0
Freezing	13 days	63.5	12.4	12.0	5.0	8.3
WA-60	Base	161.0		14.5	8.5	1.2
(Saline Control)	42 hr	127.5	37.8	17.6	15.0	6.6
Freezing	13 days	113.5	13.0	12.5	6.0	5.8
WA-63	95 hr	475.0	438.0	210.0	46.0	182.4
WA-64	95 hr	745.0	337.0	110.0	29.0	92.0
WA-67	Base	67.0		9.5	5.5	5.1
	1 hr/ 20 min.	57.5	24.6	10.6	3.0	7.7
	12 hr/ 40 min.	136.1	88.0	46.7	12.0	38.4
	30 hr	301.0	245.5	102.5	24.5	86.8
	29 hr/ 40 min.					
WA-68	Base	83.0		9.5	5.0	3.1
	1 hr	65.01	24.6	9.8	4.5	6.0
	12 hr/ 35 min.	136.1	99.0	57.5	14.0	48.4
	30 hr	305.4	215.0	127.5	23.0	111.8
	29½ hr					
	7½ days	350.2	137.5	27.5	17.5	11.5
	13 days	200.0	56.2	21.0	9.0	12.1



<u>Rabbit</u>	<u>Time</u>	<u>Cholesterol</u> mg. %		<u>Fatty</u>	<u>Phospho-</u>	<u>Trigly-</u>
		<u>Total</u>	<u>Free</u>	<u>Acids</u> meq./l.	<u>lipids</u> mg. %	<u>cerides</u> meq./l.
WA-69	Base	95.0		9.5	6.0	4.2
	1 hr/ 40 min.	79.0	34.0	10.5	5.0	6.1
	12 hr/ 25 min.	162.5	131.0	97.5	16.4	86.1
	78 hr	367.3	234.4	125.0	22.5	107.3
	7½ days	630.8	534.0	83.0	46.7	53.3
	13 days	175.5	49.9	20.0	9.0	11.6
	WA-70	Base	59.6		10.2	3.1
1½ hr		62.5	31.0	12.5	5.5	8.1
12¼ hr		124.5	95.0	50.0	13.1	41.7
78 hr		662.0	320.0	290.0	45.0	255.0
7½ days		726.3	526.3	115.0	60.0	77.0
13 days		205.0	41.5	20.0	11.0	9.5
WA-71		Base	43.0		5.0	4.0
	30 hr	388.5	235.5	125.0	24.0	109.8
	76 hr	602.5	266.4	205.0	44.0	160.6
	95 1/3 hr	635.0	507.9	190.0	52.0	156.6
	7½ days	496.3	299.0	60.0	40.0	31.7
	13 days	56.0	13.0	15.0	5.0	11.5
	WA-72	Base	85.4		7.5	6.0
28 hr/ 5 min.		388.4	306.0	155.5	29.0	136.6
76¼ hr		585.2	266.3	175.0	42.5	141.9
95¼ hr		680.0	594.3	118.0	44.0	90.3
7½ days		500.0	260.0	27.5	15.0	12.6
13 days		189.0	50.0	19.0	9.0	10.2





Chart 6



Chart 7







Chart 8







## Chart 9

## Rabbits Receiving Triton and/or Aortic Freezing

<u>Number</u> <u>Weight (lbs.)</u> <u>Sex</u>	<u>Amount (cc.)</u> <u>of Triton,</u> <u>Day Given</u>	<u>Day of</u> <u>Aortic</u> <u>Freezing</u>	<u>Gross and Microscopic Observations</u>	<u>Serum</u> <u>Grossly</u> <u>Lipemic</u>	<u>Sacrificed</u> <u>Days p</u> <u>Triton</u>	<u>Days p</u> <u>Freezing</u>
43, 8.9 ♀	8.5 1	2	Renal cortical scars.	yes	9	8
44, 8.25 ♂	8.25 1	2	Aorta - periadventitial hemorrhage, granulation and fat, medial necrosis, intimal proliferation, subintimal cellular accumulation of foam cells and monocytes. Normal heart, lung, kidney.	yes	5	4
45, 8.4 ♂	8.4 1	2	Aorta - subendothelial foam cells with little stainable lipid; medial calcification, hemorrhage and necrosis with slightly more lipid. (accessory spleen). Ventricular septal myocardial infarction; pulmonary arteriolar intimal proliferation.	yes	3 mos.	3 mos.
46, 10.4 ♂	11.8 1	-	Animal died with broken back, autolyzed. Fatty liver; gross lipemia.	yes	1	-
47, 7.9 ♂	9.0 1	-	Congested liver without lipid; pulmonary arteriolar intimal proliferation; pulmonary hemorrhage. One coronary arteriole with cells and adherent to subintimal membrane.	yes	3 mos.	3 mos.



Number Weight (lbs.) Sex	Amount (cc). of Triton, <u>Day Given</u>	Day of Aortic <u>Freezing</u>	<u>Gross and Microscopic Observations</u>	Serum		Sacrificed	
				Grossly <u>Lipemic</u>	yes	Days <u>Triton</u>	Days <u>Freezing</u>
48, 9.4 ♂	10.75 1	--	Animal missing - never found.	yes	--	--	--
49, 8.5 ♀	9.6 1	4	Aorta - extensively calcified media; some subintimal lipid and cell accumulation. Lipoid pneumonia with mononuclear cells, giant cells, hemorrhage and foam cells; pulmonary arteriolar intimal proliferation. No renal tubular lipid; renal cortical scars. Inferior vena caval phlebitis.	yes	24	20	20
50, 9.3 ♂	11.6 1	4	Aorta - subintimal proliferation; lipid deposit-ion in media; necrotic media. heart - no stainable lipid. lung - no pulmonary arteriolar lipid; some consolidation. liver - virtually no fat	yes	32	28	28
51, 8.5 ♀	9.6 1	4	Aorta - slight subintimal cellularity without lipid. heart - normal; normal amount of peritoneal fat. lungs - normal; slight subintimal cell infiltration without lipid in pulmonary arterioles.	yes	3 mos.	3 mos.	3 mos.
52 --	--	--	Died of cardiac tamponade after withdrawal of first blood sample.	---	--	--	--





<u>Number Weight (lbs.) Sex</u>	<u>Amount (cc.) of Triton, Day Given</u>	<u>Day of Aortic Freezing</u>	<u>Gross and Microscopic Observations</u>	<u>Serum Grossly Lipemic</u>	<u>Sacrificed Days <math>\bar{p}</math> Days <math>\bar{p}</math> Triton Freezing</u>
53, 8.5 ♀	9.5 1	2	Aorta - proliferated intima; slight inner medial lipid; medial necrosis. liver - slight stainable lipid. kidneys - no tubular lipid; cortical scars.	yes	13 12
54, 8.5 ♀	9.5 1	2	Aorta - without proliferated intima or medial necrosis. heart - normal. Extensive I.V.C. phlebitis; huge abdominal mass.	yes	— —
55, 7.9 ♀	9.0 1	2-sham	Venal caval phlebitis; renal cortical scars - no tubular lipid; no hepatic lipid; no aortic proliferation or lipid-periaortic arteriolar medial lipid.	yes	7 6
56, 8.4 ♀	9.5 1	2-sham	Aorta - thickened and folded inter-nal elastic lamina; very few foam cells. liver - no fat. kidneys - arterial intimal swelling; patchy parenchymal round cell infiltrates.	yes	7 6
57, 7.5 —	10.0 1	—	Fatty liver. (22 hour lipemic serum)	yes	1 —
58, 7.5 —	7.5 1	—	Fatty liver. (used for electrophoresis.)	yes	1 —
59, 6.1 ♀	(7.5 saline control)	3	Aorta - severely calcified media; subintimal proliferation without lipid. Normal heart, liver, kidney.	no	(13) 11



Number Weight (lbs.) Sex	Amount (cc.) of Triton, Day Given	Day of Aortic Freezing	Gross and Microscopic Observations	Serum		Sacrificed	
				Grossly Lipemic	no	Days p̄ Triton	Days p̄ Freezing
59 cont'd			lungs - congestion; subintimal pulmonary arteriolar swelling without lipid. Resuscitated after cardiac arrest. (2 accessory spleens)				
60, 9.25 ♀	(10 cc. 1 saline control)	3	Aorta - proliferation with some lipid deposition and foam cells; medial calcification.	no		(13)	11
61, 7.6 ♂	Corneas injected with 0.1 cc. supernat from 22 hr. pooled serum from WA-57, 58.	-	Sacrificed 8 days after corneal injection. Almost no corneal reaction except few isolated foam cells.				
62, 7.6 ♀	Corneas injected with 0.1 cc. subnate from 22 hr. pooled serum from WA-57, 58.	-	Sacrificed 8 days after corneal injection. Almost no corneal reaction except few isolated foam cells.				
63, ~8.0 ♂	10.0	-	Fatty liver.	yes		4	
64, ~8.0 ♂	8.0 subcutaneous	-	Fatty liver.	yes		4	



Number Weight (lbs.) Sex	Amount (cc.) of Triton, Day Given	Day of Aortic Freezing	Gross and Microscopic Observations	Serum		Sacrificed	
				Grossly Lipemic	Days p Triton	Days p Freezing	Days p Freezing
65, ~8.0 ♂	Corneas in- jected with 0.1 cc. super- nate from 4 day pooled sera from WA- 63, 64.	-	Sacrificed 7 days after corneal injection; fine droplets of lipid halfway through outer cornea. Corneas grossly milky.	---	---	---	---
66, ~8.0 ♂	Corneas in- jected with 0.1 cc. subnate from 4 day pooled sera from WA-63, 64.	-	Sacrificed 7 days after corneal injection; fine droplets of lipid halfway through outer cornea. Corneas only slightly cloudy. Moderately well circumscribed region of foam cellular reaction.	---	---	---	---
67	10.0 1	-	Lost in animal room.	yes	---	---	---
68, 8.75 ♀	10.0 1	-	Diminished peritoneal and perinephric fat. lungs - left upper and lower lob pneumonial, pul- monary arteriolar focal cellular infiltrate.	yes	26	---	---
69, 7.75 ♀	10.0 1	-	Abdominal wall abscess. lungs - left upper and lower lobe pneumonial, pul- monary arteriolar focal cellular infiltrate.	yes	26	---	---
70, 8.0 ♀	10.0 1	-	Abundant peritoneal fat. lungs - bilateral lower lobe pneumonia; pulmonary arterioles have foam cells and proliferated intima. (L)kidney - white 2 mm. cortical sear-superior pole.	yes	26	---	---





<u>Number</u> <u>Weight (lbs.)</u> <u>Sex</u>	<u>Amount (cc.)</u> <u>of Triton,</u> <u>Day Given</u>	<u>Day of</u> <u>Aortic</u> <u>Freezing</u>	<u>Gross and Microscopic Observations</u>	<u>Serum</u> <u>Grossly</u> <u>Lipemic</u>	<u>Sacrificed</u> <u>Days p</u> <u>Triton</u>	<u>Days p</u> <u>Freezing</u>
71, 9.1 ♀	10.0 1	-	Abundant peritoneal fat. heart and kidneys normal. lungs - bilateral lower lobe pneumonia; pulmonary arterioles have foam cells and proliferated intima.	yes	26	-
72, 6.0 ♀	10.0 1	-	lungs - bilateral lower lobe pneumonia; pulmonary arterioles have foam cells and proliferated intima.  Abundant peritoneal fat; renal cortical scarring.	yes	26	-



BIBLIOGRAPHY

1. Anitschkow, N. Uber experimentell erzeugte Ablagerungen von anisotropen Lipoidsubstanzen in der Milz und in Knochenmark. Beitr. Pathol. Anat., 57:201, 1914.
2. Anitschkow, N. and Chalатов, S. Uber experimentelle Cholesterinsteatose und ihre bedeutung fur Entstehung einiger pathologischer Prozesse. Centralbl. E. Allg. Path. U. Path. Anat., 24:1, 1913.
3. Aortic intimal lipid. Nut. Rev., 26:20, 1968.
4. Baumgartner, H.R. and Studer, A. Gezielte Überdehnung der Aorta abdominalis am normo - und hypercholesterinaemischen Kaninchen. Pathol. et Microbiol. (Basel) 26:129, 1963.
5. Bezman-Tarcher, A. Personal communication to Sheila Otway and D.S. Robinson mentioned in Otway, S., et al. The use of ... J. Physiol. (London) 190:321, 1967.
6. Bishop, Louis Faygeres (1864-1941), Arteriosclerosis: A Consideration of the Prolongation of Life and Efficiency after Forty, London, Henry Frowde, 1915, 359 pp.
7. Boggs, T.R. and Morris, R.S. Experimental lipemia in rabbits. J. Exp. Med., 11:553, 1909.
8. Brown, D.F. Evaluation and management of the patient with elevated serum lipid levels. Presented at A.M.A. Convention, June, 1967.
9. Brody, G.L., et al. Hyperlipemia and fat embolism. Am. J. Med. Sci., 247:682, 1964.
10. Brown, R.K., et al. The enzymatic transformation of lipoproteins. J. Biol. Chem., 204:423, 1953.
11. Buchwald, H., et al. Severe atherosclerotic cardiovascular disease in a 14-year-old homozygous familial hypercholesterolemic. Minn. Med. 51:477, 1968.
12. Burnstein, J., et al. Pituitary peptides with direct action on the metabolism of carbohydrates and fatty acids. Biochim. Biophys. Acta, 156:31, 1968.
13. Byers, S.O., et al. Triton hypercholesteremia: Cause or consequence of augmented cholesterol synthesis. Am. J. Physiol., 204:1100, 1963.
14. Caplan, R.M. and Block, W.D. Experimental production of hyperlipemia in rabbits by cobaltous chloride. J. Invest. Derm., 40:199, 1963.
15. Cholesterol esters and lipoproteins. Lancet, 1:902, 1968.



16. Clancy, John James. Experimental Atherosclerosis in Rabbits (and to Others). M.D. Thesis, Department of Pathology, Yale University School of Medicine, New Haven, Connecticut. 1936.
17. Clark, E. and Kaplan, B.I. Endocardial, arterial and other mesenchymal alterations associated with serum disease in man. Arch. Path., 24:458, 1937.
18. Clarkson, Thomas R. Atherosclerosis - spontaneous and induced. Adv. Lipid Res., 1:211, 1963.
19. Collins, W.S., et al. Prevention and control of experimental atherosclerosis in the rabbit. A biological factor. Arch. Surg., (Chicago) 95:871, 1967.
20. Condon, R.E. Tobias, H. and Datta, D.V. Importance of liver in release and degradation of lipoprotein lipase. Surg. Forum, 15:92, 1964.
21. Connor, W.E. Dietary sterols: Their relationship to atherosclerosis. J. Am. Diet. Ass., 52:202, 1968.
22. Constantinides, P. Experimental Atherosclerosis. New York, Elsevier, 1965, 91 pp.
23. Cornforth, J.W., et al. Antituberculosis effect of certain surface active polyoxyethylene esters in mice. Nature, 168:150, 1951.
24. Courtice, F.C. and Garlick, D.G. The permeability of the capillary wall to the different plasma lipoproteins of the hypercholesterolaemic rabbit in relation to their size. Quart. J. Exp. Physiol., 47:221, 1962.
25. Courtice, F.C. and Schmidt-Diedrichs, A. Lipid deposition in the injured wall of the carotid artery in the hypercholesterolaemic and hyperlipaemic rabbit. Quart. J. Exp. Physiol., 47:228, 1962.
26. Cowdry, E.V., Ed. Arteriosclerosis: A Survey of the Problem. New York, MacMillan, 1933. 617 pp.
27. Dately, K.K., Pinto, I.J. and Bharucha, P.E., Eds., Proceedings of the International Seminar on Arteriosclerosis, Bombay, 3,4,5 February 1962. Bombay, Asia Publishing House, 1964.
28. DeFaria, J.L. Medionekrose der grossen und mittelgrossen Arterien nach orthostischen Kollaps medionecrosis Aortae des Menschen und der spontanen Aortensklerose des Kaninchens. Beitr. Path. Anat., 115:373, 1955.
29. DeSuto-Nagy, G.I. and Waters, L.L. The effect of altered lipid metabolism on experimental lesions of the coronary arteries. Circulation, 4:468, 1951.





30. Dock, W. Atherosclerosis: The facts and the mysteries. Bull. N.Y. Acad. Med., 43:792, 1967.
31. Dole, V.P., James, A.T., Webb, J.P.W., Rizack, M.A. and Sturman, M.F. The fatty acid patterns of plasma lipids during alimentary lipaemia. J. Clin. Invest., 38:1544, 1959.
32. Duff, G., Hamilton, J.D. and Magner, D. Experimental production of arteriolonecrosis and medionecrosis of arteries by means of tyramine injections. Proc. Soc. Exp. Biol., 41:295, 1939.
33. Duff, G.L. and McMillan, G.C. Pathology of atherosclerosis. Am. J. Med., 11:92, 1951.
34. Endothelium and atherosclerosis. Lancet, 2:1239, 1967.
35. Fangman, R.J. and Hellwig, C.A. Histology of coronary arteries in newborn infants, abstracted. Am. J. Path., 23:901, 1947.
36. Fleisher, M.S., Jones, L. Serum sickness in rabbits. J. Exp. Med., 34:597, 1931.
37. Frantz, I.D., Jr., et al. Acceleration of hepatic cholesterol synthesis by Triton WR 1339. J. Exp. Med., 101:225, 1955.
38. Fredrickson, D.S. and Lees, R.S. A system for phenotyping hyperlipoproteinemia. Circulation, 31:321, 1965.
39. Fredrickson, D.S., Levy, R.I. and Lees, R.S. Fat transport in lipoproteins - an integrated approach to mechanisms and disorders. New Eng. J. Med., 276:32, 1967.
40. Friedland-unpublished observation cited by N. Nitschkow in E.V. Condry, Arteriosclerosis. A Survey of the Problem, MacMillan, New York, 1933, Chap. 10.
41. Friedman, M. and Byers, S.O. The mechanism responsible for the hypercholesterolemia induced by Triton WR 1339. J. Exp. Med., 97:117, 1953.
42. Friedman, M. and Byers, S.O. Mechanism underlying hypercholesterolemia induced by Triton WR 1339. Am. J. Physiol., 190:439, 1957.
43. Fujikura, T., et al. Unilateral thickening of fetal arteries on the placenta resembling arteriosclerosis. Am. J. Obstet. Gynec., 100:843, 1968.
44. Ghidoni, J.J., et al. Recent advances in molecular pathology: A review ultrastructure of human atheroma. Exp. Molec. Path., 7:378, 1967.





45. Gofman, J.W., et al. The role of lipids and lipoproteins in atherosclerosis. Science, 111:166, 1950.
- 45a. Gofman, J.W., Rubin, L., McGinley, J.P. and Jones, H.B. Hyperlipoproteinemia. Am. J. Med., 17:514, 1954.
46. Goldblatt, H. Studies on experimental hypertension. VIII. The production of the malignant phase of hypertension. J. Exp. Med., 67:809, 1938.
47. Goodman, D.S., et al. Turnover of plasma cholesterol in man. J. Clin. Invest., 47:321, 1968.
48. Grauer, Leonard E. Observations on the Pathogenesis of Atherosclerosis Using the Serum-Lipid Injected Rabbit Cornea as an Experimental Model. M.D. Thesis, Department of Pathology, Yale University School of Medicine, New Haven, Connecticut. 1968.
49. Gutstein, W.H., Lazzarini-Robertson, A. and LaTaillade, J.N. The role of local arterial irritability in the development of arterio-atherosclerosis. Am. J. Path., 42:61, 1963.
50. Haust, M.D. and More, R.H. "Spontaneous lesions of the aorta in the rabbit" in Comparative Atherosclerosis, James K. Roberts, Jr., Reuben Straus, Eds. Hoeber, New York, 1965.
51. Hawk, P.B., Oser, B.L. and Summersun, W.H. Practical Physiological Chemistry, 12th Ed., Philadelphia, The Blakiston Co., 1947. 541 pp.
52. Heimberg, M. Lipids and lipoproteins of human serum. J. Tenn. Med. Ass., 61:167, 1968.
53. Hellman, L., Hirsch, R.L., et al. (Unpublished observation) cited in R.L. Hirsch and A. Kellner, J. Exp. Med., 104:1, 1956.
54. Hill, M.C. Various forms of experimental arterial disease in the rabbit. Arch. Int. Med., 5:22, 1910.
55. Hirsch, R.L. and Kellner, A. The pathogenesis of hyperlipemia induced by means of surface-active agents. I. Increased total body cholesterol in mice given Triton WR 1339 parentally. J. Exp. Med., 104:1, 1956.
56. Hirsch, R.L. and Kellner, A. The pathogenesis of hyperlipemia induced by means of surface-active agents. II. Failure of exchange of cholesterol between the plasma and the liver in rabbits given Triton WR 1339. J. Exp. Med., 104:15, 1956.
57. Hirsch, R.L., Rudman, D. and Travers, R. Movement of lipids into and out of the blood during hyperlipidemia induced in rabbits by pituitary extract and fraction H. J. Lipid Res., 7:182, 1966.



58. Hoff, H.F., et al. An electron microscopic study of the rabbit aortic intima after occlusion by brief exposure to a single ligature. Brit. J. Exp. Path., 49:68, 1968.
59. Hollender, W. Recent advances in experimental and molecular pathology-influx, synthesis, and transport of arterial lipoproteins in atherosclerosis. Exp. Mol. Path., 7:248, 1967.
60. Hueper, W.C. Arteriosclerosis. Arch. Path., 30:162, 1944.
61. Hueper, W.C. Experimental studies on the therapy and the prevention of degenerative vascular diseases. II. The effects of several detergents on experimental cholesterol atheromatosis of rabbits. Arch. Path., 38:381, 1944.
62. Ignatowski, A. Zur Frage über den Einfluss der animalischen Nahrung auf den Kaninchenorganismus, Ber. der Kaiserlichen Militär-Mediz. Akad. Zu St. Petersburg, 16:174, 1908.
63. Imai, H., Lee, K.T., Pastori, S., Panlilio, E., Flörentin, R. and Thomas, W.A. Atherosclerosis in rabbits - architectural and subcellular alterations of smooth muscle cells of aortas in response to hyperlipemia. Exp. Mol. Path., 5:273, 1966.
64. Indian Council of Medical Research, Seminar on Atherosclerosis and Ischaemic Heart Disease, New Delhi, 1962.
65. Israel, O. Experimentelle Untersuchung über den Zusammenhang zwischen Nieren-Krankheiten und secundären Veränderungen des Circulationssystems. Virchow. Arch. Path. Anat., 86:299, 1881.
66. Josué, O. Atherome aortique experimental par injections repetees d'adrenaline dans les viens. Compt. Rend. Soc. DeBiol., 55:1374, 1903.
67. Katz, Louis N., and Stamler, Jeremiah. Experimental Atherosclerosis. Springfield, Illinois, Charles C. Thomas. 1953. 376 pp.
68. Kellner, A., Correll, J.W., Ladd, A.T. Sustained hyperlipemia induced in rabbits by means of intravenously injected surface-active agents. J. Exp. Med., 93:373, 1951.
69. Kellner, A., et al. The influence of intravenously administered surface-active agents on the development of experimental atherosclerosis in rabbits. J. Exp. Med., 93:385, 1951.
70. Kellner, A., Hirsch, R.L. and Freeman, E.B. Inhibition of lipoprotein lipase activity following injection of pituitary extracts into rabbits. J. Exp. Med., 12:1, 1960.





71. Kelly, F.B., Jr., Taylor, C.B. and Hass, G.M. Experimental atheroarteriosclerosis - localization of lipids in experimental arterial lesions of rabbits with hypercholesteremia. Arch. Path., 53:419, 1952.
72. Kelly, John. The Effects of Certain Enzymes on Purified Human Serum Beta Lipoproteins. M.D. Thesis, Department of Pathology, Yale University School of Medicine, New Haven, Connecticut. 1969.
73. Kesten, H.D. Early incidence of spontaneous medial degeneration ("arteriosclerosis") in the aorta of the rabbit. A. M. A. Arch. Pathol., 20:1, 1935.
74. Kline, I.K. Heart lesions due to pheochromocytomas. Am. J. Path., 38:539, 1961.
75. Klotz, O. and Manning, M.F. Fatty streaks in the intima of arteries. J. Path. and Bact., 16:211, 1912.
76. Kobernick, S.D. and Hashimoto, Y. Histochemistry of atherosclerosis. II. Spontaneous degenerative lesions of aorta of exercised and sedentary rabbits. Lab. Invest., 12:685, 1963.
77. Koshy, P. "Results in the use of hypocholesterolemic agents in atherosclerosis", in Proceedings of the International Seminar on Arteriosclerosis, Bombay, Asia Publishing House. 1964.
78. Kritchevsky, D. and Tepper, S.A. Cholesterol vehicle in experimental atherosclerosis - X. Influence of specific saturated fatty acids. Exp. Mol. Path., 6:394, 1967.
79. Kummerow, F.A., Ed., Metabolism of Lipids as Related to Atherosclerosis. A Symposium, Springfield, Illinois, Charles C. Thomas, 1965. 300 pp.
80. Lawry, E.Y., Mann, G.U., Peterson, A., Wysocki, A.P., O'Connell, R., and Stare, F.J. Cholesterol and beta-lipoproteins in the serum of Americans - well persons and those with coronary heart disease. Am. J. Med., 22:605, 1957.
81. Levy, R.I., Lees, R.S. and Fredrickson, D.S. A functional role for plasma alpha<sub>1</sub> lipoprotein. J. Clin. Invest., 44:1068, 1965.
82. Levy, R.I., Lees, R.S. and Fredrickson, D.S. The nature of pre-beta (very low density) lipoproteins. J. Clin. Invest., 45:63, 1966.
83. Levy, R.I. and Fredrickson, D.S. Diagnoses and management of hyperlipoproteinemia. Am. J. Cardiol., 22:576, 1968.
84. Lloyd, J.K. Primary disorders of lipoprotein metabolism in childhood. Postgrad. Med. J., 43:691, 1967.





85. Lucien, M. and Parisot, J. L'athérome spontané chez le lapin, sa fréquence et ses caractères généraux. Comp. Rend. Soc. de Biol., 64: 917, 1908.
86. Malkoff, G. Bedeutung der traumatischen Verletzung von arterien. Bietr. Z. Path. Anat. U. Z. Allg. Path., 25:431, 1899.
87. Man, E.B. and Gildea, E.F. A modification of the Stoddard and Drury titrimetric method for the determination of fatty acids in blood serum. J. Biol. Chem., 99:43, 1932.
88. Mandel, P., Poirel, G. and Simard-Duquesne, N. Oxygen uptake of normal and atherosclerotic rabbit aortae in mediums of normal and hyperlipaemic sera and plasmas. J. Atherosclerosis Res., 6:463, 1966.
89. Marchand, F. Ueber Athero-sklerose. Verhandl. D. Konger F. Inn. Med., 21:23, 1904
90. Matsuyama, K., et al. Experimental studies on the role of plasma proteins and lipids in the evolution of atherosclerosis. I. Tissue reaction to lipids, exp. to cholesterol and its esters. Gunma. J. Med. Sci., 15:131, 1966.
91. Matthews, R.J. Type III and IV familial hyperlipoproteinemia. Evidence that these two syndromes are different phenotypic expressions of the same mutant gene(s). Am. J. Med., 44:188, 1968.
92. Miles, A.B. and Johnstone, O.P. Spontaneous arterial degeneration in rabbits. J. Am. Med. Ass., 49:1173, 1907.
93. Minowski, W.L. The coronary arteries of infants. Am. J. Med. Sci., 214:623, 1947.
94. Moses, C. Atherosclerosis: Mechanisms as a Guide to Prevention. Philadelphia, Lea and Febiger. 1963.
95. Nazum, F.R., Elliot, A.H., Evans, R.D. and Priest, B.V. The occurrence and nature of spontaneous arteriosclerosis and nephritis in the rabbit. Arch. Path., 10:697, 1930.
96. Norum, K.R., et al. Familial serum cholesterol esterification failure. A new inborn error of metabolism. Biochem. Biophys. Acta., 114:698, 1967.
97. Ophüls, W. Spontaneous arteriosclerosis of the aorta (athaeroma) in a rabbit. J. Am. Med. Ass., 48:326, 1907.
98. Otway, S., et al. The use of a non-ionic detergent (Triton WR 1339) to determine rates of triglyceride entry into the circulation of the rat under different physiological conditions. J. Physiol. (Lon.) 190:321, 1967.

The first part of the document discusses the importance of maintaining accurate records of all transactions. It emphasizes that every entry should be supported by a valid receipt or invoice. This ensures transparency and allows for easy verification of the data.

In the second section, the author outlines the various methods used to collect and analyze data. This includes both primary and secondary research techniques. The primary research involves direct observation and interviews, while secondary research involves analyzing existing data sources.

The third part of the document focuses on the statistical analysis of the collected data. It describes the use of various statistical tests to determine the significance of the findings. The author also discusses the limitations of these tests and the potential for bias in the results.

Finally, the document concludes with a summary of the key findings and recommendations. It suggests that further research is needed to explore the long-term effects of the intervention. The author also provides a list of references for further reading on the topic.

99. Packman, M.A., Rowsell, H.C., Jorgensen, L. and Mustard, J.F. Localized protein accumulation in the wall of the aorta. Exp. Mol. Path., 7:214, 1967.
100. Page, I.H. (Ed.), Fredrickson, D.S., et al. Chemistry of Lipids as Related to Atherosclerosis. Springfield, Illinois. C.C. Thomas, 1958. 215 pp.
101. Pawar, S.S., et al. Effect of Triton ingestion on fat retention, blood lipids and growth in rats. Proc. Soc. Exp. Biol. & Med., 122:665, 1966.
102. Pearce, R.M. Occurrence of spontaneous arterial degeneration in the rabbit. J. Am. Med. Ass., 51:1056, 1908.
103. Pethica, B.A., et al. The physical chemistry of haemolysis by surface-active agents. Biochem. J. (London) 53:177, 1953.
104. Petroff, J.R. Über die Vitalfärbung der Gefässwand. Beitr. Z. Path. & Anat. U. Z. Allg. Path., 71:115, 1922.
105. Pick, R. The present state of knowledge about the prevention and therapy of atherosclerosis. Med. Clin. N. Amer., 51:97, 1967.
106. Prior, J.T., Hartmann, W.H. The effect of hypercholesterolemia upon intimal repair of the aorta of the rabbit following experimental trauma. Am. J. Path., 32:417, 1956.
107. Prior, J.T., Hutter, R.V.P. Intimal repair of the aorta of the rabbit following experimental trauma. Am. J. Path., 31:107, 1955.
108. Prior, J.T., Kurtz, D.M., Ziegler, D.D. The hypercholesteremic rabbit: An aid to understanding arteriosclerosis in man? Arch. Pathol., 71:672, 1961.
109. Quadri, G. Altérations athéromateuses de l'aorte des lapins en l'absence de toute injection de substances toxiques. J. des Practiciens, 21:519, 1907.
110. Radding, C.M. and Steinberg, D. Studies on the synthesis and secretion of serum lipoproteins by rat liver slices. J. Clin. Invest., 39:1560, 1960.
111. Rich, A.R. The role of hypersensitivity in periarteritis nodosa, (as indicated by seven cases developing during serum sickness and sulfonamide therapy). Bull. J. Hopkins Hosp., 71:123, 1942.
112. Rich, A.R. Additional evidence of the role of hypersensitivity in the etiology of periarteritis nodosa. Another case associated with a sulfonamide reaction. Bull. J. Hopkins Hosp., 71:375, 1942.





113. Rich, A.R. and Gregory, J.E. The experimental demonstration that periarteritis nodosa is a manifestation of hypersensitivity. Bull. J. Hopkins Hosp., 72:65, 1943.
114. Rich, A.R., Cochran, T.H. and McGoon, D.C. Marked lipemia resulting from the administration of cortisone. Bull. J. Hopkins Hosp., 88:101, 1951.
115. Roberts, J.C., Jr. and Straus, R. Comparative Atherosclerosis: The Morphology of Spontaneous and Induced Atherosclerotic Lesions in Animals and its Relation to Human Disease. New York, Harper and Row, Hoeber Medical Division, 1965. 538 pp.
116. Robertson, W.B., Geer, J.C., Strong, J.P. and McGill, H.C., Jr. The fate of the fatty streak. Exp. Mol. Path., Suppl. 1:28, 1963.
117. Robinson, D.S. The clearing factor lipase and its action in the transport of fatty acids between the blood and the tissues. Adv. Lipid Res., 1:133, 1963.
118. Rosenman, R.H., et al. Plasma lipid interrelationships in experimental nephrosis. J. Clin. Invest., 36:1558, 1957.
119. Rosenthal, S.R. Studies in Atherosclerosis; Chemical, Experimental and Morphologic. Chicago, (reprinted from - Arch. Path.) 134, 88 pp.
120. Rudman, D. and Seidman, F. Lipemia in the rabbit following injection of pituitary extract. Proc. Soc. Exp. Biol. & Med., 99:146, 1958.
- 120a. Saltykow, S. Die experimentelle Erzeugten Arterien-Veränderungen in ihrer Beziehung zu Atherosklerose und verwandten Krankheiten des Menschen. Zent. F. Allg. Path. U. Path. Anat., 19:321, 1908.
121. Scanu, A. and Oriente, P. Triton hyperlipemia in dogs. I. In vitro effects of the detergent on serum lipoproteins and chylomicrons. J. Exp. Med., 113:735, 1961.
122. Scanu, A., et al. Triton hyperlipemia in dogs. II. Atherosclerosis, diffuse lipidosis, and depletion of fat stores produced by prolonged administration of the non-ionic surface-active agent. J. Exp. Med., 114:279, 1961.
123. Scanu, A., et al. Plasma transport of lipids and lipoproteins in dogs treated with Triton WR-1339. J. Clin. Invest., 41:495, 1962.
124. Scanu, A.M. Factors affecting lipoprotein metabolism. Adv. Lipid Res., 3:63, 1965.
125. Schlichter, J.G. Experimental medionecrosis of the aorta. Arch. Path., 42:182, 1946.





126. Schmidtman, M. Vigantolversuche. Verhandl. Der Deutsch. Pathologischen Ellschaft, 24:75, 1929.
127. Schoenheimer, R. and Sperry, W.M. A micromethod for the determination of free and combined cholesterol. J. Biol. Chem., 106:745, 1934.
128. Schotz, M.C., et al. Effect of Triton on lipoprotein lipase of rat plasma. Am. J. Physiol., 188:399, 1957.
129. Schwartz, A.M., et al. Surface Active Agents and Detergents. New York, Interscience Publishers, Inc., 1958, 2 Vols.
130. Schwartz, C.J. The nature of the ground substance changes in experimental lathyrism and their effect on atherogenesis in cholesterol fed rabbits. Brit. J. Exp. Path., 40:44, 1959.
131. Seifter, J. and Baeder, D.H. Lipemia clearing by hyaluronidase, hyaluronate, and desoxycorticosterone, and its inhibition by cortisone, stress and nephrosis. Proc. Soc. Exp. Biol. & Med., 86:709, 1954.
132. Seifter, J. and Baeder, D.H. Lipid mobilizer (LM) from posterior pituitary hogs. Proc. Soc. Exp. Biol. & Med., 95:318, 1957.
133. Sheehan, J.F. Foam cell plaques in the intima of irradiated small arteries (one hundred to five hundred microns in external diameter). Arch. Path., 37:297, 1944.
134. Simpson, C.F. and Harms, R.H. Pathology of aortic atherosclerosis and dissecting aneurysms induced by diethylstilbestrol. Exp. & Mol. Path., 5:183, 1966.
135. Sisson, J.A., et al. Plasma lipids in maternal and fetal rabbits fed stock and peanut oil-cholesterol diets. J. Nutr., 92:435, 1967.
136. Ssolowjew, A. Experimentelle Untersuchungen über die bedeutung von lokaler Schädigung für die Lipoidablagerung in die Arterienwand. Z. Ges. Exper. Med., 69:94, 1929.
137. Ssolowjew, A. Über experimentell hervorgerufene elastic Arisse der arterien und deren bedeutung für die lipoidablagerung. Virch. Arch. F. Path. Anat., 283:213, 1932.
138. Ster, I. and Shapiro, B. A rapid and simple method for the determination of esterified fatty acids and for total fatty acids in blood. J. Clin. Path., 6:158, 1953.
139. Still, W.J.S. and Dennison, S.M. Reaction of the arterial intima of the rabbit to trauma and hyperlipemia. Exp. & Mol. Path., 6:245, 1967.
140. Szakács, J.E. and Cannon, A. 1-norepinephrine myocarditis. Am. J. Clin. Path., 30:425, 1958.



141. Szakács, J.E. and Mehlman, B. Pathologic changes induced by 1-nor-epinephrine. Am. J. Cardiol., 5:619, 1960.
142. Taylor, C.B., Baldwin, D. and Haas, G.M. Localized arteriosclerotic lesions induced in the aorta of the juvenile rabbit by freezing. Arch. Path., 49:623, 1950.
143. Tompkins, E.H. Reaction of the reticuloendothelial cells to subcutaneous injections of cholesterol. Arch. Path., 42:299, 1946.
144. Triton and fat balance. Nutr. Rev., 24:338, 1966.
145. Trygstad, O., et al. Inhibition by human serum of the adipokinetic effect of a human pituitary lipid-mobilizing factor (LMF) in rabbits. Acta Endocr. (Kobenhavn)., 56:649, 1967.
146. Trygstad, O. The lipid-mobilizing effect of some pituitary gland preparations. II. Preparation of a human pituitary lipid-mobilizing factor (LMF) with hypocalcemic and hyperglycaemic effects in rabbits. Acta Endocr. (Kobenhavn), 57:81, 1968.
147. Van der Bosch, J. Evrard, E., Billiau, A., Josseus, J.V. and Desomer, P. The role of liver and spleen in the metabolism of intravenously injected fat in rabbits. J. Exp. Med., 114:1035, 1961.
148. Vidone, R.A., Lowman, R.M., Hukill, P.B., et al. Experimental arteriosclerosis in dogs: The effects of Triton-induced hyperlipemia. Angiology, 18:204, 1967.
149. Waterman, N. Einige Bemerkungen zur Frage: Arteriosklerose nach Adrenalen-Injektionen. Virch. Arch. Path. Anat., 191:202, 1908.
150. Waters, L.L. Changes in the coronary arteries of the dog following injections of allylamine. Am. Heart J., 35:212, 1948.
151. Waters, L.L. and DeSuto-Nagy, G.I. Circulatory factors in the pathogenesis of experimental arteriolar necrosis. Yale J. Biol. & Med., 22:751, 1950.
152. Waters, L.L. The reaction of the artery wall to injury by chemicals or infection. In: National Academy of Sciences-National Research Council, Publication 338, Symposium on Atherosclerosis, Washington, D.C., 1954, pp. 91-98.
153. Waters, L.L. Studies on the pathogenesis of vascular disease. The effect of intravenously injected human plasma and of lipid-rich plasma globulins on inflammatory lesions of the coronary arteries of dogs. Yale J. of Biol. & Med., 30:57, 1957.
154. Waters, L.L. Behavior and fate of injected plasma lipids in corneal connective tissues, abstracted. Circ., 24:1107, 1961.





155. Waters, L.L. Corneal stromal reactions in rabbits following injections of hyperlipemic and of delipidized homologous serum. Am. J. Path., 47:51, 1965.
156. Waters, L.L. Experimental atherosclerosis. Yale J. Biol. & Med., 38:389, 1966.
157. Waters, L.L. Removal of serum lipoproteins by cornea in vivo. Circ., 34:I. 33, 1966.
158. Wherat, A.F. Recent advances in experimental and molecular pathology - Atherosclerosis and metabolic disorder in the arterial wall. Exp. Mol. Path., 7:233, 1967.
159. Williams, A.W. Relation of atheroma to local trauma. J. Path. Bact., 81:419, 1961.
160. Winternitz, M.C., Thomas, R.M. and LeCompte, P.M. The Biology of Arteriosclerosis. Springfield, Illinois, Charles C. Thomas, 1938, 142 pp.
161. Zarafonetis, C.J.D., et al. Lipid mobilizer in hypercholesterolemic states and in surgical stress. Abstr. J. Lab. & Clin. Med., 50:965, 1957.
162. Zarafonetis, C.J.D., et al. Occurrence of clearing factor inhibitor in pregnancy and newborn. Clin. Res., 6:625, 1958.
163. Zarafonetis, C.J.D., et al. Lipid mobilization as a consequence of surgical stress. Am. J. Med. Sci., 237:418, 1959.
164. Zarafonetis, C.J.D., et al. Lipid mobilizer hormone in cobalt chloride hyperlipemia. J. Am. Med. Ass., 191:235, 1965.
165. Zarafonetis, C.J.D., et al. Lipid mobilizer hormone in Triton hyperlipemia. Proc. Soc. Exp. Biol. & Med., 125:321, 1967.
166. Zarafonetis, C.J.D., et al. Plasma protein changes consequent to hyperlipemia induced by cobaltous chloride or Triton WR-1339. Am. J. Med. Sci., 254:506, 1967.
167. Zimmerman, M., et al. Effect of Triton WR-1339 on the metabolism of KB cells in culture. Fed. Proc., 26:513, 1967.



















YALE MEDICAL LIBRARY

Manuscript Theses

Unpublished theses submitted for the Master's and Doctor's degrees and deposited in the Yale Medical Library are to be used only with due regard to the rights of the authors. Bibliographical references may be noted, but passages must not be copied without permission of the authors, and without proper credit being given in subsequent written or published work.

This thesis by \_\_\_\_\_ has been used by the following persons, whose signatures attest their acceptance of the above restrictions.

---

---

NAME AND ADDRESS

DATE

