

1981

Histamine receptors in the airways of healthy human subjects

Steven M. Brown
Yale University

Follow this and additional works at: <http://elischolar.library.yale.edu/ymtdl>

Recommended Citation

Brown, Steven M., "Histamine receptors in the airways of healthy human subjects" (1981). *Yale Medicine Thesis Digital Library*. 2426.
<http://elischolar.library.yale.edu/ymtdl/2426>

This Open Access Thesis is brought to you for free and open access by the School of Medicine at EliScholar – A Digital Platform for Scholarly Publishing at Yale. It has been accepted for inclusion in Yale Medicine Thesis Digital Library by an authorized administrator of EliScholar – A Digital Platform for Scholarly Publishing at Yale. For more information, please contact elischolar@yale.edu.

YALE MEDICAL LIBRARY



3 9002 08627 9289

HISTAMINE RECEPTORS IN THE
AIRWAYS OF HEALTHY HUMAN SUBJECTS

STEVEN MARK BROWN

1981

YALE



MEDICAL LIBRARY

Permission for photocopying or microfilming of " _____

Histamine Receptors in the Airways of Healthy Human Subjects

(Title of thesis)


for the purpose of individual scholarly consultation or reference is hereby granted by the author. This permission is not to be interpreted as affecting publication of this work or otherwise placing it in the public domain, and the author reserves all rights of ownership guaranteed under common law protection of unpublished manuscripts.

Steven M. Jones

Signature of Author

3/8/81

Date



Digitized by the Internet Archive
in 2017 with funding from
The National Endowment for the Humanities and the Arcadia Fund

<https://archive.org/details/histaminerecepto00brow>

HISTAMINE RECEPTORS IN THE
AIRWAYS OF HEALTHY HUMAN SUBJECTS

By Steven M. Brown

A Thesis Submitted to the
Yale University School of Medicine
in Partial Fulfillment of the
Requirement of the Degree of
Doctor of Medicine

March 2, 1981

Med. Lib.

T113

Y12

4044

TO MY PARENTS

TABLE OF CONTENTS

INTRODUCTION.....	Page 1
HISTAMINE.....	Page 2
<u>Background and History</u>	Page 2
<u>Histamine and Its Receptors</u>	Page 3
THE ACTIONS AND ROLE OF HISTAMINE AND ANTI-HISTAMINES IN PHYSIOLOGIC PROCESSES.....	Page 5
<u>Respiratory System</u>	Page 6
<u>Circulatory System</u>	Page 6
<u>Endocrine System</u>	Page 8
<u>Tissue Growth and Repair</u>	Page 8
<u>Gastrointestinal System</u>	Page 9
<u>Nervous System</u>	Page 9
<u>Urinary System</u>	Page 10
<u>H₁ Antihistamines</u>	Page 10
<u>H₂ Antihistamines</u>	Page 11
THE PRESENCE AND ROLE OF H ₁ AND H ₂ RECEPTORS IN THE RESPIRATORY SYSTEM.....	Page 12
<u>Histamine Receptors in Guinea Pig Airways</u>	Page 12
<u>Histamine Receptors in the Respiratory System of the Rat and Ferret: Possible Presence of a Subclass of Receptors</u>	Page 17
<u>More Evidence for Histamine₂ Receptors: The Airways of the Rabbit</u>	Page 18
<u>Histamine Receptors in the Horse: A Possible Animal Model of Human Asthma</u>	Page 19
<u>Histamine Receptors in Sheep Airways</u>	Page 20

<u>Histamine Receptors in Cat Airways</u>	Page 21
<u>Histamine Receptors in the Canine Respiratory System</u>	Page 22
<u>Histamine Receptors in the Airways of Rhesus Monkeys</u>	Page 24
<u>Histamine Receptors in the Airways of Man</u>	Page 25
<u>Pulmonary Vascular Histamine Receptors</u>	Page 31
THE ROLE OF HISTAMINE IN PATHOLOGIC PROCESSES..	Page 34
<u>Histamine, Inflammation, and the Allergic Response</u>	Page 34
<u>Mechanisms of Histamine Interaction</u>	Page 35
<u>Step #1: Histamine Release</u>	Page 35
<u>H₁ Receptors Promote the Inflammatory Response</u>	Page 36
<u>H₂ Receptors Inhibit the Inflammatory Response: Potentiation of Anaphylaxis By H₂ Receptor Blockade</u>	Page 38
<u>Effect on Histamine Release</u>	Page 38
<u>Effect on Histamine Production, Metabolism, Uptake, Storage, and Clearance</u>	Page 40
<u>Effect on Other Participants in the Immune Response</u>	Page 41
<u>Histamine, Antihistamines, and the Immune Reaction in the Clinical Setting</u>	Page 42
Antihistamines as Preventors of the Inflammatory Response.....	Page 42
Antihistamines as Potentiators of the Inflammatory Response.....	Page 44
<u>The Role of Histamine in Asthma</u>	Page 46
Introduction.....	Page 46

The Role of Histamine in Asthma.....	Page 48
HISTAMINE RECEPTORS IN THE AIRWAYS OF HEALTHY, HUMAN SUBJECTS.....	Page 52
<u>Introduction</u>	Page 52
<u>Materials and Methods</u>	Page 53
Subject selection.....	Page 53
Histamine Preparation.....	Page 53
Drugs Used.....	Page 54
Aerosol Challenge.....	Page 55
Pulmonary Function Testing.....	Page 55
Protocol.....	Page 58
<u>Day 1</u> --Histamine Dose Response.....	Page 58
<u>Days 2, 3, and 4</u> --Premedication With Antihistamines.....	Page 59
<u>Analysis of Data</u>	Page 60
<u>Results</u>	Page 61
<u>Discussion</u>	Page 62
<u>Figures</u>	Page 74
<u>Tables</u>	Page 78
BIBLIOGRAPHY.....	Page 87
ACKNOWLEDGEMENTS.....	Page 105

INTRODUCTION

Several chemical mediators play an important role in the normal functioning of the respiratory system in man, in addition to the expression of the immune response. These mediators include histamine, prostaglandins, kallikrein, eosinophil chemotactic factor of anaphylaxis (ECF-A), slow-reacting factor of anaphylaxis (SRS-A), and platelet-activating factor (PAF). All may be released by normal respiratory tissue after passive sensitization with IgE antibody and challenge with specific antigen (8).

Mediators may be released from cells such as mast cells and basophils in response to a number of immunologic or non-specific stimuli. Airway-constricting stimuli include histamine, cholinergic agonists, SRS-A, bradykinin, prostaglandin $F_{2\alpha}$, β -adrenergic blocking drugs, cold air, dust, sulfur dioxide, exercise, and suggestion (15). In addition to their role in normal physiologic processes, mediators play an important role in the pathogenesis of disease states such as asthma and the anaphylactic response.

The following pages will focus on histamine, a

well-investigated mediator. The actions and role of histamine in both the normal and pathologic setting will be addressed. There will be particular focus on the presence of histamine in the respiratory system. The final chapter includes an original investigation of the presence of histamine receptors in the airways of healthy, human subjects.

HISTAMINE

Background and History

Histamine, β -imidazolyethylamine, was first synthesized in the laboratory in 1907. In vivo, it is formed by the decarboxylation of L-histidine, by histidine decarboxylase. Histamine was first isolated in bacteria in 1910 and was one of the first vaso-active substances to be identified (42, 43). The name is derived from the Greek word histos, meaning tissue.

Histamine is stored in mast cells in various tissues as well as in basophils in blood. The mast cell, which also contains heparin, is the major repository for histamine. Histamine is also present in as yet unidentified non-mast cells of the gastric and intestinal mucosa and the brain. In the rabbit (12), the bulk of histamine in the blood is stored in the platelets.

Histamine is released in response to various immunological stimuli as in IgE-antigen-mediated immune reactions (14). It is also released in response to nonimmunological stimuli such as trauma, toxins, or suggestion (131). The liberation of histamine by cells in response to injurious stimuli was noted as early as 1927 by Lewis (52). Mast cell liberators, such as dextran and anaphylatoxin, may also release histamine.


During the release of histamine from mast cells, the mast cell degranulates. The histamine stored in the granule, along with other preformed and stored mediators, is expelled from the cell and is followed by release of histamine from a heparin-protein complex (12). A highly polar molecule at physiologic pH, histamine does not diffuse across the blood-brain barrier or cell membranes.

The action of histamine is rapid. It quickly diffuses into tissues and is rapidly metabolized. Gut flora in man converts ingested histamine to the inactive N-acetylhistamine (52). Once released, histamine will disappear from the blood stream within minutes.

Histamine and Its Receptors

Histamine exerts its effects through interaction with at least two identifiable receptors, which are referred to as H₁ and H₂ receptors. These receptors

may be preferentially blocked by specific antihistamines. The H₁ class of antihistamines are referred to as the "classical antihistamines". The first antihistamines to be developed, Bovet and Staub discovered the first H₁ antihistamines in 1937. The structure is of the form:



where X may be C, O, or N, Ar₁ and Ar₂ are aromatic or aryl ring structures and N must be charged at physiologic pH. Antihistamines act by competitive and reversible occupation of histamine receptor sites without themselves initiating a response.

Ash and Schild (6) were the first to demonstrate the presence of more than one class of histamine receptor. They demonstrated that several effects of histamine, such as stimulation of gastric secretion and inhibition of rat uterus contractions could not be suppressed by classical antihistamines.

Black, et al. (18) were the first to develop burimamide, an H₂ receptor antagonist. It was noted that while burimamide did not have any significant H₁ receptor antagonism, it successfully prevented gastric acid secretion.

Further evidence for the existence of two separate

classes of histamine receptors has come from the synthesis of agonists specific to H₁ and H₂ receptors. For example, while 2-methylhistamine can elicit the H₁ effect of stimulating bronchial contraction, it has no significant effect on gastric acid secretion. On the other hand, 4-methylhistamine can increase gastric acid secretion, but has little effect on airway constriction (52). In one study (18), 2-methylhistamine was shown to have 16.5% of the H₁ activity of histamine as measured by ability to constrict guinea pig terminal ileum. Its H₂-mediated ability to stimulate isolated atrium was only 4.4%. While 4-methylhistamine had 43% of the ability of histamine to stimulate the guinea pig isolated atria, the relative activity on terminal ileum was only 0.23%.

Other H₁ agonists include 2-(2-aminoethyl)-pyridine and 2-(2-aminoethyl)-thiazole (125). Other H₂ agonists are clonidine (7), St 600 ([2-(5-fluoro-o-toluidine)-2-imidazoline hydrochloride])(27), and dimaprit ([S-[3-(N,N-dimethylamino)propyl]isothiourea])(121).

THE ACTIONS AND ROLE OF HISTAMINE AND ANTIHISTAMINES IN PHYSIOLOGIC PROCESSES

Histamine was recognized a half-century ago (11) as being involved in the inflammatory response and allergic phenomena. The "classic" histamine response

in man includes itching, swelling, redness (histamine flush), headache, and bronchoconstriction. Histamine will elicit the cutaneous "triple response" of a red spot and flush, wheal, and itch--symptoms which are associated with the inflammatory response and anaphylaxis. The following are brief outlines of some other actions of histamine, with emphasis, where possible, of the pharmacologic effects of histamine and anti-histamines in man. The role of histamine in the immune response and pathologic processes will be dealt with in greater detail in a subsequent chapter.

Respiratory System

In man, large concentrations of histamine have the primary effect of bronchoconstriction. Histamine causes an increase in respiratory air resistance and a decrease in airway compliance (8). Other animal species may exhibit either bronchoconstriction, bronchodilation, or both in response to histamine. The response will be determined by which part of the respiratory system is stimulated, which specific receptors are agonized, and what are the effects of those specific receptors. This will be examined in a forthcoming section.

Circulatory System

As in the respiratory system, vasodilation or vasoconstriction will be the effect of histamine agonism,

and will vary depending upon the size and location of the blood vessel, the classification of histamine receptor which is being agonized, as well as the species involved. In man, histamine will cause constriction of large arteries and veins and dilation of the smallest vessels to contain smooth muscle. Intravenous histamine in man, cats, dogs, and sheep will cause a fall in systemic blood pressure secondary to peripheral vasodilation (1, 19).

In cats, dogs, and chickens, the blood vessels dilate in response to both H_1 and H_2 agonism (19, 34, 145). In rabbit blood vessels, H_1 agonism causes vasoconstriction and H_2 agonism causes vasodilation (34). The reverse is the case for calves (34, 66).

In man, peripheral vasodilation is mediated by both H_1 and H_2 receptors (14, 99). Combinations of H_1 and H_2 antagonists are more successful in preventing the effects of histamine on skin (101) and the cardiovascular system (13), than use of any one antagonist alone.

Vasodilation in the cerebral blood vessels in man accounts for histamine-induced headache.

Histamine's evanescent effect on small blood vessel dilation leads to enhancement of the body's microcirculation, which is due, in part, to the con-

tinuous alternating vasodilation and vasoconstriction caused by changing concentrations of histamine. Histamine will increase capillary permeability with the resultant effect of formation of edema, increased blood viscosity, and hemoconcentration.

The effect on the heart is to stimulate cardiac contractility. Histamine will also increase the rate of cardiac contraction, and will decrease AV-node conduction. In vitro studies of the guinea pig heart (92) show that H₁ receptors may be involved in mediation of a negative dromotropic effect, an atrial inotropic effect, and promotion of histamine-induced arrhythmias of conduction. H₂ receptors mediate positive chronotropic effects, ventricular inotropic effects, and promote histamine-induced arrhythmias of automaticity.

Histamine will constrict smooth muscle of the spleen.

Endocrine system

The adrenal medulla will release catecholamines in response to histamine.

Tissue Growth and Repair

Histamine is found in increased concentration in areas of wound repair. Increased levels of histamine may be found in regenerating liver, granulation tissue, and the fetus. Tissue damage through local irritants

will increase histidine decarboxylase activity at the site on injury, leading to increased histamine levels (12). Histamine promotes wound healing as well as the formation of keloids and scar tissue. It also helps to provide an increased blood supply to areas of injury.

Gastrointestinal system

Acting through stimulation of H₂ receptors, histamine stimulates gastric acid secretion by the parietal cells of the stomach. H₂ receptor blockade, in addition to blocking gastric acid secretion, also blocks the effects of gastrin and pentagastrin. This finding suggests that histamine may act as a final "common messenger" for several hormones in the gastric mucosa (34). Histamine may also stimulate, via H₂ receptors, pancreatic secretion and cholecystokinesis (34). It will also contract smooth muscle of the gut via H₁ receptor stimulation.

Nervous system

The function of histamine in the brain remains obscure, although its presence has been demonstrated in a variety of structures. The binding of radioactive H₂ antagonists has been demonstrated in guinea pig midbrain, cortex, hippocampus, thalamus, brainstem, hypothalamus, cerebellum, and corpus striatum (23). Histamine has been isolated in rat, mouse, and monkey

hypothalamus as well as midbrain, cortex, and cerebellum (13). Histamine agonism with the H₂ receptor increases levels of cyclic AMP in these structures (14).

Histamine stimulates sensory nerve endings, producing pruritis. Histamine may facilitate ganglionic transmission via H₁ agonism. H₂ receptors may inhibit ganglionic transmission.

Urinary system

Histamine has a dual mode of action in the human kidney. It causes a pressor response on renal vasculature via H₁ agonism and a depressor response via H₂ agonism.

H₁ Antihistamines

H₁ antihistamines, which include chlorpheniramine, diphenhydramine, and mepyramine, are of therapeutic use in seasonal rhinitis, allergic dermatoses, itching pruritides, hay fever, insect bites, ivy poisoning, blood transfusion reactions, serum sickness, allergic reactions to drugs, Ménière's disease, and vertigo (13, 52). They inhibit the histamine-induced increase in capillary permeability during anaphylaxis. H₁ antihistamines do not completely block anaphylaxis in man because of the significant contribution of several other mediators.

H₁ antihistamines may also cause somnolence,

prevent motion sickness, provide a mild anticholinergic effect, and have a mild anesthetic effect. Because of their mild anticholinergic, antiadrenergic, and anti-serotonergic activities, the use of H₁ antihistamines in laboratory investigations has occasionally resulted in erroneously ascribing cholinergic, adrenergic, or serotonergic effects to histamine.

H₁ antihistamines are metabolized by the liver and degradation products are almost completely excreted within 24 hours. Tissues are almost totally free of H₁ antagonists within 6 hours.

H₂ Antihistamines

H₂ antihistamines have gained recent wide clinical use in the treatment of a variety of gastrointestinal disorders associated with hypersecretion of gastric acid. Of particular significance is the recent popularity of cimetidine in the treatment of duodenal ulcers.

Another class of histamine antagonist is the mast cell stabilizer, such as cromolyn sodium. Cromolyn, which can also inhibit antigen-induced production of histamine (52), has as its main effect, the prevention of release of histamine and SRS-A. Cromolyn sodium does not interact directly with either H₁ or H₂ receptors.

THE PRESENCE AND ROLE OF H₁ AND H₂ RECEPTORS IN THE RESPIRATORY SYSTEM

The laboratory investigation which will be described in a forthcoming chapter will examine the presence of histamine H₁ and H₂ receptors in the airways of healthy human subjects. The following is a detailed description of studies which have investigated the presence of histamine receptors in the respiratory systems of a variety of species, both in vivo and in vitro.

Histamine Receptors in Guinea Pig Airways

Guinea pig trachea may be caused to contract due to antigen-induced anaphylaxis (Schultz-Dale phenomenon). It was noted almost fifty years ago (11) that these contractions are associated with the release of histamine from guinea pig lung. The causative effect of histamine in producing the contractions has been demonstrated by the partial or complete block of these contractions by H₁ antihistamines (38, 85). The fact that mepyramine only partially prevents antigen-induced contractions indicates the presence of other mediators which cause tracheal contractility.

Direct stimulation of guinea pig tracheobronchial tissue by histamine results in a profound net smooth muscle contraction (52). Mepyramine totally blocks histamine-induced contraction (85). Metiamide, which

antagonizes H₂ receptors, potentiates histamine-induced contraction. However, stimulation of H₁ receptors by the specific H₁ agonist, 2-methylhistamine, causes constriction which is not potentiated by metiamide (117). It can thus be inferred that histamine causes both H₁ stimulated contraction and H₂ stimulated relaxation of guinea pig tracheobronchial muscle. Each effect can be isolated by use of specific antagonists.

The H₁ constricting effects and H₂ relaxing effects on guinea pig tracheobronchial muscle can be demonstrated indirectly through studies with antihistamines. However, more direct evidence is available through observation of the effects of histamine agonists. As previously mentioned, the effect of 2-methylhistamine is to cause tracheobronchial contraction.

Another H₁ agonist is 2-(2-pyridyl)-ethylamine (2-PEA). 2-PEA will cause contraction of guinea pig tracheal spirals, in vitro. This constriction of tracheal spirals and parenchymal strips is reversed by dimaprit, an H₂ agonist, which mediates relaxation of the smooth muscle. The effect of dimaprit may be blocked by metiamide (54).

In guinea pig bronchial smooth muscle undergoing ovalbumin antigen-induced anaphylactic contractions, dimaprit had a significant bronchodilatory effect (32).

This effect was even more profound than treatment with an H₁ antagonist, mepyramine.

Direct evidence for histamine H₁ receptor presence in guinea pig lung and other organ systems has come from studies which trace histamine receptors by radioactively labelling tissues with [³H]-mepyramine (41).

Drazan and Schneider (53) found that histamine induced constriction in both guinea pig trachea and lung parenchyma. They discovered that histamine H₁ antagonism by mepyramine was more successful in blocking constriction in lung parenchyma than in trachea. This suggests that histamine receptors are not necessarily distributed evenly in the airways. There appears to be a greater number of receptors and/or sensitivity to the effects of histamine in peripheral airways in the guinea pig. A future study involving radioactively-labelled antagonists in different parts of the respiratory system would be useful in supporting these findings.

In addition to the effect of the direct interaction of histamine with its receptors, histamine receptor agonism may also cause release of prostaglandins. These prostaglandins act as additional mediators of airway reactivity, but do not act as the final mediators of histamine agonism. The mechanism for release of prostaglandins in response to histamine stimulation involves H₁ receptor-cyclic GMP stimulated release of PGF_{2α} and H₂ receptor-cyclic AMP stimulated release of PGE.

Yen, et al. (156) used the presence of prostaglandins

PGF_{2α} and PGE as an assay for determining the distribution of histamine H₁ and H₂ receptors within the lung. They found that the H₁ antagonist, pyrilamine, blocked the histamine-induced increases in PGF_{2α}. Metiamide diminished the histamine-induced release of PGE.

The authors found that in response to histamine in guinea pig trachea, PGE was present in much greater concentrations than PGF_{2α}. In guinea pig peripheral airways, PGF_{2α} was present in higher concentrations than PGE (156). The authors conclude that there might be more H₂ receptors in central than peripheral airways. They find this conclusion consistent with the physiologic importance of maintaining the patency of central airways. The rapid shifts, on the other hand, of peripheral airway size through bronchoconstriction, has been suggested to maximize ventilation-perfusion ratios (156).

In addition to prostaglandins, levels of cyclic nucleotides are a useful assay for determining the presence of histamine receptors. In guinea pig lung anaphylaxis, the antigen-antibody reaction yields an increase in levels of cyclic AMP, cyclic GMP, and the ratio of cAMP to cGMP. Pyrilamine inhibits histamine stimulated increases in cyclic GMP (103). Burimamide had no effect on cGMP but blocked the histamine-induced increases in cAMP.

While there may exist different distributions of H₁ and H₂ receptors within guinea pig airways, H₁ and

H₂ receptors may also differ in sensitivity to histamine. Yen (155) showed that in guinea pig peripheral airway smooth muscles, concentrations of 10⁻¹⁰ - 10⁻⁷ M of histamine caused bronchorelaxation which was blocked by metiamide. H₁ receptors had the overwhelming response in concentrations of greater than 10⁻⁶ M of histamine with the resultant contraction blocked by chlorpheniramine. It appears that H₂ receptors are more sensitive to histamine than H₁ receptors, but in high concentrations, H₁ receptors win out, probably because of greater numbers.

The opposite sensitivities to histamine were observed by Martin and Fertel (102) in guinea pig tracheal rings. They noted that at concentrations of 5 x 10⁻⁵ M, guinea pig tracheal rings contracted. However, at concentrations of 10⁻⁴ M of histamine, the tissue relaxed. Consistent with these observations was the increase in cyclic GMP levels at low histamine concentrations. Increased cyclic AMP levels were noted at high concentrations of histamine. Thus, in trachea, there may be greater sensitivity for H₁ receptors, but greater numbers of H₂ receptors.

In addition to the presence of histamine receptors in guinea pig airway smooth muscle, the presence of histamine H₁ receptors in guinea pig alveolar macrophages

has been demonstrated by both agonist and antagonist studies (50).

Histamine Receptors in the Respiratory System of the Rat and Ferret: Possible Presence of a Subclass of Receptors

Chand and Eyre (34, 40) have done extensive studies on the presence and action of histamine receptors in a number of species. They determined that ferret trachea and bronchus both constricted in response to histamine in vitro. This contraction of smooth muscle was blocked by H₁ antagonism with mepyramine. Rat trachea did not contract in the presence of histamine (40).

Both rat and ferret trachea and bronchi could be made to contract in vitro by exposure to carbachol. When subsequently exposed to histamine in the presence of H₁ blockade, the muscles relaxed. This indicates that an H₂ receptor must have been mediating the tracheo-bronchorelaxation, since the H₁ receptors were blocked. Further evidence comes from studies of direct H₂-agonism by 4-methylhistamine which also caused relaxation of smooth muscle (40).

It is very interesting to note that the relaxation was not blocked by metiamide, burimamide, or cimetidine. This important finding suggests the presence of a subclass of histamine receptors which are not blocked by conventional H₂ antagonists, yet are stimulated by

4-methylhistamine and produce relaxation of airway smooth muscle. The proposed name for this subclass of receptors is H_{2B} or H_3 receptors.

More Evidence for Histamine₃ Receptors:
The Airways of the Rabbit

Using a similar methodology for in vitro studies as Chand and Eyre (40), Fleisch and Calkins (70) ascertained that histamine induces contraction of rabbit bronchus. This effect was blocked by H_1 antagonism with pyrilamine. Interestingly, under similar experimental conditions, partially contracted rabbit trachea experienced relaxation in the presence of histamine. This effect could not be prevented by pyrilamine, burimamide, metiamide, indomethacin, or propranolol. Thus, histamine-induced tracheal relaxation does not act through H_1 , H_2 , prostaglandin release, or β -adrenergic mechanisms (65, 70).

Chand and Eyre (36) also found that partially-contracted rabbit bronchus relaxed in the presence of H_1 blockade. This response was not eliminated by metiamide. It is thus proposed that rabbit trachea (70) and rabbit bronchus (36) may contain H_3 receptors which mediate smooth muscle relaxation.

It is important to note that, as in the guinea pig, the airways of the rabbit experienced different effects

of histamine stimulation, depending upon whether central or peripheral airways were being agonized. Thus, two closely associated tissues in the same system can be pharmacologically distinct. It is important, therefore, to state specifically which tissues in an airway are being stimulated, when analyzing the results of a histamine study or when designing an investigation.

Histamine Receptors in the Horse: A Possible Animal Model of Human Asthma

Using their model of carbachol-induced, partially-constricted airways, in vitro, Chand and Eyre (37, 39) determined that histamine or 2-methylhistamine causes tracheobronchial constriction in the horse. Mepyramine prevents this histamine-induced contraction. H₂ agonists such as 4-methylhistamine and dimaprit caused tracheobronchial relaxation in this experimental model. Further support of H₂ stimulation of muscle relaxation comes from treatment of partially-constricted bronchus in the presence of H₁ blockade. This results in relaxation, which could only be the result of H₂ receptor agonism.

The relaxation in horse bronchus was blocked by metiamide and burimamide. However, in horse trachea, metiamide, cimetidine, and burimamide had no inhibitory effect. Perhaps horse trachea also has H₃ receptors.

It has been suggested (39) that the horse may be

a good model for human asthma. Horses suffer from "broken wind", an obstructive respiratory syndrome, also known as equine pulmonary emphysema, which, clinically resembles asthma. Thus, the clinical use of H₂ agonists in asthma may be worth future investigation.

H₁ blockade is only minimally useful in inhibiting anaphylaxis in the sensitized horse (63). As in human anaphylaxis, histamine is only one of several mediators which are involved in the anaphylactic response in the horse. Of importance is the fact that H₂ antagonists such as metiamide and burimamide potentiate anaphylaxis in the horse (39). H₂ antagonists have also been noted to exacerbate anaphylaxis in adult domestic fowl and the calf (39). The mechanism behind this potentiation and its implications in the clinical setting will be dealt with in a forthcoming chapter.

Histamine Receptors in Sheep Airways

Using burimamide, Eyre was the first to discover (14, 61, 62, 64) the presence of H₂ receptors in isolated smooth muscle of sheep bronchi. He found that histamine causes contraction of trachea and the major bronchi in sheep. This effect may be blocked by H₁ antihistamines. In response to histamine, the lesser

bronchi and bronchioles would relax. This effect could be blocked by burimamide (14, 64). Intravenous histamine results in a net bronchoconstriction in sheep. The net effect of inhaled histamine is either bronchodilation or minimal bronchoconstriction in the sheep (61, 65).

Histamine Receptors in Cat Airways

In in vitro cat trachea partially contracted due to carbachol, histamine challenge leads to a net bronchodilation (14, 62, 64, 100). This effect may be blocked by a combination of H₁ and H₂ antagonists in addition to propranolol. This finding suggests, that in the cat trachea, histamine acts by both direct action with both receptors as well as by indirect action through local catecholamine release to produce relaxation.

The cat bronchus also relaxes in response to histamine as well as 2-methylhistamine and 4-methylhistamine (35). This effect is not reversed, however, by either H₁ or H₂ antagonism, β -blockade, or prostaglandin inhibition. thereby suggesting an H₃ mechanism.

Other investigators (96, 123) found that in the cat bronchus, in vitro, histamine challenge caused smooth muscle contraction, acting via H₁ mechanisms which could be blocked by mepyramine. Similarly, antigen-induced contraction could be abolished by mepyramine in cat bronchial tissue.

Histamine Receptors in the Canine Respiratory System

Histamine causes bronchoconstriction in vivo in the dog. When delivered histamine size is greater than 10 microns, bronchoconstriction is mostly through a reflex vagal mechanism. There is a direct effect of histamine on smooth muscle receptors if particle size is less than 0.5 microns (128).

Dixon, et al. (51) examined the effect of histamine, 4-methylhistamine, cimetidine, and chlorpheniramine on total lung resistance and dynamic lung compliance in anaesthetized dogs. While histamine caused an increase in total lung resistance and a decrease in dynamic lung compliance, the H₂ agonist 4-methylhistamine had no such effect. Accordingly, the H₁ antagonist, chlorpheniramine blocked the effect of histamine, while cimetidine had no protective effect. The authors conclude that histamine acts mostly through H₁ mechanisms in the dog and causes airway constriction. Since dynamic lung compliance and total lung resistance are particularly useful in measuring large airways pulmonary function (98), perhaps this conclusion should not be extended to include the peripheral airways of canines. The counterargument is that at 0.5 microns in diameter, the histamine particles were delivered to the peripheral airways (45). Other investigators (5, 33, 83, 132, 153) have found similar

in vitro and in vivo evidence in peripheral and central airways of dogs. The possibility of regional differences in histamine receptor distribution are similarly supported by scientific evidence (33).

Some investigators (153, 154) have found that cimetidine tended to increase airway resistance, suggesting a small H₂-mediated bronchodilatory effect of histamine in the dog lung. On the other hand, Irvin and Dempsey (82) find a minor contribution of H₂-mediated bronchoconstriction in the peripheral airways of dogs. They concur with other authors regarding the absence of H₂ receptors in canine central airways.

As in humans, mepyramine does not completely block antigen-induced anaphylaxis (33), suggesting the role of other mediators such as SRS-A and serotonin in this response. Whereas metiamide has been shown to exacerbate anaphylaxis in the horse, calf, and domestic fowl (39), the H₂ antagonist has no such in vivo effect in immediate hypersensitivity reactions (33, 153) or in vitro models of canine asthma (5, 33). In ovalbumin-sensitized dogs (5), isolated trachealis muscle contracted in response to histamine. This was inhibited by pyrilamine. H₂ blockade with metiamide had no effect.

Of interest is the effect of H₂ stimulation on respiratory secretions, rather than respiratory muscle

activity. Cimetidine blocked stimulation of canine secretory activity in the lung in histamine-induced asthma. Chlorpheniramine also had this effect, but to a smaller extent (153). The investigators conclude that histamine acts to promote canine asthma through an H₁-mediated muscular effect and an H₂-mediated effect on secretory activity of the respiratory tract (153, 154). They propose that H₂ antihistamines might have therapeutic value in decreasing respiratory secretions in the asthmatic.

Histamine Receptors in the Airways of Rhesus Monkeys

One published investigation of histamine receptors in the airways of rhesus monkeys examined the effect of H₁ and H₂ antagonism on in vivo pulmonary function (76). The authors measured peak expiratory flow, total pulmonary resistance, dynamic lung compliance, tidal volume, and respiratory frequency.

The results of the study indicated that pretreatment with the H₁ antagonist diphenhydramine inhibited histamine-induced bronchoconstriction. Metiamide potentiated histamine's bronchoconstrictory effects when given prior to histamine challenge (76). Thus, this study seems to indicate that histamine causes bronchoconstriction in the rhesus monkey via H₁ agonism. H₂ receptors are present and apparently modulate the

bronchoconstriction, perhaps via relaxation of smooth muscle or by some feedback mechanism.

Histamine Receptors in the Airways of Man

There have been a few recent studies which have investigated the role of histamine receptors in the airways of man. These preliminary reports include studies of healthy human subjects as well as human asthmatics, who represent a special class of people who are exquisitely sensitive to minute quantities of histamine.

Histamine alone causes a net bronchoconstriction in man. Normal subjects receiving histamine have been reported (110) to experience an increase in respiratory system resistance and closing volume and a decrease in vital capacity (VC), forced expiratory volume in one second (FEV_1), and the ratio of FEV_1 to VC. These parameters suggest both large and small airways constriction due to histamine.

Chlorpheniramine clearly prevents histamine-induced bronchospasm in children (137) and adults (50, 71, 99).

Maconochie, et al. (99) found that in healthy subjects, 8 mg. or oral or 5 or 10 mg. of intravenous chlorpheniramine generally prevented histamine-induced bronchoconstriction, as measured by FEV_1 .

However, 400 mg. or oral cimetidine or 100 or 200 mg. of intravenous cimetidine blockade of H₂ receptors had no effect on bronchoconstriction in the subjects studied (99). These findings indicate the primary and perhaps sole involvement of H₁ receptors in histamine-induced bronchospasm in healthy, human airways.

The authors offer (99), as an explanation for the lack of effect of cimetidine, the possibility that cimetidine did not reach the H₂ receptors in the lung in sufficient concentrations to produce an effect. However, the doses of cimetidine used were adequate to inhibit gastric acid secretion and histamine challenge did coincide with peak blood levels of cimetidine.

Eiser (59) was similarly not able to demonstrate the presence of H₂ histamine receptors in normal human airways. He found that H₂ blockade with 200 mg. of intravenous cimetidine had no effect on histamine-induced changes in airway specific conductance in the nine subjects tested. 20 mg. of intravenous chlorpheniramine, however, significantly shifted the histamine dose-response curve to the right. Eiser found no difference between chlorpheniramine administration and the administration of chlorpheniramine and cimetidine together (59).

Cimetidine has received wide clinical use since its development. In the clinical setting, it does not

appear that cimetidine promotes asthmatic bronchospasm (14).

However, Frith, et al. (71) are preparing a report on a study of ten asthmatics. The subjects were selected on the basis of histories of episodic dyspnea as well as documented reversible airflow destruction. Subjects received an increasing dose of inhaled histamine until a 20% drop in FEV₁ was reached. On subsequent days, the same protocol was followed, except subjects were pretreated with chlorpheniramine 8 mg. or cimetidine 600 mg. or both.

The findings show that H₁ blockade significantly increased the level of histamine necessary to produce a 20% drop in FEV₁ (p=0.0001)(71). Cimetidine did not differ from placebo. However, cimetidine did significantly (p=0.04) reduce the effectiveness of chlorpheniramine when the two drugs were given together. These results indicate the presence of histamine H₁ and H₂ receptors in asthmatic airways, although the H₂ effect does not seem to be very profound. The authors do not address the influence of drug-drug interactions in their findings. The dose of cimetidine used was at least twice that required to reduce food-stimulated gastric acid secretion by 50%.

Nathan, et al. (109) achieved similar results in eleven asthmatics. Chlorpheniramine significantly

raised (p less than 0.002) the level of histamine required to cause a 20 percent drop in FEV₁, a 35 percent decline in maximum expiratory flow at 50 percent below vital capacity, or a fifty percent decrease in airway conductance. Cimetidine significantly decreased the amount of histamine necessary to produce these results (p less than 0.02). The authors conclude that H₂ receptors, mediating bronchodilation, exist in asthmatic airways.

Pretreatment with aerosolized diphenhydramine in adult asthmatics significantly blocked the effects of inhaled histamine (28) as measured by the amount of histamine required to produce a 20% drop in FEV₁. This result lends further support to the theory that histamine produces bronchoconstriction by direct action on H₁ receptor sites.

Although oral antihistamines are of little clinical use in asthmatics, clemastine, an H₁ antagonist, when delivered directly to the bronchial tree by inhalation (113), caused bronchodilation in asthmatics. Pulmonary function parameters were FEV₁ and peak expiratory flow rate (PEFR). In addition, the histamine-dose response curve was significantly shifted to the right (112). These findings were comparable to the results achieved by salbutamol, a β_2 receptor agonist. The results of

this study may be due to complete blockage of H₁ receptors in asthmatics, thereby allowing endogenous histamine to react specifically with the only unblocked receptors, H₂ receptors, thereby mediating relaxation in addition to the obvious elimination of bronchoconstricting H₁ activity. An interesting follow-up study would investigate whether cimetidine reverses the bronchodilatory effects of inhaled clemastine.

Dunlop and Smith (57) as well as Kaliner and Platshon (89) have provided in vitro evidence for the presence of histamine receptors in human airways. Dunlop and Smith (57) showed, in vitro, in sensitized human bronchus, 2-3 mm. in diameter, that in the presence of H₁ receptor blockade, histamine agonism led to broncho-relaxation. This effect could be eliminated by H₂ antagonism with metiamide. Furthermore, in human bronchus caused to contract due to exposure to house dust mite antigen (57), H₁ blockade with mepyramine caused decreased contraction while H₂ blockade caused increased bronchial contraction.

Kaliner and Platshon (89) demonstrated that 2-methylhistamine produced an increase in cyclic GMP, with subsequent enhancement of mediator release (SRS-A). H₂ agonism with dimaprit in the human lung caused increased levels of cyclic AMP with subsequent decreased

release of SRS-A. This finding indicates that H₂ agonism with substances such as dimaprit might be useful in the treatment of anaphylaxis.

Histamine's action in asthmatic airways to cause bronchoconstriction is almost entirely via direct stimulation of smooth muscle receptors (29). In asthmatics receiving a dose of atropine, a parasympatholytic drug, at a dose strong enough to achieve cholinergic blockade, the histamine dose-response curve was slightly shifted to the right (29, 84, 137). Thus, in asthmatics, histamine does not act primarily through cholinergic pathways.

Similarly, the vagal-blocker SCH 1000 did not prevent histamine-induced bronchospasm in asthmatics (150) although it did prevent methacholine-induced bronchospasm. Histamine does have some reflex vagal nerve irritant receptor action, although this is a minor effect (3, 135). In large airways, it is proposed that histamine has more of a reflex vagal effect (8, 47) while in small airways, constriction is due to direct histamine H₁ antagonism.

As is the case in other species, such as the guinea pig and sheep, the distribution and sensitivity of histamine receptors may vary in different parts of the respiratory system such as in central versus peripheral

airways. This possibility makes in vivo studies in man more difficult than in vitro studies. Methodologies must deliver histamine to specific parts of the airway. Pulmonary function tests must be used to measure constriction or dilation in the specific stimulated sections of airway under investigation. Thus, one would not want to only measure large airways constriction in a protocol where histamine is delivered mostly to the respiratory bronchioles. Since large airways have a reflex vagal contribution to histamine-induced bronchospasm (47) this provides another factor which makes analysis of the presence of histamine receptors in vivo difficult.

Pulmonary Vascular Histamine Receptors

The most profound effect of histamine on airway mechanics is by its direct interaction with smooth muscle receptors. However, interaction of histamine with pulmonary vascular receptors is also very important in the regulation of respiration. Studies with a variety of results have been performed in a number of animal species.

In the guinea pig, histamine agonism leads to pulmonary vasoconstriction. This effect is most profound in the pulmonary veins (34, 115). It is proposed (64) that histamine-induced pulmonary venospasm may increase right atrial pressure and thereby contribute to the formation of pulmonary edema. The pulmonary

vasoconstriction appears to be mediated by H₁ receptors in the guinea pig. Burimamide potentiates this response. Mepyramine has a depressor action on pulmonary vasculature, indicating the presence of H₂ receptors which mediate pulmonary vasodilation. Some authors feel that pulmonary vascular dilation may increase airway resistance, through mechanical obstruction.

The rat exhibits pulmonary vasoconstriction in response to hypoxia. This effect is blocked by metiamide (10) indicating that H₂ stimulation leads to pulmonary vasoconstriction in the rat. This is the opposite effect to be observed in the cat (79, 145) where H₁ agonism causes pulmonary constriction and H₂ agonism mediates dilation of pulmonary vasculature.

Comparable results are available in the dog (74). Okpako (116) demonstrated that H₂ agonism with 4-methylhistamine had 15 times the vasodepressor potency of 2-methylhistamine. This vasodepression could be abolished by burimamide.

Tucker, et al. arrive at similar results in the dog (145, 146). Measuring pulmonary vascular resistance, they found that H₁ agonism led to an increase in resistance, while H₂ agonism caused a fall in pulmonary vascular resistance. Antagonists had the expected results of reversal of the effects of agonists. The

authors also found (146) that H₂ blockade would also potentiate pulmonary vasoconstriction during hypoxia. Okpako (116) concludes that H₂ receptor agonism may lead to "pooling of blood" in the bronchial mucosa with resultant mechanical obstruction of the airway.

The effect of histamine on pulmonary vessels seems to vary with development in dogs (111). One group of authors found that while metiamide potentiated pulmonary vasoconstriction in canine pups, this effect disappeared after 15 days of life.

In the horse (77), histamine causes pulmonary vasoconstriction. This appears to be mostly an H₁ effect, although H₂ agonism may also contribute to vasoconstriction to a lesser degree. In the nonpregnant ewe (152), histamine also mediates pulmonary artery constriction which may be blocked equally well by H₁ or H₂ antagonists (benadryl and metiamide).

THE ROLE OF HISTAMINE IN PATHOLOGIC PROCESSES

Histamine, Inflammation, and the Allergic Response

As early as 1910 (42, 43), Dale and Laidlaw appreciated the similarities between the effects of histamine and anaphylaxis. These include redness, swelling, edema, and hypotension. Twenty years later, Bartosch et al. (11) demonstrated the presence of histamine in guinea pig lung during the antigen-antibody reaction.

Sensitized human lung and skin mast cells and leukocytes will release histamine in vitro when challenged with specific antigens (17). In vivo, sensitized humans and asthmatics experience increased plasma histamine levels following allergen-induced bronchoprovocation (17, 28). The increase in histamine levels correlates well with the onset of bronchospasm.

The release is initiated by IgE antibody bound to the cell surface. IgE-induced effects are mediated by alterations in cyclic nucleotides.

The fact that a second messenger, in this instance, the cyclic nucleotides cAMP and cGMP, may act as the final common mediators for the elaboration of the different physiological responses of different tissues to antigen-antibody interaction, may have been first proposed in 1919 (43):

So that it is quite possible that the production by histamine, and by a whole group of other substances, of a complex including contraction of plain muscle with relaxation and permeability of capillaries, may depend on a common type of physical change in protoplasm produced by all of them, the result of which receives different expression in terms of the physiology of different tissues.

Mechanisms of Histamine Interaction

In cells undergoing the IgE mediated antigen-antibody reaction (Type I immediate hypersensitivity reaction), histamine, acting on H₁ receptors, participates in an exquisitely fine-tuned mechanism to promote the inflammatory process. In addition, histamine also acts on H₂ receptors to provide a negative feedback on its own actions (34).

Step #1: Histamine Release

Following the interaction of cell surface-bound IgE antibody with specific antigen, there is an influx of extracellular calcium, important for the initiation of release of histamine-containing granules. This is followed by the energy-dependent activation of microtubules and microfilaments which results in the fusion of the perigranular membrane with the cell membrane, followed by exocytosis of granules. Extracellular ions such as sodium then penetrate into the granules with the resultant release of the histamine-

heparin-protein complex by ion exchange (114).

Further release of histamine is modulated by intracellular levels of cyclic AMP and cyclic GMP, which, as will be seen, are themselves regulated by histamine. Platshon and Kaliner (124) found that cGMP and cAMP levels increased profoundly, simultaneously with the appearance of mediators in sensitized human bronchus.

H₁ Receptors Promote the Inflammatory Response

Acting on H₁ receptors, histamine's effect on smooth muscle is to facilitate contraction by increasing calcium entry. This leads to depolarization and impulse-carrying currents.

The sudden contraction, which is a result of histamine release, is followed by a brief period of partial relaxation during the anaphylactic response. This is followed by a secondary, sustained contraction, which is resistant to classical antihistamines and which is attributed to the delayed release of non-histamine mediators such as SRS-A (38).

Cyclic GMP may be implicated in this H₁ mediated contraction. Increased levels of cyclic GMP are found following stimulation with histamine (124). This increase may be blocked by H₁ antihistamines (124). The rise in cyclic GMP is calcium-dependent (14). The role of cyclic GMP may be to activate protein kinases

which in turn enhance phosphorylation of proteins which participate in calcium ion mobilization (14) which in turn leads to smooth muscle contraction.

Histamine seems to promote its own release through H_1 -mediated positive feedback (14). H_1 receptor anti-histamines inhibit the IgE induced release of histamine as well as antigen-induced release of histamine from sensitized monkey lung (14).

H_1 receptor agonism enhances eosinophil migration.

Histamine causes release of prostaglandins from the lung. Indomethacin, a prostaglandin synthetase inhibitor, diminishes allergic bronchospasm by 40%, indicating a role of prostaglandins in allergic bronchoconstriction, although the prostaglandin synthesis which accompanies anaphylaxis has been considered secondary to the action of histamine. Suppression of prostaglandin synthesis does not prevent hypersensitivity-associated rises in cAMP and cGMP.

The cyclic GMP dependent release of $PGF_{2\alpha}$ and thromboxane A_2 , which intensify bronchoconstriction, may be inhibited by H_1 receptor antagonism with pyrilamine or mepyramine (64, 125). H_1 agonism with 2-methylhistamine promotes release of $PGF_{2\alpha}$ and thromboxane A_2 (16, 124).

Histamine agonism of H_1 receptors (and to a

lesser extent H₂ receptors) also increases vascular permeability. This facilitates the migration of immunoglobulins and leukocytes to the site of inflammation.

H₂ Receptors Inhibit the Inflammatory Response:
Potentiation of Anaphylaxis By H₂ Receptor Blockade

Histamine acts through H₂ receptors in a different manner than H₁ receptors. Stimulation of histamine H₂ receptors causes a rise in cyclic AMP (124). Blockade of the H₂ receptor prevents a rise in cyclic AMP (2, 124, 134, 138), except in mice, where H₁ and not H₂ receptor blockade prevents histamine-induced cAMP accumulation (118).

Cyclic AMP facilitates calcium sequestration and extracellular transport, with resultant relaxation of contractile proteins. Elevated levels of cyclic AMP correlate with bronchial smooth muscle relaxation and bronchodilation.

Effect on Histamine Release

H₂ receptor agonism and the subsequent rise in cyclic AMP is also very important in the regulation of the immune process. Elevated levels of cyclic AMP inhibit histamine and SRS-A release from the mast cell and human leukocytes (21).

In humans allergic to ragweed, histamine is re-

leased upon exposure to ragweed antigen. In leukocytes of these human donors, in vitro, this release of histamine may be inhibited by histamine, isoproterenol, or methylxanthines. Each of these substances acts to increase levels of cyclic AMP (21). Lichtenstein and Gillespie (94) demonstrated that H₂ antagonists could prevent the inhibition of histamine release from blood basophils. H₁ antagonists did not block this response.

Ovalbumin antigen caused dose-related release of histamine from in vitro guinea pig lung. Dulabh and Vickers (56) found that H₂ antagonism with cimetidine or burimamide potentiated this release. This demonstrates the role of H₂ receptors in modulating the immune response-mediated release of histamine.

In addition, metiamide or cimetidine potentiated the antigen-induced bronchoconstriction in guinea pig lung (56). Other investigators have achieved similar results (2, 90).

It appears that metiamide also potentiates the antigen-induced release of histamine in sensitized rhesus monkey lung (30, 31), canine lung, bovine lung, and monkey skin (30) in vivo. However, Platshon and Kaliner (124) were unable to prove that H₂ blockade potentiated histamine release in the human lung in vitro. H₂ antagonism does not potentiate histamine release from passively sensitized rat lung (30, 34). This

suggests that H₂ agonism may not inhibit endogenous histamine release in the rat (30). Holyroyde and Eyre (80) showed that H₂ blockade actually prevented histamine release in sensitized bovine lung. It is interesting to note that H₂ blockade does not, in some experimental models (30, 33, 88) also enhance SRS-A release from sensitized tissues. This is a curious result, since SRS-A release is inhibited by increased levels of cyclic AMP and H₂ agonism increases cyclic AMP levels.

Effect on Histamine Production, Metabolism, Uptake, Storage, and Clearance

Not only does H₂ agonism prevent release of histamine, it also inhibits production of histamine. H₂ antagonists induce histidine decarboxylase to produce histamine (104, 125). Histamine metabolism is also promoted by H₂ receptor agonism and is inhibited by H₂ antihistamines (67, 107, 125). Taylor (144) was unable to show, however, that either the H₁ antihistamine mepyramine, or the H₂ antihistamine burimamide had any effect on histamine methyltransferase, the enzyme which is involved in the methylation of histamine, the predominant mode of inactivation of the mediator. Blockage of the H₂ receptor also inhibits histamine storage in mast cells (14, 107).

H₂ antagonists prevent histamine uptake (34, 68, 107) and delay clearance of histamine from the circulation (34, 133).

Effect on Other Participants in the Immune Response

Increased levels of cyclic AMP, the result of H₂ agonism, also inhibit T-cell induced cytotoxicity (14, 126). H₂ agonists in high concentrations will inhibit eosinophil migration (14) as well as inhibit lysosomal enzyme release from human polymorphonuclear cells (93, 125). Acting through the H₂ receptor, histamine will inhibit the production of lymphokines.

Just as histamine acts upon H₁ receptors to promote PGF_{2α} release, histamine acts upon H₂ receptors to promote release of PGE, a prostaglandin which has cyclic-AMP-mediated, bronchodilating activity. PGE may in turn inhibit further histamine release by activation of cAMP in mast cells (16). Metiamide can antagonize histamine-induced release of PGE.

In conclusion, histamine released during a hypersensitivity reaction may participate in a number of mechanisms to provide feedback inhibition in order to limit the severity of the reaction.

Histamine, Antihistamines, and the Immune Reaction in the Clinical Setting

Antihistamines as Preventors of the Inflammatory Response

Although histamine is a major mediator of anaphylaxis, antihistamines have not been effective in the complete control of this process. Several reasons for this have been suggested. Other mediators are involved in the anaphylactic response. These include SRS-A, kinins, prostaglandins, and serotonin. SRS-A has been shown to be the mediator responsible for the late, prolonged phase of bronchoconstriction in the antigen-challenged human bronchus (125). Although H₂ receptors seem to be most important in inhibition of the inflammatory response, to the effect that they cause increased vascular permeability, they contribute to the inflammatory process. Classical H₁ antihistamines have no effect on the H₂ receptor contribution to this process.

Antihistamines nevertheless do have important clinical relevance in the modification of the inflammatory reaction. Plaut (125) found that cutaneous reactivity to compound 48/80 could be inhibited by combinations of H₁ antihistamines and cimetidine. He proposes that H₁ and H₂ receptor blockade might be helpful for treating some cutaneous manifestations of allergic diseases such as chronic urticaria (125).

Similarly, Harvey and Schocket (78) found that H₂ antihistamines potentiated the effect of H₁ antihistamines in blocking the cutaneous wheal response, although H₂ antihistamines alone had no effect. It is suggested (78) that H₂ receptors are somewhat involved in the cutaneous histamine response. Goadby and Little (75) found that metiamide protected against anaphylaxis in the guinea pig, a finding which is unusual, considering the purported role of H₂ receptors in modifying the immune response.

Rocklin (129) noted that in sensitized guinea pigs, histamine in concentrations of 10⁻³M reduced the size of a delayed hypersensitivity skin test. This effect could be reversed by an H₂ antagonist. He proposed that histamine suppresses cutaneous delayed hypersensitivity in part, by inhibiting the production and proliferation of migration inhibiting factor (MIF). H₂ antagonists reversed this inhibition of MIF (129). In so doing, H₂ antagonists such as cimetidine have been shown to be of therapeutic value in allowing the expression of cutaneous cell-mediated immunity. For example, cimetidine is useful in the treatment of selected deficiencies in the immunologic response to candida antigen (86).

Antihistamines as Potentiators of the Inflammatory Response

H₂ antihistamines, by antagonizing H₂-mediated inhibition of inflammation, may potentiate anaphylaxis. Dunlop and Smith demonstrated (57) that metiamide potentiates anaphylactic bronchoconstriction in vitro. This was the observation of Drazen, et al. (55) who found that H₂ antihistamines burimamide and metiamide increased the severity of immune-initiated anaphylaxis in the guinea pig. 4-methylhistamine decreased the severity of the reaction (55). Of note is the fact that cimetidine, another class of H₂ antagonist, had no effect.

Wolfe, et al. (151) came to the same conclusion regarding cimetidine. They were unable to demonstrate a prominent modulating role for H₂ receptors in immediate or delayed skin test reactivity. Krell and Chakrin (90) were unable to find that H₂ blockade with metiamide potentiated antigen-induced or histamine-induced changes in dynamic lung compliance or pulmonary resistance in the dog at doses up to 16 times the ED₅₀ for inhibition of gastric acid secretion. They conclude that insofar as the sensitized canine might be a model for human asthma, it appears that H₂ antihistamines may not be deleterious to allergic asthmatics (90).

While in one study, cimetidine potentiated the

anaphylactic response in the guinea pig (56), required doses were 100 micromoles per kilogram, a dose between 50 and 100 times that required to inhibit maximal gastric output by 50% in the rat and dog (14). It is unlikely that the results of Dulabh and Vickers (56) can be extended to conclude that a potential effect of therapeutic doses of H₂ receptor antihistamines would be to cause intensified allergic reactions in man.

However, a recent case report in the New England Journal of Medicine (48), described the appearance of a hypersensitivity reaction which coincided with the oral administration of cimetidine. The reaction disappeared after discontinuance of the H₂ antagonist.

This finding was supported by the work of Avella, et al. (9) who found that in patients receiving cimetidine, there was a significant enhancement of the delayed hypersensitivity response to four common antigens as compared to controls. This is most likely due to the antagonism of H₂ receptor mediated stimulation of suppressor T cell function (9).

Expecting that H₂ blockade would remove inhibition of the delayed hypersensitivity reaction, Dale (44) noted that metiamide did not exacerbate the tuberculin reaction in the guinea pig. She concluded that other factors in addition to histamine must regulate the

immune response.

The Role of Histamine in Asthma

Introduction

Asthma may be defined as a disease process characterized by an increased reactivity of the trachea and bronchi to various stimuli (4). This leads to reversible narrowing of airways. Asthma is marked by increased contraction of bronchial smooth muscle and increased secretion of mucous in the bronchial tree. Asthma is manifested by the symptoms and signs of dyspnea, wheezing, cough, prolonged expiration, and responsiveness to bronchodilator drugs.

Asthmatic airways display hyperreactivity to a number of stimuli. These include specific allergens such as house dust, animal dander, pollen, and drugs (114). Other stimuli include suggestion (81, 97, 131, 140), smoke, exercise (136), respiratory infections (4), and emotional reactions (4).

The mechanism for the allergic hyperreactivity in asthmatics is via the Type I immediate hypersensitivity reaction which is mediated by the IgE antibody which reacts with surface bound antigen. Asthmatic bronchospasm is also mediated via Type III hypersensitivity, a late, gradually appearing immune-complex reaction.

The most important mediators of asthmatic broncho-spasm are histamine and SRS-A. Other mediators such as bradykinin and prostaglandins play a less important role. Bronchoconstriction is mediated by chemical mediators, as previously discussed, via a rise in intracellular levels of cyclic GMP. This in turn causes a rise in intracellular Ca^{++} levels with subsequent activation of contractile proteins.

In addition to chemical mediators, bronchoconstriction is promoted by increased parasympathetic tone. Cholinergic stimulation similarly causes a rise in intracellular calcium ion levels (114). Interaction of bronchoconstricting elements with "irritant receptors" causes reflex bronchoconstriction through vagal stimulation. Vagally-induced bronchoconstriction can not be inhibited by classical H_1 antihistamines, although it can be prevented by treatment with parasympathetic blocking agents such as atropine or ipratropium bromide (114).

Agonism with β_2 receptors on respiratory smooth muscle causes bronchorelaxation. Neurotransmitters which combine with β_2 receptors will relax bronchial smooth muscle. The mechanism of bronchorelaxation is via stimulation of cyclic AMP which causes decreased calcium ion levels. This increase in cyclic AMP

which results from catecholamine stimulation, also decreases the antigen-induced release of histamine in sensitized lung in vitro (106).

Asthmatics may experience disturbances in any of several mechanisms which lead to increased bronchial activity. They may display hypersensitivity to antigens, increased responsiveness to histamine, abnormally active parasympathetic tone, or diminished sympathetic (β_2 receptor) responsiveness (106, 110, 142). The following pages will focus on the role of histamine in asthma.

The Role of Histamine in Asthma

The findings that histamine causes bronchoconstriction and is released in immediate hypersensitivity reactions have led to a strong association of asthmatic bronchospasm with histamine (117). With regard to histamine, asthmatics may experience an increased release of histamine in response to bronchoconstricting stimuli as compared to healthy humans.

For example, in addition to the role of vagal stimulation in exercise-induced bronchospasm, histamine also seems to be involved. Simon, et al. (136) found that among some asthmatics, exercise provoked a fall in pulmonary function (FEV_1). This decline in FEV_1 correlated well with increased levels of histamine in the peripheral venous blood.

Asthmatics may also have an increased sensitivity to the histamine which is released. Indeed, asthmatics develop bronchoconstriction in response to lower concentrations of inhaled histamine than do normal persons (91, 110, 139). This fact is useful in the identification of persons with hyperreactive airways (109). There is some evidence in the mouse model for asthma, that histamine sensitivity may be in part, an inheritable trait (147).

In addition, asthmatics differ from normals in that asthmatics have detectable levels of histamine in their plasma (17) even without antigen challenge. Nonasthmatics do not have any detectable level of endogenous histamine in their plasma.

Asthmatics might suffer from an imbalance in the numbers of different types of histamine receptors which have opposing effects on bronchial smooth muscle. This theory is supported by recent experimental evidence. Busse and Sosman (26) noted that histamine, acting through H_2 receptors, inhibits serum-activated zymosan-induced lysosomal enzyme release from granulocytes. This response parallels a simultaneous increase in levels of cyclic AMP. The authors found (26) that this H_2 -mediated inhibition of lysosomal enzyme release is significantly decreased in asthmatics as well as

normal volunteers infected with rhinovirus 16 (24). This suggests that asthmatics have decreased numbers or sensitivity of H₂ receptors. Furthermore, viruses may exacerbate the asthmatic condition through their effect on H₂ activity.

The decreased responsiveness of H₂ receptors in asthmatics is important. The antigen-antibody reaction might initiate a selective increase in cyclic GMP levels in asthmatics (124). Since the H₂ receptor acts to control further histamine release from sensitized cells as well as acts to stimulate histamine metabolism and uptake, diminished H₂ reactivity would cause increased levels of endogenous histamine. In asthmatics, who are sensitive to even minute quantities of histamine, this increase in histamine levels would lead to further bronchoconstriction. For the same reason, H₂ blockade in asthmatics would be expected to be more deleterious than in non-asthmatics, because the slight increases in histamine would effect asthmatics, whose H₁ receptors are much more sensitive to slight increases in the levels of histamine.

Airway obstruction in asthma is due, in part, to inflammatory changes in the bronchi. Granulocytes may be found in the respiratory tree in increased numbers during bronchial reactivity. Important medi-

ators of this inflammatory response are the lysosomal enzymes. Decreased inhibition by the H₂ receptor of these lysosomal enzymes may contribute to the inflammatory response in asthmatics (26). It is interesting to note that persons with other defects in immune responsiveness, such as those with atopic eczema, may also have diminished H₂ receptor-mediated inhibition of the inflammatory process (25).

Some authors, however, have found no difference between H₂ receptor responsiveness in asthmatics and non-asthmatics. Gillespie et al. (73) found no significant difference between asthmatics and non-asthmatics in the effect of histamine on lymphocyte cAMP levels (an H₂ receptor-mediated process).

The finding that asthmatics have decreased H₂ receptor responsiveness is analagous to the decreased responsiveness of β_2 adrenergic receptors seen in asthmatics as compared to non-asthmatics (106, 110, 142).

HISTAMINE RECEPTORS IN THE AIRWAYS OF HEALTHY, HUMAN SUBJECTS

Introduction

While many of the previously described studies have provided excellent evidence for the presence of histamine H₁ and H₂ receptors in the airways of many species, there is little evidence for the presence of H₂ receptors in human airways. Should H₂ receptors exist in normal or asthmatic airways, and should they provide a role in bronchodilation, as in animal models, this finding might have important therapeutic implications. For example, H₂ receptor agonists might be useful as bronchodilators. Furthermore, caution might need to be followed in using H₂ receptor antagonists in persons with compromised respiratory function.

Evidence to date includes in vitro studies on the presence of H₂ receptors in sensitized human bronchus (57, 89). Two recent studies in non-asthmatics (59, 99) have found no evidence for the presence of H₂ receptors in the airways of non-asthmatics. Two recent studies (71, 109) provide preliminary evidence that H₁ and H₂ receptors are present in asthmatic airways, and H₂ receptors mediate bronchodilation.

The following experiment describes an investigation of the effects of histamine H₁ and H₂ receptor blockade

in the airways of healthy, non-asthmatic, human subjects. The goal of the experiment was to provide evidence for the presence or absence of H₁ and H₂ receptors in the small airways of normal man.

Materials and Methods

Subject Selection

Subjects were recruited as paid volunteers. They gave written, informed consent for participation in the experiment, as approved by the Yale University School of Medicine Human Investigations Committee.

Eleven healthy, non-asthmatic subjects (8 males, 3 females) ranged in age from 18 to 35. The mean age was 26.3 ± 5.1 years. All had no history of asthma or recent respiratory diseases which could simulate an asthmatic response to histamine (24, 60). Baseline pulmonary functions were at least 85% of normal predicted values for the group as a whole. (TABLE I)

Histamine preparation

Histamine dihydrochloride (Sigma, St. Louis) was prepared in normal saline as a stock solution of 128 mg. base ml.. Stock solution was kept frozen when not in use. Serial dilutions of the stock solution provided concentrations of histamine used for the study. Subjects received histamine challenges of 0.1, 2, 4, 8,

16, 32, 64, and 128 mg./ml. ($9 \times 10^{-4}M$ to 1.15M; pH=7.0 to 3.7).

Drugs Used

All drugs were administered single-blindedly.

Cimetidine 300 mg. (Smith, Kline, and French, Philadelphia) is a highly specific H_2 receptor antagonist (22). It was selected because of its recent wide use in the treatment of various illnesses, particularly gastrointestinal disorders (149). The dosage of the drug provided a blood level of cimetidine of twice that required to suppress by 80% secretion of gastric acid by the stomach (149).

Chlorpheniramine maleate 8 mg. (USV, Tuckahoe), an alkylamine derivative, was selected because of its negligible sedative and anticholinergic effects as well as its high specificity for H_1 receptors as compared to other H_1 antihistamines (119).

A lactose placebo dispensed in a pink gelatin capsule (Eli Lilly and Company, Indianapolis) was used.

Medications were administered orally, after an overnight fast, $1\frac{1}{2}$ to 2 hours prior to histamine challenge, a time designed to coincide with peak levels of each drug (127, 130, 149). No side effects of cimetidine, chlorpheniramine, or placebo were reported. Since both cimetidine (143, 149) and chlorpheniramine (122) have very short physiological and pharmacological half-lives, the possibility of

drug-drug interactions was negligible, since the drugs were administered at least 24 hours apart.

Aerosol Challenge

Histamine was delivered via a Dautrebande D-30 nebulizer (45, 46), driven by 20 pounds per square inch of compressed air (0.5 ml. liquid nebulized/min.). Particles of histamine were thus consistently aerosolized to less than 0.5 microns in diameter. This assured deposition of histamine in the respiratory bronchioles and lung alveolar spaces.

Subjects wore a noseclip and inhaled the histamine through a mouthpiece attached to a separate, valved breathing circuit. Histamine was inhaled for 30 seconds by tidal breathing. Tidal breathing controlled against the effects of deep inspiration on pulmonary function in non-asthmatics (108). Adverse symptoms to inhaled histamine at highest challenge doses included headache, cough, flushing of the skin, tachycardia, and chest tightness. Higher doses of histamine were not given to subjects after the onset of adverse symptoms.

Pulmonary Function Testing

Histamine induces large airway constriction through both direct agonism with H_1 receptors, and to a lesser extent, by vagal mechanisms. Peripheral

airway constriction is almost entirely the result of direct agonism with H₁ receptors of smooth muscle (8, 47). Keeping in mind the fact that histamine was being delivered primarily to small airways, the pulmonary function tests selected for this experiment were selected for their sensitivity in measuring small airway constriction. A particularly sensitive measure of small airways constriction is the MEF₄₀(P), the maximum expiratory flow at 60 percent below vital capacity on a partial expiratory curve (20, 105). This test measures flow during the part of expiration which is independent of expiratory effort. The flow at 60 percent of vital capacity is determined by the static recoil pressure of the lung and the flow-resistive properties of small airways (20, 105).

The particular sensitivity of the partial flow-volume curve may reflect the absence of the effects of a deep inspiration required for the generation of maximal expiratory flow volume curves (69, 72, 108).

All subjects were familiarized with the breathing maneuvers and equipment at the beginning of the study. Breathing maneuvers were repeated until reproducible curves could be obtained.

Subjects blew into a cardboard tube attached to a computerized pneumotachograph-integrator system (148)

which calculated flow and volume. The measurements were recorded on a Gould x-y recorder (slew rate of 40 inches per second). The equipment was calibrated daily.

Subjects expired to residual volume, then inspired to approximately 50-70% of their vital capacity. While wearing a noseclip, the subjects then expired into the cardboard tube as fast as possible to residual volume, thereby generating the partial expiratory flow-volume (PEFV) curve.

Next, subjects inspired to vital capacity and then expired as fast as possible to residual volume, thereby generating the maximal expiratory flow-volume (MEFV) curve. A programmable marker, set to trigger at 1 second, permitted identification of the forced expiratory volume at 1 second (FEV_1). The resultant curves allowed measurement of the forced vital capacity (FVC), peak expiratory flow rate (PEFR), and maximum expiratory flow rate at 60% below vital capacity on the MEFV curve (MEF_{40}) and PEFV curve ($MEF_{40}(P)$). (Figure 1) FEV_1 is a particularly sensitive measure of airway obstruction during a relatively effort-independent portion of the curve.

Protocol

Each day of the experiment, there were controls against factors known to influence responses to histamine (87). Each day, pre-challenge flow-volume maneuvers were performed to establish the baseline for the day. The baseline FEV₁ and MEF₄₀(P) for the subjects on each day was expressed as a percent of the baseline on Day 1. (TABLE II) Histamine challenge was performed at the same time of day to control for diurnal variations in histamine response (49, 95).

Day 1--Histamine Dose Response

On Day 1, the baseline pulmonary function was determined. Prior to each histamine challenge, each subject repeated three flow-volume maneuvers to establish a pre-challenge control value. Starting with doses of 0.1 mg./ml. of histamine, subjects received a histamine challenge as previously described. Inhalation of histamine was followed immediately by pulmonary function testing at 0, 1, 2, 4, 6, and 8 minutes. The averages of the first three and second three blows were calculated.

If a subject did not experience a 20% fall in MEF₄₀ (P) as compared to pre-challenge control values, within the first three minutes, then the next successive dose of histamine was administered after a thirty-minute

wait. The wait between successive doses of histamine challenge was designed to control against any cumulative effect of histamine (87). Pre-challenge controls expressed as a percentage of the baseline for Day 1 are recorded in TABLE III.

When a subject experienced a 20% decline in $MEF_{40}(P)$ as compared to pre-challenge control, this dosage was referred to as the "threshold" dose. (Figure 2)

Days 2, 3, and 4--Premedication With Antihistamines

On each of Days 2, 3, and 4, subjects were given, on successive days, an oral dose of H_1 antihistamine (chlorpheniramine), H_2 antihistamine (cimetidine), or placebo (lactose), $1\frac{1}{2}$ to 2 hours prior to histamine challenge.

Baseline pulmonary function was determined at the beginning of each day. Prior to each histamine challenge, each subject established a pre-challenge control value as on Day 1. Pre-challenge controls as a percent of baseline for Days 2, 3, and 4 are recorded in TABLE IV.

Each subject received a dose of histamine at one dose below the threshold dose as determined on Day 1 (T-1). Subjects also received a histamine challenge at threshold dose (T) and one dose above the threshold

(T+1). Each histamine challenge was followed by pulmonary function testing at 0, 1, 2, 4, 6, and 8 minutes after histamine challenge. The averages of the first three and second three blows were calculated for $MEF_{40}(P)$ and FEV_1 . The average of the first three blows was calculated as a percentage of the pre-challenge control.(TABLE V)(Figures 3 and 4)

Analysis of Data

The means and standard deviations for the pulmonary function tests were calculated. Using this information, a t-test for the comparison of two means (58) was used to compare changes in daily baseline or pre-challenge control pulmonary function, as well as changes in pulmonary function following histamine challenge as compared to pre-challenge control values.

Underlying the comparison of the $MEF_{40}(P)$ values, is the assumption that the air flow reflected a defined lung volume. This could be the case only if total lung capacity remained unchanged during histamine provocation. Total lung capacity has indeed been shown (20, 141) to remain constant during induced bronchoconstriction in asthmatics and normal subjects.

Results

Analysis of the baseline FEV₁ and MEF₄₀(P) for each day (TABLE II) shows that pulmonary function did not significantly vary from day to day.

A clear dose response curve (TABLE VI) could be generated for histamine inhalation. The effects of histamine on pulmonary function for the group became significant at 32 mg./ml. for both the MEF₄₀(P) and FEV₁ curves. The MEF₄₀(P) curve was more significant than the FEV₁ curve at doses of 32 mg./ml. and 64 mg./ml.. (Figure 2, TABLE VII)

There was a slight cumulative effect of histamine on pre-challenge control levels at 64 mg./ml. of histamine. (TABLE III)

Chlorpheniramine significantly prevented the decline in MEF₄₀(P), at threshold doses of histamine, as compared to Day 1. Pulmonary function values for placebo and cimetidine were not significantly different at threshold as compared to Day 1. (TABLE VIII)

Pretreatment with chlorpheniramine significantly protected against the fall in MEF₄₀(P) as compared to placebo at doses (T) and (T+1). Significant protection of chlorpheniramine against the fall in FEV₁ as compared to placebo was apparent at dose (T+1). There was no significant difference between placebo and cimetidine

for either $MEF_{40}(P)$ or FEV_1 at any test dose of histamine. (Figures 3 and 4, TABLES V, VIII, and IX)

Discussion

Various conclusions may be drawn from this study. It can be seen from the comparison of baseline pulmonary function from each day that chlorpheniramine, cimetidine, and placebo each had no bronchoconstrictory or bronchodilatory effects on baseline pulmonary function. Popa (127) studied the effects of oral and intravenous chlorpheniramine in asthmatics. He found that chlorpheniramine did improve baseline pulmonary function. However, this study examined the effect of chlorpheniramine on non-asthmatics. The difference in findings may be due to the relative absence of endogenous histamine in non-asthmatics as compared to asthmatics (17) in addition to the decreased sensitivity of non-asthmatics to the little endogenous histamine which may exist (91, 110, 139). Thus, while chlorpheniramine may block H_1 receptors in asthmatics which would otherwise be stimulated by histamine, chlorpheniramine blocks receptors in non-asthmatics, which, in the absence of exogenous histamine provocation, would otherwise not be agonized.

Subjects could not be excluded from the study if baseline pulmonary function varied from day to day, as

this variation might be attributed to the effect of the drug. Exclusion of these persons might therefore skew the results in favor of persons in whom the drugs had no effect.

The finding that baseline pulmonary function did not significantly vary daily added, in retrospect, a useful control to the study. Benson (15) found that the resting state of the airways was an important determinant of bronchial reactivity. Thus, if either chlorpheniramine, cimetidine, or placebo increased the resting bronchomotor tone in the airways, they could have potentiated subsequent bronchoconstriction due to histamine.

It may also be concluded that one dose of 300 mg. of oral cimetidine has no effect on baseline pulmonary function in non-asthmatics. It would be premature to extend this conclusion to the clinical setting of continuous cimetidine use in non-asthmatics. However, this finding suggests the possibility of conducting such a study which would examine pulmonary function in non-asthmatics before, during, and after chronic cimetidine use.

This study confirmed that the methodology used can successfully generate a dose response curve for inhaled histamine provocation. Furthermore, the usefulness of the $MEF_{40}(P)$ pulmonary function test as a

sensitive measure of bronchoconstriction has been demonstrated. The measurement of MEF₄₀(P) is particularly useful in measuring small airways constriction. In addition, the Dautrebande D-30 nebulizer is particularly useful in delivering small histamine particles to the smallest airways. It may thus be concluded that receptors mediating bronchoconstriction, which are stimulated by histamine, exist in the small airways of healthy, human subjects.

A small cumulative effect of histamine was observed to occur on Day 1 as evidenced by pre-challenge pulmonary function, upon reaching a dose of 64 mg./ml., the highest threshold dose of histamine for any subject. However, no cumulative response was apparent on Days 2, 3, and 4. The finding of a cumulative response on Day 1 is surprising. Juniper, et al. (87) found no such response after waiting only five minutes between successive histamine challenges, although they did not challenge anyone with more than 16 mg./ml. of histamine. Perhaps the time required to adequately clear 32 mg./ml. from the respiratory system, as delivered to the airways by the protocol used in this study is greater than the thirty minutes which was permitted before the 64 mg./ml. challenge. The small decline in baseline pulmonary function prior to the

64 mg./ml. challenge might also be attributed to the general fatigue of the subjects, who spent over two hours waiting and who previously performed the pulmonary function maneuver more than for any other test dose.

Chlorpheniramine had a very significant effect in preventing induced bronchoconstriction due to histamine inhalation. Blockade of the H₁ receptors with the oral medication was significantly greater than placebo. The difference between chlorpheniramine and placebo could be detected at both the threshold and superthreshold doses of histamine. As in the histamine dose response curve, MEF₄₀(P) was a more sensitive test than FEV₁ in detecting differences between chlorpheniramine and placebo, as well as detecting the protective effect of chlorpheniramine as compared to the histamine dose response curve.

Chlorpheniramine did not have a completely protective effect against the bronchoconstricting effects of histamine. This may have been due, in part, to incomplete antagonism of all the H₁ receptors in the small airway by the oral medication. Inhaled H₁ receptor antagonists, such as clemastine (113) have apparently been more successful than oral medications in providing H₁ antagonism in asthmatics. In fact, clemastine (113) causes bronchodilation in asthmatics, perhaps by leaving

endogenous histamine free to stimulate bronchodilating H₂ receptors.

Chlorpheniramine is a competitive antagonist of H₁ receptors. Thus, high concentrations of histamine, such as those which may have been by the T+1 dose, could displace the antagonist from the receptor.

A small component on the bronchoconstriction due to histamine may have been the result of vagally-mediated, reflex bronchoconstriction (52). Histamine particles, particularly those with a moderately acidic pH among the higher doses, may have stimulated irritant receptors, particularly in the larger airways (47) to cause bronchoconstriction which can not be prevented by H₁ antagonism. In this study, parasympatholytic agents, such as atropine, may have provided additional protection.

Unlike inhaled clemastine (113), chlorpheniramine did not yield any bronchodilation either at baseline or upon histamine challenge. This result does not support the presence of H₂ receptors in the normal human lung which mediate bronchodilation.

The absence of an H₂ receptor-mediated bronchodilation at baseline may be explained as due to either the absolute deficiency of H₂ receptors, or more likely, the absence of endogenous histamine in non-asthmatics which could bind with H₂ receptors. The fact that histamine challenge did not yield bronchodilation

during H₁ receptor blockade, may be due to the absolute lack of H₂ receptors. It may also be possible, that with the oral medication, even at subthreshold levels of histamine, there were still an adequate number of unbound H₁ receptors agonized by histamine to counter any effect of H₂ agonism. Thus, a net bronchoconstriction would be observed.

Further studies in the non-asthmatic might explore the possible bronchodilatory effects of an inhaled antihistamine on both baseline pulmonary function and response to histamine challenge. Smaller doses of histamine than those which were used in this study might simulate endogenous levels of histamine in the asthmatic and agonize any possibly existent H₂ receptors without displacing H₁ antagonists and causing a net H₁ receptor-mediated bronchoconstriction. Pretreatment with an H₂ antagonist in addition to an H₁ antagonist on the same day might remove H₂-mediated bronchodilation, resulting in a significantly larger net bronchoconstriction than in pretreatment with only an H₁ antagonist.

No placebo effect was observed. Placebos have been reported to influence the amount of bronchoconstriction due to histamine provocation (131). However, the placebo has to be presented to the subject as a substance which has a bronchoconstricting or broncho-

dilating effect. Subjects in this study were not informed as to what the effects of the drugs might be with regard to subsequent histamine challenge.

Cimetidine was no different than placebo with regard to pulmonary function observed upon histamine challenge. Blockade of H₂ receptors neither protected against or potentiated bronchoconstriction. If H₂ receptors, which mediated bronchoconstriction, existed in the lung, then cimetidine would be expected to have a protective effect, as was the case with chlorpheniramine.

If H₂ receptors which mediate bronchorelaxation were successfully antagonized by cimetidine, then subsequent histamine challenge would stimulate only H₁ receptors. This would promote an even greater bronchospasm than that observed with placebo, in which case, both H₁ and H₂ receptors are stimulated.

The H₂ receptor has been implicated, through cyclic AMP mechanisms, to have a variety of effects which would inhibit the effects of histamine challenge. These effects include H₂ receptor-mediated inhibition of further histamine release (94), inhibition of histamine production (104, 125), promotion of histamine metabolism (67, 107, 125), promotion of histamine clearance from the circulation (34, 133), and promotion

of histamine uptake by mast cells (34, 68, 107). Blockade of these effects, particularly the effects on metabolism, uptake, and clearance, by cimetidine would also potentiate histamine-induced bronchospasm. However, cimetidine was not successful in potentiating histamine-induced bronchospasm.

These findings are consistent with the preliminary reports of Eiser (59) and Maconochie (99) who also found no bronchoconstrictory effects of cimetidine. The most likely conclusion to be drawn from the evidence is that H_2 receptors do not exist to any significant degree in non-asthmatic airways. If H_2 receptors are present in the lung, the effects of H_2 agonism are negligible compared to the bronchoconstrictory effects of H_1 agonism.

If there are few H_2 receptors in the lung, it is possible that at the doses of histamine used for this experiment, the histamine overcame the competitive antagonism by cimetidine for H_2 receptor sites in the lung. Thus, cimetidine might have been displaced from the opportunity to potentiate bronchospasm. At lower levels of histamine agonism, H_2 blockade might have potentiated bronchospasm. While it is difficult to measure bronchospasm in non-asthmatics at low levels of histamine challenge, asthmatics are much more

sensitive to low levels of histamine (91, 110, 139). The finding that cimetidine promotes bronchospasm in asthmatics (71, 109) may be due to the fact that cimetidine is not displaced by the low levels of histamine used in the histamine challenges. In addition, acting through mechanisms which potentiate the presence and release of histamine (34, 67, 68, 94, 104, 125, 133) small levels of additional histamine which would not provoke bronchospasm in the non-asthmatic might potentiate bronchospasm in the sensitive asthmatic.

It is possible, as has been pointed out by Macanochie, et al. (99) that the cimetidine used in this study did not achieve adequate levels in the lung. Cimetidine levels in human lung following oral administration have not been studied. The effects of cimetidine on histamine metabolism, uptake and clearance have also not been investigated. However, the dosage of cimetidine used in this study provided a blood level of twice that required to inhibit gastric acid secretion in the stomach by 80%.

Perhaps future studies could investigate the effects of an inhaled H₂ agonist, such as 4-methylhistamine or dimaprit on pulmonary function. Future studies could also investigate the effects of an inhaled H₂ antagonist,

in hopes of achieving better penetration in the lung than with oral administration.

A possible reason for the findings of this investigation may be the failure of cimetidine to adequately antagonize the histamine receptors in the lung which mediate bronchodilation. A previous study (66) demonstrated that cimetidine did not potentiate the depressor actions of histamine in tissue in which burimamide and metiamide, which belong to a different class of H₂ antagonists, potentiated the depressor effects on carotid blood pressure. Similarly, a subclass of H₂ receptor not antagonized by cimetidine, might be present in the human lung.

Several other studies have demonstrated the presence of histamine receptors in the airways of the rat, ferret (40), cat (35), horse (37, 39), and rabbit (36, 70) which mediate bronchodilation and can be agonized by 4-methylhistamine (40), yet which cannot be antagonized by conventional H₂ antihistamines. Perhaps such a class of H₃ receptors may exist in human lung.

Histamine was aerosolized so as to reach the smallest airways (45, 46). The MEF₄₀(P) pulmonary function test is particularly sensitive to small airways constriction (20, 105). If H₂ receptors in the human lung exist

predominantly in the large airways, the protocol of this investigation may not have been sensitive enough to document any effects of H₂ blockade. Regional differences in the distribution of H₂ receptors has been demonstrated in several animal species (14, 64, 156). For example, H₂ histamine receptors are more prominent in the central airways than the peripheral airways in the guinea pig (156).

The regional differences in histamine receptor distribution may correlate with the physiologic importance of maintaining the patency of the large airways. An alternative study could examine the effects of larger aerosolized histamine particles on pulmonary flow resistance, a sensitive measure of central resistance (98), in order to more closely investigate the presence of H₂ receptors in large airways. Atropine would have to be used in this proposed protocol, because of the relative importance of irritant receptors in the large airways.

In conclusion, the experimental evidence presented in this investigation confirms the presence of H₁ receptors in the airways of healthy human subjects. These receptors appear to mediate bronchoconstriction and can be successfully antagonized by chlorpheniramine.

The role of H₂ receptors in healthy human airways, as either bronchodilators acting directly on smooth muscle or as inhibitors of the presence of histamine, appears to be negligible under the conditions of the protocol used in this study. Cimetidine did not potentiate histamine-induced bronchospasm. The role of H₂ receptors in asthmatic bronchoconstriction and bronchodilation remains a possibility for future investigation.

MAXIMAL AND PARTIAL EXPIRATORY FLOW-VOLUME CURVES




Figure 1

HISTAMINE DOSE RESPONSE AS CHARACTERIZED
BY TWO MEASUREMENTS OF PULMONARY FUNCTION
(FEV₁ and MEF 40% (P))




Figure 2

THE EFFECT OF ANTIHISTAMINES
ON HISTAMINE CHALLENGE AS MEASURED
BY MEF 40% (P)




Figure 3

THE EFFECT OF ANTIHISTAMINES
ON HISTAMINE CHALLENGE AS MEASURED
BY FEV₁




Figure 4

TABLE I
ANTHROPOMORPHIC DATA

	FVC	FEV ₁	FEV ₁ %	PEFR	MEF ₅₀ %	Age
Mean	4.41	3.68	83.9	8.21	4.50	26.3
Standard deviation	0.82	0.69	9.3	1.62	2.10	5.1
Percent of expected values	89%	94%		93%	102%	

TABLE II

BASELINES AT BEGINNING OF DAY
AS PERCENTAGE OF BASELINE ON DAY 1

DAY	1	CM	CH	P
FEV ₁	100	99.6	99.8	99.0
MEF ₄₀ (P)	100	98.1	104.6	109.9

CM=Cimetidine

CT=Chlorpheniramine

P =Placebo

TABLE III

AVERAGE HISTAMINE DOSE-RESPONSE
 PRE-CHALLENGE CONTROLS AS
 PERCENT OF INITIAL BASELINE

DOSE	.1	2	4	8	16	32	64
FEV ₁	100	100	98.9	98.1	98.6	99.0	100.7
MEF ₄₀ (P)	100	110	94.1	102.4	98.1	93.9	89.0*

*p= .031

TABLE IV

PRE-CHALLENGE CONTROLS ON DAYS 2, 3, AND 4
AS PERCENT OF INITIAL BASELINE

CIMETIDINE

DOSE	T-1	T	T+1
FEV ₁	100	98.45	98.6
MEF ₄₀ (P)	100	101.4	101.9

CHLORPHENIRAMINE

DOSE	T-1	T	T+1
FEV ₁	100	100.36	100.8
MEF ₄₀ (P)	100	102.3	98.4

PLACEBO

DOSE	T-1	T	T+1
FEV ₁	100	99.5	102.0
MEF ₄₀ (P)	100	97.0	95.4

T=Threshold Dose

TABLE V

THE EFFECT OF PRETREATMENT WITH ANTIHISTAMINE
UPON SUBSEQUENT RESPONSE TO HISTAMINE(Means \pm Standard Error of the Mean)

DOSE	FEV ₁		
	T-1	T	T+1
CIMETIDINE	99.0 ± 1.13	93.64 ± 2.79	86.0 ± 4.30
CHLORPHENIRAMINE	101.72 ± 1.71	98.91 ± 1.23	96.09 ± 2.60
PLACEBO	101.88 ± 4.42	95.75 ± 2.54	85.88 ± 2.49

MEF₄₀(P)

DOSE	T-1	T	T+1
CIMETIDINE	96.0 ± 4.26	79.64 ± 3.70	57.80 ± 7.39
CHLORPHENIRAMINE	99.0 ± 4.17	91.45 ± 2.54	82.18 ± 5.58
PLACEBO	91.88 ± 3.66	78.13 ± 5.72	57.88 ± 5.15

TABLE VI

HISTAMINE DOSE RESPONSE
AS PERCENTAGE OF PRE-CHALLENGE CONTROL

(Means \pm Standard Error of the Mean)

DOSE	.1	2	4	8	16	32	64
FEV ₁	101.1 ± 2.65	103 ± 0	98.63 ± 1.39	101.6 ± 2.01	99.45 ± 0.73	91.0 ± 3.59	87.57 ± 2.98
MEF ₄₀ (P)	99.90 ± 2.68	97 ± 0	100.5 ± 3.17	94.2 ± 2.46	92.0 ± 3.90	79.72 ± 4.92	73.29 ± 4.53

TABLE VII

RESPONSE TO INHALED HISTAMINE
 AS COMPARED TO RESPONSE TO
 0.1 MG./ML. OF INHALED HISTAMINE

DOSE	4	8	16	32	64
FEV ₁	NS	NS	NS	p<.05	p<.05
MEF ₄₀ (P)	NS	NS	p=.10	p<.005	p<.001

NS=No significance

TABLE VIII

COMPARISON OF THE PROTECTIVE EFFECT
OF CHLORPHENIRAMINE, CIMETIDINE, AND PLACEBO
UPON SUBSEQUENT RESPONSE TO INHALED HISTAMINE
AS MEASURED BY $MEF_{40}(P)$

	DAY 1	CHLORPHENIRAMINE(CT)	CIMETIDINE(CM)	PLACEBO(P)
<u>DAY 1</u>				
T-1	*****	NS	NS	NS
T	*****	p<.0025	NS	NS
T+1	*****	*	*	*
<u>CT</u>				
T-1	NS	*****	NS	NS
T	p<.0025	*****	p<.01	p<.05
T+1	*	*****	p<.0005	p<.005
<u>CM</u>				
T-1	NS	NS	*****	NS
T	NS	p<.01	*****	NS
T+1	*	p<.0005	*****	NS
<u>P</u>				
T-1	NS	NS	NS	*****
T	NS	p<.05	NS	*****
T+1	*	p<.005	NS	*****

*Insufficient data to make comparison
NS=No significance

TABLE IX

COMPARISON OF THE PROTECTIVE EFFECT
OF CHLORPHENIRAMINE, CIMETIDINE, AND PLACEBO
UPON SUBSEQUENT RESPONSE TO INHALED HISTAMINE
AS MEASURED BY FEV₁

	DAY 1	CHLORPHENIRAMINE(CT)	CIMETIDINE(CM)	PLACEBO(P)
<u>DAY 1</u>				
T-1	*****	p=.07	NS	NS
T	*****	p=.14	NS	NS
T+1	*****	*	*	*
<u>CT</u>				
T-1	p=.07	*****	NS	NS
T	p=.14	*****	p<.05	NS
T+1	*	*****	p<.001	p<.05
<u>CM</u>				
T-1	NS	NS	*****	NS
T	NS	p<.05	*****	NS
T+1	*	p<.001	*****	NS
<u>P</u>				
T-1	NS	NS	NS	*****
T	NS	NS	p<.05	*****
T+1	*	p<.05	p<.001	*****

*Insufficient data to make comparison
NS=No significance

BIBLIOGRAPHY

1. Alexander, F., P. Eyre, K.W. Head, and J. Sanford (1967) Effects of Histamine and 5-Hydroxytryptamine on Sheep. Journal of Comparative Pathology. 77: 1-12.
2. Allouche, G., M-C. Santais. C. Foussard, F. Ruff, and J.-L. Parrot (1977) Role of H₂ Receptor Antagonists in the Feedback of Histamine (abstract). Agents and Actions, 7: 103-104.
3. Altounyan. R.E.C. (1971) Comments in Discussion. in Identification of Asthma, Ruth Porter and Joan Birch (editors). Churchill Livingstone, Edinburgh and London. pages 160-162.
4. American Thoracic Society Committee on Diagnostic Standards for Nontuberculous Respiratory Diseases (1962) Chronic Bronchitis, Asthma, and Pulmonary Emphysema. American Review of Respiratory Disease, 85: 762-768.
5. Antonissen, L.A., R.W. Mitchell, E.A. Kroeger, and N.L. Stephens (1978) Airway Smooth Muscle Pharmacology in a Canine Asthmatic Model (abstract). Federation Proceedings, 37: 639.
6. Ash, A.S.F., and H.O. Schild (1966) Receptors Mediating Some Aspects of Histamine. British Journal of Pharmacology and Chemotherapeutics, 27: 427-439.
7. Audigier, Y., A. Viron, and J.C.-Schwartz (1977) Clonidine, a Potent Agonist of Histamine H₂-Receptors in Guinea-Pig Brain (abstract). Agents and Actions, 7: 104.
8. Austen, K. Frank, and Robert P. Orange (1975) Bronchial Asthma: The Possible Role of the Chemical Mediators of Immediate Hypersensitivity in the Pathogenesis of Subacute Chronic Disease. American Review of Respiratory Disease, 112: 423-436.

9. Avella, Jerry, John E. Madsen, Henry J. Binder, and Philip W. Askenase (1978) Effect of Histamine H₂-Receptor Antagonism on Delayed Hypersensitivity. Lancet 1: 624-626.
10. Barer, Gwenda R., Celia J. Emery, F.H. Mohammed, and I.P.F. Mungall (1978) H₁ and H₂ Histamine Actions on Lung Vessels: Their Relevance to Hypoxic Vasoconstriction. Quarterly Journal of Experimental Physiology, 63: 157-169.
11. Bartosch, R., W. Feldberg, and E. Nagel (1932) Das Freiwerden Eines Histaminähnlichen Stoffes bei der Anaphylaxie des Meerschweinchens. Pflügers Archiv Für de Gesamte Physiologie, 230: 120-153.
12. Beaven, Michael A. (1976) Histamine (First of Two Parts). New England Journal of Medicine, 294: 30-36.
13. Beaven, Michael A. (1976) Histamine (Second of Two Parts). New England Journal of Medicine, 294: 320-325.
14. Beaven, Michael A. (1978) Histamine: Its Role in Physiological and Pathological Processes. Monographs in Allergy, 13: 1-113.
15. Benson, Malcolm K. (1975) Bronchial Hyperreactivity. British Journal of Diseases of the Chest, 69: 227-239.
16. Berti, F., G.C. Folco, Simonetta Nicosia, C. Omini, and R. Pasargiklian (1979) The Role of Histamine H₁- and H₂- Receptors in the Generation of Thromboxane A₂ in Perfused Guinea-Pig Lungs. British Journal of Pharmacology, 65: 629-633.
17. Bhat, K. Naras, Carlos M. Arroyave, Samuel R. Marney, Jr., Donald D. Stevenson, and Eng M. Tan (1976) Plasma Histamine Changes During Provoked Bronchospasm in Asthmatic Patients. Journal of Allergy and Clinical Immunology, 58: 647-656.

18. Black., J.W., W.A.M. Duncan, C.J. Durant, C.R. Ganellin, and E.M. Parsons (1972) Definition and Antagonism of H₂- Receptors. Nature, 236: 385-390.
19. Black, J.W., D.A.A. Owen, and M.E. Parsons (1975) An Analysis of the Depressor Responses to Histamine in the Cat and Dog: Involvement of Both H₁- and H₂- Receptors. British Journal of Pharmacology, 54: 319-324.
20. Bouhuys, A., V.R. Hunt, B.M. Kim, and A. Zapletal (1969) Maximum Expiratory Flow Rates in Induced Bronchoconstriction in Man. Journal of Clinical Investigation, 48: 1159-1168.
21. Bourne, Henry R., Kenneth L. Melmon, and Lawrence M. Lichtenstein (1971) Histamine Augments Leukocyte Adenosine 3', 5'-Monophosphate and Blocks Antigenic Histamine Release. Science, 273: 743-745.
22. Brimblecombe, R.W., W.A.M. Duncan, G.J. Durant, J.C. Emmett, C.R. Ganellin, G.B. Leslie, and M.E. Parsons (1978) Characterization and Development of Cimetidine as a Histamine H₂- Receptor Antagonist. Gastroenterology, 74: 339-347.
23. Burkard, Willy P. (1978) Histamine H₂- Receptor Binding With ³H-Cimetidine in Brain. European Journal of Pharmacology, 50: 449-450.
24. Bush, Robert K., William Busse, Dennis Flaherty, David Warshauer, Elliot C. Dick, and Charles E. Reed (1978) Effects of Experimental Rhinovirus 16 Infection on Airways and Leukocyte Function in Normal Subjects. Journal of Allergy and Clinical Immunology, 61: 80-87.
25. Busse, William W., and Sharon D.H. Lantis (1979) Impaired H₂ Histamine Granulocyte Response in Active Atopic Eczema. Journal of Investigative Dermatology, 73: 184-187.
26. Busse, William W., and Jeffrey Sosman (1977) Decreased H₂ Histamine Response of Granulocytes of Asthmatic Patients. Journal of Clinical Investigation, 59: 1080-1087.

27. Caputi, A.P., E. Lampa, F. Rossi, G. Nistico, and E. Marmo (1978) Activation of Histamine H₂-Receptors by St 600. Agressologie, 19: 35-37.
28. Casterline, Charlotte L., and Richard Evans III (1977) Further Studies on the Mechanism of Human Histamine-Induced Asthma: The Effect of an Aerosolized H₁ Receptor Antagonist (Diphenhydramine). Journal of Allergy and Clinical Immunology, 59: 420-424.
29. Casterline, Charlotte L., Richard Evans III, and George W. Ward, Jr. (1976) The Effect of Atropine and Albuterol Aerosols on the Human Bronchial Response to Histamine. Journal of Allergy and Clinical Immunology, 58: 607-613.
30. Chakrin, L.W., R.D. Krell, J. Mengel, D. Young, C. Zaher, and J.R. Wardell, Jr. (1974) Effect of a Histamine H₂ Receptor Antagonist on Immunologically Induced Mediator Release In Vitro. Agents and Actions, 4: 297-303.
31. Chakrin, L.W., J. Mengel, D. Young, C. Zaher, R.D. Krell, and J.R. Wardell, Jr. (1974) Enhancement of Immediate-Type Hypersensitivity Reactions By N-Methyl-N'-[2[(4-Methyl-5 Imidazolyl)methylthio]ethyl]thiourea, A Histamine H₂- Receptor Antagonist (abstract). Federation Proceedings, 33: 585.
32. Chand, N. (1979) In Vitro Anaphylaxis in Guinea-Pig Lung: Evidence for the Protective Role of Histamine H₂- Receptors. European Journal of Pharmacology, 55: 337-339.
33. Chand, N., L. DeRoth, and P. Eyre (1979) Pharmacology of Schultz-Dale Reaction in Canine Lung Strip In Vitro: Possible Model for Allergic Asthma. British Journal of Pharmacology, 66: 511-516.
34. Chand, N., and P. Eyre (1975) Classification and Biological Distribution of Histamine Receptor Sub-Types. Agents and Actions, 5: 277-295.

35. Chand, N., and P. Eyre (1977) Atypical (Relaxant) Response to Histamine in Cat Bronchus. Agents and Actions, 7: 183-190.
36. Chand, N., and P. Eyre (1977) Spasmolytic Actions of Histamine in Airway Smooth Muscle--An Atypical H-Receptor (abstract). Federation Proceedings, 36: 1022.
37. Chand, N., and P. Eyre (1977) Histamine Relaxes Constricted Trachea and Bronchi of Horse. Veterinary Science Communications, 1: 85-90.
38. Chand, Naresh, and Peter Eyre (1978) The Schultz-Dale Reaction: A Review. Agents and Actions, 8: 171-184.
39. Chand, N., and P. Eyre (1978) Spasmolytic Action of Histamine in Airway Smooth Muscle of Horse. Agents and Actions, 8: 191-198.
40. Chand, N., and P. Eyre (1978) Histamine Receptors of Airway Smooth Muscle of Ferret and Rat. Research Communications in Chemical Pathology and Pharmacology, 21: 55-65.
41. Chang, Raymond S.L., Vinh Tan Tran, and Solomon H. Snyder (1979) Characteristics of Histamine H₁-Receptors in Peripheral Tissues Labeled With [³H] Mepyramine. Journal of Pharmacology and Experimental Therapeutics, 209: 437-442.
42. Dale, H.H., and P.P. Laidlaw (1910) The Physiological Action of β -Iminazolylethylamine. Journal of Physiology, 41: 318-344.
43. Dale, H.H., and P.P. Laidlaw (1919) Histamine Shock. Journal of Physiology, 52: 355-390.
44. Dale, M. Maureen (1977) The Effects of Metiamide on Cell-Mediated Immune Reactions in the Guinea-Pig. British Journal of Pharmacology, 60: 441-444.
45. Dautrebande, Lucien (1962) Microaerosols. Academic Press, New York.

46. Dautrebande, Lucien (1970) Experimental Observations on the Participation of Alveolar Spaces in Airway Dynamics. in Airway Dynamics: Physiology and Pharmacology, Arend Bouhuys, (editor), Charles C. Thomas, Springfield, pages 153-168.
47. DeKock, M.A., J.A. Nadel, S. Zwi, H.J.H. Colebatch, and C.R. Olsen (1966) New Method For Perfusing Bronchial Arteries: Histamine Bronchoconstriction and Apnea. Journal of Applied Physiology, 21: 185-194.
48. Delaunois, L. (1979) Hypersensitivity to Cimetidine. New England Journal of Medicine, 300: 1216.
49. deVries, K., J.T. Goei, H. Booy-Noord, and N.G.M. Orie (1962) Changes During 24 Hours in the Lung Function and Histamine Hyperreactivity of the Bronchial Tree in Asthmatic and Bronchitic Patients. International Archives of Allergy and Applied Immunology, 20: 93-101.
50. Diaz, Patricia, D.G. Jones, and A.B. Kay (1979) Histamine Receptors on Guinea-Pig Alveolar Macrophages: Chemical Specificity and the Effects of H₁- and H₂- Receptor Agonists and Antagonists. Clinical and Experimental Immunology, 35: 462-469.
51. Dixon, M., D.M. Jackson, and I.M. Richards (1979) The Effects of H₁- and H₂- Receptor Agonists and Antagonists on Total Lung Resistance, Dynamic Lung Compliance and Irritant Receptor Discharge in the Anaesthetized Dog. British Journal of Pharmacology, 66: 203-209.
52. Douglas, William W. (1975) Histamine and Antihistamines; 5-Hydroxytryptamine and Antagonists. in The Pharmacological Basis of Therapeutics (fifth edition), Louis S. Goodman and Alfred Gilman (editors), Macmillan Publishing Company, Inc., New York, pages 590-629.
53. Drazen, Jeffrey M., and M.W. Schneider (1978) Comparative Responses of Tracheal Spirals and Parenchymal Strips to Histamine and Carbachol In Vitro. Journal of Clinical Investigation, 61: 1441-1447.

54. Drazen, Jeffrey M., M.W. Schneider, and C.S. Venugopalan (1979) Bronchodilator Activity of Dimaprit in the Guinea Pig In Vitro and In Vivo. European Journal of Pharmacology, 55: 233-239.
55. Drazen, Jeffrey M., C.S. Venugopalan, and Nicholas A. Soter (1978) H₂ Receptor Mediated Inhibition of Immediate Type Hypersensitivity Reactions In Vivo. American Review of Respiratory Disease, 117: 479-484.
56. Dulabh, R., and M.R. Vickers (1978) The Effects of H₂- Receptor Antagonists on Anaphylaxis in the Guinea-Pig. Agents and Actions, 8: 559-565.
57. Dunlop, Lyn S., and A.P. Smith (1977) The Effect of Histamine Antagonists On Antigen-Induced Contractions of Sensitized Human Bronchus In Vitro. British Journal of Pharmacology, 59: 475P.
58. Edwards, Allen L. (1967) Statistical Methods.(second edition), Holt, Rinehart, and Winston, Inc., New York.
59. Eiser, Naomi M., A. Guz, and P.D. Snashall (1978) The Effect of H₁ and H₂ Receptor Antagonists on Bronchial Dose-Response Curves to Histamine in Normal Subjects. Clinical Science and Molecular Medicine, 54: 10P-11P.
60. Empey, D.W., L.A. Laitinen, L. Jacobs, W.M. Gold, and J.A. Nadel (1976) Mechanisms of Bronchial Hyperreactivity in Normal Subjects After Upper Respiratory Tract Infection. American Review of Respiratory Disease, 113: 131-139.
61. Eyre, P. (1969) The Pharmacology of Sheep Tracheo-bronchial Muscle: A Relaxant Effect of Histamine on the Isolated Bronchi. British Journal of Pharmacology, 36: 409-417.
62. Eyre, P. (1973) Histamine H₂-Receptors in the Sheep Bronchus and Cat Trachea: The Action of Burimamide. British Journal of Pharmacology, 48: 321-323.

63. Eyre, P. (1976) Preliminary Studies of Pharmacological Antagonism of Anaphylaxis in the Horse. Canadian Journal of Comparative Medicine, 40: 149-152.
64. Eyre, P. (1977) Pulmonary Histamine H₁ and H₂ Receptor Studies. in Asthma: Physiology, Immunopharmacology, and Treatment--Second International Symposium, Lawrence M. Lichtenstein and K. Frank Austen (editors), Academic Press, New York, pages 169-180.
65. Eyre, P., and N. Chand (1979) Preliminary Evidence for Two Subclasses of Histamine H₂-Receptors. Agents and Actions, 9: 1-3.
66. Eyre, P., and P.W. Wells (1973) Histamine H₂-Receptors Modulate Systemic Anaphylaxis: A Dual Cardiovascular Action of Histamine in Calves. British Journal of Pharmacology, 49: 364-367.
67. Fantozzi, R, Flavia Franconi, P.F. Mannaioni, Emanuela Massini, and F. Moroni (1975) Interaction of Histamine H₁- and H₂- Receptor Antagonists With Histamine Uptake and Metabolism By Guinea-Pig Isolated Atrium and Mouse Neoplastic Mast Cells In Vitro. British Journal of Pharmacology, 53: 569-574.
68. Fantozzi, R., P.F. Mannaioni, and F. Moroni (1973) The Inhibition of Histamine Uptake and Metabolism By Burimamide in the Guinea-Pig Atrium and in Mouse Neoplastic Mast Cells, In Vitro. British Journal of Pharmacology, 49: 361-363.
69. Fish, James E., Virginia I. Peterman, and David W. Cugell (1977) Effect of Deep Inspiration on Airway Conductance in Subjects With Allergic Rhinitis and Allergic Asthma. Journal of Allergy and Clinical Immunology, 60: 41-46.
70. Fleisch, Jerome H., and Patricia J. Calkins (1976) Comparison of Drug-Induced Responses of Rabbit Trachea and Bronchus. Journal of Applied Physiology, 41: 62-66.
71. Frith, P.A., R.E. Ruffin, F.A. Oldenburg, M.T. Newhouse, and F.E. Hargreave (1980) Evidence for Histamine H₂ Receptors in Asthmatic Airways (in preparation).

72. Gayrard, Pierre, Jean Orehek, Charles Grimaud, and Jacques Charpin (1975) Bronchoconstrictor Effects of a Deep Inspiration in Patients With Asthma. American Review of Respiratory Disease, 111: 433-439.
73. Gillespie, Elizabeth, Martin D. Valentine, and Lawrence M. Lichtenstein (1974) Cyclic AMP Metabolism in Asthma: Studies With Leukocytes and Lymphocytes. Journal of Allergy and Clinical Immunology, 53: 27-33.
74. Giordano, Joseph, Michael Zinner, Alexander Guba, and Noreen Lynch (1977) Effects of H₁ and H₂ Histamine Antagonists on the Pulmonary Pressor Response to Alveolar Hypoxia. Journal of Surgical Research, 22: 392-397.
75. Goadby, P., and M. Ann Little (1978) Histamine H₂-Receptors in Guinea Pig Lung and Their Role in Anaphylaxis. Agents and Actions, 8: 389-390.
76. Greenberger, Paul, Kathleen Harris, and Roy Patterson (1979) The Effect of Histamine-1 and Histamine-2 Antagonists on Airway Responses to Histamine in the Rhesus Monkey. Journal of Allergy and Clinical Immunology, 64: 189-196.
77. Hanna, C.J., and P. Eyre (1978) Pharmacological Studies on the Pulmonary Vein of the Horse. I. Effects of Selected Spasmogens. Canadian Journal of Physiology and Pharmacology, 56: 812-817.
78. Harvey, R.P., and A.L. Schocket (1979) The Effect of H₁ and H₂ Blockade on Cutaneous Histamine Response in Man (abstract). Journal of Allergy and Clinical Immunology, 63: 208-209.
79. Hoffman, Eric A., Mary L. Munroe, Alan Tucker, and John T. Reeves (1977) Histamine H₁- and H₂-Receptors in the Cat and Their Roles During Alveolar Hypoxia. Respiration Physiology, 29: 255-264.

80. Holroyde, Martin C., and Peter Eyre (1978) Histamine Enhances Anaphylactic Histamine Release From Bovine Lung and Leukocytes Via Histamine H₂-Receptor. Journal of Pharmacology and Experimental Therapeutics, 204: 183-188.
81. Horton, Douglas J., William L. Suda, Robert A. Kinsman, Joseph Souhrada, and Sheldon L. Spector (1978) Bronchoconstrictive Suggestion in Asthma: A Role For Airways Hyperreactivity and Emotions. American Review of Respiratory Disease, 117: 1029-1038.
82. Irvin, C.G., and J.A. Dempsey (1977) Relative Contribution of H₁ and H₂ Receptors to Large and Small Airway Constriction (abstract). Federation Proceedings, 36: 613.
83. Irvin, Charles G., and J.A. Dempsey (1978) Role of H₁ and H₂ Receptors in Increased Airway Resistance in the Dog. Respiration Physiology, 35: 161-176.
84. Itkin, Irving H., and Suresh C. Anand (1970) The Role of Atropine as a Mediator Blocker of Induced Bronchial Obstruction. Journal of Allergy, 45: 178-186.
85. Joiner, P.D., M. Wall, L.B. Davis, and F. Hahn (1974) Role of Amines in Anaphylactic Contraction of Guinea Pig Isolated Smooth Muscle. Journal of Allergy and Clinical Immunology, 53: 261-270.
86. Jorizzo, Joseph L., W. Mitchell Sams, Jr., Brian V. Jegasothy, and Alan L. Olansky (1980) Cimetidine as an Immunomodulator: Chronic Mucocutaneous Candidiasis as a Model. Annals of Internal Medicine, 92: 192-195.
87. Juniper, E.F., P.A. Frith, C. Dunnett, D.W. Cockcroft, and F.E. Hargreave (1978) Reproducibility and Comparison of Responses To Inhaled Histamine and Methacholine. Thorax, 33: 705-710.
88. Kaliner, Michael (1978) Human Lung Tissue and Anaphylaxis: The Effects of Histamine on the Immunologic Release of Mediators. American Review of Respiratory Disease, 118: 1015-1022.

89. Kaliner, M., and L. Platshon (1978) The Effect of Histamine on Human Lung Anaphylaxis. Journal of Allergy and Clinical Immunology, 61: 130.
90. Krell, Robert D., and Lawrence W. Chakrin (1977) The Effect of Metiamide in In Vitro and In Vivo Canine Models of Type I Hypersensitivity Reactions. European Journal of Pharmacology, 44: 35-44.
91. Laitinen, Lauri A.I. (1974) Histamine and Metacholine Challenge in the Testing of Bronchial Reactivity. Scandinavian Journal of Respiratory Diseases Supplementum, 86: 3-48.
92. Levi, Roberto, Geoffrey Allan, and James H. Zavecz (1976) Cardiac Histamine Receptors. Federation Proceedings, 35: 1942-1947.
93. Lichtenstein, Lawrence M. (1977) Mediator Release and Asthma. in Asthma: Physiology, Immunopharmacology, and Treatment--Second International Symposium, Lawrence M. Lichtenstein and K. Frank Austen (editors), Academic Press, New York, pages 169-180.
94. Lichtenstein, Lawrence M., and Elizabeth Gillespie (1973) Inhibition of Histamine Release By Histamine Controlled By H₂ Receptor. Nature, 244: 287-288.
95. Littner, Michael R., E. Neil Schachter, and Arend Bouhuys (1979) Modification of Induced Airway Constriction in Healthy Subjects. Annals of Allergy, 43: 135-143.
96. Lulich, K.M., H.W. Mitchell, and M.P. Sparrow (1976) The Cat Lung Strip as an In Vitro Preparation of Peripheral Airways: A Comparison of β -Adrenoceptor Agonists, Autoacids, and Anaphylactic Challenge on the Lung Strip and Trachea. British Journal of Pharmacology, 58: 71-79.
97. Luparello, Thomas J., Nancy Leist, Cary H. Lourie, and Pauline Sweet (1970) The Interaction of Psychologic Stimuli and Pharmacologic Agents on Airway Reactivity in Asthmatic Subjects. Psychosomatic Medicine, 32: 509-513.

98. Macklem, Peter T., and Jere Mead (1967) Resistance of Central and Peripheral Airways Measured By a Retrograde Catheter. Journal of Applied Physiology, 22: 395-401.
99. Maconochie, J.G., E.P. Woodings, and D.A. Richards (1979) Effects of H₁- and H₂- Receptor Blocking Agents on Histamine-Induced Bronchoconstriction in Non-Asthmatic Subjects. British Journal of Clinical Pharmacology, 7: 231-236.
100. Maengwyn-Davies, Gertrude D. (1968) The Dual Mode of Action in the Cat Isolated Tracheal Chain. Journal of Pharmacy and Pharmacology, 20: 572-573.
101. Marks, R., and M.W. Greaves (1977) Vascular Reactions to Histamine and Compound 48/80 in Human Skin: Suppression By a Histamine H₂-Receptor Blocking Agent. British Journal of Clinical Pharmacology, 4: 367-369.
102. Martin, G., and R.H. Fertel (1978) Guinea Pig Lung May Have Both H₁ and H₂ Receptors (abstract). Federation Proceedings, 37: 393.
103. Mathé, A.A., L. Volicer, and S.K. Puri (1974) Effect of Anaphylaxis and Histamine, Pyrilamine, and Burimamide on Levels of Cyclic AMP and Cyclic GMP in Guinea-Pig Lung. Research Communications in Chemical Pathology and Pharmacology, 8: 635-651.
104. Maudsley, David V., Yutaka Kobayashi, Ethel Williamson, and Larry Bovaird (1973) Stimulation of Histidine Decarboxylase By Burimamide in the Rat Glandular Stomach (abstract). Federation Proceedings, 32: 808.
105. Mead, Jere, James M. Turner, Peter T. Macklem, and John B. Little (1967) Significance of the Relationship Between Lung Recoil and Maximum Expiratory Flow. Journal of Applied Physiology, 22: 95-108.
106. Middleton, Elliot Jr. (1972) Autonomic Imbalance in Asthma With Special Reference to Beta Adrenergic Blockade. Advances in Internal Medicine, 18: 177-197.

107. Moroni, F., R. Fantozzi, and P.F. Mannioni (1973) The Influence of Burimamide on ¹⁴C-Histamine Uptake and Metabolism by Guinea Pig Atrium and Mouse Neoplastic Mast Cells, In Vitro. Agents and Actions, 3: 187-189.
108. Nadel, Jay A., and Donald F. Tierney (1961) Effect of a Previous Deep Inspiration on Airway Resistance in Man. Journal of Applied Physiology, 16: 717-719.
109. Nathan, Robert A., Nathan Segall, Gail C. Glover, and Alan L. Schocket (1979) The Effects of H₁ and H₂ Antihistamines on Histamine Inhalation Challenges in Asthmatic Patients. American Review of Respiratory Disease, 120: 1251-1258.
110. Newball, H.H., and H.R. Keiser (1973) Relative Effects of Bradykinin and Histamine on the Respiratory System in Man. Journal of Applied Physiology, 35: 552-556.
111. Newman, John H., Joseph F. Souhrada, John T. Reeves, Carlos M. Arroyave, and Robert F. Grover (1979) Postnatal Changes in Response of Canine Neonatal Pulmonary Arteries to Histamine. American Journal of Physiology, 237: H76-H82.
112. Nogrady, S.G., and C. Bevan (1978) Inhaled Antihistamines--Bronchodilatation and Effect of Histamine- and Methacholine-Induced Bronchoconstriction. Thorax, 33: 700-704.
113. Nogrady, S.G., J.P.R. Hartley, P.D.J. Handslip, and N.P. Hurst (1978) Bronchodilatation After Inhalation of the Antihistamine Clemastine. Thorax, 33: 479-482.
114. Norn, S., and P. Stahl Skov (1979) Pathophysiologic Mechanisms in Asthma. Molecular and Pharmacological Aspects. Scandinavian Journal of Respiratory Diseases Supplementum, 103: 25-32.
115. Okpako, D.T. (1972) A Dual Action of Histamine on Guinea-Pig Lung Vessels. British Journal of Pharmacology, 45: 311-321.

116. Okpako, D. Tinakpo'Evwan (1977) A Vasodepressor Action of Histamine Mediated By H_2 -Receptor Activation, European Journal of Pharmacology, 29: 10-14.
117. Okpako, D.T., N. Chand, and P. Eyre (1978) The Presence of Inhibitory Histamine H_2 -Receptors in Guinea-Pig Tracheobronchial Muscle. Journal of Pharmacy and Pharmacology, 30: 181-182.
118. Ortez, Richard A. (1977) Pharmacologic Effect of the Blockade of Histamine on Lung Cyclic AMP Levels in Normal and Pertussis-Vaccinated Mice. Biochemical Pharmacology, 26: 529-533.
119. Pandit, P.R., Sunita D. Kulkarni, and G.V. Joglekar (1973) Study of Relationship Between Antihistaminic, Anticholinergic and Central Depressant Properties of Various Antihistaminics. Indian Journal of Medical Sciences, 27: 920-924.
120. Parker, Charles W., and Jay W. Smith (1973) Alterations in Cyclic Adenosine Monophosphate Metabolism in Human Bronchial Asthma. I. Leukocyte Responsiveness to β -Adrenergic Agents. Journal of Clinical Investigation, 52: 48-59.
121. Parsons, M.E., D.A.A. Owen, C.R. Ganellin, and G.J. Durant (1977) Dimaprit—[S-[3-(N,N-dimethylamino)propyl]isothiourea]—A Highly Specific Histamine H_2 -Receptor Agonist. Part 1. Pharmacology. Agents and Actions, 7: 31-37.
122. Peets, E.A., M. Jackson, and S. Symchowicz (1972) Metabolism of Chlorpheniramine Maleate in Man. Journal of Pharmacology and Experimental Therapeutics, 180: 464-474.
123. Persson, Carl G.A., and Marianne Ekman (1976) Contractile Effects of Histamine in Large and Small Respiratory Airways. Agents and Actions, 6: 389-393.
124. Platshon, Leslie F., and Michael Kaliner (1978) The Effects of Histamine Upon Human Lung Cyclic Nucleotide Levels and Prostaglandin Generation. Journal of Clinical Investigation, 62: 1113-1121.

125. Plaut, Marshall (1979) Histamine H₁ and H₂ Antihistamines, And Immediate Hypersensitivity Reactions. Journal of Allergy and Clinical Immunology, 63: 371-375.
126. Plaut, Marshall, Lawrence M. Lichtenstein, and Christopher S. Henney (1975) Properties of a Subpopulation of T Cells Bearing Histamine Receptors. Journal of Clinical Investigation, 55: 856-874.
127. Popa, Valentin T. (1977) Bronchodilating Activity of an H₁ Blocker, Chlorpheniramine. Journal of Allergy and Clinical Immunology, 59: 54-63.
128. Richards, I.M., and D.M. Jackson (1976) An Investigation of Histamine Aerosol Induced Reflex Bronchoconstriction in the Anaesthetized Dog. British Journal of Pharmacology, 57: 439P-440P.
129. Rocklin, Ross E. (1976) Modulation of Cellular Immune-Responses In Vivo and In Vitro by Histamine Receptor-Bearing Lymphocytes. Journal of Clinical Investigation, 57: 1051-1058.
130. Rowley-Jones, D., W.L. Burland, and R. Griffiths (1978) Pharmacokinetics and Pharmacological Properties of Cimetidine. in Cimetidine: Proceedings of an International Symposium on Histamine H₂-Receptor Antagonists, W. Creutzfeldt (editor), Excerpta Medica, Amsterdam, pages 72-78.
131. Ruffin, R.E., P.A. Frith, R.C. Anderton, C.R. Kumana, M.T. Newhouse, and F.E. Hargreave (1979) Selectivity of Beta Adrenoreceptor Antagonist Drugs Assessed By Histamine Bronchial Provocation. Clinical Pharmacology and Therapeutics, 25: 536-540.
132. Sampson, S.R., and E.H. Vidruk (1979) The Nature of the Receptor Mediating Stimulant Effects of Histamine On Rapidly Adapting Vagal Afferents in the Lungs. Journal of Physiology, 287: 509-518.

133. Schayer, Richard W. (1973) Effect of Burimamide on Histamine Metabolism. Agents and Actions, 3: 191-192.
134. Schmidt, Michael J., Lewis L. Truex, Carol S. Hooker, Stephen M. Spaethe, and Jerome H. Fleisch (1979) Ethanol Inhibition of Antigen-Induced Histamine Release From Guinea Pig Lung: Correlation With Intracellular Levels of Cyclic Nucleotides. Life Sciences, 24: 2321-2322.
135. Sellick, Hilary, and J.G. Widdicombe (1971) Stimulation of Lung Irritant Receptors By Cigarette Smoke, Carbon Dust, and Histamine Aerosol. Journal of Applied Physiology, 31: 15-19.
136. Simon, R.A., M. Ginsberg, R.M. Timms, and D.D. Stevenson (1979) Exercise-Induced Bronchospasm: A Study of Plasma Mediators. Journal of Allergy and Clinical Immunology, 63: 153.
137. Simonsson, B.G., F.M. Jacobs, and J.A. Nadel (1967) Mechanisms of Changes in Airway Size During Inhalation of Various Substances in Asthmatics: Role of Autonomic Nervous System. American Review of Respiratory Disease, 95: 873-874.
138. Sohn, Richard J., Aleksander A. Mathé, and Crystal A. Leslie (1977) Guinea Pig Anaphylaxis: Time Course of Changes in Cyclic AMP and Cyclic GMP. Life Sciences, 21: 1365-1372.
139. Spector, Sheldon L., and Richard S. Farr (1975) A Comparison of Methacholine and Histamine Inhalations in Asthmatics. Journal of Allergy and Clinical Immunology, 56: 308-316.
140. Spector, Sheldon, Thomas J. Luparello, Michael T. Kopetzky, Joseph Souhrada, and Robert A. Kinsman (1976) Response of Asthmatics to Methacholine and Suggestion. American Review of Respiratory Disease, 113: 43-50.
141. Stanescu, Dan C., and Lucien A. Brasseur (1973) Maximal Expiratory Flow Rates and Airway Resistance Following Histamine Aerosols in Asthmatics, Scandinavian Journal of Respiratory Diseases, 54: 333-340.

142. Szentivanyi, Andor (1968) The Beta Adrenergic Theory of the Atopic Abnormality in Bronchial Asthma. Journal of Allergy, 42: 203-232.
143. Taylor, Donald C., Peter R. Cresswell, and David C. Bartlett (1978) The Metabolism and Elimination of Cimetidine, A Histamine H₂-Receptor Antagonist, in the Rat, Dog, and Man. Drug Metabolism and Disposition, 6: 21-30.
144. Taylor, Kenneth M. (1973) Comparison of the Anti-histaminic Effect of Burimamide and Mepyramine and Their Effect on the Activity of Histamine Methyltransferase. Biochemical Pharmacology, 22: 2775-2776.
145. Tucker, Alan, Eric A. Hoffman, and John T. Reeves (1977) Vascular Actions of Histamine H₁- and H₂- Receptor Agonists in Dogs and Cats. European Journal of Pharmacology, 45: 73-77.
146. Tucker, Alan, Eric A. Hoffman, and E. Kenneth Weir (1977) Histamine Receptor Antagonism Does Not Inhibit Hypoxic Pulmonary Vasoconstriction in Dogs. Chest, 71: 261-262.
147. Vaz, Nelson M., Celso M. deSouza, Margaret M. Hornbrook, Donald G. Hanson, and Neil R. Lynch (1977) Sensitivity to Intravenous Injections of Histamine and Serotonin In Inbred Mouse Strains. International Archives of Allergy and Applied Immunology, 53: 545-554.
148. Virgulto, James, and Arend Bouhuys (1973) Electronic Circuits for Recording of Maximum Expiratory Flow-Volume (MEFV) Curves. Journal of Applied Physiology, 35: 145-147.
149. Walkenstein, Sidney S., Jeffrey W. Dubb, Warren C. Randolph, Wilfred J. Westlake, Robert M. Stote, and Alfred P. Intoccia (1978) Bioavailability of Cimetidine in Man. Gastroenterology, 74: 360-365.

150. Woenne, Roland, Meyer Kattan, Robert P. Orange, and Henry Levison (1978) Bronchial Hyperreactivity to Histamine and Methacholine in Asthmatic Children After Inhalation of SCH 1000 and Chlorpheniramine Maleate. Journal of Allergy and Clinical Immunology, 62: 119-124.
151. Wolfe, James D., Marshall Plaut, Philip S. Norman, and Lawrence M. Lichtenstein (1979) The Effect of an H₂ Receptor Antagonist on Immediate and Delayed Skin Test Reactivity in Man (abstract). Journal of Allergy and Clinical Immunology, 63: 208.
152. Woods, James R., Jr., C.R. Brinkman III, A. Dandavino, K. Murayama, and N.S. Assali (1977) Action of Histamine and H₁ and H₂ Blockers on the Cardiopulmonary Circulation. American Journal of Physiology, 232: H73-H78.
153. Yamatake, Yoshikazu, Sumiko Sasakawa, Toshimitsu Yanagiura, and Noboru Kobayashi (1977) Pharmacological Research on the Respiratory Tract-- Report 13: The Role of Histamine H₁ and H₂ Receptors in Asthma Attacks in Dogs (abstract). Folia Pharmacologica Japonica, 73: 125P.
154. Yamatake, Yoshikazu, Sumiko Sasagawa, Saizo Yanaura, and Noboru Kobayashi (1977) Involvement of Histamine H₁- and H₂-Receptors in Induced Asthmas in Dogs. Japanese Journal of Pharmacology, 27: 791-797.
155. Yen, Shyue-Shong (1978) Histaminergic and Cholinergic Receptors in Guinea Pig Lung (abstract). Federation Proceedings, 37: 689.
156. Yen, S.-S., A.A. Mathe, and J.J. Dugan (1976) Release of Prostaglandins From Healthy and Sensitized Guinea-Pig Lung and Trachea by Histamine. Prostaglandins, 11: 227-239.

ACKNOWLEDGEMENTS

I wish to acknowledge the assistance and cooperation of the following persons:

Mr. and Mrs. Morton Brown, my parents, for their continued support throughout medical school and especially during the writing of this thesis. Thanks for everything.

Dr. E. Neil Schachter, my thesis advisor, and good friend, for all his guidance, enormous patience, instruction, suggestions, continued encouragement, and interest. My sincerest thanks for an exciting learning experience.

Dr. Brett Gerstenhaber and Dr. Marjorie Lee, for their technical assistance.

Dr. Gerald Beck, for his advice on statistics.

Dr. Philip Askenase for his comments and suggestions, as well as careful review of the first draft of this thesis.

The rest of my family, Liz, Linda, and Bill for their continued encouragement.

Jennifer Abod, Dr. Demento, Dave Dresner, Kate Eglee, Nora Eisenstat, Ben Engel, Carmel Garguilo, Dave Gendelman, Gwyneth Irwin, Eli Lach, Don Regula, and many others, without whom, this thesis would have been completed many weeks earlier. Thanks for keeping things interesting.

YALE MEDICAL LIBRARY

Manuscript Theses

Unpublished theses submitted for the Master's and Doctor's degrees and deposited in the Yale Medical Library are to be used only with due regard to the rights of the authors. Bibliographical references may be noted, but passages must not be copied without permission of the authors, and without proper credit being given in subsequent written or published work.

This thesis by _____ has been used by the following persons, whose signatures attest their acceptance of the above restrictions.

NAME AND ADDRESS

DATE

