

10-2010

Dorsal Eye Selector Pannier (pnr) Suppresses the Eye Fate to Define Dorsal Margin of the Drosophila Eye


Sarah M. Oros
University of Dayton

Meghana Tare
University of Dayton

Madhuri Kango-Singh
University of Dayton, mkangosingh1@udayton.edu

Amit Singh
University of Dayton, asingh1@udayton.edu

Follow this and additional works at: https://ecommons.udayton.edu/bio_fac_pub

 Part of the [Biology Commons](#), [Computational Biology Commons](#), [Genetics Commons](#), [Genomics Commons](#), [Molecular Genetics Commons](#), [Organismal Biological Physiology Commons](#), and the [Other Genetics and Genomics Commons](#)

eCommons Citation

Oros, Sarah M.; Tare, Meghana; Kango-Singh, Madhuri; and Singh, Amit, "Dorsal Eye Selector Pannier (pnr) Suppresses the Eye Fate to Define Dorsal Margin of the Drosophila Eye" (2010). *Biology Faculty Publications*. 118.
https://ecommons.udayton.edu/bio_fac_pub/118

This Article is brought to you for free and open access by the Department of Biology at eCommons. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of eCommons. For more information, please contact frice1@udayton.edu, mschlange1@udayton.edu.

Published in final edited form as:

Dev Biol. 2010 October 15; 346(2): 258–271. doi:10.1016/j.ydbio.2010.07.030.

Dorsal eye selector *pannier* (*pnr*) suppresses the eye fate to define dorsal margin of the *Drosophila* eye

Sarah M. Oros^{1,2,*}, Meghana Tare^{2,*}, Madhuri Kango-Singh^{1,2,3}, and Amit Singh^{1,2,3,4}

¹ Premedical Programs, University of Dayton, Dayton, OH 45469

² Department of Biology, University of Dayton, Dayton, OH 45469

³ Center for Tissue Regeneration and Engineering at Dayton (TREND), University of Dayton, Dayton, OH 45469

Abstract

Axial patterning is crucial for organogenesis. During *Drosophila* eye development, dorso-ventral (DV) axis determination is the first lineage restriction event. The eye primordium begins with a default ventral fate, on which the dorsal eye fate is established by expression of the GATA-1 transcription factor *pannier* (*pnr*). Earlier, it was suggested that loss of *pnr* function induces enlargement in the dorsal eye due to ectopic equator formation. Interestingly, we found that in addition to regulating DV patterning, *pnr* suppresses the eye fate by downregulating the core retinal determination genes *eyes absent* (*eya*), *sine oculis* (*so*) and *dacshund* (*dac*) to define the dorsal eye margin. We found that *pnr* acts downstream of Ey and affect the retinal determination pathway by suppressing *eya*. Further analysis of the “eye suppression” function of *pnr* revealed that this function is likely mediated through suppression of the homeotic gene *teashirt* (*tsh*) and is independent of *homothorax* (*hth*), a negative regulator of eye. Pnr expression is restricted to the peripodial membrane on the dorsal eye margin, which gives rise to head structures around the eye, and *pnr* is not expressed in the eye disc proper that forms the retina. Thus, *pnr* has dual function, during early developmental stages *pnr* is involved in axial patterning whereas later it promotes the head specific fate. These studies will help in understanding the developmental regulation of boundary formation of the eye field on the dorsal eye margin.

Keywords

Drosophila eye; Dorso-ventral eye patterning; *pannier*; GATA-1; retinal determination; *homothorax*; *teashirt*; Wg Signaling

INTRODUCTION

Axial patterning is required for the transition of a single sheet of cells into a three-dimensional organ. Axial patterning involves generation of two different cell populations or compartments. Compartments are the fundamental units of patterning that are generated by localized expression of transcription factors called selectors. The selectors, when expressed

⁴Corresponding Author: Amit Singh, amit.singh@notes.udayton.edu, Tel: 937-229-2894, Fax: 937-229-2120.

*Equal contribution

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

in a group of cells, can confer compartment-specific properties to these cells (Curtiss et al., 2002; Mann and Carroll, 2002). Signaling between the cells of two compartments is crucial for the patterning, growth and differentiation of a developing field (Blair, 2001). The developing eye of the fruit fly, *Drosophila melanogaster*, has been extensively used to study patterning and growth. The compound eye of the adult fly develops from an epithelial bilayer called the eye-antennal imaginal disc that is derived from the embryonic ectoderm (reviewed by Cohen, 1993, Held, 2002). The imaginal disc is a sac-like structure present inside the larva, which is the product of two different layers: the **peripodial membrane (PM)** and the **disc proper (DP)**. The *Drosophila* retina develops from the DP while the PM of the eye-antennal imaginal disc contributes to the adult head structures (Milner et al., 1983; Haynie and Bryant, 1986; Atkins and Mardon, 2009). Morphogenesis during animal development involves signaling between different layers of the tissue (Furuta and Hogan, 1998; Obara-Ishihara et al., 1999). Similarly, signaling between the PM and the DP of the eye disc is essential for **Dorso-ventral (DV)** axis establishment and patterning (Cho et al., 2000; Gibson and Schubiger, 2000; Atkins and Mardon, 2009).

The adult eye is a highly precise hexagonal array of ~800 ommatidial clusters or unit eyes (Ready et al., 1976; Wolff and Ready, 1993). The ommatidia are arranged in two chiral forms, which are in a mirror image asymmetry along the DV midline called the equator. The equator demarcates the boundary between the dorsal and the ventral eye, and is the site for upregulation of Notch (N) signaling, which triggers cell proliferation and differentiation (Cho and Choi, 1998; Dominguez and de Celis, 1998; Papayannopoulos et al., 1998; Singh et al., 2005b). Although the mirror image asymmetry is generated during the third instar stage of larval eye development, the subdivision of the eye into dorsal and ventral compartments takes place even earlier by domain specific expression and function of DV patterning genes (Cho and Choi, 1998; Dominguez and de Celis, 1998; Papayannopoulos et al., 1998; Cavodeassi et al., 1999; Maurel-Zaffran and Treisman, 2000; Singh et al., 2005b). In antenna, wing and leg imaginal discs, DV boundary formation takes place after the Antero-Posterior (AP) lineage restriction is generated (Blair, 2001; Garcia-Bellido and Santamaria, 1972; Milan and Cohen, 2003; Morata and Lawrence, 1975; Tabata et al., 1995). However, in the eye disc, the AP pattern is established dynamically in the third larval instar stage (after DV lineage is established) when the morphogenetic furrow (MF) is initiated (Singh et al., 2005b). The MF is a wave of retinal differentiation, which progresses anteriorly resulting in the transformation of undifferentiated retinal precursor cells (anterior to MF) into differentiated photoreceptor neurons (posterior to MF) of the eye (Ready et al., 1976; Wolff and Ready, 1993; Heberlein and Moses, 1995; Lee and Treisman, 2001). Therefore, the DV lineage, which is established at late first instar or early second instar larval stages, is the first lineage restriction event in the eye, and is crucial for the growth and differentiation of the eye (Singh and Choi, 2003; Singh et al., 2005b).

During genesis of eye, the entire early eye imaginal primordium initiates from the default ventral fate, which depends on the functions of *Lobe (L)* and *Serrate (Ser)* genes (Chern and Choi, 2002; Singh and Choi, 2003; Singh et al., 2005a, b; 2006). The onset of expression of the dorsal selector gene *pnr*, a member of the GATA-1 family of transcription factors, at the dorsal margin of early second instar larval eye discs establishes the DV lineage in the eye (Maurel-Zaffran and Treisman, 2000; Singh and Choi, 2003; Singh et al., 2005a). It has been shown that *pnr* (Maurel-Zaffran and Treisman, 2000) and members of the *Iro-C* homeodomain genes *viz.*, *araucan (ara)*, *caupolican (caup)* (Cavodeassi et al., 1999;) and *mirror (mirr)* (Kehl et al., 1998; McNeill et al., 1997) are expressed in the dorsal region of the prospective eye (Dominguez and de Celis, 1998; McNeil et al., 1997) and act as the dorsal eye fate selectors. *pnr*, the most upstream gene known in the dorsal eye gene hierarchy, regulates the expression of downstream *Iro-C* genes through Wingless (Wg) signaling (Heberlein et al., 1998; Maurel-Zaffran and Treisman, 2000). Wg, which encodes

a secreted protein, is expressed along the antero-lateral margins of the third instar eye-antennal imaginal disc (Baker, 1988), and prevents ectopic initiation of retinal differentiation from these positions (Ma and Moses, 1995; Treisman and Rubin, 1995). Wg signaling promotes the growth of cells in the eye-antennal disc and is sufficient to maintain cells in an undifferentiated state such that these cells continue to express anterior head specific markers (Lee and Treisman, 2001). In the dorsal eye, Wg promotes expression of *Iro-C* genes during early eye development. The dorsal eye genes and the genes involved in ventral eye development act antagonistically to each other (Singh et al., 2005a, b). These genetic interactions define a signaling pathway that contributes toward the positioning of the equator (Cho and Choi, 1998; Dominguez and de Celis, 1998; Papayannopoulos et al., 1998; Maurel-Zaffran and Treisman, 2000). Thus, *pnr* is known to specify dorsal eye fate. However, the role of *pnr* during retinal differentiation of the eye is not known. Thus, it is important to discern the role of DV patterning gene, *pnr*, during later stages of eye development.

Interestingly, loss-of-function of DV patterning genes manifest defects in the eye growth and patterning but the mechanism by which the DV patterning genes contribute to retinal determination is unknown. It is known that eye specification, and determination depends on a core of retinal determination (hereafter, **RD**) genes. These RD genes include PAX-6 homolog *eyeless* (*ey*), *twin of eyeless* (*toy*), *eyes absent* (*eya*), *sine oculis* (*so*) *dachshund* (*dac*), *optix* (*opt*), and *eyegone* (*eyg*) (Pappu & Mardon, 2004; Dominguez and Casares, 2005; Kumar, 2009). *Ey* is one of the early expressed gene which is required for eye field specification and is reported to induce expression of *eya* and *so* to promote the eye growth and specification (for review Pappu and Mardon, 2004; Silver and Rebay, 2005; Kumar, 2009). The multiple feedback and cross regulatory interactions among RD genes lead to formation of the eye. Loss-of-function of RD genes results in the loss of eye field whereas ectopic expression of RD genes results in the induction of ectopic eyes (Halder et al., 1995; Pappu & Mardon, 2004; Silver and Rebay, 2005; Kumar, 2009). Several genes other than RD genes contribute towards eye development. A homeotic gene, *tsh*, which encodes a C₂H₂ Zinc finger transcription factor with three widely spaced Zinc finger domains (Fasano et al., 1991), has been suggested to act upstream of *eya*, *so* and *dac* during eye development (Pan and Rubin, 1998; Kumar, 2009). Interestingly, *tsh* also exhibits asymmetric DV response in the eye (Singh et al., 2002). Misexpression of *tsh* suppresses the eye fate in the ventral eye and promotes ectopic dorsal eye enlargement (Singh et al., 2002). It has been shown that *tsh* collaborates with the genes that express in a domain specific manner to exhibit DV asymmetric response in the developing eye disc (Singh et al., 2004). Interestingly, in the dorsal eye, the gain-of-function phenotype of *teashirt* (*tsh*) is similar to the loss-of-function phenotype of *pnr*. However, the mechanism of their interaction during eye development is not fully understood. In the ventral eye, *tsh* suppresses the eye by induction of a Meis class of homeotic gene, *homothorax* (*hth*) (Reichkof et al., 1997; Singh et al., 2002). *Tsh* has been shown to physically bind *Hth* anterior to the eye field (Bessa et al., 2002). *Hth*, is known to act as a negative regulator of eye development. Loss-of-function of *hth* results in induction of the ventral eye or enlargement of the ventral eye domain (Pai et al., 1998). However, loss-of-function of *hth* in the dorsal eye has no effect even though *hth* is expressed in the dorsal eye (Pichaud & Casares, 2000; Jaw et al., 2000). Further, *tsh* does not affect *hth* expression in the dorsal eye whereas *tsh* acts upstream of *hth* in the ventral eye (Singh et al., 2002). Therefore, the mechanism of *hth* regulation in the dorsal eye remains unknown. Interestingly, the mechanism of genetic regulation of dorsal eye field growth is not very clear.

pnr plays an important role in the dorsal eye development. Onset of *pnr* expression in the dorsal eye margin is associated with DV lineage restriction (Maurel-Zaffran and Treisman, 2000; Cavodeassi et al., 2000; Singh et al., 2005b). The loss-of-function phenotypes of *pnr*

result in ectopic eye enlargement of the dorsal eye. Here we report that in addition to its earlier reported role of dorsal selector during axial (DV) patterning, *pnr* plays an important role in defining the dorsal eye margin by regulating retinal determination. We have found that gain of function of *pnr* suppresses the retinal determination whereas the loss of *pnr* results in the ectopic induction of retinal determination genes. Our data suggests that *pnr* suppresses the retinal determination by downregulation of homeotic gene *tsh*, and is independent of *hth*. Interestingly, *pnr* is expressed only in the peripodial membrane and not in the disc proper, which gives rise to the retina. Thus, a late function of *pnr* is to block retinal determination in the peripodial membrane to define the dorsal eye margin.

MATERIALS AND METHODS

Stocks

Fly stocks used in this study are described in Flybase (<http://flybase.bio.indiana.edu>). We used *y, w, eyFLP* (Newsome et al., 2000), *y, w; FRT82B pnr^{vx6}/CyO*, (Heitzler et al. 1996), *UAS-pnr^{D4}* (Haenlin et al. 1997), *UAS-pnr^{ENR}* (Klinedinst and Bodmer 2003), *FRT82D hth^{P2}* (Noro et al., 2006), *UAS-ara* (Gomez-Skarmeta et al. 1996), *UAS-hth¹²* (Pai et al., 1998), *UAS-hth-ENR* (Inbal et al., 2001), *y, w; tsh⁸/CyO* (Fasano et al., 1991); *UAS-tsh* (Gallet et al., 1998), *UAS-ds tsh* (Bessa and Casares, 2005), *y w; tsh^{A8}* (Sun et al., 1995); *y,w; UAS-NLS-GFP^{S65T}* (Ito et al., 1997) *UAS-wg* (Azpiazu and Morata, 1998), *pnr-Gal4*, *UAS-GFP* (Singh et al., 2005a), *ey-Gal4* (Hazelett et al., 1998), *bi-Gal4* (Calleja et al., 1996). We have used Gal4/UAS system for targeted misexpression studies (Brand and Perrimon, 1993). All Gal4/UAS crosses were done at 18°C, 25°C and 29°C, unless specified, to sample different induction levels.

Genetic Mosaic Analysis

Loss-of-function clones were generated using the FLP/FRT system of mitotic recombination (Xu and Rubin, 1993). To generate loss-of-function clones of *pnr* in the eye, *eyFLP*; *FRT82B Ubi-GFP* females were crossed to *y, w; FRT82B pnr^{vx6}* males. Gain-of-function clones of *pnr* were generated using *hs-FLP* method where *y, w, hsFLP¹²²; P(Act>y⁺>Gal4) 25 P(UAS-GFP^{S65T})/CyO* (Struhl and Basler, 1993) flies were crossed to *UAS-pnr^{D4}* flies.

Immunohistochemistry

Eye-antennal imaginal discs were dissected from wandering third instar larvae and stained following the standard protocol (Singh et al., 2002). Antibodies used were mouse and rabbit anti-β galactosidase (1:200) (Cappel); chicken anti-GFP (1:200) (Upstate biotechnology), rat anti-Elav (1:100); mouse anti-WG (1:50) (Developmental Studies Hybridoma Bank); rabbit anti-Dlg (a gift from K. Cho), rabbit anti-Ey (a gift from Uwe Walldorf and Patrick Callaerts), anti-Hth (a gift from H. Sun and R. Mann) mouse anti-So (1:100), mouse anti-Dac (1:100), mouse anti-Eya (1:100) (Developmental Studies Hybridoma Bank), Rat anti-Tsh (1:50) (Gallet et al., 1998). Secondary antibodies (Jackson Laboratories) were goat anti-rat IgG conjugated with Cy5 (1:200), donkey anti-rabbit IgG conjugated to Cy3 (1:250), donkey anti-rabbit IgG conjugated to FITC, donkey anti-mouse IgG conjugated to Cy3 (1:200). Pnr expression was detected using *pnr-Gal4> UAS-GFP* (Pichaud and Casares, 2000; Singh and Choi, 2003; Singh et al., 2005a). Immunofluorescent images were analyzed using the Olympus Fluoview 1000 Laser Scanning Confocal Microscope.

Scanning Electron Microscopy (SEM)

The flies were prepared for scanning electron microscopy by dehydration through a series of increasing concentrations of acetone. Dehydrated flies were then stored in 1:1 mixture of acetone and Hexa Methyl Di Silazane (HMDS, Electron Microscopy Sciences), and then

stored in 100% HMDS. The flies were allowed to air dry in HMDS. Dehydrated flies were mounted on a carbon conductive tape on EM stubs. Fly samples were coated with gold using a Denton vacuum sputter coater and analyzed using a Hitachi S-4800 High Resolution Scanning Electron Microscope (HRSEM).

RESULTS

Pnr expression is restricted to the peripodial membrane (PM) of the dorsal eye margin

In the *Drosophila* embryo, *pnr* is expressed in the dorsal most embryonic cells in a domain of presumptive notum surrounding the dorsal midline, and at the dorsal anterior margin of the eye disc (Heitzler et al., 1996; Ramain et al., 1993; Maurel-Zaffran and Treisman, 2000). Pnr is not expressed in the first instar larval eye-antennal imaginal disc (Fig. 1A). In the eye-antennal disc, *pnr* expression begins either in the late first instar stage or in the early second instar stage (Singh and Choi, 2003). In the early second instar eye-antennal imaginal disc, *pnr* expression begins in 5–7 cells in the dorsal margin of antenna and head region of the eye-antennal disc (Fig. 1A). During mid- to late- second instar of larval eye development, *pnr* begins to express in 30–35 cells (Fig. 1B). During the late second instar stage of larval eye development, *pnr* expression spreads to 80–100 cells in the dorsal eye domain (Fig. 1C). Interestingly, *pnr* expression is not seen in any disc proper cells, which differentiate to retinal photoreceptor cells in the eye (Fig. 1C'). Furthermore, *pnr* expression is restricted to the peripodial membrane on the dorsal eye margin (Pereira et al., 2006). Rarely a few disc proper cells (anterior to the morphogenetic furrow) at the border with peripodial membrane in the eye disc show *pnr* expression (Fig. 1C''). As the larva progresses into its third instar stage, *pnr* extends throughout the dorsal-most region of the head and antenna. The *pnr* expression in the third instar eye-antennal imaginal evolves into 4–5 rows of cells on the dorsal eye margin (Fig. 1D). Pnr expression overlaps with the MF but does not coincide with retinal cells as it is expressed only in the peripodial membrane.

Wg expression, which acts downstream to Pnr, is localized to the dorsal as well as the ventral eye margins in the disc proper cells (Fig. 1B–E). Wg is also expressed throughout all larval stages in the PM (Cho et al., 2000). Wg is expressed in both the dorsal and ventral compartments, but expression in the dorsal is constant throughout all larval stages in the peripodial membrane. Wg is controlled by *pnr* in the peripodial membrane only; its disc proper expression is not influenced by any identified gene as of yet. In the first instar and early second instar eye disc, *hth* is expressed in the entire eye disc (Fig. 1B; Singh et al., 2002; Bessa et al., 2002). The expression of *hth* in the disc proper (DP) begins to retract in late second instar (Fig. 1C') whereas *hth* is expressed in the entire peripodial membrane (PM). In late third instar stage, *hth* expression stays anterior to the MF in the late third instar stage (Fig. 1D).

Loss-of-function clones of *pnr* show four different phenotypes

We employed the genetic mosaic approach to generate loss-of-function clones of *pnr* in the developing eye-antennal imaginal disc (Xu and Rubin, 1993). We used the *pnr^{vxb}* mutant, a null allele which has a deletion of all but 9 amino acids of the coding region (Ramain et al., 1993; Heitzler et al., 1996), to generate the genetic mosaic clones. Mutant clones were generated in the eye using the FLP/FRT system where Flippase is under the control of an eye-specific enhancer of *eyeless*, (*ey*), (Quiring et al., 1994). Since *pnr* is expressed in the dorsal eye, the loss-of-function clones of *pnr* that are located only in the dorsal eye margin exhibit phenotypes. The loss-of-function clonal phenotypes of *pnr* can be classified into four different categories:

(I) Non-autonomous dorsal eye enlargement—Loss-of-function clones of *pnr* in the dorsal eye result in an ectopic eye field or enlargement of the existing eye field comprising of differentiating photoreceptor neurons. These eye field enlargements or ectopic eye fields can even extend anterior to the furrow only on the dorsal eye margin (Fig. 2A; marked by white dotted line). Further, these eye enlargement phenotypes are non-autonomous, which include both mutant cells (lack GFP reporter, marked by dotted boundary in Fig. 2A) as well as the adjoining wild-type cells (GFP positive). In the adult eyes, the *pnr* clones were marked by absence of the mini-*white* reporter gene, which is involved in the pigment uptake in the eye (Sun et al., 1995). These clones resulted in either enlargement of the pre-existing dorsal eye field or generation of a *de novo* ectopic eye field in the dorsal head cuticle (Fig. 2B; marked by black dotted line). These ectopic eye fields did not arise exclusively within the *pnr* mutant clones, but also contained a domain of the wild-type cells (marked by dark red pigment). The phenotype of these clones resembled the loss-of-function clone phenotypes of *pnr* described earlier (Maurel-Zaffran and Treisman, 2000; Singh and Choi, 2003). These phenotypes were explained to be due to generation of a *de novo* equator between the *pnr*⁻ and *pnr*⁺ cells. The frequency of these clones is around ~8.3% of the total *pnr* loss-of-function clones (Table 1). Strikingly, we observe only the bigger size clones in this category.

(II) Autonomous dorsal eye enlargement—Loss-of-function clones of *pnr* in this category showed an ectopic field of differentiating photoreceptors anterior to the morphogenetic furrow in the dorsal eye domain. Unlike the clones of previous category, the ectopic eye field in these clones was autonomous (restricted within the *pnr* loss-of-function clones) (Fig. 2C, clonal boundary marked by white dotted line). In the adult flies, these clones resulted in the formation of an ectopic eye field in the dorsal head cuticle anterior to the eye field (Fig. 2D, black dotted line). The ectopic eye field in the clones of this category was devoid of any wild-type *pnr*⁺ cells, clearly suggesting that *pnr* loss-of-function led to the generation of ectopic eyes. The frequency of these clones was nearly 12.6%, which comprises of both the smaller (7.0%) as well as the bigger (5.6%) clones (Table 1). Some of these clones were accompanied by cuticle enlargement in the head.

(III) Absence of dorsal eye enlargement—Unlike the previous two categories of *pnr* clones (Fig. 2A–D), loss-of-function clones in this category does not result in any ectopic dorsal eye enlargement (Fig. 2E). Interestingly, even though these clones span both the anterior as well as the posterior regions of the morphogenetic furrow in the dorsal eye-antennal imaginal disc but did not result in any ectopic eyes in the eye disc (Fig. 2E) as well as in adult flies (Fig. 2F). The frequency of these clones is nearly 75.3% (Table 1). We found that these clones unlike the clones from the previous two categories, are restricted only to the disc proper in the dorsal eye; a domain where *pnr* is normally not expressed (Fig. 1).

(IV) Antennal duplication—Loss-of-function clones of *pnr* in the antenna region of the eye-antennal imaginal disc, results in the duplication of antennal field (Fig. 2G). Interestingly, most of these antennal duplications were accompanied with ectopic eye enlargements (Fig. 2H). However, in some of these clones only antennal duplication were observed. These clones led to duplication of the ventral head structures such as antenna and maxillary palps in the dorsal head (Fig. 2H; Maurel-Zaffran and Treisman, 2000; Pichaud and Casares, 2000; Singh et al., 2005b). The frequency of these clones was ~3.6% (Table 1). Thus, our analysis of loss-of-function clones suggests that *pnr* may be involved in the suppression of eye fate.

Pnr suppresses the eye fate in the dorsal eye

To test the role of *pnr* in eye fate determination, we used the target system of Gal4/UAS to misexpress *pnr* in the eye (Brand and Perrimon, 1993). We used a *UAS-pnr^{D4}* construct that behaves like wild-type *pnr* in the absence of *U-shaped (ush)* function (Haenlin et al., 1997; Maurel-Zaffran and Treisman, 2000; Fossett et al., 2001; Singh & Choi, 2003). *Ush* encodes a zinc finger protein that dimerizes with Pnr and acts as a negative regulator of *pnr* transcriptional activity (Haenlin et al., 1997). Since *ush* is not expressed in the eye-antennal imaginal disc, the *UAS-pnr^{D4}* construct behaves in a wild-type fashion in the eye (Maurel-Zaffran and Treisman, 2000; Fossett et al., 2001). We used an *ey-Gal4* driver that drives the expression of UAS-GFP transgene in the entire eye disc, which comprises of differentiating retinal neurons (posterior to MF, marked by white arrowhead) as well as the region forming the prospective head cuticle (anterior to MF) (Fig. 3A; Singh et al., 2005a). Misexpression of *pnr* in the entire eye disc using *ey-Gal4 (ey>pnr^{D4})* results in the complete loss of eye field as evident from the absence of neuronal marker *Elav* whereas the size of antennal field is not affected (Fig. 3C). The misexpression of *pnr* in the entire eye (*ey>pnr^{D4}*) results in the adult flies with highly reduced eye field or what we refer to as the “no-eye” phenotypes (Fig. 3D) as compared to the wild-type eyes (Fig. 3B). To test if there is any domain specific response of *pnr* misexpression, we employed *bi-Gal4* driver, which drives the expression of UAS- GFP transgene (*bi>GFP*) on both dorsal and ventral margins of the developing eye-antennal imaginal disc (Fig 3E; Calleja et al., 1996; Singh et al., 2002; 2004). Misexpression of *pnr* using *bi-Gal4 (bi>pnr^{D4})* suppressed the eye fate on both dorsal and ventral eye margins as evident from the absence of *Elav* expression (Fig. 3F; white arrows). This suggests that *pnr* upon misexpression suppresses the eye fate, irrespective of dorsal or ventral domains. Random gain-of-function clones of *pnr* in the eye using *UAS-pnr^{D4}* (marked by GFP reporter) caused suppression of photoreceptors as evident from the absence of *Elav* positive cells in the eye disc (Fig. 3G, G') as well as in the adult eye (Fig. 3H). In addition to the small eye phenotypes seen in the gain-of-function clone of *pnr*, we observed necrosis as evident from presence of dark spots in the adult eye (Fig. 3H). These results suggest *pnr* can suppress the eye fate. The frequency of *pnr* gain-of-function clones was extremely low in the eye disc as well as the adults probably due to issues with cell survival.

We then disrupted Pnr function using dominant negative *pnr (UAS-pnr^{ENR}})* (Fu et al., 1998; Klinedinst and Bodmer, 2003) where the construct contains the repressor domain from the Engrailed transcription factor (EnR, amino acid 2-298) (Jaynes and O'Farrell, 1991) and the two N-terminal zinc-finger domains from *pnr* (amino acids 153-293) (Ramain et al., 1993). Disrupting *pnr* function in the entire eye-antennal imaginal disc all along during eye development by misexpression of *pnr^{ENR} (ey> pnr^{ENR}})* resulted in a small group of *Elav* positive retinal cells in the eye field (Fig. 3G), and a highly reduced eye in the adult fly (Fig. 3H). However, there was no affect on the developing antennal field (Fig. 3G, H). This suggests that during early eye development when DV patterning is being established *pnr* function is crucial for eye development. However, misexpression of *pnr^{ENR}}* on both dorsal and ventral eye margins using *bi-Gal4 (bi>pnr^{ENR}})* caused enlargement of only the dorsal eye (Fig. 3H, arrow). In *bi>pnr^{ENR}}* background, *pnr* function was abolished only in a few subset of cells within the endogenous *pnr* expression domain in the peripodial membrane of the dorsal eye. These dorsal eye enlargement phenotypes further substantiated the possibility that *pnr* may suppress retinal determination in the dorsal eye margin.

Pnr downregulates the retinal determination genes to suppress the eye

To address the role of *pnr* in retinal determination, we checked the expression of members of the **retinal determination (RD)** gene pathway in the loss-of-function clones of *pnr* in eye-antennal imaginal disc. The loss-of-function clones of *pnr* (marked by absence of the GFP reporter) that exhibit enlargement of the dorsal eye showed no ectopic induction of *Ey* (Fig.

4A, A', A''). It has been shown that Ey is expressed in undifferentiated retinal precursor cells early in eye development and after the onset of photoreceptor differentiation; Ey continues to express in the undifferentiated retinal precursor cells anterior to the MF (Quiring et al., 1994; Lee and Treisman, 2001; Singh et al., 2002; Bessa et al., 2002). Although Ey expression was not affected, the loss-of-function clones of *pnr* which caused ectopic dorsal eye enlargements showed ectopic induction of Eya (Fig. 4B, B', B''), which acts downstream to Ey. The loss-of-function clones of *pnr* in the ventral domain of the eye did not exhibit any effect on wild-type Eya expression. During the late first instar stage, *eya* begins expression in the eye region of the disc. A short time after *eya* expression, *so*, *dac*, and *eyg* are expressed in the late second instar stage in the region posterior to the MF (Bonini et al. 1993; Cheyette et al. 1994; Jang et al. 2003; Kenyon et al. 2003; Mardon et al., 1994). After the MF begins expression of *eya* and *so* are restricted to the area within and posterior to the MF (Bonini et al. 1993; Cheyette et al. 1994). The expression of *dac* is restricted to the MF in the area that directly precedes the MF and continues in R1, R6 and R7 for a few columns posterior to the MF and sharply disappears after that domain (Mardon et al., 1994; Tavsanlı et al. 2004). The loss-of-function clones of *pnr* in the dorsal eye showed ectopic induction of So (Fig. 4C, C', C''; white arrows) and Dac (Fig. 4D, D', D''; white arrows). Dac is expressed downstream to *eya* and *so* (Chen et al., 1997). The ventral eye clones did not exhibit any effect on the expression of RD genes. These results suggest that *ey*, a gene expressed in undifferentiated cells, is not induced in ectopic dorsal eye enlargement whereas the downstream RD genes like *eya*, *so*, *dac* that are expressed in differentiating photoreceptor neurons are upregulated in the *pnr* loss-of-function clones. The *pnr* loss-of-function clones which only result in antennal duplications does not affect RD genes expression (data not shown). These results suggest that *pnr* may act downstream of *ey* to suppress retinal determination on the dorsal eye margin. Furthermore, since *pnr* is expressed in the peripodial membrane on the dorsal eye margin, these loss-of-function clonal phenotypes of *pnr* suggests that *pnr* blocks retinal determination in the peripodial membrane of the dorsal eye margin. Thus, *pnr* may promote head specific fate by blocking retinal determination in the peripodial membrane.

We tested this hypothesis by checking RD gene expression in *bi>pnr^{D4}* background where *pnr* misexpression suppresses the eye both on dorsal and ventral eye margins (Fig. 3F). Interestingly, Ey was present on both dorsal and ventral margins (Fig. 4E, E'; white arrowheads). However, the expression of other RD genes like Eya (Fig. 4F, F'; white arrowheads), So (Fig. 4G, G'; white arrowheads), and Dac (Fig. 4I, I'; white arrowheads) was down regulated on both the dorsal as well as the ventral eye margins of highly reduced eye-antennal imaginal disc. Dac expression anterior to the MF is not affected (Fig. 4I, I'). These results strongly suggest that *pnr* suppresses retinal determination by blocking Eya, So, and Dac expression.

Pnr suppresses eye by induction of its downstream target Wg

In order to test if, the misexpressed *pnr* is functional in the eye disc; we tested the levels of Wg as a functional read out of *pnr* in the eye. Pnr acts upstream to the signaling molecule Wg, that suppresses the eye fate (Ma and Moses, 1995; Treisman and Rubin, 1995; Lee and Treisman, 2001). Wg is expressed on the antero-lateral margin of both dorsal and ventral eye (Fig. 1D). Misexpression of *pnr* in the eye disc (*ey>pnr^{D4}*) results in the suppression of eye (Fig. 3B). We found that misexpression of *pnr* on both the dorsal and the ventral margin (*bi>pnr^{D4}*) of eye disc results in robust induction of Wg along with a strong suppression of the eye on both the dorsal as well as the ventral margins (Fig. 5A, A'). *bi>pnr^{D4}* showed similar eye suppression phenotypes on both the dorsal and ventral eye margins in the adult eye (Fig. 5B). This phenotype is similar to the misexpression of Wg on both dorsal and ventral margins (*bi>wg*) that results in suppression of the eye on both dorsal as well as

ventral margins (Singh et al., 2002). Targeted misexpression of *pnr* using (*ey>pnr^{D4}*) results in a “no-eye” phenotype by induction of Wg in the entire eye (Fig. 5C). These phenotypes are comparable to the ectopic induction of Wg in the eye (*ey>wg*) (data not shown, Singh et al., 2002). Interestingly, the ectopic eye induction and Wg downregulation did not cover the entire loss-of-function clone of *pnr* in the dorsal eye (Fig. 5D, marked by the white dotted line). But the Wg expression was present within the clone juxtaposed to the wild-type *wg* expression domain in the head region (Fig. 5D). This Wg expression phenotype can be explained as rescue of Wg within clone from the wild-type cells due to the secretory nature of Wg. The loss-of-function clones of *pnr* in the disc proper (DP) did not result in ectopic eye enlargements and showed no affect on Wg expression (Fig. 5F, F'). In some of the larger loss-of-function clones of *pnr*, which extend from the dorsal eye margin into the antennal field, we observed a surprising result where the entire clone did not show ectopic eye enlargement. The area of the clone, which did not have eye enlargement, showed strong Wg expression even though the expression domain of this Wg was not close to wild-type expression domain of Wg. This result suggests that *pnr* is not the sole regulator of *wg* expression in the dorsal eye. Since *wg* acts in a feedback loop with *hth* in the ventral eye (Pichaud and Casares, 2000; Singh et al., 2002), there is a possibility of the role of *hth* in *wg* regulation in the dorsal eye.

Pnr eye suppression function in the dorsal eye margin is independent of *hth*

In order to understand the mechanism by which *pnr* suppresses the eye fate, we tested whether loss of *pnr* function induces *hth*. Misexpression of *hth* on both dorsal and ventral eye margins (*bi>hth*) results in suppression of the eye on both dorsal and ventral eye margins (Fig. 6A, A', arrows) as also seen in *bi>pnr^{D4}* eye-antennal discs (Fig. 3F). Therefore, we tested levels of Hth in *bi>pnr^{D4}* eye-antennal disc. Interestingly, Hth was induced both on the dorsal as well as the ventral eye margins (Fig. 6B, B', arrows). This raises the possibility that *pnr* may suppress the eye development by inducing downstream *hth*. Therefore, if *hth* is downstream to *pnr* in the dorsal eye, then loss-of-function of *hth* must be similar to *pnr* loss-of-function phenotypes. We generated loss-of-function clones of *hth* in the eye disc. We found that the loss-of-function clones of *hth* induced at any time during larval development autonomously induced ectopic eyes only in the ventral head capsule (Fig. 6C, C'; marked by white dotted line, Pai et al., 1998; Pichaud and Casares, 2000). However, *hth* loss-of-function clones did not show any ectopic dorsal eye enlargements as seen in the *pnr* loss-of-function clones (Fig. 2). Thus, unlike gain-of-function of *hth* that corresponds to the gain-of-function of *pnr* (Fig. 3), the loss-of-function of *hth* does not match *pnr* loss-of-function clonal phenotypes. This result rules out the possibility of *pnr* acting upstream of *hth* in the dorsal eye. We studied the expression of Hth in the *pnr* loss-of-function clones and found that Hth expression was not affected in loss-of-function clones of *pnr* showing ectopic eye enlargements (Fig. 6D, D'). Hth marks the undifferentiated retinal precursor cells anterior to the furrow (Fig. 1; Pai et al., 1998; Bessa et al., 2002). In *pnr* loss-of-function clones, where no ectopic eye enlargements were seen, Hth expression was not affected (Fig. 6E, E'). Interestingly, Hth expression was induced in *pnr* loss-of-function clones where the duplication of antennal region took place (Fig. 6F, F'). The duplication of antennal field represents the ventral structures in head capsule (Casares and Mann, 1998). Since *hth* is expressed in the proximal domains of antennal field there is ectopic induction of *hth* when duplication of the antennal field and cuticle enlargement occurs. Thus, our results suggest that *pnr* does not directly affect the *hth* expression in the dorsal eye.

Pnr suppresses the eye by downregulating *tsh*

We employed a candidate gene approach and looked for the genes which might affect the dorsal eye patterning. Homeotic gene *teashirt* (*tsh*) shows an asymmetric response on the

dorsal and ventral eye margins. In the early eye, *tsh* is expressed in the entire disc (Singh et al., 2002; Bessa et al., 2002). In the second instar eye-antennal imaginal disc, Tsh expression begins to retract anteriorly (Fig. 7A), and in the third instar eye disc Tsh is expressed in the retinal precursor cells anterior to the MF (Fig. 7A, B; Bessa et al., 2002; Singh et al., 2002). We tested expression of *tsh* in the loss-of-function clones of *pnr* in the dorsal eye. We found that *pnr* loss-of-function clones, which exhibit dorsal eye enlargement also exhibit ectopic induction of Tsh (Fig. 7C, C') as well as the *tsh* reporter $y, w; tsh^{A8}$ (Fig. 7C, C"). It suggests that *pnr* might suppress *tsh* expression at the transcription level. In order to test whether *pnr* suppresses the eye by downregulating *tsh*, we generated *pnr* loss-of-function clones where *tsh* levels were reduced to 50% using a heterozygous combination of *tsh⁸*, a null allele of *tsh* (Fasano et al., 1991). Interestingly, in these *pnr* loss-of-function clones where *tsh* function was reduced to half, we did not see any ectopic eye enlargement (Fig. 7D). Interestingly, although the dorsal clones did induce overgrowths/enlargement, these clones did not show any ectopic Elav expression. However, these clones exhibited strong Ey expression (Fig. 7D', D"). The adult eye phenotype of these clones is similar to the eye disc phenotype of lack of any dorsal eye enlargements (Fig. 7E). To further test our hypothesis that *pnr* affects *tsh* expression at the transcription level, we misexpressed *pnr* on the dorsal and ventral eye margins (*bi>pnr^{D4}*) and checked the expression of the *tsh* reporter. We found that *tsh* expression was downregulated on both the dorsal and ventral eye margins (Fig. 7F, arrows). Since endogenous expression of *pnr* is restricted to the dorsal eye margin, our results suggest that *pnr* suppresses the *tsh* expression in the dorsal eye. We further tested *pnr* and *tsh* interaction, by misexpressing *pnr* on both dorsal and ventral eye margins in a *tsh^{8/+}* heterozygous background (*bi>pnr^{D4}; tsh^{8/+}*). In this *tsh* heterozygous background, misexpression of *pnr* strongly enhances the eye suppression phenotype on both the dorsal and the ventral eye margins (Fig. 7G) as compared to *bi>pnr^{D4}* alone (Fig. 7F). These results suggest that *pnr* may suppress *tsh* at the dorsal eye margin. Misexpression of *tsh* on the dorsal and the ventral eye margin (*bi>tsh*) results in dorsal eye enlargement and ventral eye suppression (Fig. 7H; Singh et al., 2002). This phenotype is complementary to the *pnr* loss-of-function phenotype. Misexpression of *tsh* RNAi using *bi*-Gal4 results in complementary to *bi>tsh* phenotype in the dorsal eye (data not shown). We therefore tested whether misexpression of *tsh* can rescue the *pnr* misexpression phenotype. Misexpression of both *pnr* and *tsh* (*bi>tsh+pnr^{D4}*) resulted in the lethality as early as the first instar larval stage. We therefore, misexpressed *ara*, a downstream target of *pnr* and a member of Iro-C complex, with *tsh* on the dorsal and ventral eye margins (*bi>tsh+ara*), which resulted in the enlargement of the eye on the dorsal eye margin (Fig. 7C; Singh et al., 2004). We further tested the hypothesis that *pnr* down regulates *tsh* in the dorsal eye to suppress eye fate. Misexpression of *tsh* in the dorsal eye margin using a *pnr*-Gal4 driver (*pnr>tsh*) resulted in the enlargement of eye on the dorsal margin (Fig. 7J). Lastly, we tested whether reducing *pnr* levels affects the *bi>tsh* phenotype. In heterozygous *pnr* background we misexpressed *tsh* (*bi>tsh; pnr^{xx6/+}*) and found that it results in the dorsal eye enlargements (Fig. 7K). These eye enlargements were similar to that of the *bi>tsh* alone (Fig. 7H). This suggests that *tsh* acts downstream to *pnr* and therefore levels of *tsh* are crucial for the dorsal eye enlargement phenotype. Thus, *pnr* suppresses the eye development on the dorsal eye margin by suppressing *tsh*.

DISCUSSION

We have addressed a basic question pertaining to regulation of patterning, growth and differentiation of the developing eye field. Our results provide an important insight into the role of *pnr*, a gene known to confer dorsal eye identity during axial patterning of the eye (Maurel-Zaffran and Treisman, 2000; Pichaud and Casares, 2000; Singh and Choi, 2003). We and others have shown that the onset of *pnr* expression during early eye development results in the generation of dorsal lineage in the eye. It results in the formation of a DV

boundary (equator), which triggers N signaling at the border of the dorsal and ventral compartments to initiate growth and differentiation (Cho and Choi, 1998; Dominguez and deCelis, 1998, Papayannopoulos et al., 1998; Singh et al., 2005b).

Earlier, we have tested the spatial as well as temporal requirement for the genes controlling ventral eye growth and development (Singh and Choi, 2003). During early eye development, prior to the onset of *pnr* expression in the dorsal eye, entire early eye primordium is ventral in fate (Singh and Choi, 2003). Removal of function of genes controlling ventral eye development prior to the onset of *pnr* expression, result in complete elimination of the eye field whereas later when *pnr* starts expressing, the eye suppression phenotype gets restricted only to the ventral eye (Singh and Choi, 2003; Singh et al., 2005a). These studies suggested that *pnr* plays an important role in dorso-ventral (axial) patterning. However, the role of dorsal selector *pnr* in retinal determination was unknown.

Pnr suppresses the eye fate

Loss-of-function clones of *pnr* in the dorsal eye exhibit eye enlargement (Fig. 2A; Maurel-Zaffran and Treisman, 2000; Pichaud and Casares, 2000; Singh and Choi, 2003). It was suggested that when *pnr* function was abolished in the dorsal eye using loss-of-function clones, it results in the change of dorsal eye fate to ventral. This results in generation of a *de novo* equator, the border between dorsal and ventral half of the eye, which triggers ectopic N signaling to promote growth and cell proliferation. The same premise was used to explain the gain-of-function phenotype of *pnr* in the eye. Misexpression of *pnr* in the entire eye (*ey>pnr^{D4}*) generates a completely dorsalized eye field as *pnr* acts as the dorsal fate selector. The fully dorsalized eye lack DV polarity (equator), which results in the complete loss of eye field due to lack of N upregulation (Fig. 3C, D; Maurel-Zaffran and Treisman, 2000). Here, we addressed another possibility to see if *pnr* suppresses the eye fate upon misexpression in the entire eye as evident from the “no-eye” phenotype.

To test if *pnr* suppresses the eye fate, we misexpressed *pnr* both in the dorsal and ventral (DV) eye margins of the eye using a *bi*-Gal4 driver (*bi>pnr^{D4}*) (Fig. 3F). The rationale was if *pnr* is only required to assign the dorsal eye fate, in that case *pnr* misexpression (*bi>pnr^{D4}*) will assign a dorsal fate on the margin of ventral eye. Thus, by this logic, it would result in generation of a *de novo* equator on the ventral eye margin, which should manifest as eye enlargements in the ventral eye. The argument was based on the similar premise that was employed to explain that loss-of-function clones of *pnr* in the dorsal eye generated a new equator and led to the dorsal eye enlargement. We did not observe any ventral eye enlargements in *bi>pnr^{D4}* eye disc (Fig. 3F). Instead we saw suppression of the eye on both the dorsal as well as ventral margins (Fig. 3F). The suppression of eye fate on both the dorsal and the ventral margins suggests that *pnr*, irrespective of the domain where it is expressed, can suppress the eye fate.

Pnr suppresses the Retinal Determination (RD) genes function

Since *pnr* suppresses the eye fate, it is possible that it may be involved in regulation of expression of genes of the core retinal determination machinery. Loss-of-function of *pnr* in the dorsal eye clones results in the eye enlargements as evident from Elav positive cells but it does not induce ectopic Ey (Fig. 4A). Ey expression evolves during eye development and is localized anterior to the morphogenetic furrow in retinal precursor cells and is downregulated and degraded posterior to the furrow in differentiation retinal neurons (Quiring et al., 1994; Halder et al., 1995; 1998; Baonza and Freeman, 2002; Kango-Singh et al., 2003; Lee and Treisman, 2001). We found that in the loss-of-function clones of *pnr* in the eye, the expression of retinal determination pathway members like Eya, So and Dac, which act downstream to Ey, was ectopically induced (Fig. 4B–D). In the converse situation, where

pnr was misexpressed on both dorsal and ventral eye margins (*bi>pnr^{D4}*), we observed ectopic induction of *Ey* on both the dorsal and the ventral margins (Fig. 4E) whereas the expression of *Eya* (Fig. 4F), *So* (Fig. 4G) and *Dac* (Fig. 4H) were suppressed. Thus, misexpression of *pnr* in the eye prevents the photoreceptor differentiation irrespective of the dorsal or ventral domain. Based on these results we can propose that *pnr* suppresses the eye fate on the dorsal eye margin by downregulating RD genes like *eya*, *so* and *dac* (Fig. 8).

Since *ey* is responsible for the specification of the eye field and marks the retinal precursor cells, it suggests that *pnr* does not affect the eye field formation or specification. In fact *pnr* affects expression of RD genes *eya*, *so* and *dac* (Fig. 4), which acts downstream to *ey*, and are involved in retinal determination (Kango-Singh et al., 2003; Pappu and Mardon, 2004; Silver and Rebay, 2005; Kumar, 2009). Our results suggest that *pnr* suppress retinal determination genes. Since endogenous expression of *pnr* is restricted to the peripodial membrane of the dorsal eye margin (Fig. 1), it suggests that *pnr* may be involved in suppression of retinal determination on the dorsal peripodial membrane. Thus, our results suggest that *pnr* generally acts at the stage when photoreceptor differentiation is initiated with the formation of the morphogenetic furrow (MF) and promotes the dorsal head cuticle fate by suppressing retinal differentiation (Fig. 8).

Dual function of *pnr* during eye development

Based on our new findings and other previously published results, we propose that *pnr* may be required for two different functions during eye development: (1) axis determination during DV patterning and (2) suppression of the retinal determination process to define the dorsal eye field margin. These functions of *pnr* appear to be temporally controlled as DV axis determination takes place in late first- or early second- instar of eye development (Singh et al., 2003; Singh et al., 2005b) while suppression of the eye fate is evident in late second instar of larval development.

The axis determination function of *pnr* is required in the earlier time window. This is further validated by the loss-of-function clones of *pnr* of first category, which are bigger and exhibits non –autonomous dorsal eye enlargement phenotypes (Fig. 2A, B). These dorsal eye enlargements that are spanning both wild-type and *pnr* mutant cells in the eye disc conforms to the notion that when *pnr* is lost in a group of cells during early development, it fails to confer dorsal identity over the default ventral state. As a consequence *de novo* equator is generated which results in the ectopic dorsal eye enlargements (Maurel-Zaffran and Treisman, 2000; Pichaud and Casares, 2000; Singh et al., 2005b). Since these clones are always bigger (Table 1), suggesting that they might be formed earlier. The late function of *pnr* in suppression of retinal determination is validated both by the gain-of-function studies (Fig. 3) as well as the loss-of-function clones of second category, which are both bigger as well as smaller in size and are autonomous in nature (Fig. 2C, D; Table 1). These clones have ectopic dorsal eyes, which are restricted within the clones, thereby suggesting that absence of *pnr* function promotes ectopic eye formation in the dorsal eye margin. Thus, during the early second instar of development, before the onset of retinal differentiation, *pnr* is required for defining the dorsal lineage by inducing *Wg* and members of the *Iro-C* complex (Maurel-Zaffran and Treisman, 2000; Singh et al., 2005b). However, later during the late second- instar stage of eye development, when the morphogenetic furrow (MF) is initiated, *pnr* suppresses the photoreceptor differentiation at the dorsal eye margin. The endogenous expression of *pnr* in only the peripodial membrane of the dorsal eye margin further confirms this notion (Fig. 1; Pereira et al., 2006). Lack of phenotypes in *pnr* clones which are restricted to the disc proper (DP) alone verifies *pnr* localization (Fig. 2E, F). Thus, *pnr* defines the boundary between the eye field and the head cuticle on the dorsal margin. An interesting question will be to identify which gene is responsible for defining the ventral eye margin. In the ventral eye where *pnr* is not expressed, *hth* is known to suppress the eye

fate (Pai et al., 1998; Pichaud and Casares, 2000). There is a strong possibility that *hth* may be involved in defining the boundary of eye field on the ventral eye margin between the disc proper and peripodial membrane.

Pnr induces Wg to suppress the eye development independent of *hth*

Since *pnr* induces Wg, it is expected that *pnr* may suppress the eye by induction of Wg (Fig. 8). Even though Wg signaling is responsible for suppression of photoreceptor differentiation on both dorsal and ventral eye margins (Ma and Moses, 1995; Treisman and Rubin, 1995; Lee and Treisman, 2001; Baonza and Freeman, 2002), its regulation is different on both dorsal and ventral eye margins (Pichaud and Casares, 2000; Maurel-Zaffran and Treisman, 2000). In the ventral eye, *wg* is involved in a feedback loop with *hth* to suppress the eye fate (Pichaud and Casares, 2000; Singh et al., 2002). In the dorsal eye, Pnr induces Wg signaling, which in turn induces the members of Iro-C complex, and ultimately these signaling interactions define the dorsal eye fate. Interestingly, it seems Pnr is not the sole regulator of Wg in the dorsal eye (Fig. 5). Since loss-of-function of *hth* does not exhibit the phenotypes similar to the loss-of-function of *pnr* (Fig. 6) or *wg* (Treisman and Rubin, 1995; Ma and Moses, 1995), it is expected that the positive feedback loop regulation of *wg* and *hth* as seen in the ventral eye margin does not hold true on the dorsal eye margin. We found that *hth* is not affected in the *pnr* clones that exhibit dorsal eye enlargements. Furthermore, when we made clones of *hth* in *pnr* heterozygous condition, we did not see any dorsal eye enlargements suggesting that *hth* and *pnr* do not interact. Thus, like others, our results also verified that *hth* is not involved in *pnr* mediated eye suppression on the dorsal eye margin (Fig. 8; Pichaud and Casares, 2000).

Eye suppression function of *pnr* is mediated through the suppression of *tsh*

It is known that the gain of function of *tsh* in the dorsal eye results in ectopic eye enlargement whereas gain of function of *tsh* in the ventral eye result in suppression of ventral eye (Singh et al., 2002; 2004). The dorsal eye enlargements seen in *tsh* gain-of-function is a phenotype similar to *pnr* loss-of-function in the dorsal eye. In *pnr* loss-of-function clones, we found that *tsh* was ectopically induced (Fig. 7C). Furthermore, when *pnr* loss-of-function clones were generated in a heterozygous background of the *tsh* null allele *tsh*⁸/CyO (Fasano et al., 1991), the dorsal eye enlargement phenotype was dramatically suppressed and no longer observed. Interestingly, we found that dorsal enlargements were there but were not accompanied with ectopic eyes as evident from absence of Elav expression (Fig. 7D). All these dorsal enlargements were showing strong Ey expression. Among 500 flies counted we found only two flies that showed subtle dorsal eye enlargements. Interestingly, we also found that in *pnr* loss-of-function clones the mini-*white* reporter gene under *tsh* was ectopically induced (data not shown). The loss-of-function phenotypes of *pnr* were more pronounced when *tsh* was misexpressed in the clones. Thus, *pnr* expressed in the dorsal peripodial membrane may suppress *tsh* in the dorsal eye to suppress the eye fate (Fig. 8). Interestingly, *tsh* is known to act upstream of *eya*, *so* and *dac* (Pan and Rubin, 1998). Thus, the dorsal eye enlargement observed in the *pnr* mutant is due to ectopic induction of *tsh* in the dorsal eye, which in turn can induce the RD genes (Fig. 8).

Functional conservation of dorsal selector Pnr

The *Drosophila* eye is similar to the vertebrate eye in several features (Sanes and Zipursky, 2010) like: (i) the morphogenetic furrow in the fly eye is analogous to the wave of neurogenesis in the vertebrate eye (Neumann and Nuesslein-Volhard, 2000; Hartenstein and Reh, 2002), (ii) Like *Drosophila*, in higher vertebrates dorsal eye genes like *Bmp4* and *Tbx5* act as 'dorsal selectors' and restrict the expression of ventral eye genes *Vax2* and *Pax2* (Koshiba-Takeuchi et al., 2000; Peters and Cepko, 2002). These DV expression domains or developmental compartments (Peters, 2002) lead to formation of DV lineage restriction as

seen in the *Drosophila* eye (Singh and Choi, 2003; Singh et al., 2005a), (iii) The DV lineage in the vertebrate eye also develops from a ventral-equivalent initial state (for review see Singh et al., 2005b). The dorsal genes *pnr* and *iro-C* are highly conserved across the species, and are involved in organogenesis and neural development (Gomez-Skarmaeta and Modolell, 2002; Singh et al., 2005b). Therefore, it would be interesting to see whether the dorsal selectors in the vertebrate eye play a role in defining the boundary of the eye by suppressing retinal differentiation.

Acknowledgments

We thank Kwang-Wook Choi, J. Treisman, Y Henry Sun, Justin Kumar, Fernando Casares, H. McNeill, S. Campuzano, Nancy Fossett, and the Bloomington Stock Center for the *Drosophila* strains; U. Walldorf, Patrick Callaerts, K. Cho and the Developmental Studies Hybridoma Bank (DSHB) for the antibodies; and members of Singh and Kango-Singh lab for critical comments on the manuscript. We thank Shiranee Gunasekera for fly food and stock maintenance. Confocal microscopy was supported by Biology Department central core facility. SMO has been supported by the Fight for Sight Inc. Summer research fellowship and an award from the Szabo Endowment fund. MKS is supported by Knights Templar Eye Foundation Inc., and start-up support from University of Dayton. This work is supported by NIH grant (1R15 HD064557-01), start-up support from University of Dayton, and Ohio Cancer Research Associates seed grant to AS.

References

- Atkins M, Mardon G. Signaling in the third dimension: the peripodial epithelium in eye disc development. *Dev Dyn.* 2009; 238:2139–48. [PubMed: 19623613]
- Azpiazua N, Morata G. Functional and regulatory interactions between Hox and extradenticle genes. *Genes Dev.* 1998; 12:261–273. [PubMed: 9436985]
- Baker NE. Embryonic and imaginal requirements for wingless, a segment-polarity gene in *Drosophila*. *Dev Biol.* 1988; 125:96–108. [PubMed: 3119404]
- Baonza A, Freeman M. Control of *Drosophila* eye specification by Wingless signalling. *Development.* 2002; 129:5313–5322. [PubMed: 12403704]
- Bessa J, Casares F. Restricted teashirt expression confers eye-specific responsiveness to Dpp and Wg signals during eye specification in *Drosophila*. *Development.* 2005; 132:5011–5020. [PubMed: 16221726]
- Bessa J, Gebelein B, Pichaud F, Casares F, Mann RS. Combinatorial control of *Drosophila* eye development by eyeless, homothorax, and teashirt. *Genes Dev.* 2002; 16:2415–27. [PubMed: 12231630]
- Blair SS. Cell lineage: compartments and Capricious. *Curr Biol.* 2001; 11:R1017–21. [PubMed: 11747838]
- Bonini NM, Leiserson WM, Benzer S. The eyes absent gene: genetic control of cell survival and differentiation in the developing *Drosophila* eye. *Cell.* 1993; 72(3):379–95. [PubMed: 8431945]
- Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development.* 1993; 118:401–15. [PubMed: 8223268]
- Calleja M, Moreno E, Pelaz S, Morata G. Visualization of gene expression in living adult *Drosophila*. *Science.* 1996; 274:252–5. [PubMed: 8824191]
- Casares F, Mann RS. Control of antennal versus leg development in *Drosophila*. *Nature.* 1998; 392(6677):723–6. [PubMed: 9565034]
- Cavodeassi F, Diez Del Corral R, Campuzano S, Domínguez M. Compartments and organizing boundaries in the *Drosophila* eye: the role of the homeodomain Iroquois proteins. *Development.* 1999; 126:4933–42. [PubMed: 10529412]
- Cavodeassi F, Modolell J, Campuzano S. The Iroquois homeobox genes function as dorsal selectors in the *Drosophila* eye. *Development.* 2000; 127:1921–9. [PubMed: 10751180]
- Chen R, Amoui M, Zhang Z, Mardon G. Dachshund and eyes absent proteins form a complex and function synergistically to induce ectopic eye development in *Drosophila*. *Cell.* 1997; 91(7):893–903. [PubMed: 9428513]

- Chern JJ, Choi KW. Lobe mediates Notch signaling to control domain-specific growth in the *Drosophila* eye disc. *Development*. 2002; 129:4005–13. [PubMed: 12163404]
- Cheyette BN, Green PJ, Martin K, Garren H, Hartenstein V, Zipursky SL. The *Drosophila sine oculis* locus encodes a homeodomain-containing protein required for the development of the entire visual system. *Neuron*. 1994; 12(5):977–996. [PubMed: 7910468]
- Cho K, Choi K. Fringe is essential for mirror symmetry and morphogenesis in the *Drosophila* eye. *Nature*. 1998; 396:272–6. [PubMed: 9834034]
- Cho K, Chern J, Izaddoost S, Choi K-W. Novel Signaling from the peripodial membrane is essential for eye disc patterning in *Drosophila*. *Cell*. 2000; 103:331–342. [PubMed: 11057905]
- Cohen, SM. Imaginal disc development. Bate, Martinez Arias; 1993. p. 747-841.
- Curtiss J, Halder G, Mlodzik M. Selector and signaling molecules cooperate in organ patterning. *Nat Cell Biol*. 2002; 4:E48–51. [PubMed: 11875444]
- Czerny T, Halder G, Kloter U, Souabni A, Gehring WJ, Busslinger M. twin of eyeless, a second Pax-6 gene of *Drosophila*, acts upstream of eyeless in the control of eye development. *Mol Cell*. 1999; 3(3):297–307. [PubMed: 10198632]
- Diaz-Benjumea FJ, Cohen SM. Interaction between dorsal and ventral cells in the imaginal disc directs wing development in *Drosophila*. *Cell*. 1993; 75:741–52. [PubMed: 8242746]
- Domínguez M, Casares F. Organ specification-growth control connection: New in-sights from the *Drosophila* eye-antennal disc. *Dev Dynamics* Vol. 2005; 232 (3):673–684.
- Domínguez M, de Celis JF. A dorsal/ventral boundary established by Notch controls growth and polarity in the *Drosophila* eye. *Nature* 1998. 1998; 396:276–8.
- Fasano L, Roder L, Core N, Alexandre E, Vola C, Jacq B, Kerridge S. The gene *teashirt* is required for the development of *Drosophila* embryonic trunk segments and encodes a protein with widely spaced zinc finger motifs. *Cell*. 1991; 64:63–79. [PubMed: 1846092]
- Fossett N, Tevosian SG, Gajewski K, Zhang Q, Orkin SH, Schulz RA. The Friend of GATA proteins U-shaped, FOG-1, and FOG-2 function as negative regulators of blood, heart, and eye development in *Drosophila*. *Proc Natl Acad Sci USA*. 2001; 98:7342–7. [PubMed: 11404479]
- Fu Y, Yan W, Mohun TJ, Evans SM. Vertebrate tinman homologues XNkx2–3 and XNkx2–5 are required for heart formation in a functionally redundant manner. *Development*. 1998; 124:4439–4449. [PubMed: 9778503]
- Furuta Y, Hogan BL. BMP4 is essential for lens induction in the mouse embryo. *Genes Dev*. 1998; 23:3764–3775. [PubMed: 9851982]
- Gallet A, Erkner A, Charroux B, Fasano L, Kerridge S. Trunk-specific modulation of wingless signaling in *Drosophila* by teashirt binding to *armadillo*. *Curr Biol*. 1998; 8:893–902. [PubMed: 9707400]
- García-Bellido A, Santamaria P. Developmental analysis of the wing disc in the mutant engrailed of *Drosophila melanogaster*. *Genetics*. 1972; 72:87–104. [PubMed: 4627463]
- Gibson M, Scubiger G. Peripodial cells regulate proliferation and patterning of *Drosophila* Imaginal Discs. *Cell*. 2000; 103:343–350. [PubMed: 11057906]
- Gómez-Skarmeta JL, Modolell J. Araucan and caupolican provide a link between compartment subdivisions and patterning of sensory organs and veins in the *Drosophila* wing. *Genes Dev*. 1996; 10:2935–45. [PubMed: 8918894]
- Gómez-Skarmeta JL, Modolell J. Iroquois genes: genomic organization and function in vertebrate neural development. *Curr Opin Genet Dev*. 2002; 12:403–8. [PubMed: 12100884]
- Haenlin M, Cubadda Y, Blondeau F, Heitzler P, Lutz Y, Simpson P, Romain P. Transcriptional Activity of pannier is regulated negatively by heteroimerization of the GATA DNA-binding domain with a cofactor encoded by *u-shaped* gene of *Drosophila*. *Genes Dev*. 1997; 11:3096–3108. [PubMed: 9367990]
- Halder G, Callaerts P, Gehring WJ. Induction of ectopic eyes by targeted expression of the *eyeless* gene in *Drosophila*. *Science*. 1995; 267(5205):1788–1792. [PubMed: 7892602]
- Halder G, Callaerts P, Flister S, Walldorf U, Kloter U, Gehring WJ. Eyeless initiates the expression of both *sine oculis* and *eyes absent* during *Drosophila* compound eye development. *Development*. 1998; 125:2181–2191. [PubMed: 9584118]

- Hartenstein, V.; Reh, TA. Homologies between vertebrate and invertebrate eyes. In: Moses, K., editor. *Drosophila Eye Development*. Vol. 37. Berlin: Springer-Verlag; 2002. p. 219-51.
- Haynie JL, Bryant PJ. Development in the eye-antennal imaginal disc and morphogenesis of the adult head in *Drosophila melanogaster*. *J Exp Zool*. 1986; 3:293-308. [PubMed: 3084703]
- Hazelett DJ, Bourouis M, Walldorf U, Treisman JE. *decapentaplegic* and *wingless* are regulated by *eyes absent* and *eyegone* and interact to direct the pattern of retinal differentiation in the eye disc. *Development*. 1998; 125:3741-3751. [PubMed: 9716539]
- Heberlein U, Moses K. Mechanisms of *Drosophila* retinal morphogenesis: the virtues of being progressive. *Cell*. 1995; 81:987-90. [PubMed: 7600585]
- Heberlein U, Borod ER, Chanut FA. Dorsoroventral patterning in the *Drosophila* retina by *wingless*. *Development*. 1998; 125:567-577. [PubMed: 9435278]
- Heitzler P, Haenlin M, Ramain P, Calleja M, Simpson P. A genetic analysis of *pannier*, a gene necessary for viability of dorsal tissues and bristle positioning in *Drosophila*. *Genetics*. 1996; 143:1271-86. [PubMed: 8807299]
- Held, LI, Jr. *Imaginal Discs: The Genetic and Cellular Logic of Pattern Formation*. Cambridge University Press; New York: 2002.
- Inbal A, Halachmi N, Dibner C, Frank D, Salzberg A. Genetic evidence for the transcriptional activating function of Homothorax during adult fly development. *Development*. 2001; 128:3405-13. [PubMed: 11566847]
- Ito K, Awano W, Suzuki K, Hiromi Y, Yamamoto D. The *Drosophila* mushroom body is a quadruple structure of clonal units each of which contains a virtually identical set of neurones and glial cells. *Development*. 1997; 124:761-771. [PubMed: 9043058]
- Jang CC, Chao JL, Jones N, Yao LC, Bessarab DA, Kuo YM, Jun S, Desplan C, Beckendorf SK, Sun YH. Two Pax genes, *eye gone* and *eyeless*, act cooperatively in promoting *Drosophila* eye development. *Development*. 2003; 130(13):2939-51. [PubMed: 12756177]
- Jaw TJ, You LR, Knoepfler PS, Yao LC, Pai CY, Tang CY, Chang LP, Berthelsen J, Blasi F, Kamps MP, Sun YH. Direct interaction of two homeoproteins, homothorax and extradenticle, is essential for EXD nuclear localization and function. *Mech Dev*. 2000; 91(1-2):279-91. [PubMed: 10704852]
- Jaynes JB, O'Farrell PH. Active repression of transcription by the engrailed homeodomain protein. *EMBO J*. 1991; 10:1427-1433. [PubMed: 1673924]
- Jurgens; Hartenstein. The terminal regions of the body pattern. In: Bate, M.; Martinez-Arias, A., editors. *The Development of Drosophila melanogaster*. Vol. I. Cold Spring Harbor: Cold Spring Harbor Laboratory Press; 1993. p. 687-746.
- Kango-Singh M, Singh A, Sun YH. *Eyeless* collaborates with Hedgehog and Decapentaplegic signaling in *Drosophila* eye induction. *Developmental Biology*. 2003; 256 (1):49-60. [PubMed: 12654291]
- Kehl BT, Cho KO, Choi KW. *mirror*, a *Drosophila* homeobox gene in the Iroquois complex. Is required for sensory organ and alula formation. *Development*. 1998; 125:1217-27. [PubMed: 9477320]
- Kenyon KL, Ranade SS, Curtiss J, Mlodzik M, Pignoni F. Coordinating proliferation and tissue specification to promote regional identity in the *Drosophila* head. *Dev Cell*. 2003; 5(3):403-14. [PubMed: 12967560]
- Klinedinst SL, Bodmer R. Gata factor *pannier* is required to establish competence for heart progenitor formation. *Development*. 2003; 130:3027-38. [PubMed: 12756184]
- Koshiba-Takeuchi K, Takeuchi JK, Matsumoto K, Momose T, Uno K, Hoepker V, Ogura K, Takahashi N, Nakamura H, Yasuda K, Ogura T. *Tbx5* and the retinotectum projection. *Science*. 2000; 287:134-7. [PubMed: 10615048]
- Kumar JP. The molecular circuitry governing retinal determination. *Biochim Biophys Acta*. 2009; 1789:306-14. [PubMed: 19013263]
- Lee JD, Treisman JE. *Sightless* has homology to Transmembrane acyltransferases and is required to generate active Hedgehog protein. *Curr Biol*. 2001; 11:1147-52. [PubMed: 11509241]
- Lee JD, Treisman JE. The role of *Wingless* signaling in establishing the anteroposterior and dorsoventral axes of the eye disc. *Development*. 2001; 128:1519-29. [PubMed: 11290291]

- Lee T, Luo L. Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. *Neuron*. 1999; 22:451–61. [PubMed: 10197526]
- Ma C, Moses K. Wingless and patched are negative regulators of the morphogenetic furrow and can affect tissue polarity in the developing *Drosophila* compound eye. *Development*. 1995; 121:2279–2289. [PubMed: 7671795]
- Mann R, Carroll S. Molecular mechanisms of selector gene function and evolution. *Curr Opin Genet Dev*. 2002; 12:592–600. [PubMed: 12200165]
- Mardon G, Solomon NM, Rubin GM. *dachshund* encodes a nuclear protein required for normal eye and leg development in *Drosophila*. *Development*. 1994; 120:3473–3486. [PubMed: 7821215]
- Maurel-Zaffran C, Treisman J. *pannier* acts upstream of wingless to direct dorsal eye development in *Drosophila*. *Development*. 2000; 127:1007–16. [PubMed: 10662640]
- McNeil H, Yang CH, Brodsky M, Ungos J, Simon MA. *mirror* encodes a novel PBX-class homeodomain that functions in the definition of the dorsal-ventral border in the *Drosophila* eye. *Genes Dev*. 1997; 11:1073–82. [PubMed: 9136934]
- Mílan M, Cohen SM. A re-evaluation of the contributions of *Apterous* and *Notch* to the dorsoventral lineage restriction boundary in the *Drosophila* wing. *Development*. 2003; 130:553–62. [PubMed: 12490561]
- Milner MJ, Bleasby AJ, Pyott A. The role of the peripodial membrane in the morphogenesis of the eye antennal disc of *Drosophila melanogaster*. *Roux Arch Dev Biol*. 1983; 192:164–170.
- Morata G, Lawrence P. Control of compartment development by the engrailed gene in *Drosophila*. *Nature*. 1975; 255:614–7. [PubMed: 1134551]
- Neumann CJ, Nüsslein-Volhard C. Patterning of the zebrafish retina by a wave of sonic hedgehog activity. *Science*. 2000; 289:2137–9. [PubMed: 11000118]
- Newsome TP, Asling B, Dickson BJ. Analysis of *Drosophila* photoreceptor guidance in eye specific mosaics. *Development*. 2000; 127:851–60. [PubMed: 10648243]
- Noro B, Culi J, McKay DJ, Zhang W, Mann RS. Distinct functions of homeodomain-containing and homeodomain-less isoforms encoded by *homothorax*. *Genes Dev*. 2006; 20(12):1636–50. [PubMed: 16778079]
- Obara-Ishihara T, Kuhlman J, Niswander L, Herzlinger D. The surface ectoderm is essential for nephric duct formation in the intermediate mesoderm. *Development*. 1999; 6:1103–1108. [PubMed: 10021330]
- Pai CY, Kuo TS, Jaw TJ, Kurant E, Chen CT, Bessarab DA, Salzberg A, Sun YH. The *Homothorax* homeoprotein activates the nuclear localization of another homeodomain, extradenticle, and suppresses eye development in *Drosophila*. *Genes Dev*. 1998; 12:435–46. [PubMed: 9450936]
- Pan D, Rubin GM. Targeted expression of *teashirt* induces ectopic eyes in *Drosophila*. *Proc Natl Acad Sci USA*. 1998; 95:15508–12. [PubMed: 9860999]
- Papayannopoulos V, Tomlinson A, Panin V, Rauskolb C, Irvine K. Dorsal-ventral signaling in the *Drosophila* eye. *Science*. 1998; 281:2031–2034. [PubMed: 9748163]
- Pappu KS, Mardon G. Genetic control of retinal specification and determination in *Drosophila*. *Int J Biol Dev*. 2004; 48:418–424.
- Pereira PS, Pinho S, Johnson K, Couso JP, Casares F. A 3' cis-regulatory region controls wingless expression in the *Drosophila* eye and leg primordia. *Dev Dyn*. 2006; 235(1):225–234. [PubMed: 16261625]
- Peters MA. Patterning the neural retina. *Curr Opin Neurobiol*. 2002; 12:43–8. [PubMed: 11861163]
- Peters MA, Cepko CL. The dorsal-ventral axis of the neural retina is divided into multiple domains of restricted gene expression which exhibit features of lineage compartments. *Dev Biol*. 2002; 251:59–73. [PubMed: 12413898]
- Pichaud F, Casares F. *homothorax* and *iroquois-C* genes are required for the establishment of territories within the developing eye disc. *Mech Dev*. 2000; 96:15–25. [PubMed: 10940621]
- Quiring R, Walldorf U, Kloter U, Gehring WJ. Homology of the *eyeless* Gene of *Drosophila* to the Small eye Gene in Mice and Aniridia in Humans. *Science*. 1994; 265:785–789. [PubMed: 7914031]

- Ramain P, Heitzler P, Haenlin M, Simpson P. *pannier*, a negative regulator of *achaete* and *scute* in *Drosophila*, encodes a zinc finger protein with homology to the vertebrate transcription factor GATA-1. *Development*. 1993; 119:1277–91. [PubMed: 7916679]
- Ready DF, Hanson TE, Benzer S. Development of the *Drosophila* retina, a neurocrystalline lattice. *Dev Biol*. 1976; 53:217–40. [PubMed: 825400]
- Rieckhof GE, Casares F, Ryoo HD, Abu-Shaar M, Mann RS. Nuclear translocation of extradenticle requires homothorax, which encodes an extradenticle-related homeodomain protein. *Cell*. 1997; 91(2):171–83. [PubMed: 9346235]
- Sanes JR, Zipursky SL. Design principles of insect and vertebrate visual systems. *Neuron*. 2010; 66:15–36. [PubMed: 20399726]
- Silver SJ, Rebay I. Signaling circuitries in development: insights from the retinal determination gene network. *Development*. 2005; 132(1):3–13. [PubMed: 15590745]
- Singh A, Choi KW. Initial state of *Drosophila* eye before dorso-ventral specification is equivalent to ventral. *Development*. 2003; 130:6351–6360. [PubMed: 14623824]
- Singh A, Kango-Singh M, Sun YH. Eye suppression, a novel function of *teashirt*, requires *Wingless* signaling. *Development*. 2002; 129:4217–80.
- Singh A, Kango-Singh M, Choi KW, Sun YH. Dorso-ventral asymmetric functions of *teashirt* in *Drosophila* eye development depend on spatial cues provided by early DV patterning genes. *Mech Dev*. 2004; 121:365–370. [PubMed: 15110046]
- Singh A, Chan J, Chern JJ, Choi KW. Genetic interaction of *Lobe* with its modifiers in dorsoventral patterning and growth of the *Drosophila* eye. *Genetics*. 2005a; 171:169–83. [PubMed: 15976174]
- Singh, A.; Lim, J.; Choi, K-W. Dorso-ventral boundary is required for organizing growth and planar polarity in the *Drosophila* eye. In: Mlodzik, M., editor. *Planar Cell Polarization during Development: Advances in Developmental Biology and Biochemistry*. Elsevier Science & Technology Books; 2005b. p. 59-91.
- Singh A, Shi X, Choi KW. *Lobe* and *Serrate* are required for cell survival during early eye development in *Drosophila*. *Development*. 2006; 133:4771–81. [PubMed: 17090721]
- Struhl G, Basler K. Organizing activity of *wingless* protein in *Drosophila*. *Cell*. 1993; 72:527–40. [PubMed: 8440019]
- Sun YH, Tsai CJ, Green MM, Chao JL, Yu CT, Jaw TJ, Yeh JY, Bolshakov VN. *white* as a reporter gene to detect transcriptional silencers specifying position-specific gene expression during *Drosophila melanogaster* eye development. *Genetics*. 1995; 141(3):1075–1086. [PubMed: 8582614]
- Tabata T, Schwartz C, Gustavson E, Ali Z, Kornberg T. Creating a *Drosophila* wing de novo, the role of *engrailed*, and the compartment border hypothesis. *Development*. 1995; 121:3359–69. [PubMed: 7588069]
- Tavsanli BC, Ostrin EJ, Burgess HK, Middlebrooks BW, Pham TA, Mardon G. Structure-function analysis of the *Drosophila* retinal determination protein *Dachshund*. *Dev Biol*. 2004; 272(1):231–47. [PubMed: 15242803]
- Treisman JE, Rubin GM. *wingless* inhibits the morphogenic furrow movement in the *Drosophila* eye disc. *Development*. 1995; 11:3519–3527. [PubMed: 8582266]
- Wolff, T.; Ready, DF. Pattern formation in the *Drosophila* retina. In: Bate, M.; Martinez-Arias, A., editors. *The Development of Drosophila melanogaster*. Vol. II. Cold-Spring Harbor: Cold Spring Harbor Laboratory Press; 1993. p. 1277-1325.
- Xu T, Rubin GM. Analysis of genetic mosaics in developing and adult *Drosophila* tissues. *Development*. 1993; 117:1223–1237. [PubMed: 8404527]




Fig. 1. Pnr expression is restricted to the peripodial membrane (PM) of the dorsal eye margin (A) *pnr* expression (*pnr* Gal4 drive UAS-GFP, Singh and Choi, 2003; Singh et al., 2005) is absent in the first instar eye-antennal imaginal disc whereas Wg (red) is expressed in the entire eye disc. Note that *pnr* expression in the brain at this stage is seen. (B) In the second instar eye-antennal imaginal disc, *pnr* expression (green) is initiated in 15–20 cells on the dorsal eye margin and Wg (red) is expressed laterally on both dorsal and ventral eye margins. At this stage, Hth (blue) expression is present in the entire eye disc. (C, C', C'') In the early third instar eye-antennal disc, *pnr* expression in the dorsal eye margin is restricted only to the peripodial membrane (PM) whereas Hth (blue) is also expressed in peripodial membrane (PM) of the eye-antennal disc. (C') *pnr* (green) expression at this stage is absent in the disc proper (DP). Hth (blue) expression begins to retract with the initiation of MF and stays anterior to the furrow (Pai et al., 1998; Bessa et al., 2002; Singh et al., 2002). (C'') Pnr expression is restricted to the peripodial membrane (PM) specific cells on the dorsal eye margin. (D) In the late third instar eye-antennal imaginal disc, *pnr* (green) expression is restricted to the dorsal eye margin whereas Wg (red) expression is restricted to the dorsal and ventral eye margins. Hth (Blue) is expressed in rings in the proximal region of antenna and expressed both in the dorsal and ventral part of the disc proper anterior to the furrow. Dashed lines indicate the approximate midline, the border between D (Dorsal) and V (ventral) eye. All the eye-antennal imaginal discs and the adult eyes are organized as Dorsal (D) up and the ventral (V) down. Markers for immunostaining are shown in color labels. (AN: Antenna)




Fig. 2. Loss-of-function of *pnr* exhibits a range of eye enlargements and antennal duplications in the dorsal eye

(A, B) Loss-of-function clones of *pnr* in the dorsal eye margin {marked by the absence of GFP (green) in the eye-antennal imaginal disc and absence of the *mini-white* reporter (red) in the adult eye} results in a non-autonomous ectopic eye enlargement as seen in the eye-antennal imaginal disc and in the adult eye. The ectopic eye enlargements are not restricted within the clone. However, they extend both in the wild-type as well as in the *pnr* mutant cells of the eye-antennal disc. Note that the dorsal clone boundary is marked by white dotted line in the eye disc and by black dotted line in the adult eye. (C, D) Loss-of-function of *pnr* in the dorsal eye results in an autonomous ectopic dorsal eye anterior to the normal eye field. These ectopic eyes are restricted to within the clones. Note that not all the cells of the *pnr* loss-of-function clone differentiate to the photoreceptors. (E, F) Loss-of-function clones of *pnr* in the dorsal eye have no effect on the eye field as seen in the eye disc and the adult eye. All these clones were restricted to the disc proper. (G, H) Loss-of-function clones of *pnr* in the antenna results in duplication of the antennal field as seen in (G) the eye-antennal disc and (H) the adult head. (H) Scanning electron microscopy (SEM) of the adult head showing antennal duplication and dorsal eye enlargement (Magnification X180). Note that only a few *pnr* loss-of-function clones show both dorsal eye enlargements along with the antennal duplication.




Fig. 3. Pnr suppresses the eye fate

(A) Eye-antennal imaginal disc showing domain of expression of GFP reporter (green) under the *ey* Gal4 (*ey>GFP*). Note that the *ey* Gal4 drives the expression of GFP reporter in the entire eye-antennal imaginal disc (both anterior as well as posterior to the morphogenetic furrow (MF) marked by white arrowhead). (B) Wild-type adult eye. (C, D) Misexpression of *pnr* in the eye using the *ey*-Gal4 driver (*ey>pnr^{D4}*) results in the suppression of eye fate and leads to a “no-eye” phenotype in the (C) eye-antennal disc (Elav, a pan-neural marker, which marks the photoreceptors) as well as in the (D) adult eye. The white dotted line in 3C marks the possible outline of the eye disc. There is no effect on the antennal field both in the eye-antennal imaginal disc as well as the adult head. (E) Another Gal4 driver, *bi*-Gal4 drives expression of a GFP reporter (*bi>GFP*) both on the dorsal and the ventral eye disc margin. (F) Misexpression of *pnr* using *bi*-Gal4 (*bi>pnr^{D4}*) results in the suppression of eye fate on both the dorsal and the ventral eye margin as evident from the loss of Elav expression (white arrows). (G, G', H) Gain-of-function clones of *pnr* (marked by GFP, white arrowhead) generated by random “flp-out” approach in the eye using the *heat shock*-FLP showed the suppression of eye as evident from the absence of (G') Elav in the eye disc as well as (H) in the adult eye. Note that the eye suppression in the *pnr* heat shock “flp out” clones was seen only in the larger clones. Further, necrosis (black spots) is also seen in the adult eye upon misexpression of *pnr* in the eye. (I, J) Blocking *pnr* function in the entire eye using *pnr^{ENR}* construct (*ey>pnr^{ENR}*) results in a “small eye” phenotype as seen (I) in the eye imaginal disc as well as (J) in the adult eye. (K) However, blocking *pnr* function both on the dorsal and the ventral eye disc margin (*bi>pnr^{ENR}*) results in the dorsal eye enlargement whereas there was no effect on the ventral eye margin. This data suggests that *pnr* suppresses eye on the dorsal eye margin.




Fig. 4. Pnr suppresses the expression of retinal differentiation genes in the eye
 (A, A', A'') Loss-of-function clones of *pnr* in the eye exhibit dorsal eye enlargement by (A'') ectopic Elav expression. In these clones where dorsal eye enlargement is seen, (A') the expression of retinal precursor marker Ey is restricted anterior to the furrow (white arrow). The dorsal eye enlargement, marked by Elav (blue) is the outcome of the *pnr* loss-of-function clone. Note that Ey is absent in the differentiating photoreceptors. Therefore, Ey is not seen in these clones. (B-D) Loss-of-function clones of *pnr* showing an ectopic dorsal eye phenotype with ectopic induction of retinal determination genes like (B, B', B'') Eya (white arrow), (C, C', C'') So (white arrow), and (D, D', D'') Dac (white arrow). Note that these retinal determination genes, which act downstream to Ey, and unlike Ey are expressed in the differentiating photoreceptor neurons. (E- H) Gain-of-function of *pnr* in the eye suppresses the retinal determination genes. (E, E') Misexpression of *pnr* on dorsal and ventral eye

margins by using a *bi*-Gal4 driver (*bi>pnr^{D4}*), results in strong upregulation of *Ey* on both dorsal and ventral eye margins (marked by a white arrowhead). (F-H) However, misexpression of *pnr* (*bi>pnr^{D4}*) suppresses the downstream retinal differentiation genes (F, F') *Eya*, (G) *So*, (H) *Dac* on both dorsal and ventral margins (marked by arrow heads). The anterior *Dac* expression (anterior to furrow) went all the way down to the posterior margin in *bi>pnr^{D4}* misexpression. *Ey* marks retinal precursor cells and is required for the specification of eye field. Our results suggest that *pnr* may not affect early eye specification function of *Ey*, whereas *pnr* suppresses the retinal determination genes like *eya*, *so* and *dac*, which acts downstream to *Ey*.




Fig. 5. *pnr* induces downstream target *Wg* to suppress the eye

Wg is known to act as a negative suppressor of eye fate. *Wg* is expressed laterally both on the dorsal and the ventral eye margins (Fig. 1B). (A, A') Misexpression of *pnr* on both dorsal and ventral eye margin using *bi-Gal4* results in the suppression of eye on both DV margins along with ectopic induction of *Wg* (marked by white arrows). (B) *bi>pnr^{D4}* results in the reduction of eye both on the dorsal and ventral eye margins. This phenotype is similar to *bi>wg* (Singh et al., 2002). (C) Misexpression of *Wg* in the entire eye using *ey-Gal4* (*ey>wg*) results in “no-eye”. (D) Loss-of-function clones of *pnr* in the dorsal eye (marked by absence of GFP reporter and white dotted line) result in the ectopic eye enlargement along with the suppression of *Wg* expression. (E, E') Loss-of-function clones of *pnr* in the DP (marked by white dotted line) caused no effect on *Wg* expression as *Pnr* is not expressed in the DP. (E') Higher magnification of the clone showing its location restricted to the DP. (F) Interestingly, some of the bigger loss-of-function clones of *pnr* (marked by white dotted line) exhibit the dorsal eye enlargement. However, this eye enlargement do not cover the entire clone. The part of the clone anterior to the eye enlargement show robust *Wg* expression. These clones show some overgrowth in the eye disc, which will form head specific structures, suggesting that within dorsal eye margin *pnr* is not the sole *Wg* regulator.




Fig. 6. *pnr* suppresses the eye fate at dorsal eye margin independent of *hth*

hth, a Meis class of gene (Rieckhof et al., 1997), acts as a negative regulator of the eye (Pai et al., 1998). *Hth* expression is restricted anterior to furrow in 10–15 cell wide domain and in entire peripodial membrane (Fig. 1). (A, A') Misexpression of *hth* on both dorsal and ventral eye margin (*bi>hth*) results in the suppression of eye fate on both dorsal and ventral margin of the eye disc as evident from (A') suppression of *Elav* (marked by white arrows). (B, B') Misexpression of *pnr* on both dorsal and ventral eye margin (*bi>pnr^{D4}*) results in suppression of eye on both dorsal and ventral eye margin (marked by white arrows), which is accompanied by induction of *Wg* (green) as well as *Hth* (red; white arrows). (C, C') Loss-of-function clone of *hth* in the eye has DV asymmetric phenotypes. The loss-of-function clone of *hth* in the ventral eye results in the eye enlargement as evident from *Elav* expression (marked by white dotted line). Note that the dorsal eye clones do not exhibit any phenotype. (D–F) In loss-of-function clones of *pnr*, (D, D') which result in dorsal eye enlargement (marked by white dotted line) or (E, E') which do not exhibit dorsal eye enlargement (marked by white dotted line), *Hth* (red) expression stays anterior to the furrow as seen in the wild-type eye disc. Loss-of-function clones of *pnr* in the antennal disc which results in the duplication of antennal field exhibit ectopic *Hth* expression in the duplicated antennal disc (marked by white dotted line). Note that *hth* is expressed in the proximal region of the antennal disc.




Fig. 7. Pnr suppresses the eye fate by downregulating *teashirt* (*tsh*) in the dorsal eye margin
Tsh, a Hox gene (Fasano et al., 1991), exhibits Dorso-ventral (DV) asymmetric function in the eye (Singh et al., 2002). *pnr* expression initiates in early second instar eye-antennal imaginal disc (Singh & Choi, 2003). (A) In the late second instar, *pnr* expression evolves and is restricted to 50–100 cells of the dorsal eye margin. At this stage when MF has just initiated, *Tsh* is expressed anterior to the furrow (MF). (B) In the third instar eye imaginal disc, *pnr* is expressed on the dorsal eye margin whereas *tsh* is expressed in the eye disc anterior to the furrow. (C–C'') Loss-of-function clone of *pnr* in the dorsal eye marked by the loss of GFP reporter (marked by white dotted line) exhibit (C') ectopic localization of *Tsh* protein, and (C'') ectopic expression of *tsh* reporter (*tsh^{A8}/CyO*) in the dorsal eye. (D–D'') Loss-of-function clones of *pnr* (marked by loss of GFP), where *tsh* function is reduced to half using a heterozygous background of *tsh* null allele (*tsh^{8/+}*), (D') exhibit outgrowth on the dorsal eye margin which is positive for *Ey* expression but there is no ectopic eye enlargement as evident from (D'') absence of neuronal marker *Elav* expression. The dorsal overgrowth exhibits robust expression of *Ey*, a marker for undifferentiated retinal precursor cells. (E) Loss-of-function clone of *pnr* in the *tsh* heterozygous background marked by the loss of *mini-white* reporter (red: clonal boundary marked by black dotted line) results in the absence of eye enlargement in the adult eye. These results suggest that *pnr* eye suppression function is mediated through down regulation of *tsh*. (F) Misexpression of *pnr* on the dorsal and the ventral eye margin, *bi>pnr^{D4}*, results in the suppression of *tsh* reporter on both dorsal and ventral eye margin (white arrowhead) along with the suppression of eye as evident from *Elav* (blue) expression. Note that eye size is reduced on both margins. (G) Misexpression of *pnr^{D4}* both on dorsal and ventral eye margin in *tsh* heterozygous background (*tsh^{8/+}; bi>pnr^{D4}*) exhibits strong suppression of eye resulting in a highly reduced eye. Note that *bi>pnr^{D4}* alone (F) shows suppression of eye both on the dorsal and the ventral eye margin. However, the size of *bi>tsh^{8/+}; pnr^{D4}* eye imaginal disc size is extremely reduced as compared to *bi>pnr^{D4}* alone. (H) Misexpression of *tsh* on DV margin (*bi>tsh*) results in the suppression of eye on the ventral margin whereas eye enlargement in the dorsal eye (Singh et al., 2002). Misexpression of both *tsh* and *pnr* on DV margin results in early lethality. Therefore, we misexpressed *pnr* downstream target *ara* with *tsh*. (I) Misexpression of *tsh* with dorsal eye selector *ara*, a downstream target of *pnr*, on DV margin using *bi-Gal4* (*bi>tsh+ara*) results in the enlargement on both dorsal and ventral eye

margins. Misexpression of *tsh* and *ara* on DV margin results in strong dorsal eye enlargements. (J) Misexpression of *tsh* using *pnr*-Gal4 driver (*pnr>tsh*) results in the enlargement of the dorsal eye. (K) Misexpression of *tsh* in the heterozygous *pnr* background results in the dorsal eye enlargement. However, these eye enlargements are not bigger than what is seen in (H) *bi>tsh* or (J) *pnr>tsh*, suggesting that *pnr* acts upstream of *tsh*.




Fig. 8. Pnr suppresses the eye fate by downregulating *tsh* which results in suppression of retinal determination genes at the dorsal eye margin

GATA-1 transcription factor *pnr*, which is expressed in the peripodial membrane (PM) at the dorsal eye margin, suppresses the retinal determination. The suppression of retinal determination genes by *pnr* can be mediated by two possible ways: (i) *pnr* directly suppresses the retinal determination genes to suppress the eye. *pnr* may act downstream to *ey* and suppress the downstream retinal determination target *eya* and other downstream genes *so* and *dac*. During eye development, *ey* is required for eye specification and other downstream targets are required for retinal determination. Our studies suggest that *pnr* acts on retinal determination process, which corresponds to the onset of *pnr* expression in the eye. (ii) Alternatively, *pnr* suppresses the eye by downregulating homeotic gene *teashirt* (*tsh*) in the dorsal eye. Interestingly, the *tsh* gain-of-function in the dorsal eye (Singh et al., 2002) is complementary to the loss-of-function of *pnr* in the dorsal eye. *tsh* is known to act upstream of *eya*, *so* and *dac* (Pan and Rubin, 1998). Thus, the dorsal eye enlargement observed in *pnr* mutant is due to ectopic induction of *tsh* in the dorsal eye, which in turn can induce the RD genes. Lastly, Pnr mediated suppression of the eye fate is independent of Meis class of homeotic gene, *homothorax* (*hth*) function.

Table 1
 Classification of the loss-of-function phenotypes of *pannier* (*pnr*) clones in the eye and antenna.

Phenotype	Large clones with dorsal eye enlargements	Small clones with dorsal eye enlargements	No dorsal eye enlargements	Antennal duplication	Total flies (with clones) counted
Clones with Non-autonomous dorsal eye enlargement Class I	34 (8.3%)				34
Clones with autonomous dorsal eye enlargement Class II	23 (5.6%)	29 (7.0%)			52
No dorsal eye enlargement Class III			308 (75.3%)		308
Antennal Duplication Class IV	(10)*	(4)*		15 (3.6%)	15
Grand Total					409

* The flies in this category showed both antennal duplications dorsal eye enlargements of large and small size.