

University of Dayton eCommons

Health and Sport Science Faculty Publications

Department of Health and Sport Science

3-2007

Derivation of an Age and Weight Handicap for the 5K Run

Paul M. Vanderburgh

University of Dayton, pvanderburgh1@udayton.edu

Lloyd L. Laubach

University of Dayton, llaubach1@udayton.edu

Follow this and additional works at: https://ecommons.udayton.edu/hss_fac_pub

 Part of the [Exercise Physiology Commons](#), [Kinesiology Commons](#), and the [Sports Sciences Commons](#)

eCommons Citation

Vanderburgh, Paul M. and Laubach, Lloyd L., "Derivation of an Age and Weight Handicap for the 5K Run" (2007). *Health and Sport Science Faculty Publications*. 27.

https://ecommons.udayton.edu/hss_fac_pub/27

This Article is brought to you for free and open access by the Department of Health and Sport Science at eCommons. It has been accepted for inclusion in Health and Sport Science Faculty Publications by an authorized administrator of eCommons. For more information, please contact frice1@udayton.edu, mschlangen1@udayton.edu.

DERIVATION OF AN AGE AND WEIGHT HANDICAP FOR THE 5K RUN

**(Final Revision, Accepted for Publication in Measurement in Physical Education
and Exercise Science)**

Published in 2007

Paul M. Vanderburgh, EdD*

Department of Health and Sport Science

University of Dayton, Dayton OH, 45469-1210

TEL: 937.229.4213

FAX: 937.229.4244

Email: vanderburgh@udayton.edu

and

Lloyd L. Laubach, PhD

Department of Health and Sport Science

University of Dayton, Dayton OH, 45469-1210

***Corresponding Author**

Running Title: 5K Run Handicap

1 Abstract

2

3 The adverse effect of increasing age and/or body weight on distance run performance has
4 been well documented. Accordingly, nearly all five kilometer (5K) road races employ
5 age categories and, sometimes, a heavier body weight classification. Problems with such
6 conventions include small numbers of runners within older age categories and the
7 advantage given to the lightest runners within each weight category. We developed a 5K
8 Handicap (5KH), a model that calculates an adjusted run time based on the inputs of
9 actual 5K run time, age, and body weight for men and women. This adjusted time, then,
10 can be compared between runners of different ages and body weights. The purpose of
11 this paper was to explain, in detail, the derivation of the 5KH formula using published
12 theoretical and empirical findings on age, body weight and distance run time
13 relationships. To our knowledge, the 5KH is the first such model and overcomes the
14 problems associated with being heavier within one weight class and having too few
15 runners in certain age categories. We are currently undertaking large-scale validation
16 studies and evaluation of its race day implementation.

17

18 Key Words: allometry, age handicap, weight handicap, running

19

1 Participation in distance running road races suggests their increasing popularity.
2 According to the Running USA’s Road Running Information Center
3 (www.runningusa.org), 2004 U.S. road race participation grew by an average of 4% from
4 the previous year, with increases evidenced at every standard race distance, and nearly
5 eight million finishers. Furthermore, recent research advances offer new ways to
6 evaluate race performance that may enhance such participation especially for older and
7 heavier runners.

8 While many factors contribute to distance run performance, the notion that
9 runners of advanced age and body weight (BW) post slower times is widely accepted.
10 Accordingly, nearly all races have age categories and, in some cases, body weight
11 divisions. The primary limitation of the age categories is that there are often very few
12 competitors in the older categories, sometimes fewer runners than awards. When BW
13 divisions are used, typically only two heavier weight divisions are added such that
14 runners above the minimum weight can compete against each other for the additional
15 awards beyond those of the open division. The divisions, named “Clydesdale” for men
16 and “Fillies” or “Athena” for women, have different BW cut-offs depending on the
17 choice of organization affiliation (www.clydesdale.org or www.usa-clydesdale.com).
18 The minimum BW established for these divisions is arbitrarily assigned and the use of
19 such a category gives competitors close to the minimum weight an advantage over
20 heavier ones (based on physiological laws of similarity discussed in the Methods section
21 below). These limitations could be mitigated by using an age and BW handicapping
22 system that more precisely handicaps age and weight as continuous variables. More

1 importantly, the modeling of distance run performance has more recently included both
2 theoretical and physiological evidence defending various handicap systems.

3 The World Association of Veteran Athletes (WAVA) has popularized an age
4 handicap model for many different track and field events including distance runs up to
5 the marathon. Though never published, and because the organization no longer exists,
6 the WAVA age handicap model appears to be available only on an independent website
7 (<http://www.howardgrubb.co.uk/athletics/wavalookup.html>, Nov 2005) which also
8 includes links to the development of its algorithms. According to this source, the WAVA
9 model is based on world best performances which determine the slopes of the age vs. run
10 time curves. Comparisons of distance run times between individuals of different age,
11 then, are based on distance from the curve. This can be problematic because the slope of
12 any such curve is very sensitive to data at the limits of age. The fewer elderly
13 competitors at the world class level would likely contribute to world bests not being age-
14 comparable to those of younger runners. This, in turn, would lead to larger than
15 appropriate age handicaps for older runners.

16 To our knowledge, however, there exists no published, physiologically defensible
17 model that adjusts distance run time by age and BW. Recently, however, research has
18 been published examining the relationship between body size and age on various indices
19 of physical performance, including distance running (Nevill, Ramsbottom, & Williams,
20 1992; Vanderburgh & Mahar, 1995). These data and the application of related
21 theoretical models of allometry, also receiving attention in recent literature, can be used
22 in combination to better quantify the contributions of age or BW on distance run times.

1 The first step in developing an age and BW handicap model for distance runs is to
2 quantify the physiological effect of age on distance run performance. The pathway of
3 assimilating empirical findings is based on two key relationships: age vs. maximal
4 oxygen uptake (VO_{2max}) and VO_{2max} vs. distance run time. The former has been well
5 documented for men (Jackson et al., 1995) and women (Jackson et al., 1996). In these
6 large scale studies, Jackson and colleagues assessed VO_{2max} , percent body fat, and self-
7 reported physical activity levels for 1499 men (ages 20-75) and 409 women (ages 20-64).
8 Using cross-sectional and longitudinal analyses, they determined that, for both genders,
9 50% of the age-associated decline in VO_{2max} was due to the influence of percent body fat
10 and self-reported physical activity. Furthermore, they quantified the change in VO_{2max}
11 with age independent of these two factors as 0.26 and 0.25 $ml \cdot kg^{-1} \cdot min^{-1} \cdot year^{-1}$ for men
12 and women, respectively. These slopes are key in the consideration of an age handicap
13 model for distance runs because they represent how functional capacity, a key
14 determinant of distance run speed, should change due to age, independent of body fatness
15 and activity level.

16 The link between VO_{2max} and distance run speed has been well-established for the
17 5K distance. Nevill and colleagues assessed the relationship between 5K run speed and
18 VO_{2max} and BW for 308 recreationally active men and women (Nevill et al., 1992) and
19 developed the following equation:

$$20 \qquad \qquad \qquad 5K \text{ Run Speed} = 84.3(VO_{2max})^{1.01}(BW)^{-1.03} \qquad (1)$$

21 where run speed, VO_{2max} and BW are expressed in $m \cdot sec^{-1}$, $l \cdot min^{-1}$ and kg, respectively.

22 This equation can then be used in combination with the findings of Jackson and

1 This relationship has empirical support. Crowder and Yunker (1996), in a sample
2 of 238 fit service academy males, reported a BW exponent of 0.26 ± 0.06 for the two-
3 mile run. Similarly, Vanderburgh and Mahar (1995), with a sample of 59 male cadets
4 from the same service academy, found the BW exponent to be 0.40 ± 0.09 . Because
5 these were active, fit, lean, and young cadets, the effect of BW on these run times can be
6 considered somewhat independent of body fatness and physical activity level.

7 The purpose of this paper, then, was to use published theoretical and empirical
8 evidence to develop an age and BW handicap system for the 5K distance run that could
9 be used to evaluate run performance in a novel way.

10

11 METHODS

12 The basis of our methodology was to use these theoretically based and empirically
13 supported findings to develop the 5K Handicap (5KH), an age and BW handicap model
14 for male and female 5K runners. The model calculates an adjusted run time, RT_{adj} , the
15 score that can be used to compare against those of other runners of different BW or age,
16 within each gender. We modeled our approach after that of the Wilks Formula of the
17 International Powerlifting Federation (<http://www.powerlifting-ipf.com>, Nov 2005),
18 validated by Vanderburgh and Batterham (1999). This formula computes a correction
19 factor, F , for BW which is multiplied by the total weight lifted to yield an adjusted score
20 that determines the overall best lifter among all BW categories. For the 5KH, however,
21 two correction factors are needed, F_{age} and F_{BW} , such that:

$$22 \qquad \qquad \qquad RT_{adj} = F_{BW} F_{age} RT \qquad (6)$$

1 But each F is equal to the ratio of the adjusted run time for that variable (RT_{BW}
2 and RT_{age}) to the actual run time. Therefore:

$$3 \quad RT_{adj} = RT \cdot (RT_{BW} \cdot RT^{-1}) \cdot (RT_{age} \cdot RT^{-1})$$

4 or

$$5 \quad RT_{adj} = RT_{age} \cdot RT_{BW} \cdot RT^{-1} \quad (7)$$

6 Obtaining one's RT_{adj} , then, requires the calculation of RT_{age} and RT_{BW} . These can be
7 thought of as the RT the runner would obtain if he/she were a “scale model” of
8 him/herself but younger and lighter, at a standard age and BW for all runners. In other
9 words, the 5KH statistically makes each runner the same age and BW and computes the
10 resulting performance.

11 Distance run time, widely acknowledged to increase with increasing age and BW,
12 presents an interesting modeling dilemma. One could mathematically infer that lower
13 BW and age result in improved run performance. Clearly, though, as age and BW
14 approach zero, run performance cannot possibly improve. To account for this, we
15 established limits for both age and BW, below which RT would not be adjusted. These
16 limits could be thought of as the “optimal” distance running age and BW for each gender.
17 We chose 25 years, 50 kg for women, and 65 kg for men as these lower limits. The age
18 value of 25 was based on the mean age 25 (male) and 26 (female) for the top 20 5K male
19 and female runners in the world, according to the International Association of Athletics
20 Federations (IAAF) rankings as of November, 2005 (www.iaaf.org, Nov 2005). The
21 BWs of 50 kg for women and 65 kg for men were selected based on the mean BWs of
22 samples of national-class female (Morikawa et al., 2001, $N = 26$, mean BW = 51.8 kg)
23 and male (Saunders et al., 2004, $N = 81$, mean BW = 66.6 kg) middle-distance runners.

1 Using these elite lower limit values is actually more advantageous to novice runners than
2 using those of the reference man and woman because this affords everyone above 50 kg
3 (women), 65 kg (men), and/or 25 years of age a handicap.

4 Fundamentally, RT_{age} is the equivalent RT if the runner were “made younger” to
5 the age of 25 years. This entails computing the change in VO_{2max} with the age change via
6 Jackson’s slope values (1995, 1996) as well as the resulting RT using Equation 1.

7 Though the equation becomes a bit lengthy, it can be broken down into several steps.

8 First, the equivalent VO_{2max} is computed at the specified RT and BW with Nevill’s
9 Equation 1 (1992). This involves both the conversion of run speed to run time and
10 absolute ($l \cdot min^{-1}$) to relative VO_{2max} ($ml \cdot kg^{-1} \cdot min^{-1}$), both simple arithmetic operations.

11 Second, the resultant relative VO_{2max} is increased by adding actual minus target age (25
12 years) and multiplying that number by the slopes of Jackson’s (1995,1996) age vs.
13 VO_{2max} slopes, 0.25 and 0.26 $ml \cdot kg^{-1} \cdot min^{-1} \cdot year^{-1}$ for women and men, respectively.

14 Third, this adjusted VO_{2max} is then used to recompute the RT_{age} using Equation 1.

15 RT_{BW} , the adjusted RT due to BW alone, is computed by applying Equation 5:

16 $RT \propto BW^{1/3}$. Similar to RT_{age} , RT_{BW} is the functional equivalent of creating a scale

17 model of the runner at the standard weight of 50 and 65 kg, for women and men,

18 respectively and recomputing the resulting RT. If Equation 5 dictates that a change in

19 $BW^{1/3}$ is directly proportional to RT, then:

20
$$RT_{BW} = RT \cdot (X \cdot BW^{-1})^{1/3} \quad (8)$$

21 where X = 50 and 65 kg, for women and men, respectively. Substituting into Equation

22 7, this yields the final basic formula to compute RT_{adj} :

23
$$RT_{adj} = RT_{age} \cdot (X \cdot BW^{-1})^{1/3} \quad (9)$$

1 Armed with RT_{age} and RT_{BW} , and, of course, the actual RT, one can now apply
2 Equation 7 to compute the final RT_{adj} , which is that adjusted for BW and age. Age and
3 BW below the standard values should be entered as the standard values (e.g., “25” would
4 be entered for an age of 18 and “65” would be entered for a male BW of 59 kg) because
5 of the unknown nature of the physiological and allometric modeling for run times with
6 children and young adults. Furthermore, the 5KH is designed to overcome the
7 unavoidable RT increase due to age and BW, which primarily occurs only in adulthood.
8 The formula is not designed to handicap young, pre-pubertal runners for being below the
9 age and BW standards.

10

11 RESULTS

12 Table 1 shows four hypothetical examples of male and female runners and the
13 adjusted run times associated with age alone, RT_{age} . Runner A receives minor credit for
14 age because he is close to the age standard of 25 years. The 9 s of handicap he receives is
15 due to the three extra years above 25 that would reduce his VO_{2max} . Note that runners B
16 and D, who are 18 and 14 years, respectively, away from the standard of 25 years, receive
17 the most handicap for age which shows up in the RT minus RT_{age} .

18 Table 1 also shows RT_{BW} for the same four runners. Runners B and D receive a
19 larger handicap for BW than age because, physiologically, each experiences a greater
20 handicap for being heavier compared to being older. Incidentally, Nevill’s Equation 1
21 (1992) could have been used to compute the RT_{BW} , yielding approximately a 1%
22 difference from the RT_{adj} obtained using Equation 8. This is because the 5KH model was
23 predicated on Equation 4 which is slightly different from that derived from Equation 1

1 where the ratio of the VO_{2max} and BW exponents was not exactly 1.0. Our decision to
 2 use Equation 4 rather than Equation 1 for BW adjustment was based on our preference to
 3 defer to that which is theoretically more defensible rather than empirical. Nevertheless,
 4 either approach yields very similar results.

5 Finally, Table 1 shows the RT_{adj} column. Runners B and D, who run
 6 considerably slower than their counterparts of the same gender, actually score lower RT_{adj}
 7 values because they are older and heavier. Furthermore, for each runner, the handicap
 8 due to age (RT minus RT_{age}) plus that due to BW (RT minus RT_{BW}) is greater than the
 9 the total handicap (RT minus RT_{adj}) because of Equation 7. In other words Equation 7
 10 shows that the individual handicaps are not additive. The resulting equations, in the form
 11 that can be quickly adapted to a common spreadsheet software program, for BW (kg),
 12 AGE (years), and RT (sec), are:

$$13 \quad \text{Women's } RT_{adj} = (59.31 * BW^{1.03} / ((((((59.31 * BW^{1.03} / RT)^{(1/1.01)) * 1000 / BW +$$

$$14 \quad (AGE - 25) * 0.25)) * BW) / 1000)^{1.01} * (50 / BW)^{(1/3)} \quad (10)$$

$$16 \quad \text{Men's } RT_{adj} = (59.31 * BW^{1.03} / ((((((59.31 * BW^{1.03} / RT)^{(1/1.01)) * 1000 / BW + (AGE -$$

$$17 \quad 25) * 0.26)) * BW) / 1000)^{1.01} * (65 / BW)^{(1/3)} \quad (11)$$

19 DISCUSSION

20 To our knowledge, the 5KH is the first age and BW distance run handicap model
 21 developed using documented relationships between age and BW on distance run time.
 22 The formula is based on the scale model approach that essentially recalculates a runner's
 23 time if he/she were 25 years of age and 50 or 65 kg BW, for women and men,

1 respectively. This adjusted time, RT_{adj} , can be used to compare against any other runner
2 of the same gender but of different BW and age. Figure 1 shows four hypothetical men
3 and women of varying ages and BW and the effect of the overall handicap. Clearly, an
4 older, heavier and slower runner can “compete” with younger, lighter and faster runners.

5 One might posit that the 5KH rewards excess body fat. Detailed analysis of some
6 “what-if” scenarios suggests otherwise. Consider a man, 45 years, 85 kg, 15% body fat,
7 who runs a 1500 s (25:00) 5K. His RT_{adj} would be 1227 s (20:27). If, however, he
8 gained 3 kg of fat weight with no change in functional capacity, then his RT, according to
9 Equation 1, would change to 1556 s (25:56), 56 s slower. His new RT_{adj} would be 1253 s
10 (20:53), for a net slower RT_{adj} of 26 sec. Conversely, if he gained 3 kg of only muscle
11 mass, assuming that functional capacity remained proportionally the same with respect to
12 lean body mass, he would show a net improvement of 13 s in RT_{adj} . Finally, if the 3 kg
13 gain in weight maintained the same percent body fat with similar functional capacity
14 assumptions, the calculations reveal that his RT_{adj} would remain unchanged. This is
15 expected because he would essentially be a larger scale model of himself. In short,
16 excess fat weight is disadvantageous for the 5KH.

17 Validation of the 5KH is an important objective associated with any such model,
18 but should be done only after careful consideration of the model’s characteristics. From a
19 simplistic perspective, a valid model means that the bias against heavier and older
20 runners is eliminated. Statistically, this could be ascertained by correlating age and BW
21 each with RT_{adj} ; both correlations should be near zero. For a large sample of 5K runners,
22 however, this is not likely even with a valid model. For example, few runners will give
23 maximum effort. This will lead to slower RT performance and RT_{adj} scores not

1 commensurate with age or BW, yielding a spurious effect on the correlation.
2 Furthermore, as mentioned previously, runners who are heavier due to excess body fat,
3 will have slower than expected RT_{adj} scores. The 5KH's advantage, however, is that
4 while effort and percent body fat are not directly measured, the model does penalize
5 runners who give less than a maximum effort and/or carry excess body fat, a desirable
6 outcome from a health perspective. When attempting to validate the 5KH, therefore,
7 researchers need to control for the effects of effort and body fatness or at least consider
8 them in the interpretation of the validation data. Additional information regarding the
9 practice application of the 5KH will be available upon completion of our ongoing
10 validation and implementation studies.

11 We compared the 5KH standards to those of the aforementioned WAVA
12 (<http://www.howardgrubb.co.uk/athletics/wavalookup.html>, Nov 2005), and their age-
13 only grading formula for the 5K. Because their data include the world best performance
14 for different age categories, and the older categories have fewer competitors, we expected
15 the WAVA model to provide a larger handicap for increasing age. Indeed, as shown in
16 Figure 2, for an average RT, male and female, the WAVA standards show a steeper slope
17 than the 5KH for either gender, indicating a larger handicap for older runners than
18 physiological modeling supports. Furthermore, they more closely coincide with those of
19 the 5KH at the lighter weights, which is to be expected because world bests across all age
20 groups are most likely from light runners. Finally, the WAVA model, for heavier,
21 younger runners, yields a significantly smaller handicap than the 5KH because the
22 WAVA model makes no adjustment for body weight.

1 In conclusion, we have detailed the derivation of the 5KH, an age and BW
2 handicap model for 5K runners, which yields an adjusted race time, RT_{adj} , based on the
3 contribution of these two factors. This contribution has a theoretical and physiological
4 basis that appears defensible. We recommend rigorous large scale validation studies of
5 this model, which we are currently conducting; however, careful interpretation of results
6 will be important given the likelihood that the handicap is not likely to eliminate all
7 effects of age and BW due to the influences of effort and/or body fatness.

References

- Astrand, P.O., & Rodahl, K. (1986). *Textbook of work physiology*. New York: McGraw-Hill.
- Bourdin, M., Pastene, J., Germain, M., & Lacour, J.R. (1993). Influence of training, sex, age, and body-mass on the energy-cost of running. *European Journal of Applied Physiology and Occupational Physiology*, 66, 439-444.
- Crowder, T.A., & Yunker, C. (1996). Scaling of push-up, sit-up and two-mile run performances by body weight and fat-free weight in young, fit men. *Medicine and Science in Sports and Exercise*, 28, S183.
- Jackson, A.S., Beard, E.F., Weir, L.T., Ross, R., & Blair, S. (1995). Changes in aerobic power of men, ages 25-70 years. *Medicine and Science in Sports and Exercise*, 27, 113-120.
- Jackson, A.S., Weir, L.T., Ayers, G.W., Beard, E.F., Stuteville, J., & Blair, S. (1996). Changes in aerobic power of women, ages 20-64. *Medicine and Science in Sports and Exercise*, 28, 884-891.
- Morikawa, T.S., Sagawa, S., Torii, R., Endo, Y., Yamazaki, F., & Shiraki, K. (2001). Hypovolemic intolerance to lower body negative pressure in female runners. *Medicine and Science in Sports and Exercise*, 33, 2058-2064.
- Nevill, A.M., Ramsbottom, R., & Williams, C. (1992). Scaling physiological measurements for individuals of different body size. *European Journal of Applied Physiology*, 65, 110-117.

- Saunders, P.U., Pyne, D.B., Telford, R.D., Peltola, E.M., Cunningham, R.B., & Hawley, J.A.. (2004). Reliability and variability of running economy in elite distance runners. *Medicine and Science in Sports and Exercise*, 36, 1972-1976.
- Vanderburgh, P.M., & Batterham, A.M. (1999). Validation of the Wilks Powerlifting Formula. *Medicine and Science in Sports and Exercise*, 31, 1869-1875.
- Vanderburgh, P.M., & Mahar, M.T. (1995). Scaling of 2-mile run times by body weight and fat-free weight in college-age men. *Journal of Strength and Conditioning Research*, 9, 67-70.
- Walker, J.L., Murray, T.D., Jackson, A.S., Morrow, J.R., & Michaud, T.J. (1999). The energy cost of horizontal walking and running in adolescents. *Medicine and Science in Sports and Exercise*, 31, 311-322.

Table 1. Four Hypothetical Examples of Runners with Handicaps for Age, BW, or Both

| Subject | Gender | RT (s) | Age (years) | BW (kg) | RT_{age} (s) | RT_{BW} (s) | RT_{adj} (s) |
|----------------|---------------|-------------------|------------------------|--------------------|---------------------------------|--------------------------------|---------------------------------|
| A | M | 1080 | 27 | 68 | 1071 | 1064 | 1055 |
| B | M | 1212 | 43 | 89 | 1128 | 1091 | 1005 |
| C | F | 1104 | 28 | 55 | 1090 | 1069 | 1056 |
| D | F | 1315 | 39 | 79 | 1229 | 1129 | 1055 |

RT (s) = actual run time

BW (kg) = body weight

RT_{age} = RT adjusted by Age only


RT_{BW} = RT adjusted by BW only

RT_{adj} = Run Time (RT) adjusted by Age and BW

Figure Caption

Figure 1. The 5K Handicap (5KH) for four hypothetical women and men of varying age and body weight. Run time values are shown adjacent to each data point.

Women



Men





Figure Caption

Figure 2. Comparison of World Association of Veteran Athletes (WAVA) age handicap standards with those of the 5K Handicap which handicaps both age and weight.

Men with a 22:00 (1320 sec) 5K Actual Run Time



Women with a 25:00 (1500 sec) 5K Actual Run Time

