

1989

Pixley-Roy Hyperspaces of ω -graphs

Joe Mashburn

Follow this and additional works at: https://ecommons.udayton.edu/mth_fac_pub

 Part of the [Applied Mathematics Commons](#), [Mathematics Commons](#), and the [Statistics and Probability Commons](#)

eCommons Citation

Mashburn, Joe, "Pixley-Roy Hyperspaces of ω -graphs" (1989). *Mathematics Faculty Publications*. 28.
https://ecommons.udayton.edu/mth_fac_pub/28

This Article is brought to you for free and open access by the Department of Mathematics at eCommons. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of eCommons. For more information, please contact frice1@udayton.edu, mschlange1@udayton.edu.

PIXLEY-ROY HYPERSPACES OF ω -GRAPHS

J. D. MASHBURN

ABSTRACT. The techniques developed by Wage and Norden are used to show that the Pixley-Roy hyperspaces of any two ω -graphs are homeomorphic. The Pixley-Roy hyperspaces of several subsets of \mathbb{R}^n are also shown to be homeomorphic.

I. INTRODUCTION

Since it was introduced in 1969, the Pixley-Roy hyperspace, $\text{PR}[X]$, of a topological space X has been intensely studied with the hope of establishing how the properties of X affect those of $\text{PR}[X]$. This study has met with some success, especially in the area of cardinal functions. However, there is a class of questions which, until recently, eluded investigators: For which spaces X and Y will $\text{PR}[X]$ be homeomorphic to $\text{PR}[Y]$? For several years the only results in this area were some embedding results obtained by van Douwen [vD] and Lutzer [L]. In 1985 Wage [W] achieved a breakthrough by developing a technique for breaking up neighborhoods around points in certain spaces which allowed him to define homeomorphisms between those neighborhoods. Using this technique he was able to show that Pixley-Roy hyperspaces of spaces like R or $[0, 1]$ are homogeneous. In 1986 Norden [N] extended Wage's technique to one which broke up an entire space. With this he was able to show that the Pixley-Roy hyperspaces of any two P -graphs (one-dimensional polyhedra with a finite number of points removed) are homeomorphic. It follows that the Pixley-Roy hyperspaces of spaces like R , $[0, 1]$, and the circle are all homeomorphic. It is the purpose of this paper to use Norden's technique to show that Pixley-Roy hyperspaces of infinite, as well as finite, graphs are all the same.

Definition. A T_2 space X with no isolated points is an ω -graph if there is a countable discrete subset D of X and a countable collection \mathbf{I} of pairwise disjoint copies of $(0, 1)$ such that $X \setminus D = \bigcup \mathbf{I}$, \mathbf{I} is locally finite on X , and for every $x \in D$, $\{x\} \cup (\bigcup \{I \in \mathbf{I} : x \in \bar{I}\})$ is a neighborhood of x which can be embedded in \mathbb{R}^2 . The set D is called a dividing set for X .

The main result of this paper can be stated as follows.

Received by the editors March 10, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 54B20; Secondary 54F65.

Supported by a Summer Research Grant from the University of Dayton.

Theorem 1. *If X and Y are ω -graphs then $\text{PR}[X]$ is homeomorphic to $\text{PR}[Y]$.*

§II will consist of preliminary definitions, notation, and observations necessary for the proof of the Theorem 1. Theorem 1 will be proved in §III, and §IV will contain some related results.

We will use $\text{PR}[X]$ to denote the Pixley-Roy hyperspace of X . Our notation for the open subsets of $\text{PR}[X]$ will be standard. We will use $F[A]$ to denote the set of nonempty finite subsets of a set A , and $F'[A]$ to denote the set of all finite subsets of A . The notation " $X \approx Y$ " will mean that X is homeomorphic to Y .

II. PRELIMINARY MATTERS

Let X be an ω -graph and let X_0 be a dividing set for X . Enumerate X_0 as $\{x_n : n < \omega\}$. Let I_0 be the countable collection of pairwise disjoint copies of $(0, 1)$ whose union makes up $X \setminus X_0$. We may assume that every element of I_0 has at least one endpoint in X_0 . For each $n < \omega$ let $\mu(n)$ be the number of elements of $X \setminus X_0$ having x_n as an endpoint. For each $I \in I_0$, fix a linear structure and orientation for I . Let Q_0 be the set of all midpoints of elements of I_0 and, for each $p \in X_0$, let O_p be the component of $X \setminus Q_0$ containing p . Then Q_0 is a discrete subset of X and $O_p \cap O_q = \emptyset$ if $p \neq q$.

For each $p \in X_0$ and each $I \in I_0$ having p as an endpoint, choose a sequence of points in $I \cap O_p$ converging monotonically to p . This can be done because each element of X_0 is the endpoint of at least one element of I_0 . Let Q_1 be the set of all points of X which are elements either of Q_0 or of the sequences just chosen. Call Q_1 the 1st cut-set of X . Set $\widehat{Q}_1 = Q_1$. Let I_1 be the countable collection of pairwise disjoint copies of $(0, 1)$ whose union makes up $X \setminus (\widehat{Q}_1 \cup X_0)$. Call I_1 the set of intervals in X derived from \widehat{Q}_1 .

Assume that $n < \omega$, that Q_n is a discrete subset of $X \setminus X_0$, and that I_n is a countable collection of pairwise disjoint intervals in X . Let Q_{n+1} , the $(n+1)$ th cut-set of X , be the set of midpoints of elements of I_n and let $\widehat{Q}_{n+1} = \widehat{Q}_n \cup Q_{n+1}$. Let I_{n+1} , the set of intervals in X derived from \widehat{Q}_{n+1} , be the countable collection of pairwise disjoint copies of $(0, 1)$ whose union makes up $X \setminus (\widehat{Q}_{n+1} \cup X_0)$. Set $Q = \bigcup_{n < \omega} Q_n$.

For every $1 \leq m < \omega$ and every $n < \omega$, let $I_{m,n} = \{I \in I_m : I \subset O_{x_n}\}$. This is the set of those elements of I_m which "cluster" around x_n .

For every $1 \leq n < \omega$ let $\Sigma(n)$ be the set of sequences, σ , defined on $n+1$ such that $\sigma(0), \sigma(1) \in \omega$ and $\sigma(m) \in \{0, 1\}$ for all $1 < m \leq n$. Let $m < \omega$. Since $I_{1,m}$ is countable, it can be enumerated as $\{I_{\langle m,n \rangle} : n < \omega\}$. In this way the set I_1 is indexed by $\Sigma(1)$. Assume that the elements of $\Sigma(n)$ have been used to index the elements of I_n . Let $I \in I_{n+1}$. There is a unique $\sigma \in \Sigma(n)$ such that $I \subset I_\sigma$. If I is the left-hand half of I_σ , then let τ be the element of $\Sigma(n+1)$ such that $\tau|n+1 = \sigma$ and $\tau(n+1) = 0$ and set $I_\tau = I$. If I is the right-hand half of I_σ , then let τ be the element of $\Sigma(n+1)$ such that $\tau|n+1 = \sigma$ and $\tau(n+1) = 1$ and set $I_\tau = I$. Let $\Sigma = \bigcup_{1 \leq n < \omega} \Sigma(n)$.

The following lemma consists of observations which are immediate consequences of the previous definitions and its proof is omitted.

Lemma 2. *Let $1 \leq m \leq n < \omega$.*

1. *If $I \in \mathbf{I}_n$ then $I \cap Q_m \neq \emptyset$.*
2. *If $p \in Q_m$ then there are exactly two elements, I_1 and I_2 , of \mathbf{I}_n such that p is an endpoint of both I_1 and I_2 . Furthermore, $I_1 \cup I_2 \cup \{p\}$ is open in X .*
3. *If $I \in \mathbf{I}_m$ then there are exactly two elements of \mathbf{I}_{m+1} that are subintervals of I .*
4. *If $I_\sigma \in \mathbf{I}_n$ then there is exactly one element, $I_{\sigma|_{m+1}}$, of \mathbf{I}_m that contains I_σ .*
5. *If $\sigma \in \Sigma(1)$, $\sigma(0) = k$, and $\sigma(1) = 1$, then I_σ is the l th element of $\mathbf{I}_{1,k}$.*
6. *If $J_\sigma \in \mathbf{I}_{n,k}$ then $\sigma \in \Sigma(n)$ and $\sigma(0) = k$.*
7. *For any $n, k < \omega$, $\{\text{Int}[\text{Cl}(\cup\{I_\sigma \in \mathbf{I}_{n,k} : \sigma(1) > a\})] : a < \omega\}$ forms a local base for x_k .*

For each $p \in X$ and each $1 \leq n < \omega$ let $A_n(p) = \{I \in \mathbf{I}_n : p \in \bar{I}\}$ and let $A_n^*(p) = \cup A_n(p)$. If $p \in Q_n$ then $A(p)$ and $A^*(p)$ will denote $A_{n+1}(p)$ and $A_{n+1}^*(p)$ respectively. If $B \in \text{PR}[X]$ then set $A_n(B) = \cup_{p \in B} A_n(p)$ and $A_n^*(B) = \cup_{p \in B} A_n^*(p)$. If $B \in F[Q_n]$ then set $A(B) = \cup_{p \in B} A(p)$ and $A^*(B) = \cup_{p \in B} A^*(p)$.

Set $M_0 = \{\emptyset\}$ and, for each $1 \leq n < \omega$, let $M_n = \{E \in F(\widehat{Q}_n) : E \cap Q_m \neq \emptyset \text{ for all } 1 \leq m \leq n\}$. For $1 \leq n < \omega$ call M_n the set of elements of $\text{PR}[X]$ compatible with \widehat{Q}_n . Note that if $m > n$ and $E \in M_n$ then $E \cap Q_m = \emptyset$. Also, if $k \neq l$ then $M_k \cap M_l = \emptyset$. For each $n < \omega$ and each $E \in M_n$, let $S_E = \{A \in \text{PR}[X] : A \cap \widehat{Q}_{n+1} = E\}$. Thus, if $A \in S_E$ and $E \in M_n$, then $A \cap Q_{n+1} = \emptyset$. The set $\{S_E : E \in M\}$ where $M = \cup_{n < \omega} M_n$ is a partition of $\text{PR}[X]$ and is called the fundamental partition of $\text{PR}[X]$ based on M . If $E \in M_n$ then S_E can be written as $\{A \cup B \cup E : A \in F'[X_0] \text{ and } B \in F'[X \setminus (\widehat{Q}_{n+1} \cup X_0)]\}$. Recall that $X \setminus (\widehat{Q}_{n+1} \cup X_0) = \cup \mathbf{I}_{n+1}$.

For each $E \in M_n$ let $\widehat{F}_E = \{I \in \mathbf{I}_{n+1} : I \subset A^*(E)\}$. If $n \geq 2$, let $E' = E \setminus Q_n = E \cap \widehat{Q}_{n-1}$. If $n \geq 3$ then E'' is $E \cap \widehat{Q}_{n-2}$. If $n = 2$ then set $E'' = \emptyset$.

Now let Y be another ω -graph and let Y_0 be a dividing set for Y . Enumerate Y_0 as $\{y_n : n < \omega\}$. Then the function $\lambda : X_0 \rightarrow Y_0$ given by $\lambda(x_n) = y_n$ is a bijection. Let J_0 be a countable collection of pairwise disjoint copies of $(0, 1)$ whose union is $Y \setminus Y_0$. We may again assume that every element of J_0 has at least one endpoint in Y_0 . Let R_0 be the set of midpoints of elements of J_0 . Let $\{R_n : 1 \leq n < \omega\}$ be the collection of cut-sets for Y and set $R = \cup_{n > \omega} R_n$. Let P_n be the component of $Y \setminus R_0$ that contains y_n . For each $0 < n < \omega$ let J_n be the set of intervals of $\text{PR}[Y]$ derived from R_n , each indexed as before by the elements of Σ . Let $\{N_k : k < \omega\}$ be the collection of

sets of elements of $\text{PR}[Y]$ compatible with $\{\widehat{R}_k : k < \omega\}$ and let $\{T_E : E \in N\}$ be the fundamental partition of $\text{PR}[Y]$ based on $N = \bigcup_{k < \omega} N_k$. If $E \subset Q$ and $f : E \rightarrow R$, then f is level preserving if $f(E \cap Q_n) \subset R_n$ for all $n < \omega$.

For each $I \in \mathbf{I}_n$ and $J \in \mathbf{J}_n$ there is a unique linear homeomorphism between I and J that preserves orientation. Denote this homeomorphism by $\eta_{I,J}$. If $\sigma, \tau \in \Sigma(n)$, $I = I_{\sigma \upharpoonright m+1}$, and $J = J_{\tau \upharpoonright m+1}$ for some $m < n$, then $\eta_{I,J}(I_\sigma) = J_\tau$ if and only if $\sigma(k) = \tau(k)$ for all $m < k \leq n$. If $\Gamma : \mathbf{I}_n \rightarrow \mathbf{J}_n$ is a bijection, then $\Gamma^* : \bigcup \mathbf{I}_n \rightarrow \bigcup \mathbf{J}_n$ is the function $\bigcup_{I \in \mathbf{I}_n} \eta_{I, \Gamma(I)}$. Γ^* is a homeomorphism that is linear and orientation preserving on each element of \mathbf{I}_n .

Now order each \mathbf{I}_n and \mathbf{J}_n lexicographically using the indices of their elements. These collections then have order-type ω^2 . Let $F \subset \mathbf{I}_n$ and $G \subset \mathbf{J}_n$ be equipotent finite sets and let $\gamma : F \rightarrow G$ be a bijection. Then $\mathbf{I}_n \setminus F$ and $\mathbf{J}_n \setminus G$ still have order-type ω^2 , so there is a unique order isomorphism $\Delta_F : \mathbf{I}_n \setminus F \rightarrow \mathbf{J}_n \setminus G$. Define $\Gamma : \mathbf{I}_n \rightarrow \mathbf{J}_n$ by $\Gamma = \gamma \cup \Delta_F$. Then Γ is a bijection.

In those situations where more than one F is being considered and subscripts are used to distinguish the various set, the same subscripts will be used to distinguish the corresponding γ, Δ , and Γ functions. For example, the functions associated with F_1 will be γ_1, Δ_1 , and Γ_1 .

It will be necessary in what follows to compare the index of I_σ with that of $\gamma(I_\sigma)$ or $\Gamma(I_\sigma)$. In order to facilitate this, we will use $\gamma(\sigma)$ and $\Gamma(\sigma)$ to denote the indices of $\gamma(I_\sigma)$ and $\Gamma(I_\sigma)$ respectively.

The next lemma is obvious and its proof is omitted.

Lemma 3. *Let $m \leq n < \omega$ and let $F_1 \subset \mathbf{I}_m$ and $F_2 \subset \mathbf{I}_n$ with $\{I \in \mathbf{I}_n : I \subset F_1\} \subset \bigcup F_2$. If $\gamma_1 : F_1 \rightarrow \mathbf{J}_m$ is a one-to-one function and $\gamma_2 : F_2 \rightarrow \mathbf{J}_n$ is defined by $\gamma_2(I) = \Gamma_1^*(I)$, then $\Gamma_1^*(I) = \Gamma_2^*(I)$ for all $I \in \mathbf{I}_n$.*

Lemma 4. *Let $F \subset \mathbf{I}_k$ be finite and let $\gamma : F \rightarrow \mathbf{J}_k$ be a one-to-one function. Assume that there are $b, c, m < \omega$ such that*

1. $c - m > b$;
2. if $I_\sigma \in F$ then either $\sigma(1) \leq b$ or $\sigma(1) > c$;
3. if $I_\sigma \in F \cap \mathbf{I}_{k,n}$ and $m \leq \sigma(1) \leq b$ then $\gamma(I_\sigma) \in \mathbf{J}_{k,n}$ and $\gamma(\sigma)(1) \leq b$;
and
4. if $I_\sigma \in F \cap \mathbf{I}_{k,n}$ and $\sigma(1) > c$ then $\gamma(I_\sigma) \in \mathbf{J}_{k,n}$ and $\gamma(\sigma)(1) > b$.

Then $\Gamma(I_\sigma) \in \mathbf{J}_{k,n}$ and $\Gamma(\sigma)(1) > b$ for all $I_\sigma \in \mathbf{I}_{k,n}$ with $\sigma(1) > c$.

Proof. Let $n < \omega$. The elements of $\mathbf{J}_{k,n} \setminus \gamma(F)$ are the images under Δ_F of $\mathbf{I}_{k,n} \setminus F$. By conditions 2 and 3,

$$\begin{aligned} |\mathbf{F} \cap \{I_\sigma \in \mathbf{I}_{k,n} : m \leq \sigma(1) \leq c\}| &= |\mathbf{F} \cap \{I_\sigma \in \mathbf{I}_{k,n} : m \leq \sigma(1) \leq b\}| \\ &= |\{\gamma(I_\sigma) : I_\sigma \in \mathbf{I}_{k,n} \text{ and } m \leq \sigma(1) \leq b\}| \\ &\leq |\{J_\sigma \in \mathbf{J}_{k,n} : J_\sigma \in \gamma(F) \text{ and } \sigma(1) \leq b\}| \\ &= |\gamma(\mathbf{F}) \cap \{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\}|. \end{aligned}$$

Also, $|\{I_\sigma \in \mathbf{I}_{k,n} : m \leq \sigma(1) \leq c\}| \geq |\{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\}|$ because $c - m > b$. Therefore,

$$\begin{aligned} & |\{I_\sigma \in \mathbf{I}_{k,n} : m \leq \sigma(1) \leq c\} \setminus \mathbf{F}| \\ &= |\{I_\sigma \in \mathbf{I}_{k,n} : m \leq \sigma(1) \leq c\} \setminus (\mathbf{F} \cap \{I_\sigma \in \mathbf{I}_{k,n} : m \leq \sigma(1) \leq c\})| \\ &\geq |\{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\} \setminus (\gamma(\mathbf{F}) \cap \{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\})| \\ &= |\{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\} \setminus \gamma(\mathbf{F})|. \end{aligned}$$

Thus, if $J_\tau \in \mathbf{J}_{k,n}$ and $\tau(1) \leq b$ then there is $I_\sigma \in \mathbf{I}_k$ such that either $I_\sigma \in \mathbf{F}$ or $I_\sigma \in \mathbf{I}_{k,n}$ and $\sigma(1) \leq c$, and $\Gamma(I_\sigma) = J_\tau$. It follows from this and condition 4 that if $I_\sigma \in \mathbf{I}_{k,n}$ and $\sigma(1) > c$, then $\Gamma(I_\sigma) \in \mathbf{J}_{k,n}$ and $\Gamma(\sigma)(1) > b$.

Lemma 5. Let $\mathbf{F}_1, \mathbf{F}_2 \subset \mathbf{I}_k$ be finite and let $\gamma_1: \mathbf{F}_1 \rightarrow \mathbf{J}_k$ and $\gamma_2: \mathbf{F}_2 \rightarrow \mathbf{J}_k$ be one-to-one functions. Let $a, b, m < \omega$ such that

1. $b - a > m$;
2. $\{I_\sigma \in \mathbf{F}_1 : \sigma(1) \leq a\} = \{I_\sigma \in \mathbf{F}_2 : \sigma(1) \leq a\} = \mathbf{G}$; and
3. $\gamma_1(I_\sigma) = \gamma_2(I_\sigma)$ for all $I_\sigma \in \mathbf{G}$;

and that for $i = 1$ or 2 ,

4. if $J_\sigma \in \gamma_i(\mathbf{F}_i)$ then either $\sigma(1) \leq a$ or $\sigma(1) > b$;
5. if $I_\sigma \in \mathbf{F}_i$ and $\sigma(1) > b$ then $\gamma_i(\sigma)(1) > a$; and
6. for all $n < \omega$, if $J_\sigma \in \gamma_i(\mathbf{F}_i) \cap \mathbf{J}_{k,n}$ and $\gamma_i^{-1}(J_\sigma) \notin \mathbf{I}_{k,n}$ then $\sigma(1) < m$.

Then $\Gamma_1(I_\sigma) = \Gamma_2(I_\sigma)$ for all $I_\sigma \in \mathbf{I}_n$ with $\sigma(1) \leq a$.

Proof. Let $n < \omega$. By condition 2,

$$\{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \cap \mathbf{F}_1 = \mathbf{I}_{k,n} \cap \mathbf{G} = \{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \cap \mathbf{F}_2$$

and

$$\begin{aligned} \{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \setminus \mathbf{F}_1 &= \{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \setminus \mathbf{G} \\ &= \{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \setminus \mathbf{F}_2. \end{aligned}$$

By conditions 2, 3, and 4,

$$\begin{aligned} \{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\} \setminus \gamma_1(\mathbf{F}_1) &= \{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\} \setminus \gamma_1(\mathbf{G}) \\ &= \{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\} \setminus \gamma_2(\mathbf{F}_2). \end{aligned}$$

If $I_\sigma \in \mathbf{I}_{k,n} \cap \mathbf{G}$ then $\Gamma_1(I_\sigma) = \gamma_1(I_\sigma) = \gamma_2(I_\sigma) = \Gamma_2(I_\sigma)$. The values of Γ_1 and Γ_2 on $\{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \setminus \mathbf{G}$ are determined by Δ_1 and Δ_2 respectively. We can establish the equality of Γ_1 and Γ_2 on $\{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \setminus \mathbf{G}$ by showing that this set is no larger than $\{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\} \setminus \gamma_1(\mathbf{G})$. Then, since both Δ_1 and Δ_2 take the α th element of $\{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \setminus \mathbf{G}$ to the

α th element of $\{J_\sigma \in \mathbf{G}J_{k,n} : \sigma(1) \leq b\} \setminus \gamma_1(\mathbf{G})$, they must be equal.

$$\begin{aligned}
 & |\{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq b\} \setminus \gamma_1(\mathbf{G})| \\
 &= |\{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq a\} \setminus \gamma_1(\mathbf{G})| + |\{J_\sigma \in \mathbf{J}_{k,n} : a < \sigma(1) \leq b\}| \\
 & \hspace{15em} \text{(by condition 4)} \\
 &= |\{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq a\} \setminus (\{\gamma_1(I_\sigma) \in \mathbf{J}_{k,n} : I_\sigma \in \mathbf{G} \setminus \mathbf{I}_{k,n}\} \\
 & \quad \cup \{\gamma_1(I_\sigma) \in \mathbf{J}_{k,n} : I_\sigma \in \mathbf{G} \cap \mathbf{I}_{k,n}\})| \\
 & \quad + |\{J_\sigma \in \mathbf{J}_{k,n} : a < \sigma(1) \leq b\}| \\
 &\geq |\{J_\sigma \in \mathbf{J}_{k,n} : \sigma(1) \leq a\} \setminus \{\gamma_1(I_\sigma) \in \mathbf{J}_{k,n} : I_\sigma \in \mathbf{G} \cap \mathbf{I}_{k,n}\}| \\
 & \hspace{15em} \text{(by conditions 1 and 6)} \\
 &= |\{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \setminus \{I_\sigma \in \mathbf{G} \cap \mathbf{I}_{k,n} : \gamma_1(I_\sigma) \in \mathbf{J}_{k,n}\}| \\
 &\geq |\{I_\sigma \in \mathbf{I}_{k,n} : \sigma(1) \leq a\} \setminus \mathbf{G}|.
 \end{aligned}$$

III. PROOF OF THEOREM 1

Let X and Y be ω -graphs with dividing sets X_0 and Y_0 . We will use the structures and definitions developed in §II. Let $g: Q_1 \rightarrow R_1$ be a bijection such that $g(Q_1 \cap O_n) = R_1 \cap P_n$ for all $n < \omega$. Then $g(Q_0) = R_0$. For our convenience later in the proof, we will assume that the first $\mu(n)$ elements of any $\mathbf{I}_{m,n}$ are those elements of $\mathbf{I}_{m,n}$ having an element of Q_0 as an endpoint.

The homeomorphism we will define is essentially that defined by Norden in [N].

Define $\Gamma_\phi: \mathbf{I}_1 \rightarrow \mathbf{J}_1$ by $\Gamma_\phi(I_\sigma) = J_\sigma$, and $h_\phi: \bigcup \mathbf{I}_1 \rightarrow \bigcup \mathbf{J}_1$ by $h_\phi = \Gamma_\phi^*$. Then h_ϕ is a homeomorphism. Set $\theta(\phi) = \phi$.

Let $E \in M_1$. Set $f_E = g \upharpoonright E$ and $\theta(E) = f_E(E)$. Let $\mathbf{F}_E = \widehat{\mathbf{F}}_E$ and $\mathbf{F}_{\theta(E)} = \widehat{\mathbf{F}}_{\theta(E)}$. Each $I \in \mathbf{F}_E$ is adjacent to exactly one element of E and each element of E is the endpoint of exactly two elements of \mathbf{F}_E . Similarly, each element of $\mathbf{F}_{\theta(E)}$ is adjacent to exactly one element of $\theta(E)$ and each element of $\theta(E)$ is the endpoint of exactly two elements of $\mathbf{F}_{\theta(E)}$. Define $\gamma_E: \mathbf{F}_E \rightarrow \mathbf{F}_{\theta(E)}$ as follows. Let $I \in \mathbf{F}_E$ and let $p \in E$ be an endpoint of I . If p is the right-hand endpoint of I , then set $\gamma_E(I)$ equal to the element of $\mathbf{F}_{\theta(E)}$ which has $g(p)$ for its right-hand endpoint. If p is the left-hand endpoint of I , then set $\gamma_E(I)$ equal to the element of $\mathbf{F}_{\theta(E)}$ which has $g(p)$ for its left-hand endpoint. Then γ_E is a bijection. Define $h_E: (\bigcup \mathbf{I}_2) \cup E \rightarrow (\bigcup \mathbf{J}_2) \cup \theta(E)$ by $h_E = \Gamma_E^* \cup f_E$. Both Γ_E^* and f_E are bijections so h_E is a bijection. It is also a homeomorphism on $\bigcup \mathbf{I}_2$ because Γ_E^* is. Let $x \in E$ and let V be a neighborhood of $f_E(x)$ in Y . By the definition of γ_E there is a neighborhood U of x in $A^*(x) \cup \{x\}$ such that $h_E(U) \subset V$. Thus h_E is continuous at x . A similar argument shows that h_E^{-1} is continuous at $h_E(x)$, so h_E is a homeomorphism.

Let $2 \leq l < \omega$ and assume that for all $k < l$ and all $E \in M_k$,

1. $f_E: E \rightarrow \widehat{R}_k$ is a level preserving one-to-one function and $\theta(E) = f_E(E)$;
2. $F_E \subset I_{k+1}$ and $F_{\theta(E)} \subset J_{k+1}$ are finite and $\gamma_E: F_E \rightarrow F_{\theta(E)}$ is a bijection; and
3. the function $h_E: (\cup I_{k+1}) \cup E \rightarrow (\cup J_{k+1}) \cup \theta(E)$ given by $h_E = \Gamma_E^* \cup f_E$ is a homeomorphism.

Fix $E \in M_l$. Each element of $E \cap Q_l$ is the midpoint of some element of I_{l-1} and $h_{E''}$, which is defined on $\cup I_{l-1}$, takes midpoints to midpoints. Thus $h_{E''}(p) \in R_l$ for all $p \in E \cap Q_l$. Define $f_E: E \rightarrow \widehat{R}_l$ by

$$f_E(p) = \begin{cases} h_{E'}(p) & \text{if } p \in E \cap \widehat{Q}_{l-1}, \\ h_{E''}(p) & \text{if } p \in E \cap Q_l. \end{cases}$$

Then f_E is a one-to-one level preserving function. Note that if $p \in E \cap \widehat{Q}_{l-1}$ then $f_E(p) = h_{E'}(p) = f_{E'}(p)$. Extending this backward, we can see that if $1 \leq k < l$ and $p \in E \cap \widehat{Q}_k$ then $f_E(p) = f_{E \cap \widehat{Q}_k}(p)$.

Let $F_{E1} = A(E \cap Q_l)$ and $F_{\theta(E)1} = A(\theta(E) \cap R_l)$. Let $I \in F_{E1}$ and let $p \in E \cap Q_l$ be an endpoint of I . Then $f_E(p) = h_{E''}(p) \in R_l$ and $h_{E''}(p)$ is an endpoint of $h_{E''}(I)$ because $h_{E''}$ is continuous. Thus $h_{E''}(I) \in F_{\theta(E)1}$. A similar argument shows that if $h_{E''}(I) \in F_{\theta(E)1}$ then $I \in F_{E1}$.

Let $F_{E2} = \{I \in \widehat{F}_E \setminus F_{E1} : h_{E'}(I) \in \widehat{F}_{\theta(E)} \setminus F_{\theta(E)1}\}$ and let $F_{\theta(E)2} = \{J \in \widehat{F}_{\theta(E)} \setminus F_{\theta(E)1} : h_{E'}^{-1}(J) \in \widehat{F}_E \setminus F_{E1}\}$. Clearly $I \in F_{E2}$ if and only if $h_{E'}(I) \in F_{\theta(E)2}$. Set $F_E = F_{E1} \cup F_{E2}$ and $F_{\theta(E)} = F_{\theta(E)1} \cup F_{\theta(E)2}$. Define $\gamma_E: F_E \rightarrow F_{\theta(E)}$ by

$$\gamma_E(I) = \begin{cases} h_{E''}(I) & \text{if } I \in F_{E1}, \\ h_{E'}(I) & \text{if } I \in F_{E2}. \end{cases}$$

Then γ_E is a bijection.

Define $h_E: (\cup I_{l+1}) \cup E \rightarrow (\cup J_{l+1}) \cup \theta(E)$ by $h_E = \Gamma_E^* \cup f_E$. The function h_E is a bijection because Γ_E^* and f_E are bijections and is a homeomorphism on $\cup I_{l+1}$ because Γ_E^* is. If $p \in E \cap Q_l$ then $A(p) \subset F_{E1}$ and $h_E(A^*(p) \cup \{p\}) = h_{E''}(A^*(p) \cup \{p\})$. Now let $p \in E'$. If $I \in A_{l+1}(p)$ then $I \in \widehat{F}_E$. Since p is an endpoint of I and $p \in \widehat{Q}_{l-1}$, the other endpoint of I must be an element of Q_{l+1} . Hence $I \notin F_{E1}$. To show that $h_{E'}(I) \in \widehat{F}_{\theta(E)} \setminus F_{\theta(E)1}$, note that $p \in E'$ and $h_{E'}$ is continuous on $(\cup I_l) \cup E'$. So $f_E(p) = F_{E'}(p)$ is an endpoint of $h_{E'}(I)$. But $f_{E'}$ is level preserving, so $f_{E'}(p) \in \widehat{R}_{l+1}$. Again, the other endpoint of $h_{E'}(I)$ must be an element of R_{l+1} . Hence $h_{E'}(I) \in \widehat{F}_{\theta(E)} \setminus F_{\theta(E)1}$. It follows that $A_{l+1}(p) \subset F_{E2}$ and $h_E(A_{l+1}^*(p) \cup \{p\}) = h_{E'}(A_{l+1}^*(p) \cup \{p\})$. But $h_{E'}$ is a homeomorphism on $(\cup I_l) \cup E'$ and $h_{E''}$ is a homeomorphism on $\cup I_{l-1}$, so h_E is a homeomorphism on $(\cup I_{l+1}) \cup E$.

Notice that for any $k < \omega$, $E \in M_k$, $x_n \in X_0$, and $I_\sigma \in \mathbf{I}_{k,n}$, if $\Gamma_E(I_\sigma) \notin \mathbf{J}_{k,n}$ then $\sigma(1) < \mu(n)$ because only the first $\mu(n)$ elements of $\mathbf{I}_{1,n}$ have endpoints in Q_0 .

For all $n < \omega$ and all $E \in M_n$, define $H_E: S_E \rightarrow T_{\theta(E)}$ by $H_E(A) = \lambda(A \cap X_0) \cup h_E(A \setminus X_0)$. Finally, define $H: \text{PR}[X] \rightarrow \text{PR}[Y]$ by $H = \bigcup_{E \in M} H_E$. To show that H is a bijection it is sufficient to show that θ is a bijection. Let $E, D \in M$ and $E \neq D$. Then $\theta(E) = f_E(E)$ and $\theta(D) = f_D(D)$. Both f_E and f_D are level-preserving one-to-one functions, so $\theta(E) \neq \theta(D)$ if $E \in M_k$ and $D \in M_l$ and $k \neq l$. Assume that $E, D \in M_1$. Then $\theta(E) = g(E) \neq g(D) = \theta(D)$ since g is a bijection. Assume that $E, D \in M_k$ for some $k > 1$. Either $E \cap Q_k \neq D \cap Q_k$ or $E' \neq D'$. But the functions $h_{E'}$, $h_{E''}$, $h_{D'}$, and $h_{D''}$ are all one-to-one, so either $h_{E''}(E \cap Q_k) \neq h_{D''}(D \cap Q_k)$ or $h_{E'}(E') \neq h_{D'}(D')$. In either case, $\theta(E) \neq \theta(D)$.

Let $A \in S_E$ where $E \in M_k$ and let V be a neighborhood of $H(A)$ in Y . Pick $a < \omega$ such that if $I_\sigma \in \mathbf{A}_1(A)$ then $\sigma(1) \leq a$ and if $J_\sigma \in \mathbf{A}_1(H(A))$ then $\sigma(1) \leq a$. Let $m = \max\{\mu(n): \mathbf{A}_1(A) \cap \mathbf{I}_{1,n} \neq \emptyset \text{ or } \mathbf{A}_1(H(A)) \cap \mathbf{J}_{1,n} \neq \emptyset\} + 1$. Pick $b \in \omega$ such that $b - m > a$ and

$$\text{Int} \left[\text{Cl} \left(\bigcup \{J_\sigma \in \mathbf{J}_{1,n}: \sigma(1) > b\} \right) \right] \subset V$$

for all $y_n \in H(A) \cap Y_0$. Set

$$V_{y_n} = \text{Int} \left[\text{Cl} \left(\bigcup \{J_\sigma \in \mathbf{J}_{1,n}: \sigma(1) > b\} \right) \right]$$

and set $V_0 = \bigcup_{p \in H(A) \cap Y_0} V_p$. Pick $c \in \omega$ such that $c - m > b$ and if $x_n \in A \cap X_0$ and $p \in Q_1 \cap \text{Int}[\text{Cl}(\bigcup \{I_\sigma \in \mathbf{I}_{1,n}: \sigma(1) > c\})]$, then $g(p) \in V_{y_n}$. For each $x_n \in A \cap X_0$ set $U_{x_n} = \text{Int}[\text{Cl}(\bigcup \{I_\sigma \in \mathbf{I}_{1,n}: \sigma(1) > c\})]$. Let $U_0 = \bigcup_{p \in A \cap X_0} U_p$. If $A \cap X_0 = \emptyset$ then set $U_0 = \emptyset$. Pick $r \geq k + 1$ such that $h_E(\mathbf{A}_r^*(p)) \subset V$ for all $p \in A \setminus X_0$. Set $U_p = \mathbf{A}_r^*(p) \cup \{p\}$ for $p \in A \setminus X_0$ and set $U_1 = \bigcup_{p \in A \setminus X_0} U_p$. Let $U = U_0 \cup U_1$. Note that:

1. if $I_\sigma \cap U_1 \neq \emptyset$ then $\sigma(1) \leq a$;
2. if $\bar{J}_\sigma \cap (H(A) \setminus Y_0) \neq \emptyset$ then $\sigma(1) \leq a$;
3. if $I_\sigma \cap U_{x_n} \neq \emptyset$ for some $x_n \in A \cap X_0$ then $I_{\sigma+1} \in \mathbf{I}_{1,n}$ and $\sigma(1) > c$;
4. if $J_\sigma \cap V_{y_n} \neq \emptyset$ for some $y_n \in H(A) \cap Y_0$ then $J_{\sigma+1} \in \mathbf{J}_{1,n}$ and $\sigma(1) > b$;
5. if $p \in A \setminus X_0$ then $U_p \cap \hat{Q}_{k+1} \subset \{p\}$.
6. a, b, c and m satisfy condition 1 in Lemmas 4 and 5; and
7. if $I_\sigma \in \mathbf{I}_{1,n}$ and $m \leq \sigma(1)$ then $H_D(I_\sigma) \subset \bigcup \mathbf{J}_{1,n}$ for any $0 < 1 < \omega, n < \omega$, and $D \in M$.

The heart of the proof that $H([A, U]) \subset [H(A), V]$ is contained in Lemmas 6 and 7.

Lemma 6. Let $D \in M_j$ where $1 \leq j \leq k$, $D \subset U$, and $D \cap U_1 = E \cap \hat{Q}_j$. Let $C = E \cap \hat{Q}_j$. Then

1. if $p \in D \cap U_q$ for some $q \in A \cap X_0$ then $f_D(p) \in V_{\lambda(q)}$;

2. if $p \in D \cap U_q$ for some $q \in E$ then $p = q$ and $f_D(p) = F_E(p)$;
3. if $I_\sigma \in I_{j+1}$ and $\sigma(1) \leq a$ then $\Gamma_C(I_\sigma) = \Gamma_D(I_\sigma)$; and
4. if $I_\sigma \in I_{j+1,n}$, $x_n \in A \cap X_0$, and $\sigma(1) > c$, then $\Gamma_D(I_\sigma) \in J_{j+1,n}$ and $\Gamma_D(\sigma)(1) > b$.

Proof. To begin with, let us take note of three useful facts. First, since $\Gamma_\phi(I_\sigma) = J_\sigma$ for all $I_\sigma \in I_1$, if $I_\sigma \in I_{1,n}$ and $\sigma(1) > c$, then $\Gamma_\phi(I_\sigma) = J_\sigma \in J_{1,n}$ and $\Gamma_\phi(\sigma)(1) = \sigma(1) > c > b$. Also, for any j , if $p \in C$ then $f_C(p) = f_E(p)$. Furthermore, if $I_\sigma \in F_D$ then either $\sigma(1) \leq a < b$ or $\sigma(1) > c$.

Let $j = 1$. Then $D \subset Q_1$ and $D \cap U_1 = E \cap Q_1$. Let $p \in D$. If $p \in U_q$ for some $q \in A \cap X_0$, then $f_D(p) = g(p) \in V_{\lambda(q)}$. If $p \in U_q$ for some $q \in A \setminus X_0$, then $q \in E, p = q$, and $f_D(p) = g(p) = f_C(p)$.

Let $n < \omega$ and let $I_\sigma \in I_{2,n} \cap F_D$ with $\sigma(1) > c$. Let $p \in D$ be an endpoint of I_σ . Since $\sigma(1) > c$, p must be in U_{x_n} . Then $f_D(p)$, which is an endpoint of $\gamma_D(I_\sigma)$, is in V_{y_n} . Thus $\gamma_D(I_\sigma) \in J_{2,n}$ and $\gamma_D(\sigma)(1) > b > a$.

It follows from $D \cap U_1 = C$ that $F_C = \{I_\sigma \in F_D : \sigma(1) \leq a\}$. Let $I_\sigma \in F_C$. Let $p \in D$ be an endpoint of I_σ . Then p must be an element of U_1 , so $f_D(p) = f_E(p) = f_C(p)$. Thus $f_E(p)$ is an endpoint for both $\gamma_C(I_\sigma)$ and $\gamma_D(I_\sigma)$. Since both γ_C and γ_D preserve orientation, it must be true that $\gamma_C(I_\sigma) = \gamma_D(I_\sigma)$. Also, $\gamma_D(\sigma)(1) \leq a < b$ because $f_D(p) \in H(A) \setminus Y_0$.

By Lemma 4, if $I_\sigma \in I_{2,n}$ and $\sigma(1) > c$, then $\Gamma_D(I_\sigma) \in J_{2,n}$ and $\Gamma(\sigma(1)) > b$. By Lemma 5, if $I_\sigma \in I_2$ and $\sigma(1) \leq a$, then $\Gamma_D(I_\sigma) = \Gamma_C(I_\sigma)$.

Let $2 \leq j \leq k$ and assume that the lemma is valid for all $1 \leq i < j$ and all $D \in M_i$ with $D \subset U$ and $D \cap U_1 = E \cap \hat{Q}_i$. Let $D \in M_j$ with $D \subset U$ and $D \cap U_1 = E \cap \hat{Q}_j$. Then $D' \in M_{j-1}, D' \subset U$, and $D' \cap U_1 = E \cap \hat{Q}_{j-1} = C'$, so the lemma is valid for D' . If $j = 2$, then $D'' = C'' = \emptyset$. If $j > 2$, then $D'' \in M_{j-2}, D'' \subset U$, and $D'' \cap U_1 = E \cap \hat{Q}_{j-2} = C''$. Thus the lemma is valid for D'' .

Let $p \in D \cap U_{x_n}$ for some $x_n \in A \cap X_0$. If $p \in \hat{Q}_{j-1}$ then $f_D(p) = f_{D'}(p) \in V_{y_n}$. If $p \in Q_j$ then $f_D(p) = h_{D''}(p)$. Now p is the midpoint of some element I_σ of $I_{j-1,n}$ where $\sigma(1) > c$. But $\Gamma_{D''}(I_\sigma) \in J_{j-1,n}, \Gamma_{D''}(\sigma)(1) > b$, and $h_{D''}(p)$ is the midpoint of $\Gamma_{D''}(I_\sigma)$. Hence $f_D(p) \in V_{y_n}$.

Let $p \in D \cap U_q$ for some $q \in A \setminus X_0$. Then $q \in E$ and $q = p$. If $p \in \hat{Q}_{j-1}$ then $f_D(p) = f_{D'}(p) = f_E(p)$. If $p \in Q_j$ then

$$f_D(p) = h_{D''}(p) = \Gamma_{D''}^*(p) = \Gamma_{C''}^*(p) = h_{C''}(p) = f_{C''}(p) = f_E(p).$$

Let $n < \omega$ and let $I_\sigma \in F_D \cap I_{j+1,n}$ with $\sigma(1) > c$. Either $\gamma_D(I_\sigma) = \Gamma_{D'}^*(I_\sigma)$ or $\gamma_D(I_\sigma) = \Gamma_{D''}^*(I_\sigma)$. In either case, $\gamma_D(I_\sigma) \in J_{j+1,n}$ and $\gamma_D(\sigma)(1) > b > a$.

It follows from the inductive hypotheses that $F_{C1} = \{I_\sigma \in F_{D1} : \sigma(1) \leq a\}$ and $F_{C2} = \{I_\sigma \in F_{D2} : \sigma(1) \leq a\}$. Thus $F_C = \{I_\sigma \in F_D : \sigma(1) \leq a\}$. Let $I_\sigma \in F_C$. If $I_\sigma \in F_{D1}$ then $\gamma_D(I_\sigma) = \Gamma_{D''}^*(I_\sigma)$. But $\Gamma_{D''}^*(I_\sigma) = \Gamma_{C''}^*(I_\sigma)$ so $\gamma_D(I_\sigma) = \gamma_C(I_\sigma)$. If $I_\sigma \in F_{D2}$ then $\gamma_D(I_\sigma) = \Gamma_{D'}^*(I_\sigma)$. But $\Gamma_{D'}^*(I_\sigma) = \Gamma_{C'}^*(I_\sigma)$ so $\gamma_D(I_\sigma) = \gamma_C(I_\sigma)$. In either case, $\gamma_D(\sigma)(1) \leq a < b$.

By Lemma 4, if $I_\sigma \in \mathbf{I}_{j+1,n}$ and $\sigma(1) > c$, then $\Gamma_D(I_\sigma) \in \mathbf{J}_{j+1,n}$ and $\Gamma_D(\sigma)(1) > b$. By Lemma 5, if $I_\sigma \in \mathbf{I}_{j+1}$ and $\sigma(1) \leq a$, then $\Gamma_D(I_\sigma) = \Gamma_C(I_\sigma)$.

Lemma 7. *If $k \leq l, D \in M_l$, and $E \subset D \subset U$, then*

1. *if $p \in D \cap U_q$ for some $q \in A \cap X_0$ then $f_D(p) \in V_{\lambda(q)}$;*
2. *if $p \in D \cap U_q$ for some $q \in A \setminus X_0$ then $f_D(p) \in V$;*
3. *if $I_\sigma \in \mathbf{I}_{l+1,n}$ for some $x_n \in A \cap X_0$ and $\sigma(1) > c$, then $\Gamma_D(I_\sigma) \in \mathbf{J}_{l+1,n}$ and $\Gamma_D(\sigma)(1) > b$; and*
4. *if $I_\sigma \in \mathbf{I}_{l+1}$ and $\sigma(1) \leq a$ then $\Gamma_D(I_\sigma) = \Gamma_E^*(I_\sigma)$.*

Note that condition 4 implies that $\gamma_D(\sigma)(1) \leq a$ for all $I_\sigma \in \mathbf{F}_D$ with $\sigma(1) \leq a$.

Proof. The case $k = 1$ is given by Lemma 6.

Assume that $l = k + 1$. Then $D' \in M_k, D' \subset U$, and $D' \cap U_1 = E$. Also, $D'' \in M_{k-1}, D'' \subset U$, and $D'' \cap U_1 = E'$. So Lemma 6 holds for D' and D'' .

Let $p \in D \cap U_{x_n}$ for some $x_n \in A \cap X_0$. If $p \in \widehat{Q}_k$ then $f_D(p) = f_{D'}(p) \in U_{x_n}$. Let $p \in Q_1$. Then p is the midpoint of some element I_σ of $\mathbf{I}_{k,n}$ where $\sigma(1) > c$. Also, $f_D(p) = h_{D''}(p)$ and $h_{D''}(p)$ is the midpoint of $\Gamma_{D''}(I_\sigma)$. But $\Gamma_{D''}(I_\sigma) \in \mathbf{J}_{k,n}$ and $\Gamma_{D''}(\sigma)(1) > b$. Thus $f_D(p) \in U_{x_n}$.

Let $p \in D \cap U_q$ for some $q \in A \setminus X_0$. Now $U_q \cap \widehat{Q}_1 \subset \{q\}$ so $p = q$ and $p \in \widehat{Q}_k$. Thus $f_D(p) = f_{D'}(p) = f_E(p) \in V$.

Let $I_\sigma \in \mathbf{F}_D \cap \mathbf{I}_{l+1,n}$ for some $x_n \in A \cap X_0$ and let $\sigma(1) > c$. Either $\gamma_D(I_\sigma) = \Gamma_{D'}^*(I_\sigma)$ or $\gamma_D(I_\sigma) = \Gamma_{D''}^*(I_\sigma)$. In either case, $\gamma_D(I_\sigma) \in \mathbf{J}_{l+1,n}$ and $\gamma_D(\sigma)(1) > b > a$.

To show that conditions 3 and 4 hold, consider the sets $\mathbf{F} = \{I \in \mathbf{I}_{l+1} : I \subset \cup \mathbf{F}_E\}$ and $\mathbf{G} = \{I_\sigma \in \mathbf{F}_D : \sigma(1) \leq a\}$. Define γ on \mathbf{G} by $\gamma(I) = \Gamma_E^*(I)$. We will show that $\mathbf{F} \subset \mathbf{G}$. Let $I_\sigma \in \mathbf{F}$. Then $\sigma(1) \leq a$ and $I_{\sigma tk+1} \in \mathbf{F}_E$. Now $A(E) \subset A(D)$ because $E \subset D$. Also, $A(\theta(E)) \subset A(\theta(D))$. Thus $I_\sigma \in \widehat{\mathbf{F}}_D$ and $h_{D'}(I_\sigma) = h_E(I_\sigma) \in \widehat{\mathbf{F}}_{\theta(D)}$. If $I_\sigma \in \mathbf{F}_{D_1}$ then there is $p \in D \cap Q_l$ such that p is an endpoint of I_σ . Then, since $\sigma(1) \leq a$, $p \in U_1$. But $U_1 \cap Q_l = \emptyset$, so $I_\sigma \notin \mathbf{F}_{D_1}$. If $p \in D \cap Q_l$ then $p \in U_0$ and $f_D(p) \in V_0$. But $\Gamma_{D'}(\sigma)(1) \leq a$ so $h_{D'}(I_\sigma)$ cannot have an endpoint in $\theta(D) \cap R_l$. Therefore $h_{D'}(I_\sigma) \in \widehat{\mathbf{F}}_{\theta(D)} \setminus \mathbf{F}_{\theta(D)_1}$, and $I_\sigma \in \mathbf{G}$. By Lemma 3, $\Gamma(I) = \Gamma_E^*(I)$ for all $I \in \mathbf{I}_{l+1}$. If $I \in \mathbf{G}$ then $I \in \mathbf{F}_{D_2}$ so $\gamma_D(I) = \Gamma_{D'}^*(I) = \Gamma_E^*(I) = \gamma(I)$. Thus $\gamma_D(I_\sigma) \in \mathbf{J}_{l+1,n}$ and $\gamma_D(\sigma)(1) \leq a < b$ for all $I_\sigma \in \mathbf{F}_D \cap \mathbf{I}_{l+1,n}$ with $m \leq \sigma(1) \leq b$. By Lemma 4, if $I_\sigma \in \mathbf{I}_{l+1,n}$ for some $x_n \in A \setminus X_0$ and $\sigma(1) > c$, then $\Gamma_D(I_\sigma) \in \mathbf{J}_{l+1,n}$ and $\Gamma_D(\sigma)(1) > b$. By Lemma 5, $\Gamma_D(I_\sigma) = \Gamma(I_\sigma) = \Gamma_E^*(I_\sigma)$ for all $I_\sigma \in \mathbf{I}_{l+1}$ with $\sigma(1) \leq a$.

Let $l \geq k + 2$ and assume that if $j = l - 1$ or $j = l - 2$, $C \in M_j$, and $E \subset C \subset U$, then the lemma holds for C . Let $D \in M_l$ with $E \subset D \subset U$. Then $D \cap U_1 \cap \widehat{Q}_{k+1} = E$. Furthermore $D' \in M_{l-1}, E \subset D' \subset U, D'' \in M_{l-2}$, and $E \subset D'' \subset U$. Thus the lemma holds for D' and D'' .

Let $p \in D \cap U_{x_n}$ for some $x_n \in A \cap X_0$. If $p \in \widehat{Q}_{l-1}$ then $f_D(p) \subset f_{D'}(p) \in V_{y_n}$. If $p \in Q_l$ then p is the midpoint of some $I_\sigma \in \mathbf{I}_{l-1,n}$ with $\sigma(1) > c$. But $f_D(p) = h_{D''}(p)$ is the midpoint of $\Gamma_{D''}(I_\sigma)$ and $\Gamma_{D''}(I_\sigma) \in \mathbf{J}_{l-1,n}$ with $\Gamma_{D''}(\sigma)(1) > b$. Hence $f_D(p) \in V_{y_n}$.

Let $p \in D \cap U_q$ for some $q \in A \setminus X_0$. If $p \in \widehat{Q}_{l-1}$ then $f_D(p) = f_{D'}(p) \in V$. If $p \in Q_l$ then $f_D(p) = h_{D''}(p) = \Gamma_{D''}^*(p) = \Gamma_E^*(p) \in V$ because $h_E(U_q) \subset V$.

Let $I_\sigma \in \mathbf{F}_D \cap \mathbf{I}_{l+1,n}$ for some $x_n \in A \cap X_0$ and let $\sigma(1) > c$. Either $\gamma_D(I_\sigma) = \Gamma_{D'}^*(I_\sigma)$ or $\gamma_D(I_\sigma) = \Gamma_{D''}^*(I_\sigma)$. In either case, $\gamma_D(I_\sigma) \in \mathbf{J}_{l+1,n}$ and $\gamma_D(\sigma)(1) > b > a$.

To show that conditions 3 and 4 hold, consider the sets $\mathbf{F} = \{I \in \mathbf{I}_{l+1} : I \subset \bigcup \mathbf{F}_E\}$ and $\mathbf{G} = \{I \in \mathbf{F}_D : \sigma(1) \leq a\}$. Define γ on \mathbf{G} by $\gamma(I) = \Gamma_E^*(I)$. Let $I_\sigma \in \mathbf{F}$. Then $I_\sigma \in \widehat{\mathbf{F}}_D$ because $E \subset D$ and $h_{D'}(I_\sigma) = h_E(I_\sigma) \in \widehat{\mathbf{F}}_{\theta(D)}$ because $\theta(E) \subset \theta(D)$. Assume that $I_\sigma \notin \mathbf{F}_{D_1}$. Let $p \in D \cap Q_1$. We will show that $f_D(p)$ cannot be an endpoint of $h_{D'}(I_\sigma)$. If $p \in U_0$, then $f_D(p) \in V_0$. But $\Gamma_{D'}(\sigma)(1) \leq a$ so $f_D(p)$ is not an endpoint of $h_{D'}(I_\sigma)$. If $p \in U_1$ then $p \in I_\tau$ for some $I_\tau \in \mathbf{I}_{k+2}$ with $\tau(1) \leq a$. By the induction hypotheses, $f_D(p) = h_{D''}(p) = h_E(p) \in h_E(I_\tau)$. If $\sigma \upharpoonright k+2 \neq \tau$ then $I_{\sigma \upharpoonright k+2} \cap I_\tau = \emptyset$ so p cannot be an endpoint of any subinterval of $I_{\sigma \upharpoonright k+2}$. If $\sigma \upharpoonright k+2 = \tau$ then p is not an endpoint of I_σ because $I_\sigma \notin \mathbf{F}_{D_1}$. The assumption that $I_\sigma \notin \mathbf{F}_{D_1}$ also implies that $h_D(I_\sigma) = h_{D'}(I_\sigma) = h_E(I_\sigma)$. But h_E^{-1} is continuous at $h_E(p)$, so $h_E(p)$ cannot be an endpoint of $h_E(I_\sigma)$. Therefore $h_D(I_\sigma) \in \widehat{\mathbf{F}}_{\theta(D)} \setminus \mathbf{F}_{\theta(D)_1}$ and $I_\sigma \in \mathbf{F}_{D_2}$. By Lemma 3, $\gamma(I) = \Gamma_E^*(I)$ for all $I \in \mathbf{I}_{l+1}$. If $I \in \mathbf{G}$ then either $\gamma_D(I) = \Gamma_{D'}^*(I)$ or $\gamma_D(I) = \Gamma_{D''}^*(I)$. In either case, $\gamma_D(I) = \Gamma_E^*(I) = \gamma(I)$. Thus $\gamma_D(I_\sigma) \in \mathbf{J}_{l+1,n}$ and $\gamma_D(\sigma)(1) \leq a < b$ for all $I_\sigma \in \mathbf{F}_D \cap \mathbf{I}_{l+1,n}$ with $m \leq \sigma(1) \leq b$. By Lemma 4, if $I_\sigma \in \mathbf{I}_{l+1,n}$ for some $x_n \in A \cap X_0$ and $\sigma(1) > c$, then $\Gamma_D(I_\sigma) \in \mathbf{J}_{l+1,n}$ and $\Gamma_D(\sigma)(1) > b$. By Lemma 5, if $I_\sigma \in \mathbf{I}_{l+1}$ and $\sigma(1) \leq a$, then $\Gamma_D(I_\sigma) = \Gamma_E^*(I_\sigma)$.

Now let $B \in [A, U]$ and let $B \in S_D$. Then $D \in M_l$ for some $l \geq k$ and $E \subset D \subset U$. Also, $B \cap X_0 = A \cap X_0$ so $\lambda(B \cap X_0) = \lambda(A \cap X_0) \subset V$. Let $p \in B \setminus X_0$. If $p \in D$ then $f_D(p) \in V$ by Lemma 7. Assume that $p \notin D$. There is $I_\sigma \in \mathbf{I}_{l+1}$ such that $p \in I_\sigma$. If $p \in U_{x_n}$ for some $x_n \in A \cap X_0$ then $I_\sigma \in \mathbf{I}_{l+1,n}$ and $\sigma(1) > c$. By Lemma 7, $h_D(I_\sigma) = \Gamma_D^*(I_\sigma) \in \mathbf{J}_{l+1,n}$ and $\Gamma_D^*(\sigma)(1) > b$. Thus $h_D(p) \in V$. If $p \in U_q$ for some $q \in A \setminus X_0$ then $\sigma(1) \leq a$. By Lemma 7, $h_D(I_\sigma) = \Gamma_D^*(I_\sigma) = \Gamma_E^*(I_\sigma)$. Thus $h_D(p) \in V$ because $h_E(U_q) \subset V$. Therefore $H(B) \in [H(A), V]$ and H is continuous. A similar argument shows that H^{-1} is continuous.

IV. RELATED RESULTS

Corollary 8. *If X and Y are ω -graphs and D and E are equipotent discrete subsets of X and Y respectively, then $\bigcup_{p \in D}[p, X]$ is homeomorphic to $\bigcup_{p \in E}[p, Y]$.*

Proof. Extend D and E to dividing sets X_0 and Y_0 of X and Y . Order the sets X_0 and Y_0 so that $\lambda(D) = E$. Then the homeomorphism defined in the proof of Theorem 1 takes $\bigcup_{p \in D}[p, X]$ to $\bigcup_{p \in E}[p, Y]$, so these two sets are homeomorphic.

The finally results are about spaces other than graphs or ω -graphs. Theorem 2 of [N] shows that points may be removed from certain T_1 spaces without affecting its Pixley-Roy hyperspace. The next three lemmas generalize this result. Theorem 12 applies this procedure to \mathbf{R}^n .

Lemma 9. *If $\langle Z_n : n < \omega \rangle$ is a sequence of disjoint homeomorphic open and closed subsets of $\text{PR}[X]$ such that $\bigcup_{n < \omega} Z_n$ is open and closed in $\text{PR}[X]$, then $\text{PR}[X] \setminus Z_0 \approx \text{PR}[X]$.*

Proof. For each $n < \omega$ let $H_n : Z_n \rightarrow Z_{n+1}$ be a homeomorphism. Define $H : \text{PR}[X] \rightarrow \text{PR}[X] \setminus Z_0$ by

$$H(A) = \begin{cases} A & \text{if } A \notin \bigcup_{n < \omega} Z_n, \\ H_n(A) & \text{if } A \in Z_n. \end{cases}$$

Then H is a homeomorphism.

Lemma 10. *If U is an open subset of space X and C is closed in U then $\bigcup_{p \in C}[p, U]$ is open and closed in $\text{PR}[X]$.*

Proof. Clearly $\bigcup_{p \in C}[p, U]$ is an open subset of $\text{PR}[X]$. Let

$$A \in U \setminus \bigcup_{p \in C}[p, U].$$

If $A \notin U$ then $[A, X]$ is a neighborhood of A that misses $\bigcup_{p \in C}[p, U]$. If $A \subset U$ then $A \cap C = \emptyset$, so $[A, U \setminus C]$ is a neighborhood of A in $\text{PR}[X]$ that misses $\bigcup_{p \in C}[p, U]$.

Lemma 11. *Let $\langle U_n : n < \omega \rangle$ be a sequence of disjoint open subsets of a space X and let $\langle C_n : n < \omega \rangle$ be a sequence of subsets of X such that $C_n \subset U_n$ and C_n is closed in U_n for all $n < \omega$. Then $\bigcup_{n < \omega} \bigcup_{p \in C_n}[p, U_n]$ is open and closed in $\text{PR}[X]$.*

Proof. It is clear that $\bigcup_{n < \omega} \bigcup_{p \in C_n}[p, U_n]$ is open in $\text{PR}[X]$. By Lemma 10, each $\bigcup_{p \in C_n}[p, U_n]$ is closed in $\text{PR}[X]$. Let $A \in \text{PR}[X]$. Since A is finite and the U_n 's are disjoint, there is a finite subset B of ω such that $A \cap U_n \neq \emptyset$ if and only if $n \in B$. Then $(\bigcup_{m \in B}[A, U_m]) \cap (\bigcup_{p \in U_n}[p, U_n]) \neq \emptyset$ only if $n \in B$. Thus $\{\bigcup_{p \in C_n}[p, U_n] : n < \omega\}$ is locally finite, and $\bigcup_{n < \omega} \bigcup_{p \in C_n}[p, U_n]$ is closed.

Theorem 12. *Let $0 < n < \omega$ and let $X = \{\bar{x} \in \mathbf{R}^n : 0 < |\bar{x}| < 1\}$ where $|\bar{x}|$ denotes the Euclidean norm. For any $0 < m < \omega$,*

$$\text{PR}[\mathbf{R}^n] \approx \text{PR}[m \times \mathbf{R}^n] \approx \text{PR}[\omega \times \mathbf{R}^n] \approx \text{PR}[m \times X] \approx \text{PR}[\omega \times X].$$

Proof. We will show that each of these spaces is homeomorphic to $\text{PR}[\mathbf{R}^n]$. Let D be a discrete subset of $\{x \in \mathbf{R} : x \geq 0\}$ which contains 0 and let $\pi : \mathbf{R}^n \rightarrow \mathbf{R}$

be the projection onto the first coordinate. Let $L = \{\bar{x} \in \mathbb{R}^n : \pi(\bar{x}) \in D\}$ and let $C = \{\bar{x} \in \mathbb{R}^n : |\bar{x}| \in D\}$. If D is finite then $\mathbb{R}^n \setminus L = (|D| + 1) \times \mathbb{R}^n$ and $\mathbb{R}^n \setminus C = |D| \times X$. If D is infinite then $\mathbb{R}^n \setminus L \approx \omega \times \mathbb{R}^n$ and $\mathbb{R}^n \setminus C \approx \omega \times X$. Let $U_0 = \mathbb{R}^n$ and let $\langle U_k : 0 < k < \omega \rangle$ be a sequence of disjoint open balls in \mathbb{R}^n , each of which has empty intersection with L and C .

Set $C_0 = L$. For every $0 < k < \omega$ let C_k be a subset of U_k which is homeomorphic to L . Then C_k is closed in U_k for all $k < \omega$. For each $k < \omega$ set $Z_k = \bigcup_{p \in C_k} [p, U_k]$. By Lemma 10, each Z_k is open and closed in $\text{PR}[\mathbb{R}^n]$. By Lemma 11, $\bigcup_{0 < k < \omega} Z_k$ is open and closed in $\text{PR}[\mathbb{R}^n]$, so $\bigcup_{k < \omega} Z_k$ is open and closed in $\text{PR}[\mathbb{R}^n]$. Clearly each Z_k is homeomorphic to every other Z_k , so $\text{PR}[\mathbb{R}^n] \approx \text{PR}[\mathbb{R}^n] \setminus Z_0 \approx \text{PR}[\mathbb{R}^n \setminus L]$. If D is finite then $\text{PR}[\mathbb{R}^n] \approx \text{PR}[(|D| + 1) \times \mathbb{R}^n]$. If D is infinite then $\text{PR}[\mathbb{R}^n] \approx \text{PR}[\omega \times \mathbb{R}^n]$.

Now let $C_0 = C$ and for every $k < \omega$ let C_k be a subset of U_k homeomorphic to C . Set $Z_k = \bigcup_{p \in C_k} [p, U_k]$ for all $k < \omega$. Again, $\langle Z_k : k < \omega \rangle$ is a sequence of disjoint homeomorphic open and closed subsets of $\text{PR}[\mathbb{R}^n]$ so $\text{PR}[\mathbb{R}^n] \approx \text{PR}[\mathbb{R}^n] \setminus Z_0 \approx \text{PR}[\mathbb{R}^n \setminus C]$. If D is finite then $\text{PR}[\mathbb{R}^n] \approx \text{PR}[|D| \times X]$. If D is infinite then $\text{PR}[\mathbb{R}^n] \approx \text{PR}[\omega \times X]$.

BIBLIOGRAPHY

- [D] Peg Daniels, *Pixley-Roy spaces over subsets of the reals*, *Topology Appl.* **29** (1988), 93–106.
- [vD] E. K. van Douwen, *The Pixley-Roy topology on spaces of subsets*, *Set-Theoretic Topology*, Academic Press, New York, 1977, pp. 111–134.
- [L] D. J. Lutzer, *Pixley-Roy topology*, *Topology Proc.* **3** (1978), 139–158.
- [N] Jeffrey Norden, *Pixley-Roy spaces over graphs*, *Topology Appl.* (to appear).
- [W] M. L. Wage, *Homogeneity of Pixley-Roy spaces*, *Topology Appl.* **28** (1987), 45–57.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DAYTON, DAYTON, OHIO 45469