

Andrews University Digital Commons @ Andrews University

Faculty Publications

Biology

2014

Feeding Interactions Between Juvenile and Adult Flightless Cormorants.

James Hayward Andrews University, hayward@andrews.edu

Libby C. Megna Andrews University

Brianna G. Payne Andrews University

Follow this and additional works at: http://digitalcommons.andrews.edu/biology-pubs Part of the <u>Biology Commons</u>

Recommended Citation

Hayward, James; Megna, Libby C.; and Payne, Brianna G., "Feeding Interactions Between Juvenile and Adult Flightless Cormorants." (2014). *Faculty Publications*. 4. http://digitalcommons.andrews.edu/biology-pubs/4

This Article is brought to you for free and open access by the Biology at Digital Commons @ Andrews University. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Digital Commons @ Andrews University. For more information, please contact repository@andrews.edu.

FEEDING INTERACTIONS BETWEEN JUVENILE AND ADULT FLIGHTLESS CORMORANTS

JAMES L. HAYWARD, LIBBY C. MEGNA & BRIANNA G. PAYNE

Department of Biology, Andrews University, 4280 Administration Drive, Berrien Springs, MI 49104-0410, USA (hayward@andrews.edu)

Submitted 1 July 2013, accepted 8 Oct 2013

We report observations on chick feedings by adult Flightless Cormorants Phalacrocorax harrisi, indicating that, contrary to the literature, the sequence of interaction is similar to that of other Pelecaniformes. This species is among the rarest of seabirds, breeding in scattered colonies along the coastlines of Isla Fernandina and Isla Isabela, Galápagos, Ecuador (Harris 1974; Rosenberg et al. 1990). It is listed as vulnerable on the IUCN Red List of Threatened Species (IUCN 2013), and is the only flightless species of its 27-member genus (Livezey 1992, Johnsgard 1993). Flightless Cormorants exhibit more than twice the mass of other cormorants (Wilson 2008). Despite their large size and locally conspicuous presence, they were overlooked by Darwin (1845) during his celebrated visit to the Galápagos Islands in 1835. The first published descriptions of Flightless Cormorants were provided over a half century later by Rothschild (1898) and Rothschild & Hartert (1899, 1902).

Snodgrass & Heller (1904) reported on feeding interactions between juvenile and adult Flightless Cormorants: "A large, immature bird may often be seen pursuing an adult through the surf with loud cries and savage thrusts of the beak, until the latter comes to terms, thrusts its beak into the open mouth of the young and disgorges into it a mass of partially digested food." Such an occurrence would be surprising, however: a young cormorant, like other juvenile Pelecaniformes, typically feeds by reaching "directly into its parent's mouth for food" (Johnsgard 1993, p. 116; see drawings pp. 116–119).

We studied environmental factors influencing the behavior of Flightless Cormorants at a colony of over 30 individuals at Cabo Douglas, Isla Fernandina, from 28 April to 17 May 2011 (Hayward *et al.* 2013). Several times each day, we observed feeding events involving adults and large juveniles. When an adult cormorant caught prey, a juvenile swam rapidly toward the adult, flapping its wings and screaming loudly. Upon reaching the adult, the smaller and darker juvenile (see Snow 1966 for descriptions of plumage differences between adults and juveniles) thrust its head and upper

neck down the throat of the forager (Fig. 1), but not *vice versa* as reported by Snodgrass & Heller (1904). Often two pursuers, perhaps siblings, competed to accomplish this feat. Magnificent Frigatebirds *Fregata magnificens* and Brown Pelicans *Pelecanus occidentalis*, attracted by these frenzied events, often pinched the head or neck of the adult from overhead while the head and neck of the juvenile were down the throat of the adult (Fig. 1C). Sometimes these piracy attempts were successful.

Our observations and photographs clearly demonstrate that it was the juveniles that thrust their bills down the throats of the larger adults. We have found only two references to juvenile feeding published since Snodgrass & Heller's (1904) description, both of which are uninformative on this point: Snow (1966) noted that the young she observed were "fed by a parent." She described food-begging by juveniles, but did not indicate whether the juvenile thrust its head down the adult's throat or *vice versa*. Harris (1979) noted only that juveniles up to at least nine months of age are fed by adults, although juveniles are capable of feeding themselves by that age.

Snodgrass & Heller's (1904) mistaken interpretation is not surprising, given that much splashing accompanies these frenzied feedings. Only digital photographs and repeated observations enabled us to determine that adult-to-juvenile feeding behavior by these rarest of all cormorants is consistent with that of other Pelecaniformes.

ACKNOWLEDGEMENTS

We thank Galápagos National Park Service for permission to work on Isla Fernandina; Sonia Cisneros, Roberto Pepolas, and other personnel at the Charles Darwin Research Station for logistical support; and Lenin Cruz and his crew for safe passage to and from Isla Fernandina on *La Pirata*. Financial support was provided by Andrews University Faculty Grants to JLH and Shandelle M. Henson. This publication is contribution number 2077 of the Charles Darwin Foundation for the Galápagos Islands.



Fig. 1. Three feeding events between adult and juvenile Flightless Cormorants. A) A juvenile (right) thrusts its head and neck down the throat of adult (middle), and a second juvenile (left) looks on. B) Note the light-colored ventral surface of the adult (right) visible during this second event. C) A Magnificent Frigatebird pinches the united heads of juvenile (left) and adult (right) during a third event.

REFERENCES

- IUCN. 2013. IUCN red list of threatened species. Version 2013.1. [Available online at: http://www.iucnredlist.org/; accessed 1 July 2013]
- DARWIN, C. 1845. Journal of researches into the natural history and geology of the countries visited during the voyage of the H.M.S. Beagle round the world under the command of Capt. Fitz Roy, R.A. Second edition, corrected, with additions. London: John Murray.
- HARRIS, M.P. 1974. A complete census of the Flightless Cormorant (*Nannopterum harrisi*). *Biological Conservation* 6: 188–191.
- HARRIS, M.P. 1979. Population dynamics of the Flightless Cormorant *Nannopterum harrisi*. *Ibis* 121: 135–146.
- HAYWARD, J.L., MEGNA, L.C., PAYNE, B.G., VELASTEGUI CHÁVEZ, S.R. & HENSON, S.M. 2013. Temporal and environmental effects on the behavior of Flightless Cormorants. *Wilson Journal of Ornithology* 125: 790–799.
- JOHNSGARD, P.A. 1993. Cormorants, Darters, and Pelicans of the World. Washington, DC: Smithsonian Institution Press.
- LIVEZEY, B.C. 1992. Flightlessness in the Galápagos Cormorant (*Compsohalieus* [*Nannopterum*] *harrisi*)—heterochrony, giantism and specialization. *Zoological Journal of the Linnean Society* 105: 155–224.

- ROSENBERG, D.K, VALLE, C.A., COULTER, M.C. & HARCOURT, S.A. 1990. Monitoring Galapagos Penguins and Flightless Cormorants in the Galápagos Islands. *Wilson Bulletin* 102: 525–532.
- ROTHSCHILD, W. 1898. Phalacrocorax harrisi. Bulletin of the British Ornithological Club 7: 52.
- ROTHSCHILD, W. & HARTERT, E. 1899. A review of the ornithology of the Galápagos Islands. With notes on the Webster-Harris expedition. *Novitates Zoologicae* 6: 5–205.
- ROTHSCHILD, W. & HARTERT, E. 1902. Further notes on the fauna of the Galápagos Islands. *Novitates Zoologicae* 9: 373–418.
- SNODGRASS, R. & HELLER, E. 1904. Papers of the Hopkins-Stanford Galapagos Expedition, 1899, XVI: Birds. *Proceedings* of the Washington Academy of Sciences 5: 231–372.
- SNOW, B.K. 1966. Observations on the behavior and ecology of the Flightless Cormorant Nannapterum harrisi. Ibis 108: 265–280.
- WILSON, R.P., VARGAS, F.H., STEINFURTH, A., RIORDAN, P., ROPERT-COUDERT, Y. & MACDONALD, D.W. 2008. What grounds some birds for life? Movement and diving in the sexually dimorphic Galápagos cormorant. *Ecological Monographs* 78: 633–652.