



# Effects of 1-year anti-TNF- $\alpha$ therapies on bone mineral density and bone biomarkers in rheumatoid arthritis and ankylosing spondylitis

Katalin Gulyás<sup>1,2</sup> · Ágnes Horváth<sup>1</sup> · Edit Végh<sup>1</sup> · Anita Pusztai<sup>1</sup> · Ágnes Szentpétery<sup>1,3,4</sup> · Zsófia Pethö<sup>1</sup> · Andrea Váncsa<sup>1</sup> · Nóra Bodnár<sup>1</sup> · Péter Csomor<sup>1</sup> · Attila Hamar<sup>1</sup> · Levente Bodoki<sup>1</sup> · Harjit Pal Bhattoa<sup>5</sup> · Balázs Juhász<sup>6</sup> · Zoltán Nagy<sup>1</sup> · Katalin Hodosi<sup>1</sup> · Tamás Karosi<sup>7</sup> · Oliver FitzGerald<sup>4</sup> · Gabriella Szücs<sup>1</sup> · Zoltán Szekanecz<sup>1</sup> · Szilvia Szamosi<sup>1</sup> · Sándor Szántó<sup>1,2</sup>

Received: 17 May 2019 / Revised: 2 September 2019 / Accepted: 3 September 2019  
© The Author(s) 2019

## Abstract

**Objectives** Rheumatoid arthritis (RA) and ankylosing spondylitis (AS) have been associated with generalized and localized bone loss. We conducted a comprehensive study using imaging (dual-energy X-ray absorptiometry, DXA) and laboratory biomarkers in order to determine bone health and to study the effects of anti-tumor necrosis factor (TNF) biologics in RA and AS.

**Patients and methods** Thirty-six RA and 17 AS patients undergoing 1-year etanercept (ETN) or certolizumab-pegol (CZP) therapy were studied. Bone density was assessed by DXA at baseline and after 12 months. Serum C-reactive protein (CRP), calcium, phosphate, parathyroid hormone (PTH), vitamin D3, osteocalcin, procollagen type I N-propeptide (P1NP), C-terminal telopeptide ( $\beta$ CTX), osteoprotegerin, sclerostin (SOST), Dickkopf-1 (DKK-1), soluble receptor activator nuclear kappa B ligand (sRANKL), and cathepsin K (cathK) levels were determined at baseline and after 6 and 12 months.

**Results** TNF- $\alpha$  inhibition was clinically effective. Anti-TNF- $\alpha$  halted further bone loss over 1 year. In general, anti-TNF therapy significantly increased P1NP, SOST levels, and the P1NP/ $\beta$ CTX ratios, while decreased DKK-1 and CathK production at different time points in most patient subsets. In the full cohort and in RA, baseline and/or 12-month bone mineral density (BMD) at multiple sites exerted inverse relationships with CRP and  $\beta$ CTX, and positive correlation with SOST. In AS, L2-4 BMD after 1-year biologic therapy inversely correlated with baseline  $\beta$ CTX, while femoral neck BMD rather showed inverse correlations with CRP.

**Conclusions** Anti-TNF therapy slowed down generalized bone loss, in association with clinical improvements, in both diseases. TNF blockade may enhance bone formation and suppress joint destruction. Anti-TNF therapy may act inversely on DKK-1 and SOST. Independent predictors of BMD were SOST and  $\beta$ CTX in RA, whilst CRP in AS.

## Key Points

- One-year anti-TNF therapy halted generalized bone loss in association with clinical improvement in arthritides.
- Anti-TNF therapy may inversely act on DKK-1 and SOST.
- Independent predictors of BMD were SOST and  $\beta$ CTX in RA, while CRP in AS.

**Keywords** Biologics · Bone loss · DKK-1 · Erosion · JAK inhibitors · Osteoporosis · Osteoprotegerin · RANKL · Rheumatoid arthritis · Sclerostin · Spondyloarthritis · Syndesmophyte

Katalin Gulyás and Ágnes Horváth Shared first authorship with equal contribution.

Szilvia Szamosi and Sándor Szántó Shared last authorship with equal contribution.

✉ Zoltán Szekanecz  
szekanecz.zoltan@med.unideb.hu; <http://www.rheumatology.hu>

Extended author information available on the last page of the article

## Introduction

Rheumatoid arthritis (RA) and ankylosing spondylitis (AS) have been associated with osteoporosis, as well as localized inflammatory bone resorption and/or formation [1, 2]. Low bone mineral density (BMD) and increased fracture risk have been observed in two-third of these patients [1, 2]. The RANK-RANKL system is highly involved in inflammatory bone resorption [3, 4]. Among cytokines, tumor necrosis

factor  $\alpha$  (TNF- $\alpha$ ) induces bone loss via the stimulation of RANKL [1, 3–5]. TNF- $\alpha$  also directly induces osteoclast function [6, 7]. Interleukin-1 (IL-1), IL-6, and IL-17 also have similar effects [4]. Osteoprotegerin (OPG), a decoy receptor of RANKL, influences bone erosions in arthritides [8]. A low OPG/RANKL ratio has been associated with increased radiographic damage in RA [9, 10]. In clinical trials, TNF- $\alpha$  inhibitors inhibited bone destruction and reduced radiological progression in RA [11–13].

With respect to bone formation, wingless-related integration site (Wnt) proteins have emerged as central regulators. Upon osteoblast activation, a multi-molecular complex is formed.  $\beta$ -catenin becomes free, it stabilizes and it is transported to the nucleus resulting in the transcription of Wnt-induced genes regulating bone formation. This process can be inhibited by Dickkopf-1 (DKK-1) and sclerostin (SOST). DKK-1 and SOST are able to block the formation of the molecular complex mentioned above and this complex will fall into pieces [4, 14]. There is a direct interaction between DKK-1 and SOST. Blockade of DKK-1 in animal models suppresses SOST production [15]. TNF- $\alpha$  induces DKK-1 and DKK-1 stimulates SOST production by osteocytes. Both DKK-1 and SOST block bone formation [14]. An interaction between DKK-1 and OPG was proposed in experimental arthritis models suggesting that the antiresorptive and bone formation-inducing property of DKK-1 inhibition also involves OPG [16]. As TNF- $\alpha$  induces DKK-1 and SOST and thus inhibits bone formation, in contrast to RA, there have been lots of controversies with regard to the efficacy of TNF- $\alpha$  inhibitors in halting bone formation and the development of syndesmophytes in axial AS [12, 14]. In animal models of arthritis, SOST promoted, while anti-SOST attenuated the formation of erosions [17]. In contrast, another study suggested that SOST may act differently by inhibiting rather than promoting TNF- $\alpha$ -mediated inflammatory bone loss [18]. SOST may exert diverse effects on bone turnover under inflammatory and non-inflammatory conditions.

In addition to their effects on localized bone turnover, anti-TNF therapies also influence the development and progression of generalized osteoporosis in conjunction with RA and AS [12].

Anti-TNF therapies may also influence bone turnover and the production of bone biomarkers. Anti-TNF- $\alpha$  agents were shown to increase serum osteocalcin (OC) and procollagen type I N-propeptide (P1NP) levels, which are markers of bone formation, and suppressed C-terminal telopeptide ( $\beta$ CTX) and receptor activator nuclear kappa B ligand (RANKL) levels, which are markers of bone resorption, in RA and AS (reviewed in [12]). These findings were not always consistent as some studies with TNF- $\alpha$  inhibitors showed no effects on OC, P1NP and  $\beta$ CTX levels [19, 20]. On the other hand, anti-TNF agents increased OPG/RANKL, OC/ $\beta$ CTX, and

P1NP/ $\beta$ CTX ratios [20–22]. Furthermore, TNF- $\alpha$  inhibitors also suppress DKK-1 leading to increased bone formation in RA [23, 24]. Interestingly, in one study, anti-TNF treatment increased SOST production in RA [25]. In most studies, changes in bone biomarkers by anti-TNF therapy were associated with improvements in disease activity and inflammatory markers (e.g. CRP) in RA and AS [12].

As most studies in the literature have been conducted in either RA or AS/SpA, here, we conducted a comprehensive study on the effects of anti-TNF biologics on bone density and bone markers in both RA and AS. Moreover, we assessed more than 10 bone markers in order to elucidate the complex effects of biologics on bone metabolism. Our aim was to draw a complex picture of, as well as to demonstrate the effects of anti-TNF therapy on bone health in RA and AS.

## Patients and methods

### Patients

Fifty-three patients with inflammatory arthritis (36 RA and 17 AS) were enrolled in the study. Patient characteristics are seen in Table 1. The cohort included 34 women and 19 men with mean age of  $52.0 \pm 12.1$  (range 24–83) years. Mean disease duration was  $8.5 \pm 7.9$  (range 1–44) years. Patients with active disease were recruited prior to initiating a biological therapy. Inclusion criteria included definitive diagnosis of RA or AS; high disease activity (DAS28 > 5.1, BASDAI > 4 after at least 3 months of combined conventional DMARD therapy); clinical indication of biological therapy. Exclusion criteria included acute/recent infection, standard contraindications to anti-TNF therapy, chronic renal or liver failure. None of the patients had known primary osteoporosis prior to the diagnosis of RA or AS. None of the patients received replacement vitamin D therapy at the time of inclusion. At baseline, RA patients had a mean DAS28 of  $5.00 \pm 0.86$ , while AS patients exerted mean BASDAI of  $5.79 \pm 1.19$ . All patients were biologic-naïve, started on an anti-TNF therapy at baseline and received the same biological treatment at 1 year. Clinical assessments were performed at baseline, and after 3, 6, and 12 months of therapy. Among the 36 RA patients, 20 received etanercept (ETN) 50 mg/week subcutaneous (SC) and 16 received certolizumab pegol (CZP) (400 mg at 0, 2, and 4 weeks, and thereafter 200 mg twice weekly SC). Altogether 28 RA patients were administered the TNF inhibitor in combination with methotrexate (MTX). These patients had been on MTX prior to the initiation of biologics and the MTX dose was not changed. All 17 AS patients received ETN 50 mg/week SC in monotherapy. Although most RA patients and some AS patients may have received corticosteroids prior to the study, none of the patients were on corticosteroids for at least 3 months prior to and during the study. The study was

**Table 1** Patient characteristics

	RA	AS	Total
<i>n</i>	36	17	53
Female:male	31:5	3:14	34:19
Age (mean ± SEM) (range), years	55.9 ± 9.8 (35–83)	43.6 ± 12.4 (24–72)	52.0 ± 12.1 (24–83)
Postmenopausal female patients	26	1	27
Disease duration (mean ± SEM) (range), years	9.1 ± 8.3 (1–44)	7.2 ± 7.0 (1–26)	8.5 ± 7.9 (1–44)
RF positivity, <i>n</i> (%)	26 (72)	–	–
ACPA positivity, <i>n</i> (%)	21 (58)	–	–
DAS28 (baseline) (mean ± SEM)	5.00 ± 0.86	–	–
BASDAI (baseline) (mean ± SEM)	–	5.79 ± 1.19	–
Treatment (ETN, CZP)	20 ETN, 16 CZP	17 ETN	37 ETN, 16 CZP

approved by the Hungarian Scientific Research Council Ethical Committee (approval no. 14804-2/2011/EKU). Written informed consent was obtained from each patient and assessments were carried out according to the declaration of Helsinki.

### Bone densitometry assessments

Dual energy X-ray absorptiometry examination was performed using the LUNAR Prodigy (GE-Lunar Corp., Madison, WI, USA) densitometer by a single technician during the study period. The coefficient of variation (CV) of the technique at our institute was 0.8% using the anatomical spine phantom measured daily and there was absence of machine drift during the study. The short-term in vivo precision error for L2-L4 lumbar spine is 0.012 g/cm<sup>2</sup> (LSC = 0.034 g/cm<sup>2</sup> at 95% confidence level) and femur neck is 0.013 g/cm<sup>2</sup> (LSC = 0.035 g/cm<sup>2</sup> at 95% confidence level).

### Laboratory measurements and assessment of disease activity

Blood was drawn after overnight fasting. Serum high sensitivity C reactive protein (hsCRP; normal, ≤ 5 mg/l) and IgM rheumatoid factor (RF; normal, ≤ 50 IU/ml) were measured by quantitative nephelometry (Cobas Mira Plus-Roche), using CRP and RF reagents (both Dialab). ACPA (anti-CCP) auto-antibodies were detected in serum samples using a second-generation Immunoscan-RA CCP2 ELISA test (Euro Diagnostica; normal, ≤ 25 IU/ml). The assay was performed according to the manufacturer's instructions. Disease activity of RA and AS was calculated as DAS28-ESR (3 variables) and BASDAI, respectively [26].

### Bone biomarkers

Blood was drawn after overnight fasting. Serum calcium (Ca; Roche Diagnostics; normal, 2.1–2.6 mmol/l) and

phosphate (P; Roche Diagnostics; normal, 0.8–1.45 mmol/l); parathyroid hormone (PTH; Roche Diagnostics; normal, 1.6–6.9 pmol/l); 25-hydroxy-vitamin D3 (DiaSorin; normal, ≥ 75 nmol/l); osteocalcin (OC; Roche Diagnostics; normal, < 41 µg/l), P1NP (Roche Diagnostics; normal, < 75 µg/l), βCTX (Roche Diagnostics; normal, < 0.57 µg/l), osteoprotegerin (OPG; Biomedica; median, 2.7 pmol/l), sclerostin (SOST; Biomedica; median, 24.14 pmol/l), DKK-1 (Biomedica; median, 36 pmol/l), soluble RANKL (Ampli-sRANKL; Biomedica; median, 0.14 pmol/l), and cathepsin K (cathK; Biomedica; median, 8.7 pmol/l) were determined by ELISA at baseline, 6 and 12 months after treatment initiation. The above-mentioned reference values for biochemical data were generated by the local laboratory.

### Statistical analysis

Statistical analysis was performed using SPSS (IBM 22 software). Data are expressed as the mean ± SD for continuous variables and percentages for categorical variables. Continuous variables were evaluated by paired two-tailed *t* test and the Wilcoxon test. Nominal variables were compared between groups using the chi-squared or Fisher's exact test, as appropriate. Correlations were determined by Spearman's analysis. Univariate and multiple regression analysis using the stepwise method was applied to investigate independent associations between DXA measurements (dependent variables) and laboratory parameters (independent variables). Sex and menopausal status were included in the regression analyses. The β standardized linear coefficients showing linear correlations between two parameters were determined. The B (+ 95% CI) regression coefficient indicated independent associations between dependent and independent variables during changes. *P* values < 0.05 were considered significant.

## Results

### Clinical response to anti-TNF therapy in RA and AS

TNF- $\alpha$  inhibition was highly effective in RA and AS patients. In the RA cohort ( $n = 36$ ), ETN and CZP treatment resulted in significant decreases in DAS28 after 3 months ( $3.52 \pm 0.79$ ;  $p < 0.001$ ), 6 months ( $3.13 \pm 0.84$ ;  $p < 0.001$ ) and 12 months of treatment ( $3.02 \pm 0.96$ ;  $p < 0.001$ ) compared with baseline ( $5.00 \pm 0.86$ ) (data not shown). In RA, CRP at baseline ( $16.0 \pm 19.1$  mg/l) was also significantly higher than after 3 months ( $8.5 \pm 11.3$  mg/l;  $p < 0.001$ ), 6 months ( $7.0 \pm 7.1$  mg/l;  $p = 0.005$ ) and 12 months ( $7.5 \pm 7.9$  mg/l;  $p = 0.011$ ) (data not shown).

In AS ( $n = 17$ ), BASDAI significantly decreased from  $5.79 \pm 1.19$  at baseline to  $2.04 \pm 0.89$  ( $p < 0.001$ ),  $2.00 \pm 1.03$  ( $p < 0.001$ ), and  $1.86 \pm 1.04$  ( $p < 0.001$ ) at 3, 6, and 12 months, respectively (data not shown). Moreover, CRP was also higher at baseline ( $12.5 \pm 12.0$  mg/l) than after 3 months ( $5.7 \pm 13.6$  mg/l;  $p = 0.026$ ), 6 months ( $6.3 \pm 13.5$  mg/l;  $p = 0.041$ ), and 12 months of therapy ( $4.4 \pm 6.6$  mg/l;  $p = 0.003$ ) (data not shown).


Fortunately, all RA and AS patients were responders and no biologic switch was necessary in any of these patients during the 1-year treatment period.

### Effects of anti-TNF therapy on bone mineral density

While anti-TNF- $\alpha$  treatment halted further generalized bone loss over 1 year in the whole cohort, there was no significant difference in mean L2–4 vertebral and femoral neck BMD or T-scores at baseline compared with 12 months (L2–4 vertebral BMD  $0.890 \pm 0.027$  vs  $0.889 \pm 0.025$  g/cm<sup>2</sup> and T-score  $-0.51 \pm 0.21$  vs  $-0.46 \pm 0.21$ ; femoral neck BMD  $0.842 \pm 0.020$  vs  $0.838 \pm 0.020$  g/cm<sup>2</sup> and T-score  $-0.84 \pm 0.17$  vs  $-0.89 \pm 0.17$ , respectively). Similar observations were made in the RA and AS subsets (data not shown).

### Effects of biologics on bone biomarkers


Among the bone turnover (formation and resorption) markers, in the full RA+AS cohort, one-year anti-TNF therapy significantly increased P1NP after 3 months ( $51.8 \pm 22.5$   $\mu$ g/l;  $p = 0.042$ ) and 6 months ( $53.5 \pm 27.1$   $\mu$ g/l;  $p = 0.040$ ) compared with baseline ( $46.7 \pm 19.3$   $\mu$ g/l) (Fig. 1). Similar differences in P1NP levels could be observed in the RA subset after 3 months ( $52.2 \pm 24.0$   $\mu$ g/l;  $p = 0.041$ ) and 6 months ( $56.4 \pm 29.0$   $\mu$ g/l;  $p = 0.026$ ) compared with baseline ( $45.6 \pm 19.9$   $\mu$ g/l) (Fig. 1). In AS, P1NP levels significantly increased after 12 months ( $56.9 \pm 28.8$   $\mu$ g/l;  $p = 0.035$ ) versus baseline ( $49.2 \pm 18.4$   $\mu$ g/l) (Fig. 1). At the same time, OC and  $\beta$ CTX had normal levels at all time points. Neither OC (baseline,  $20.3 \pm 8.8$   $\mu$ g/l; 12 months,  $20.5 \pm 9.8$   $\mu$ g/l) nor  $\beta$ CTX (baseline,  $0.34 \pm$



**Fig. 1** Effects of 1-year anti-TNF therapy on P1NP production. TNF inhibition significantly increased P1NP levels in the RA+AS cohort after 3 and 6 months of treatment; in RA patients after 3 and 6 months and in AS, after 12 months

$0.18$   $\mu$ g/l; 12 months,  $0.35 \pm 0.18$   $\mu$ g/l) changed significantly in the full RA+AS cohort.

With respect to the Wnt- $\beta$ -catenin pathway and its regulators, in the RA+AS cohort, DKK-1 levels were higher at baseline ( $59.7 \pm 28.6$  pmol/l) compared with the reference value described above. DKK-1 levels significantly decreased after 6 months of treatment ( $51.6 \pm 25.5$  pmol/l;  $p = 0.045$ ) compared with baseline (Fig. 2A). SOST levels were also higher at baseline ( $94.6 \pm 45.3$  pmol/l) compared with the normal



**Fig. 2** Effects of anti-TNF therapy on DKK-1 and SOST levels. Anti-TNF treatment significantly decreased DKK-1 production in the full cohort and in the RA subset after 6 months (A). TNF inhibition also increased SOST levels in the RA+AS, RA, and AS subsets after 12 months (B)

reference value. Conversely to DKK-1, SOST levels significantly increased after 12 months ( $112.4 \pm 76.1$  pmol/l;  $p = 0.035$ ) versus baseline (Fig. 2B). In RA, DKK-1 also significantly decreased after 6 months ( $52.1 \pm 26.1$  pmol/l;  $p = 0.042$ ) versus baseline ( $60.6 \pm 28.9$  pmol/l) (Fig. 2A). On the other hand, SOST significantly increased after 12 months ( $128.5 \pm 83.7$  pmol/l;  $p = 0.038$ ) versus baseline ( $107.0 \pm 47.5$  pmol/l) (Fig. 2B). In AS, SOST significantly increased after 12 months ( $81.3 \pm 46.9$  pmol/l;  $p = 0.034$ ) compared with baseline ( $70.6 \pm 29.0$  pmol/l) (Fig. 2B).

In the full cohort, the baseline concentrations of CathK ( $27.4 \pm 6.8$  pmol/l) were higher compared with the normal reference value above. There was a significant decrease in CathK production after 12 months ( $25.8 \pm 5.5$  pmol/l;  $p = 0.006$ ) compared with baseline (Fig. 3). In RA, CathK also significantly decreased after 12 months ( $26.9 \pm 5.6$  pmol/l;  $p = 0.012$ ) versus baseline ( $28.7 \pm 6.2$  pmol/l) (Fig. 3).

With respect to the RANKL pathway, in the full cohort or in RA or AS patients, sRANKL and OPG levels did not show any differences during anti-TNF therapy. In addition, Ca, P, vitamin D3 and PTH levels also did not change during the 1-year period (data not shown).

When calculating bone formation/resorption ratios (PINP/ $\beta$ CTX, OC/ $\beta$ CTX and OPG/RANKL), which better reflect bone turnover changes, in the full cohort, the PINP/ $\beta$ CTX ratio significantly increased at 6 months ( $187.5 \pm 85.5$ ;  $p = 0.032$ ) compared with baseline ( $160.8 \pm 56.5$ ) (Fig. 4). In RA, the PINP/ $\beta$ CTX ratio significantly increased after 6 months ( $190.2 \pm 87.2$ ;  $p = 0.035$ ) (Fig. 4). In addition, the OC/ $\beta$ CTX and OPG/RANKL ratios did not change significantly during the 12 months in the full cohort, RA or AS (data not shown).

### Correlations between bone mineral density and laboratory biomarkers


Univariate and multiple regression analyses were performed in order to determine associations between BMD at various

sites (dependent variables) and laboratory parameters (independent variables) (Table 2). In the full cohort and in RA patients, baseline and/or 12-month BMD at multiple sites exerted inverse relationships with CRP and  $\beta$ CTX, as well as positive correlations with SOST. In the AS subset, L2–4 BMD after 1-year biologic therapy inversely correlated with baseline  $\beta$ CTX. Femoral neck BMD rather showed inverse correlations with CRP. The actual data and  $p$  values are included in Table 2.


### Discussion

RA and AS have been associated with secondary osteoporosis and increased fracture risk [1, 2]. Inflammatory bone formation is also a hallmark of AS [12, 14]. Anti-TNF biologics may inhibit bone loss, but they exert limited effects on inflammatory bone formation [12]. Furthermore, biological therapy may affect the production of bone markers in RA and AS [12]. With respect to mode of action, TNF inhibitors decrease RANKL and increase OPG expression in inflammatory diseases [12]. TNF inhibition also results in blockade of DKK-1 and stimulation of Wnt-dependent bone formation [12].

In this study, we assessed the effects of ETN or CZP therapy on disease activity, bone loss and bone biomarkers in RA and AS. As expected, 1-year anti-TNF treatment was effective in both RA and AS patients as it significantly decreased DAS28 and BASDAI, respectively. These effects of ETN and CZP were observed as early as after 3 months. The anti-inflammatory effects of anti-TNF therapy were accompanied by inhibition of bone loss in both diseases. L2–4 and femoral neck BMD remained unchanged after 1-year ETN or CZP therapy in the entire cohort, as well as in RA and AS. These effects have previously been reported by others [12, 13]. Baseline CRP inversely correlated not only with baseline L2–4 and femoral neck BMD, but also with BMD after 1-year treatment. Thus, at baseline, high-grade inflammation correlated with low BMD. Moreover, we found that baseline



**Fig. 3** Effects of 1-year anti-TNF therapy on cathepsin K (CathK) production. TNF inhibition significantly decreased CathK levels in the RA+AS cohort and in RA patients after 12 months of treatment



**Fig. 4** Effects of 1-year anti-TNF therapy on PINP/ $\beta$ CTX ratios. TNF inhibition significantly increased PINP/ $\beta$ CTX in the RA+AS cohort and in RA patients after 6 months of treatment

**Table 2** Regression analysis of CRP, bone density and biomarker values

Dependent variable	Independent variable	Univariate			Multiple		
		B (95% CI)	$\beta$	<i>p</i>	B (95% CI)	$\beta$	<i>p</i>
<b>A. All patients (<i>n</i> = 53)</b>							
L2–4 BMD (baseline)	CRP (baseline)	– 0.004 (– 0.006 to – 0.001)	– 0.332	0.020			
	$\beta$ CTX (baseline)	– 0.498 (– 0.809 to – 0.187)	– 0.426	0.002	– 0.498 (– 0.809 to – 0.187)	– 0.426	0.002
	SOST (baseline)	0.001 (0–0.003)	0.342	0.025			
L2–4 BMD (12 months)	CRP (baseline)	– 0.003 (– 0.006–0)	– 0.285	0.047			
	$\beta$ CTX (baseline)	– 0.449 (– 0.750 to – 0.147)	– 0.400	0.004			
Femoral neck BMD (baseline)	CRP (baseline)	– 0.002 (– 0.005–0)	– 0.303	0.033			
	SOST (baseline)	0.001 (0–0.002)	0.308	0.042	0.001 (0–0.002)	0.308	0.042
Femoral neck BMD (12 months)	CRP (baseline)	– 0.003 (– 0.005 to – 0.001)	– 0.396	0.004	– 0.002 (– 0.005–0)	– 0.299	0.036
	$\beta$ CTX (baseline)	– 0.282 (– 0.528 to – 0.037)	– 0.317	0.025	– 0.278 (– 0.536 to – 0.020)	– 0.303	0.035
	SOST (baseline)	0.001 (0–0.002)	0.343	0.024			
<b>B. RA (<i>n</i> = 36)</b>							
L2–4 BMD (baseline)	CRP (baseline)	– 0.003 (– 0.006–0)	– 0.331	0.049			
	$\beta$ CTX (baseline)	– 0.524 (– 0.853 to – 0.194)	– 0.485	0.003	– 0.342 (– 0.666 to – 0.017)	– 0.375	0.040
	SOST (baseline)	0.002 (0.001–0.003)	0.517	0.003	0.001 (0–0.002)	0.374	0.041
L2–4 BMD (12 months)	$\beta$ CTX (baseline)	– 0.478 (– 0.081 to – 0.145)	– 0.448	0.006	– 0.335 (– 0.669 to – 0.001)	– 0.362	0.049
	SOST (baseline)	0.001 (0–0.002)	0.360	0.047	0.001 (0–0.002)	0.384	0.038
Femoral neck BMD (baseline)	SOST (baseline)	0.001 (0–0.002)	0.375	0.038	0.001 (0–0.002)	0.402	0.021
Femoral neck BMD (12 months)	CRP (baseline)	– 0.003 (– 0.005–0)	– 0.353	0.035			
	$\beta$ CTX (baseline)	– 0.324 (– 0.617 to – 0.031)	– 0.359	0.031	– 0.434 (– 0.738 to – 0.130)	– 0.457	0.007
	SOST (baseline)	0.001 (0–0.002)	0.422	0.018			
<b>C. AS (<i>n</i> = 17)</b>							
L2–4 BMD (12 months)	$\beta$ CTX (baseline)	– 0.858 (– 1.600 to – 0.117)	– 0.609	0.027			
Femoral neck BMD (baseline)	CRP (baseline)	– 0.006 (– 0.011 to – 0.001)	– 0.589	0.027	– 0.006 (– 0.011 to – 0.001)	– 0.589	0.027
Femoral neck BMD (12 months)	CRP (baseline)	– 0.006 (– 0.011 to – 0.001)	– 0.597	0.024	0.001 (– 0.007–0)	– 0.359	0.034

$\beta$ , standardized linear coefficient; B (+ 95% CI), regression coefficient

CRP may predict further bone loss after 1-year anti-TNF therapy.

The first publications on the effects of TNF inhibitors on bone and osteoporosis were published in the 2000s. In most studies, effects on the bone have been associated with favorable clinical responses. A great amount of data have been published on infliximab, ETN, adalimumab and some data also on golimumab (reviewed in [12]). We have not found any reports on the bone effects of CZP.

Regarding bone biomarkers, 1-year anti-TNF therapy significantly increased P1NP over time in our full cohort, in RA and AS. Other markers of bone turnover, such as OC or  $\beta$ CTX did not change upon anti-TNF therapy. The P1NP/ $\beta$ CTX ratio increased over time in the full cohort and in RA. Bone resorbing activity as indicated by  $\beta$ CTX at baseline was associated with low baseline BMD and we showed that baseline  $\beta$ CTX may also predict further bone loss after 1-year anti-TNF therapy. In line with our findings, previous studies found that TNF inhibitors increased serum P1NP levels and, in some cohorts,

suppressed  $\beta$ CTX levels in RA and AS [20, 21, 23, 27–32]. These findings were not always consistent as some other studies showed no effects of TNF- $\alpha$  inhibitors on OC, P1NP, and  $\beta$ CTX levels [19, 20]. Furthermore, anti-TNF agents have been shown to increase OPG/RANKL, OC/ $\beta$ CTX, and P1NP/ $\beta$ CTX ratios [20–22]. In most studies, changes in bone biomarkers by anti-TNF therapy were associated with improvements in disease activity in RA and AS [12].

With respect to DKK-1 and SOST, upon 1-year ETN or CZP treatment, DKK-1 levels transiently decreased in the whole cohort and in the RA subset. On the other hand, SOST levels significantly increased in the entire cohort, as well as in the RA and AS subsets after 12 months of treatment. When comparing RA and AS, SOST levels were lower at all time points in AS compared with RA. Baseline SOST correlated with L2–4 and femoral neck BMD at baseline, as well as after 12 months in the whole cohort and in RA. This association was further confirmed by regression analyses. Interestingly, higher baseline SOST was

associated with higher BMD at all sites. Moreover, high baseline SOST predicted higher BMD after 12 months of anti-TNF treatment. Similar to our observations, other studies in RA also reported that TNF- $\alpha$  inhibitors suppressed DKK-1 leading to increased bone formation [23, 24]. In addition, in one study, anti-TNF treatment increased SOST production in RA [25]. The seemingly controversial effects of anti-TNF biologics on DKK-1 and SOST need further discussion. In general, DKK-1 and SOST were thought to have unidirectional effects on the bone. Both molecules were described to block Wnt- $\beta$ -catenin-dependent bone formation. TNF- $\alpha$  may induce both DKK-1 and SOST production [14]. DKK-1 may directly drive SOST expression by osteocytes and DKK-1 blockade also inhibits SOST production [14, 15]. Indeed, Chen et al. [17] administered anti-SOST antibody to human TNF transgenic mice that develop inflammatory arthritis. In this model, anti-SOST did not affect synovitis. However, anti-SOST, but not anti-TNF antibodies halted the progression of bone erosions [17]. On the other hand, more recently, Wehmeyer et al. [18] reported that in a TNF- $\alpha$ -dependent murine arthritis model, where synovial fibroblasts abundantly produce SOST, SOST inhibition leads to accelerated synovitis and joint destruction. SOST blocked TNF- $\alpha$ -dependent signaling pathways and SOST inhibition ameliorated arthritis in an animal model [18]. Moreover, low SOST and DKK-1 production have been associated with syndesmophyte formation in AS [33]. SOST levels remain low after TNF- $\alpha$  blockade, which may partly explain the inefficacy of TNF inhibitors on bone formation in AS [34, 35]. Interestingly, low SOST and DKK-1 levels in AS result in increased CRP production [35]. Indeed, SOST levels increased in our study in both AS and RA; however, the absolute serum concentrations of SOST were lower in AS vs RA at all time points. Our findings that TNF- $\alpha$  inhibitors may differentially regulate DKK-1 and SOST suggest that, apart from the direct stimulatory effect of DKK-1 on SOST [14, 15], there may be other indirect regulatory mechanisms between these two molecules. Finally, in our cohort, high SOST correlated with high BMD at all sites. This finding supports previous observations that, while under non-inflammatory conditions, such as osteoporosis, SOST inhibits Wnt-mediated bone formation and contributes to bone loss and joint destruction [18, 36, 37], under TNF-dependent inflammatory conditions, SOST may preserve bone [18] and attenuate joint damage [18]. Furthermore, anti-TNF treatment may increase SOST levels [25] and, as we demonstrated in this study, high SOST may correlate with high BMD.

When comparing bone resorption indicated by  $\beta$ CTX with bone formation associated with SOST, only  $\beta$ CTX, but not OC, and only SOST, but not DKK-1 correlated with BMD. Thus, our data suggest that  $\beta$ CTX and SOST, in addition to CRP, may be the most important biomarkers and independent predictors of

BMD. Moreover, most of these associations were seen in the entire cohort, where two-third of the patients had RA, and in the RA subset alone. Interestingly only  $\beta$ CTX and SOST, but not CRP associated with BMD in RA. In contrast, in our relatively small AS subset, only CRP, but not  $\beta$ CTX or SOST correlated with femoral neck BMD. In conclusion, in RA, SOST and  $\beta$ CTX, but not CRP are independent predictors of BMD (at baseline and after 12 months), whilst, in AS, CRP may serve as the best biomarker of inflammatory bone metabolism. As mentioned above, low SOST and DKK-1 levels in AS may stimulate CRP production [35], this may further explain the outstanding importance of CRP in AS-related bone changes.

CathK is a matrix-degrading enzyme that effectively digests collagen. CathK has been implicated in osteoporosis and a CathK inhibitor, odanacatib, has been tried in postmenopausal osteoporosis [38]. In RA, increased CathK levels correlated with joint damage [39]. CathK is strongly expressed at different spinal regions in AS [40]. In our entire cohort and in the RA subset, but not in AS, TNF inhibition resulted in significantly decreased CathK levels after 1 year. In a study of 13 AS patients, infliximab, ETN, or adalimumab treatment did not change CathK levels [41]. We did not find any reports in the literature on the effects of biologics on CathK production.

Our study has advantages and limitations. We included RA and AS patients and compared the two disease. To our knowledge, this is the first study that assesses the effects of CZP on bone loss. Moreover, we determined a high number of bone biomarkers in these patients and correlated inflammatory and bone biomarkers with BMD. Finally, we developed a potential explanation on the differing effects of SOST in inflammatory vs non-inflammatory bone loss. This study certainly has limitations, primarily the relatively low number of patients. However, we assessed BMD, as well as a great number of biomarkers in a prospective manner, which would have been more difficult in a larger patient cohort.

In summary, despite its diverse effects on bone formation in RA and AS, anti-TNF therapy (ETN and CZP) slowed down bone loss in both diseases. Independent predictors of BMD were SOST and  $\beta$ CTX in RA, whilst CRP in AS. Further studies are needed to evaluate the potential beneficial effects of biologics on inflammatory bone loss.

**Funding information** Open access funding provided by University of Debrecen (DE). This research was supported by the Hungarian National Scientific Research Fund (OTKA) grant No. K 105073 (H.P.B. and Z.S.); by the European Union and the State of Hungary and co-financed by the European Social Fund in the framework of TAMOP-4.2.4.A/2-11/1-2012-0001 'National Excellence Program'(Z.S.); and by the European Union grants GINOP-2.3.2-15-2016-00015 and GINOP-2.3.2-15-2016-00050 (Z.S.).

## Compliance with ethical standards

**Disclosures** None.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Deal C (2012) Bone loss in rheumatoid arthritis: systemic, periarticular, and focal. *Curr Rheumatol Rep* 14(3):231–237
- Magrey M, Khan MA (2010) Osteoporosis in ankylosing spondylitis. *Curr Rheumatol Rep* 12(5):332–336
- Schett G, Hayer S, Zwerina J, Redlich K, Smolen JS (2005) Mechanisms of disease: the link between RANKL and arthritic bone disease. *Nat Clin Pract Rheumatol* 1(1):47–54
- Takayanagi H (2009) Osteoimmunology and the effects of the immune system on bone. *Nat Rev Rheumatol* 5(12):667–676
- Daoussis D, Andonopoulos AP, Liossis SN (2010) Wnt pathway and IL-17: novel regulators of joint remodeling in rheumatic diseases. Looking beyond the RANK-RANKL-OPG axis. *Semin Arthritis Rheum* 39(15):369–383
- Kobayashi K, Takahashi N, Jimi E, Udagawa N, Takami M, Kotake S, Nakagawa N, Kinoshita M, Yamaguchi K, Shima N, Yasuda H, Morinaga T, Higashio K, Martin TJ, Suda T (2000) Tumor necrosis factor alpha stimulates osteoclast differentiation by a mechanism independent of the ODF/RANKL-RANK interaction. *J Exp Med* 191(2):275–286
- Lam J, Takeshita S, Barker JE, Kanagawa O, Ross FP, Teitelbaum SL (2000) TNF-alpha induces osteoclastogenesis by direct stimulation of macrophages exposed to permissive levels of RANK ligand. *J Clin Invest* 106(12):1481–1488
- Schett G, Redlich K, Smolen JS (2003) The role of osteoprotegerin in arthritis. *Arthritis Res Ther* 5(5):239–245
- van Tuyl LH, Voskuyl AE, Boers M, Geusens P, Landewe RB, Dijkmans BA et al (2010) Baseline RANKL:OPG ratio and markers of bone and cartilage degradation predict annual radiological progression over 11 years in rheumatoid arthritis. *Ann Rheum Dis* 69(9):1623–1628
- Geusens PP, Landewe RB, Garnero P, Chen D, Dunstan CR, Lems WF et al (2006) The ratio of circulating osteoprotegerin to RANKL in early rheumatoid arthritis predicts later joint destruction. *Arthritis Rheum* 54(6):1772–1777
- Smolen JS, Landewe R, Bijlsma J, Burmester G, Chatzidionysiou K, Dougados M et al (2017) EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis* 76(6):960–977
- Szentpetery A, Horvath A, Gulyas K, Petho Z, Bhattoa HP, Szanto S et al (2017) Effects of targeted therapies on the bone in arthritides. *Autoimmun Rev* 16(3):313–320
- Haugeberg G, Helgetveit KB, Forre O, Garen T, Sommerseth H, Proven A (2014) Generalized bone loss in early rheumatoid arthritis patients followed for ten years in the biologic treatment era. *BMC Musculoskelet Disord* 15:289
- Schett G, Gravalles E (2012) Bone erosion in rheumatoid arthritis: mechanisms, diagnosis and treatment. *Nat Rev Rheumatol* 8(11):656–664
- Heiland GR, Zwerina K, Baum W, Kireva T, Distler JH, Grisanti M, Asuncion F, Li X, Ominsky M, Richards W, Schett G, Zwerina J (2010) Neutralisation of Dkk-1 protects from systemic bone loss during inflammation and reduces sclerostin expression. *Ann Rheum Dis* 69(12):2152–2159
- Diarra D, Stolina M, Polzer K, Zwerina J, Ominsky MS, Dwyer D, Korb A, Smolen J, Hoffmann M, Scheinecker C, van der Heide D, Landewe R, Lacey D, Richards WG, Schett G (2007) Dickkopf-1 is a master regulator of joint remodeling. *Nat Med* 13(2):156–163
- Chen XX, Baum W, Dwyer D, Stock M, Schwabe K, Ke HZ, Stolina M, Schett G, Bozec A (2013) Sclerostin inhibition reverses systemic, periarticular and local bone loss in arthritis. *Ann Rheum Dis* 72(10):1732–1736
- Wehmeyer C, Frank S, Beckmann D, Bottcher M, Cromme C, Konig U et al (2016) Sclerostin inhibition promotes TNF-dependent inflammatory joint destruction. *Sci Transl Med* 8(330):330ra35
- Marotte H, Pallot-Prades B, Grange L, Gaudin P, Alexandre C, Miossec P (2007) A 1-year case-control study in patients with rheumatoid arthritis indicates prevention of loss of bone mineral density in both responders and nonresponders to infliximab. *Arthritis Res Ther* 9(3):R61
- Chopin F, Garnero P, le Henanff A, Debiais F, Daragon A, Roux C, Sany J, Wendling D, Zarnitsky C, Ravnaud P, Thomas T (2008) Long-term effects of infliximab on bone and cartilage turnover markers in patients with rheumatoid arthritis. *Ann Rheum Dis* 67(3):353–357
- Vis M, Havaardsholm EA, Haugeberg G, Uhlig T, Voskuyl AE, van de Stadt RJ, Dijkmans BAC, Woolf AD, Kvien TK, Lems WF (2006) Evaluation of bone mineral density, bone metabolism, osteoprotegerin and receptor activator of the NF kappa B ligand serum levels during treatment with infliximab in patients with rheumatoid arthritis. *Ann Rheum Dis* 65(11):1495–1499
- Woo JH, Lee HJ, Sung IH, Kim TH (2007) Changes of clinical response and bone biochemical markers in patients with ankylosing spondylitis taking etanercept. *J Rheumatol* 34(8):1753–1759
- Wang SY, Liu YY, Ye H, Guo JP, Li R, Liu X et al (2011) Circulating Dickkopf-1 is correlated with bone erosion and inflammation in rheumatoid arthritis. *J Rheumatol* 38(5):821–827
- Garnero P, Tabassi NC, Voorzanger-Rousselot N (2008) Circulating dickkopf-1 and radiological progression in patients with early rheumatoid arthritis treated with etanercept. *J Rheumatol* 35(12):2313–2315
- Lim MJ, Kwon SR, Joo K, Son MJ, Park SG, Park W (2014) Early effects of tumor necrosis factor inhibition on bone homeostasis after soluble tumor necrosis factor receptor use. *Korean J Intern Med* 29(6):807–813
- Prevoe ML, van't Hof MA, Kuper HH, van Leeuwen MA, van de Putte LB, van Riel PL (1995) Modified disease activity scores that include twenty-eight-joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum* 38(1):44–48
- Torikai E, Kageyama Y, Takahashi M, Suzuki M, Ichikawa T, Nagafusa T, Nagano A (2006) The effect of infliximab on bone metabolism markers in patients with rheumatoid arthritis. *Rheumatology (Oxford)* 45(6):761–764
- Visvanathan S, van der Heijde D, Deodhar A, Wagner C, Baker DG, Han J, Braun J (2009) Effects of infliximab on markers of inflammation and bone turnover and associations with bone mineral density in patients with ankylosing spondylitis. *Ann Rheum Dis* 68(2):175–182
- Visvanathan S, Wagner C, Marini JC, Baker D, Gathany T, Han J et al (2008) Inflammatory biomarkers, disease activity and spinal disease measures in patients with ankylosing spondylitis after treatment with infliximab. *Ann Rheum Dis* 67(4):511–517
- Barnabe C, Hanley DA (2009) Effect of tumor necrosis factor alpha inhibition on bone density and turnover markers in patients with rheumatoid arthritis and spondyloarthritis. *Semin Arthritis Rheum* 39(2):116–122



31. Vis M, Wolbink GJ, Lodder MC, Kostense PJ, van de Stadt RJ, de Koning MH et al (2003) Early changes in bone metabolism in rheumatoid arthritis patients treated with infliximab. *Arthritis Rheum* 48(10):2996–2997
32. Serio B, Paolino S, Sulli A, Ferretti V, Cutolo M (2006) Bone metabolism changes during anti-TNF-alpha therapy in patients with active rheumatoid arthritis. *Ann N Y Acad Sci* 1069:420–427
33. Heiland GR, Appel H, Poddubnyy D, Zwerina J, Hueber A, Haibel H, Baraliakos X, Listing J, Rudwaleit M, Schett G, Sieper J (2012) High level of functional dickkopf-1 predicts protection from syndesmophyte formation in patients with ankylosing spondylitis. *Ann Rheum Dis* 71(4):572–574
34. Saad CG, Ribeiro AC, Moraes JC, Takayama L, Goncalves CR, Rodrigues MB et al (2012) Low sclerostin levels: a predictive marker of persistent inflammation in ankylosing spondylitis during anti-tumor necrosis factor therapy? *Arthritis Res Ther* 14(5):R216
35. Klingberg E, Nurkkala M, Carlsten H, Forsblad-d'Elia H (2014) Biomarkers of bone metabolism in ankylosing spondylitis in relation to osteoproliferation and osteoporosis. *J Rheumatol* 41(7):1349–1356
36. PETERS E, Boudin E, Van Hul W (2008) Wnt signaling: a win for bone. *Arch Biochem Biophys* 473(2):112–116
37. Lerner UH, Ohlsson C (2015) The WNT system: background and its role in bone. *J Intern Med* 277(6):630–649
38. Perez-Castrillon JL, Pinacho F, De Luis D, Lopez-Menendez M, Duenas LA (2010) Odanacatib, a new drug for the treatment of osteoporosis: review of the results in postmenopausal women. *J Osteoporos* 2010:1–5
39. Skoumal M, Haberhauer G, Kolarz G, Hawa G, Woloszczuk W, Klingler A (2005) Serum cathepsin K levels of patients with longstanding rheumatoid arthritis: correlation with radiological destruction. *Arthritis Res Ther* 7(1):R65–R70
40. Neidhart M, Baraliakos X, Seemayer C, Zelder C, Gay RE, Michel BA, Boehm H, Gay S, Braun J (2009) Expression of cathepsin K and matrix metalloproteinase 1 indicate persistent osteodestructive activity in long-standing ankylosing spondylitis. *Ann Rheum Dis* 68(8):1334–1339
41. Wendling D, Cedoz JP, Racadot E (2008) Serum levels of MMP-3 and cathepsin K in patients with ankylosing spondylitis: effect of TNF alpha antagonist therapy. *Joint Bone Spine* 75(5):559–562

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Affiliations

Katalin Gulyás<sup>1,2</sup> · Ágnes Horváth<sup>1</sup> · Edit Végh<sup>1</sup> · Anita Pusztai<sup>1</sup> · Ágnes Szentpétery<sup>1,3,4</sup> · Zsófia Pethő<sup>1</sup> · Andrea Vánca<sup>1</sup> · Nóra Bodnár<sup>1</sup> · Péter Csomor<sup>1</sup> · Attila Hamar<sup>1</sup> · Levente Bodoki<sup>1</sup> · Harjit Pal Bhattoa<sup>5</sup> · Balázs Juhász<sup>6</sup> · Zoltán Nagy<sup>1</sup> · Katalin Hodosi<sup>1</sup> · Tamás Karosi<sup>7</sup> · Oliver FitzGerald<sup>4</sup> · Gabriella Szücs<sup>1</sup> · Zoltán Szekanez<sup>1</sup> · Szilvia Szamosi<sup>1</sup> · Sándor Szántó<sup>1,2</sup>

<sup>1</sup> Division of Rheumatology, Department of Internal Medicine, Faculty of Medicine, University of Debrecen, Nagyerdei str 98, Debrecen 4032, Hungary

<sup>2</sup> Department of Sports Medicine, Debrecen, Hungary

<sup>3</sup> Department of Rheumatology, Uppsala University Hospital, Uppsala, Sweden

<sup>4</sup> Department of Rheumatology, St. Vincent's University Hospital and Conway Institute for Biomolecular Research, University College Dublin, Dublin, Ireland

<sup>5</sup> Department of Laboratory Medicine, Faculty of Medicine, University of Debrecen, Debrecen, Hungary

<sup>6</sup> Department of Oncology, Faculty of Medicine, University of Debrecen, Debrecen, Hungary

<sup>7</sup> Department of Otolaryngology, Borsod County Teaching Hospital, Miskolc, Hungary