

Manuscript version: Published Version

The version presented in WRAP is the published version (Version of Record).

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/127742>

How to cite:

The repository item page linked to above, will contain details on accessing citation guidance from the publisher.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution 4.0 International license (CC BY 4.0) and may be reused according to the conditions of the license. For more details see: <http://creativecommons.org/licenses/by/4.0/>.



Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

The Widom-Rowlinson model on the Delaunay graph

Stefan Adams*

Michael Eyers†

Abstract

We establish phase transitions for continuum Delaunay multi-type particle systems (continuum Potts or Widom-Rowlinson models) with a repulsive interaction between particles of different types. Our interaction potential depends solely on the length of the Delaunay edges. We show that a phase transition occurs for sufficiently large activities and for sufficiently large potential parameter proving an old conjecture of Lebowitz and Lieb extended to the Delaunay structure. Our approach involves a Delaunay random-cluster representation analogous to the Fortuin-Kasteleyn representation of the Potts model. The phase transition manifests itself in the mixed site-bond percolation of the corresponding random-cluster model. Our proofs rely mainly on geometric properties of Delaunay tessellations in \mathbb{R}^2 and on recent studies [DDG12] of Gibbs measures for geometry-dependent interactions. The main tool is a uniform bound on the number of connected components in the Delaunay graph which provides a novel approach to Delaunay Widom Rowlinson models based on purely geometric arguments. The interaction potential ensures that shorter Delaunay edges are more likely to be open and thus offsets the possibility of having an unbounded number of connected components.

Keywords: Delaunay tessellation; Widom-Rowlinson; Gibbs measures; random cluster measures; mixed site-bond percolation; phase transition; coarse graining; multi-body interaction.

AMS MSC 2010: 60G55; 60G57; 82B21; 82B05; 82B26; 82B43.

Submitted to EJP on May 24, 2018, final version accepted on October 4, 2019.

1 Introduction and results

1.1 Introduction

This paper is a continuation of the study in [AE16] concerning phase transition in continuum systems. Phase transition results in the continuum setting are much harder to obtain, e.g., the proof of a liquid-vapor phase transition in [LMP99], or the spontaneous breaking of rotational symmetry in two dimensions for a Delaunay hard-equilaterality like

*University of Warwick, Mathematics Institute, Gibbet Hill Road, CV4 7AL Coventry, United Kingdom.
E-mail: s.adams@warwick.ac.uk

†London, United Kingdom.

E-mail: m.j.eyers@googlemail.com

interaction [MR09], for an overview see [AE16]. The focus of this research is on another specific model, the Widom and Rowlinson model [WR70], for which a phase transition is known to occur, and on a novel counting technique for connected components of the corresponding Delaunay random cluster model. A Widom and Rowlinson model is a multi-type particle system in \mathbb{R}^d , $d \geq 2$, with hard-core exclusion between particles of different type. The phase transition in this model was established by Ruelle [Rue71]. Lebowitz and Lieb [LL72] extended his result by replacing the hard-core exclusion by a soft-core repulsion between unlike particles. The phase transitions for large activities in all these systems reveal themselves in breaking of the symmetry in the type-distribution.

In this paper we establish the existence of a phase transition for a class of continuum Delaunay Widom-Rowlinson (Potts) models in \mathbb{R}^2 . The repulsive interaction between unlike particles is of finite range, and it depends on the geometry of the Delaunay tessellation, i.e., the length of the edges. The potential is formally given as

$$\phi_\beta(\ell) = \log \left(\frac{\ell^4 + \beta}{\ell^4} \right) \mathbb{1}\{\ell \leq R\}, \quad \beta > 0, \ell \geq 0,$$

where $\ell \geq 0$ is the length of an Delaunay edge and $\beta > 0$ is the potential parameter and $R > 0$ is finite range condition of the potential. There are two novelties in this research: We obtain a phase transition for soft repulsion (no hard-core repulsion) between unlike particles on the Delaunay structure with no additional constraints on the distribution of the underlying particle system. Secondly, the main novelty of our paper is a uniform bound on the number of connected components in the Delaunay random cluster model which is purely based on geometrical properties of Delaunay tessellations in two dimensions. The potential ensures that for large values of the parameter $\beta > 0$, Delaunay edges with shorter lengths are more likely to be connected than those with longer edges, enabling us to bound the number of connected components for clouds of points with vanishing point-wise distances. This paper is an extensive further development of the recent work [AE16] where all models had an additional background hard-core potential introducing a length scale for the configurations. Gibbs models on Delaunay structures have been studied in [BBD99, BBD02, BBD04, Der08, DDG12, DG09, DL11], and our results rely on the existence of Gibbs measures for the geometry-dependent interaction using methods in [DDG12]. Our approach is based on a Delaunay random-cluster representation. A phase transition for our Delaunay Potts model follows if we can show that the corresponding percolation process contains an infinite cluster. A similar program was carried out by Chayes et al. in [CCK95] for the hard-core Widom-Rowlinson model. In that case, the existence of infinite clusters follows from a stochastic comparison with the Poisson Boolean model of continuum percolation, while our framework uses a coarse graining method to derive a stochastic comparison with mixed site-bond percolation on \mathbb{Z}^2 . Our results are extension of [LL72, Rue71] and [CCK95] to the Delaunay structure replacing hard-core constraint by our soft-core repulsion. In particular we obtain phase transition for all activities once the interaction parameter $\beta > 0$ (inverse temperature) is sufficiently large depending on the activity. We note that our random-cluster representation requires the symmetry of the type interaction. In the non-symmetric Widom-Rowlinson models, the existence of a phase transition has been established by Bricmont et al. [BKL84], and recently by Suhov et al. [MSS15].

1.2 Remarks on Delaunay tessellations

We add some remarks on models defined on Delaunay hypergraph structures. There are differences between geometric models on the Delaunay hypergraph structure and point particle models such as the Widom-Rowlinson model. The first is that edges and triangles in the Delaunay hypergraph are each proportional in number to the number

of points in the configuration. However, in the case of the complete hypergraph the number of edges is proportional to the number of points squared and the number of triangles is proportional to the number of points cubed. Secondly, in complete graphs of all classical models, the neighbourhood of a given point depends only on the distance between points and so the number of neighbours increases with the activity parameter z of the underlying point process. This means that the system will become strongly connected for high values of z . This is not the case for the Delaunay hypergraphs which exhibit a self-similar property. Essentially, as the activity parameter z increases, the expected number of neighbours to a given point in the Delaunay hypergraph remains the same, see [Mø94]. Therefore, in order to keep a strong connectivity, we use a type interaction between points of Delaunay edges with a non-constant mark. Finally, and perhaps most importantly, is the question of additivity. Namely, suppose we have an existing point configuration ω and we want to add a new particle x to it. It is well-known that classical many-body interactions are additive, see [AE16] for details and references. On the other hand, in the Delaunay framework, the introduction of a new point to an existing configuration not only creates new edges and triangles, but destroys some too. The Delaunay interactions are therefore not additive, and for this reason, attractive and repulsive interactions are indistinct. In the case of a hard exclusion interaction, we arrive at the possibility that a configuration ω is excluded, but for some x , $\omega \cup x$ is not. This is called the non-hereditary property [DG09], which seems to rule out using techniques such as stochastic comparisons of point processes [GK97].

1.3 Setup

We consider configurations of points in \mathbb{R}^2 with internal degrees of freedom, or marks. Let $M_q = \{1, \dots, q\}$, $q \in \mathbb{N}$, $q \geq 2$, be the finite set of different marks. That is, each marked point is represented by a position $x \in \mathbb{R}^2$ and a mark $\sigma(x) \in M_q$, and each marked configuration ω is a countable subset of $\mathbb{R}^2 \times M_q$ having a locally finite projection onto \mathbb{R}^2 . We denote by Ω the set of all marked configurations with locally finite projection onto \mathbb{R}^2 . We will sometimes identify ω with a vector $\omega = (\omega^{(1)}, \dots, \omega^{(q)})$ of pairwise disjoint locally finite sets $\omega^{(1)}, \dots, \omega^{(q)}$ in \mathbb{R}^2 (we write Ω for the set of all locally finite configurations in \mathbb{R}^2). Any ω is uniquely determined by the pair (ω, σ) , where $\omega = \cup_{i=1}^q \omega^{(i)}$ is the set of all occupied positions, and where the mark function $\sigma: \omega \rightarrow M_q$ is defined by $\sigma(x) = i$ if $x \in \omega^{(i)}$, $i \in M_q$. For each measurable set B in $\mathbb{R}^2 \times M_q$ the counting variable $N(B): \omega \rightarrow \omega(B)$ on Ω gives the number of marked particles such that the pair (position, mark) belongs to B . We equip the space Ω with the σ -algebra \mathcal{F} generated by the counting variables $N(B)$ and the space Ω of locally finite configurations with the σ -algebra \mathcal{F} generated by the counting variables $N_\Delta = \#\{\omega \cap \Delta\}$ for $\Delta \in \mathbb{R}^2$ where we write $\Delta \in \mathbb{R}^2$ for any bounded $\Delta \subset \mathbb{R}^2$. As usual, we take as reference measure on (Ω, \mathcal{F}) the marked Poisson point process Π^z with intensity measure $z \text{Leb} \otimes \mu_u$ where $z > 0$ is an arbitrary activity, Leb is the Lebesgue measure in \mathbb{R}^2 , and μ_u is the uniform probability measure on M_q .

For each $\Lambda \subset \mathbb{R}^2$ we write $\Omega_\Lambda = \{\omega \in \Omega: \omega \subset \Lambda \times M_q\}$ for the set of configurations in Λ , $\text{pr}_\Lambda: \omega \rightarrow \omega_\Lambda := \omega \cap \Lambda \times M_q$ for the projection from Ω to Ω_Λ (similarly for unmarked configurations), $\mathcal{F}'_\Lambda = \mathcal{F}|_{\Omega_\Lambda}$ for the trace σ -algebra of \mathcal{F} on Ω_Λ , and $\mathcal{F}_\Lambda = \text{pr}_\Lambda^{-1} \mathcal{F}'_\Lambda \subset \mathcal{F}$ for the σ -algebra of all events that happen in Λ only. The reference measure on $(\Omega_\Lambda, \mathcal{F}'_\Lambda)$ is $\Pi_\Lambda^z := \Pi^z \circ \text{pr}_\Lambda^{-1}$. In a similar way we define the corresponding objects for unmarked configurations, Π^z , Π_Λ^z , Ω_Λ , pr_Λ , \mathcal{F}'_Λ , and \mathcal{F}_Λ . Finally, let $\theta = (\theta_x)_{x \in \mathbb{R}^2}$ be the shift group, where $\theta_x: \Omega \rightarrow \Omega$ is the translation of the spatial component by the vector $-x \in \mathbb{R}^2$. Note that by definition, $N_\Delta(\theta_x \omega) = N_{\Delta+x}(\omega)$ for all $\Delta \subset \mathbb{R}^2$.

We outline the definitions for the unmarked configurations first with obvious adaptations to the case of marked point configurations. The set of Delaunay hyperedges

of a given configuration $\omega \in \Omega$ consist of all $\eta \subset \omega$ for which there exists an open ball $B(\eta, \omega) \subset \mathbb{R}^2$ with $\partial B(\eta, \omega) \cap \omega = \eta$ that contains no points of ω . For $m = 1, 2, 3$, and $\omega \in \Omega$, we write $\text{Del}_m(\omega) = \{\eta \subset \omega : \#\eta = m\}$ for the set of Delaunay simplices with m vertices. It is possible that a Delaunay hyperedge η consists of four or more points on a sphere with no points inside. In fact, for this not to happen, we must consider configurations in general position as in [Mø94]. More precisely, this means that no four points lie on the boundary of a circle, no point lies inside a circumcircle for a triangle, and every half-plane contains at least one point. Fortunately, this occurs with probability one for our Poisson reference measure, and in fact, for any stationary point process. Note that the open ball $B(\eta, \omega)$ is only uniquely determined when $\#\eta = 3$ and η is affinely independent. Henceforth, for each configuration ω we have an associated Delaunay triangulation

$$\{\tau \subset \omega : \#\tau = 3, B(\tau, \omega) \cap \omega = \emptyset\} \tag{1.1}$$

of the plane, where $B(\tau, \omega)$ is the unique open ball with $\tau \subset \partial B(\tau, \omega)$ with B being the circumcircle of the triangle τ . The set in (1.1) is uniquely determined and defines a triangulation of the convex hull of ω whenever ω is in general position ([Mø94]). In a similar way one can define the marked Delaunay hyperedges $\mathbf{Del}_m(\omega)$, where the Delaunay property refers to the spatial component only.

Given a configuration $\omega \in \Omega$ (or ω) we write $\Omega_{\Lambda, \omega} = \{\zeta \in \Omega : \zeta \setminus \Lambda = \omega\}$ (resp. $\Omega_{\Lambda, \omega}$) for the set of configurations which equal ω off Λ . For any edge $\eta \in \text{Del}_2$ we denote its length by $\ell(\eta)$, i.e., $\ell(\eta) = |x - y|$ if $\eta = \{x, y\}$, and we write $\eta \equiv \eta_{x,y}$. The interaction is given by the following Hamiltonian in Λ with boundary condition $\omega \in \Omega$,

$$H_{\Lambda, \omega}(\zeta) := \sum_{\substack{\eta \in \text{Del}_2(\zeta) \\ \eta \in \text{Del}_{2, \Lambda}(\zeta)}} \phi_\beta(\ell(\eta))(1 - \delta_\sigma(\eta)), \quad \zeta \in \Omega_{\Lambda, \omega}, \tag{1.2}$$

where $\text{Del}_{2, \Lambda}(\zeta) := \{\eta \in \text{Del}_2(\zeta) : \exists \tau \in \text{Del}_3(\zeta), \eta \subset \tau, \partial B(\tau, \zeta) \cap \Lambda \neq \emptyset\}$. Here ϕ_β is a measurable function of the length $\ell(\eta)$ of an edge defined for any $\beta \geq 0$,

$$\phi_\beta(\ell) = \log\left(\frac{\ell^4 + \beta}{\ell^4}\right) \mathbb{1}\{\ell \leq R\}, \tag{1.3}$$

and

$$\delta_\sigma(\eta) = \begin{cases} 1 & , \text{ if } \sigma_\eta(x) = \sigma_\eta(y) \text{ for } \eta = \{x, y\}, \\ 0 & , \text{ otherwise.} \end{cases}$$

Note the following scaling relation for the potential

$$\phi_\beta(L\ell) = \phi_{\beta/L^4}(\ell), \quad \text{for any } L > 0 \text{ with } L\ell \leq R, \ell \leq R. \tag{1.4}$$

Following [DDG12] we define the partition function as

$$Z_\Lambda(\omega) = \int_{\Omega_{\Lambda, \omega}} e^{-H_{\Lambda, \omega}(\zeta)} \Pi^z(d\zeta^{(1)}) \dots \Pi^z(d\zeta^{(q)}).$$

The Gibbs distribution for ϕ_β , and $z > 0$ in Λ with boundary condition ω is defined as

$$\gamma_{\Lambda, \omega}(A) = \frac{1}{Z_\Lambda(\omega)} \int_{\Omega_{\Lambda, \omega}} \mathbb{1}_A(\zeta \cup \omega) e^{-H_{\Lambda, \omega}(\zeta)} \Pi_\Lambda^z(d\zeta), \quad A \in \mathcal{F}. \tag{1.5}$$

It is evident from (1.5) that, for fixed $\zeta \in \Omega_\Lambda$, the conditional distribution of the marks of $\zeta = (\zeta^{(1)}, \dots, \zeta^{(q)})$ relative to $\gamma_{\Lambda, \omega}$ is that of a discrete Potts model on ζ embedded in the Delaunay triangulation with position-dependent interaction between the marks. This justifies calling our model *Delaunay Potts model* or *Delaunay Widom-Rowlinson model*.

Definition 1.1. A probability measure μ on Ω is called a Gibbs measure for the Delaunay Potts model with activity $z > 0$ and interaction potential ϕ_β if

$$\mathbb{E}_\mu[f] = \int_\Omega \frac{1}{Z_\Lambda(\omega)} \int_{\Omega_{\Lambda,\omega}} f(\zeta \cup \omega) e^{-H_{\Lambda,\omega}(\zeta)} \Pi_\Lambda^z(d\zeta) \mu(d\omega) \quad (1.6)$$

for every $\Lambda \in \mathbb{R}^2$ and every measurable function f .

The equations in (1.6) are the DLR equations (after Dobrushin, Lanford, and Ruelle). They ensure that the Gibbs distribution in (1.5) is a version of the conditional probability $\mu(A|\mathcal{F}_{\Lambda^c})(\omega)$. The measurability of all objects is established in [Eye14, DDG12].

1.4 Results and remarks

Proposition 1.2 (Existence of Gibbs measures). For any $z > 0$ there exists at least one Gibbs measure for the Delaunay Widom-Rowlinson (Potts) model with parameter $\beta > 0$.

Remark 1.3 (Gibbs measures). The proof is using the so-called pseudo-periodic configurations (see Appendix A or [DDG12]) and properties of the potential ϕ_β . Existence of Gibbs measures for related different Delaunay models have been obtained in [BBD99, Der08, DG09]. Note that for $q = 1$ our models have no marks and Gibbs measures do exist as well ([DDG12]). \diamond

A *phase transition* is said to occur if there exists more than one Gibbs measure for the Delaunay Potts model. The following theorem shows that this happens for all activities z greater than some lower bound depending on the range $R > 0$ and the number of types and sufficiently large parameter β depending on z . Note that β is a parameter for the type interaction and not the usual inverse temperature. The statement of the following phase transition depends on the novel upper bound for the number of connected components in Theorem 2.1 below.

Theorem 1.4 (Phase transition). For $\alpha^* = \alpha^*(R, q)$ defined in (2.5) there exists $z^*(\alpha^*, R, q) > 0$ such that the following holds. For all $z > R^{-2}z^*(\alpha^*, R, q)$ there exists $\beta_0 = \beta_0(q, R, z)$ such that for all $\beta \geq \beta_0 \vee q$ there exist at least q different Gibbs measures for the Delaunay Widom Rowlinson (Potts) model.

Remark 1.5. (a) Theorem 1.4 actually establishes a break of the symmetry in the type distribution.

(b) The condition on the activity says that $zR^2 > z^*(\alpha^*, R, q)$ for some constant $z^*(\alpha^*, R, q)$ (see Lemma 2.4) depending solely on R and q . This is expected as the activity scales with the square of the given finite range length scale of the potential. Furthermore, the constant $\beta_0(q, R, z)$ is chosen to increase the quotient $\frac{R^4}{\beta}$, this is due to the scale invariance of the potential (1.4).

(c) Theorem 1.4 also holds for any potential depending on the length ℓ of Delaunay edges

$$\phi_\beta^{(\gamma)}(\ell) := \log \left(\frac{\ell^{3+\gamma} + \beta}{\ell^{3+\gamma}} \right) \mathbb{1}\{\ell \leq R\}, \quad \gamma > 0.$$

We need higher than cubic powers in the potential to balance the number of edges which grows quadratically with the edge length, see Section 3.4 for details. \diamond

Remark 1.6 (Free energy and Uniqueness of Gibbs measures). (a) One may wonder if the phase transition manifest itself thermodynamically by a non-differentiability (“discontinuity”) of the free energy (pressure). We refer the interested reader to [AE16] for more details and references.

- (b) To establish uniqueness of the Gibbs measure in our Delaunay Potts model one can use the Delaunay random-cluster measure $C_{\Lambda_n, \omega}$, to be defined in (2.2) below. Details are outlined in [AE16]. In contrast to the models studied in [AE16], in our model the interaction parameter β plays the role of the inverse temperature, that is, if β is chosen sufficiently small, then there is no percolation in our Delaunay random-cluster measure and therefore uniqueness of the Gibbs measure. \diamond

The study for Widom-Rowlinson or Potts models with geometry-dependent interaction is by far not complete, one may wish to extend the single edge (or triangle) interaction to mutual adjacent Voronoi cell interaction. The common feature of all these “ferromagnetic” systems is that phase transitions are due to breaking the symmetry of the type distribution.

The rest of the paper is organised as follows. In Section 2.1 we define the Delaunay random-cluster measure for edge configurations, and in Section 2.2 we establish percolation in this model for certain parameters. The main novelty is the extensive and elaborate proof of the uniform bound on the number of connected components using purely geometric properties in Section 3. Finally, in Section 4 we give details of our remaining proofs.

2 The random cluster method

In Section 2.1 we introduce the Delaunay random cluster model and show percolation for this model in Section 2.2 via comparison with mixed site-bond percolation on \mathbb{Z}^2 . We conclude in Section 2.3 with our proof of Theorem 1.4. The key step is our novel uniform estimate of the number of connected components in Section 3. The proof of this bound uses solely geometric arguments and constitutes a major part of this work.

2.1 Delaunay random cluster measure

For $\Lambda \in \mathbb{R}^2$ and parameters z and ϕ_β we define a joint distribution of the Delaunay Potts model and an edge process which we call Delaunay random-cluster model. The basic idea is to introduce random edges between points in the plane. Let

$$E_{\mathbb{R}^2} = \{\eta = \{x, y\} \subset \mathbb{R}^2 : x \neq y\}$$

be the set of all possible edges of points in \mathbb{R}^2 , likewise, let E_Λ be the set of all edges in Λ and E_ζ for the set of edges in $\zeta \in \Omega_{\Lambda, \omega}$. We identify ω with $\omega^{(1)}$ and $\omega = (\omega^{(1)}, \emptyset, \dots, \emptyset)$. This allows only monochromatic boundary conditions whereas the general version involves the so-called Edwards-Sokal coupling (see [GHM00] for lattice Potts models). We restrict ourselves to the former case for ease of notation. We write

$$\mathcal{E} = \{E \subset E_{\mathbb{R}^2} : E \text{ locally finite}\}$$

for the set of all locally finite edge configurations.

The joint distribution is built from the following three components.

The *point distribution* is given by the Poisson process Π^{zq} for any boundary condition $\omega \in \Omega$ and activity zq .

The *type picking mechanism* for a given configuration $\zeta \in \Omega_{\Lambda, \omega}$ is the distribution $\lambda_{\zeta, \Lambda}$ of the mark vector $\sigma \in M_q^\zeta$. Here $(\sigma(x))_{x \in \zeta}$ are independent and uniformly distributed random variables on M_q with $\sigma(x) = 1$ for all $x \in \zeta_{\Lambda^c} = \omega$. The latter condition ensures that all points outside of Λ carry the given fixed mark.

The *edge drawing mechanism*. Given a point configuration $\zeta \in \Omega_{\Lambda, \omega}$, we let $\mu_{\zeta, \Lambda}$ be the distribution of the random edge configuration $\{\eta \in E_\zeta : v(\eta) = 1\} \in \mathcal{E}$ with the edge

configuration $v \in \{0, 1\}^{E_\zeta}$ having probability

$$\prod_{\eta \in E_\zeta} p(\eta)^{v(\eta)} (1 - p(\eta))^{1-v(\eta)}$$

with

$$p(\eta) := \mathbb{P}(v(\eta) = 1) = \begin{cases} (1 - e^{-\phi_\beta(\eta)}) \mathbb{1}_{\text{Del}_2(\zeta)}(\eta) & \text{if } \eta \in E_{\mathbb{R}^2} \setminus E_{\Lambda^c}, \\ \mathbb{1}_{\text{Del}_2(\zeta)}(\eta) & \text{if } \eta \in E_{\Lambda^c}. \end{cases} \quad (2.1)$$

The measure $\mu_{\zeta, \Lambda}$ is a point process on $E_{\mathbb{R}^2}$. Note that $\zeta \rightarrow \lambda_{\zeta, \Lambda}$ and $\zeta \rightarrow \mu_{\zeta, \Lambda}$ are probability kernels (see [Eye14, AE16]). Let the measure

$$P_{\Lambda, \omega}^{zq}(\text{d}\zeta, \text{d}E) := \frac{1}{Z_\Lambda(\omega)} \Pi_{\Lambda, \omega}^{zq}(\text{d}\zeta) \lambda_{\zeta, \Lambda}(\text{d}\zeta) \mu_{\zeta, \Lambda}(\text{d}E)$$

be supported on the set of all (ζ, E) with $\zeta \in \Omega_{\Lambda, \omega}$ and $E \subset E_\zeta$. We shall condition on the event that the marks of the points are constant on each connected component in the graph $(\zeta, E \cap E_\zeta)$. Two distinct vertices x and y are adjacent to one another if there exists $\eta \in E_\zeta$ such that $\{x, y\} = \eta$. A path in the graph $(\zeta, E \cap E_\zeta)$ is an alternating sequence $v_1, e_1, v_2, e_2, \dots$ of distinct vertices v_i and edges e_j such that $\{v_i, v_{i+1}\} = e_i$ for all $i \geq 1$. We write

$$A = \{(\zeta, E) \in \Omega \times \mathcal{E} : \sum_{\eta \in E} (1 - \delta_\sigma(\eta)) = 0\}$$

for the set of marked point configurations such that all vertices of the edges carry the same mark. The set A is measurable which one can see from writing the condition in the following way

$$\sum_{\eta = \{x, y\} \in E} \sum_{i=1}^q (\mathbb{1}_{\zeta^{(i)}}(x)(1 - \mathbb{1}_{\zeta^{(i)}}(y))) = 0$$

and using the fact that $(\zeta, x) \mapsto \mathbb{1}_{\zeta^{(i)}}(x), i = 1, \dots, q$, are measurable (see [GH96, Chapter 2]). Furthermore, $\Pi_{\Lambda, \omega}^{zq}(A) > 0$, which follows easily observing $\Pi_{\Lambda, \omega}^{zq}(A) \geq \Pi_{\Lambda, \omega}^{zq}(\{\tilde{\omega}\}) = e^{-zq|\Lambda|}/Z_\Lambda(\omega)$, where $\tilde{\omega}$ is the configuration which equals ω outside of Λ and which is empty inside Λ . Henceforth, the random-cluster measure

$$\mathbf{P} = P_{\Lambda, \omega}^{zq}(\cdot | A)$$

is well-defined. We obtain the following two measures from the random-cluster measure \mathbf{P} , namely if we disregard the edges we obtain the Delaunay Gibbs distribution $\gamma_{\Lambda, \omega}$ in (1.5) (see [Eye14]). For the second measure consider the mapping $\text{sp}: (\zeta, E) \rightarrow (\zeta, E)$ from $\Omega \times \mathcal{E}$ onto $\Omega \times \mathcal{E}$ where $\zeta \mapsto \zeta = \cup_{i=1}^q \zeta^{(i)}$. For each (ζ, E) with $E \subset E_\zeta$ we let $K(\zeta, E)$ denote the number of connected components in the graph (ζ, E) . The Delaunay random-cluster distribution on $\Omega \times \mathcal{E}$ is defined by

$$C_{\Lambda, \omega}(\text{d}\zeta, \text{d}E) = \frac{1}{Z_\Lambda(\omega)} q^{K(\zeta, E)} \Pi_{\Lambda, \omega}^z(\text{d}\zeta) \mu_{\zeta, \Lambda}(\text{d}E), \quad (2.2)$$

where $\Pi_{\Lambda, \omega}^z$ is the Poisson process with activity z replacing zq and where

$$Z_\Lambda(\omega) = \int_{\Omega_{\Lambda, \omega}} \int_{\mathcal{E}} q^{K(\zeta, E)} \Pi_{\Lambda, \omega}^z(\text{d}\zeta) \mu_{\zeta, \Lambda}(\text{d}E)$$

is the normalisation. It is straightforward to show that $\mathbf{P} \circ \text{sp}^{-1} = C_{\Lambda, \omega}$.

For our main proofs we need to investigate the geometry of the Delaunay tessellation, and in particular what happens when we augment $\zeta \in \Omega_{\Lambda, \omega}$ with a new point $x_0 \notin \zeta$. Some edges (triangles) may be destroyed, some are created, and some remain. This

process is well described in [Lis94]. We give a brief account here for the convenience of the reader. We insert the point x_0 into one of the triangles τ in $\text{Del}_3(\zeta)$. We then create three new edges that join x_0 to each of the three vertices of τ . This creates three new triangles, and destroys one. We now need to verify that the new triangles each satisfy the Delaunay condition (1.1), that is, that their circumscribing balls contain no points of ζ . If this condition is satisfied the new triangle remains, if it is not satisfied, then there is a point $x_1 \in \zeta$ inside the circumscribing ball. We remove the edge not connected to x_0 , and replace it by an edge connecting x_0 and x_1 . This results in the creation of two new triangles. Each of these triangles must be checked as above and the process continues. Once all triangles satisfy the Delaunay condition, we arrive at the Delaunay triangulation $\text{Del}_3(\zeta \cup \{x_0\})$ and their Delaunay edges $\text{Del}_2(\zeta \cup \{x_0\})$. Let

$$\begin{aligned} E_{x_0, \zeta}^{(\text{ext})} &= \text{Del}_2(\zeta) \cap \text{Del}_2(\zeta \cup \{x_0\}), \\ E_{x_0, \zeta}^{(+)} &= \text{Del}_2(\zeta \cup \{x_0\}) \setminus \text{Del}_2(\zeta) = \text{Del}_2(\zeta \cup \{x_0\}) \setminus E_{x_0, \zeta}^{(\text{ext})}, \\ E_{x_0, \zeta}^{(-)} &= \text{Del}_2(\zeta) \setminus \text{Del}_2(\zeta \cup \{x_0\}) = \text{Del}_2(\zeta) \setminus E_{x_0, \zeta}^{(\text{ext})}, \end{aligned} \tag{2.3}$$

be the set of exterior, created, and destroyed Delaunay edges respectively, see Figure 1.




Figure 1: The Delaunay sets $\text{Del}_2(\zeta)$, $\text{Del}_2(\zeta \cup \{x_0\})$, and the exterior Delaunay set $E_{x_0, \zeta}^{(\text{ext})}$.

Note that any new edge must contain x_0 , i.e.,

$$E_{x_0, \zeta}^{(+)} = \{\eta = \eta_{x,y} \in \text{Del}_2(\zeta \cup \{x_0\}) : x = x_0, y \neq x_0 \text{ or } y = x_0, x \neq x_0\}.$$

We let $\mu_{x_0, \zeta}^{(-)}$, $\mu_{x_0, \zeta}^{(+)}$, and $\mu_{x_0, \zeta}^{(\text{ext})}$ be the edge drawing mechanisms on $E_{x_0, \zeta}^{(\text{ext})}$, $E_{x_0, \zeta}^{(+)}$, and $E_{x_0, \zeta}^{(-)}$, respectively, which are derived from the edge drawing measure $\mu_{\zeta, \Lambda}$ above. The crucial

step is an estimate on the number of connected components in the neighbourhood of the point x_0 . For any $\zeta \in \Omega_{\Lambda, \omega}$ the neighbourhood of a point $x_0 \in \Lambda, x_0 \notin \zeta$, is the following random graph $G_{x_0, \zeta} = (V_{x_0, \zeta}, E_{x_0, \zeta}^{(\text{nbdd})})$ where $V_{x_0, \zeta}$ is the set of points that share an edge with x_0 in $\text{Del}_2(\omega \cup \{x_0\})$ and $E_{x_0, \zeta}^{(\text{nbdd})}$ is the set of edges in $\text{Del}_2(\omega \cup \{x_0\})$ that have both endpoints in $V_{x_0, \zeta}$, more precisely,

$$V_{x_0, \zeta} = \{x \in \zeta : \eta_{x, x_0} \in E_{x_0, \zeta}^{(+)}\} \quad \text{and} \quad E_{x_0, \zeta}^{(\text{nbdd})} = \{\eta_{x, y} \in E_{x_0, \zeta}^{(\text{ext})} : x, y \in V_{x_0, \zeta}\}.$$




Figure 2: The exterior graph $E_{x_0, \zeta}^{(\text{ext})}$ and the boundary graph $G_{x_0, \zeta}$.

The graph $G_{x_0, \zeta}$ splits the plane into two regions. The region containing x_0 is called the neighbourhood of x_0 whereas $G_{x_0, \zeta}$ is called the boundary (graph) of the

neighbourhood of x_0 , see Figure 2. Having the edge drawing mechanism $\mu_{x_0, \zeta}^{(\text{ext})}$ define

$$\mu_{(\text{ext}), \zeta}^{(q)}(dE) := \frac{q^{K(\zeta, E)} \mu_{x_0, \zeta}^{(\text{ext})}(dE)}{\int q^{K(\zeta, E)} \mu_{x_0, \zeta}^{(\text{ext})}(dE)}. \tag{2.4}$$

The main task is the find an uniform upper bound (independent of ζ) for the expected number of connected components of (ζ, E) that intersect the boundary $G_{x_0, \zeta}$, where E is sampled from $\mu_{(\text{ext}), \zeta}^{(q)}$. The bound will enable us to estimate certain conditional Papangelou intensities from below and above. This in turn allows to dispense background potentials used in recent work ([BBD04, AE16]). Given $\zeta \in \Omega_{\Lambda, \omega}$ and $x_0 \in \Lambda, x_0 \notin \zeta$, and the Delaunay graph $\text{Del}_2(\zeta)$ we denote $N_{x_0}^{(\text{cc})}(\zeta, E)$ for any $E \subset E_{x_0, \zeta}^{(\text{ext})}$ the number of connected components that intersect $G_{x_0, \zeta}$.

Theorem 2.1 (Number of connected components). *Let $\beta > q$. For all $\Lambda \in \mathbb{R}^2$ and $\alpha = \alpha(R, q, \beta) := 1 + 6R^2\pi^2r^{-2}(1 + \frac{2q\pi^2r^2}{3\beta}) < \infty$ with $r = 1 \wedge \frac{R\pi}{2}$, the following estimate*

$$\int N_{x_0}^{(\text{cc})}(\zeta, E) \mu_{(\text{ext}), \zeta}^{(q)}(dE) \leq \alpha$$

holds for all $\omega \in \Omega$ and $\zeta \in \Omega_{\Lambda, \omega}$ and for all $x_0 \in \Lambda$ with $x_0 \notin \zeta$. Note that, as $\beta > q$,

$$\alpha(R, q, \beta) \leq \alpha^*(R, q) := 1 + 6R^2\pi^2r^{-2}(1 + \frac{2\pi^2r^2}{3}). \tag{2.5}$$

Moreover,

$$\lim_{\beta \rightarrow \infty} \alpha(R, q, \beta) = 1 + 6R^2\pi^2r^{-2} = \begin{cases} 25 & \text{if } R < 2/\pi, \\ 1 + 6R^2\pi^2 & \text{if } R \geq 2/\pi. \end{cases} \tag{2.6}$$

Proof. The proof is in Section 3. □

We need to study the change of $K(\zeta, E), E \subset E_\zeta \subset \mathcal{E}$ when adding the point $x_0 \notin \zeta$. Adding a point x_0 to ζ without changing E will always increase the number of connected components by one as the new point may not be connected to any other point. On the other hand, the addition of a single edge η to E can result in the connection of two different connected components, leaving one. Therefore,

$$\begin{aligned} K(\zeta \cup \{x_0\}, E) - K(\zeta, E) &= 1, \\ -1 \leq K(\zeta, E \cup \eta) - K(\zeta, E) &\leq 0. \end{aligned}$$

2.2 Delaunay edge-percolation

We establish the existence of edge percolation for the Delaunay random-cluster measure $C_{\Lambda, \omega}$ when z and the parameter β are sufficiently large. Note that for any $\Delta \in \mathbb{R}^2$ we write

$$N_{\Delta \leftrightarrow \infty}(\zeta, E) = \#\{x \in \zeta \cap \Delta : x \text{ belongs to an } \infty\text{-cluster of } (\zeta, E \cap E_\zeta)\},$$

where an ∞ -cluster is an unbounded open connected component in $(\zeta, E \cap E_\zeta)$. The key step in our results is the following percolation result.

Proposition 2.2. *Suppose all the assumptions hold and that z and β are sufficiently large. Suppose that Λ is a finite union of cells $\Delta_{k,l}$ defined in (A.1) Appendix A. Then there exists $\varepsilon > 0$ such that*

$$\int N_{\Delta \leftrightarrow \infty} dC_{\Lambda, \omega} \geq \varepsilon$$

for any cell $\Delta = \Delta_{k,l}$, any finite union Λ of cells and any pseudo-periodic boundary condition $\omega \in \Omega_\Lambda^*$.

Proof of Proposition 2.2. We split the proof in several steps and Lemmata below. Our strategy to establish percolation in the Delaunay random-cluster model is to compare it to mixed site-bond percolation on \mathbb{Z}^2 (see appendix C on mixed site-bond percolation). First we employ a coarse-graining strategy to relate each site $(k, l) \in \mathbb{Z}^2$ to a cell which is a union of rhombuses. The second step is to consider the links (bonds) of two good cells. In order to establish mixed site-bond percolation we need to define when cells are good (open) and when two neighbouring cells are linked once they are open which happens when the corresponding link (bond) is open as well. This link establishes an open connection in our Delaunay graph $\text{Del}_2(\zeta)$. We extend the coarse graining method recently used in [AE16].

Step 1: Coarse graining.

Let $\Lambda \equiv \Lambda_n \subset \mathbb{R}^2$ be the rhombus given as the finite union of cells (A.1) with side length ℓ , i.e.,

$$\Lambda_n = \bigcup_{(k,l) \in \{-n, \dots, n\}^2} \Delta_{k,l} \quad \text{and} \quad \Delta_{k,l} = \bigcup_{i,j=0}^7 \Delta_{k,l}^{i,j}, \tag{2.7}$$

where $\Delta_{k,l}^{i,j}$ are rhombuses with side length $\ell/8$ and each rhombus $\Delta_{k,l}$ has side length ℓ where the coordinate systems is the canonical one, that is, $\Delta_{k,l}^{0,0}$ is the rhombus in the bottom right corner. We denote (2.7) the ℓ -partition of the rhombus Λ , The union of the 16 smaller rhombuses towards the centre of $\Delta_{k,l}$ is denoted

$$\Delta_{k,l}^{(-)} = \bigcup_{i,j=2}^5 \Delta_{k,l}^{i,j}$$

see Figure 3.




Figure 3: The square version of ℓ -partition of Λ , i.e., the ℓ -partition of Λ into rhombuses is the sheared version by 60° (application of the matrix M defined in Appendix A). The shaded boxes are the link boxes.

These cells with index (k, l) will act as the sites in the mixed site-bond percolation model on \mathbb{Z}^2 . Finally, we define the link boxes between $\Delta_{k,l}$ and $\Delta_{k+1,l}$ as

$$\Delta_{(\text{link})}^{k:k+1,l} = \left\{ \bigcup_{j=0}^3 \Delta_{k,l}^{6,j+2} \right\} \cup \left\{ \bigcup_{j=0}^3 \Delta_{k,l}^{7,j+2} \right\} \cup \left\{ \bigcup_{j=0}^3 \Delta_{k+1,l}^{0,j+2} \right\} \cup \left\{ \bigcup_{j=0}^3 \Delta_{k+1,l}^{1,j+2} \right\} \quad (2.8)$$

which act as the bonds in the mixed site-bond percolation model on \mathbb{Z}^2 , see Figure 3.

We shall choose

$$\ell \in \left(0, \frac{R}{2\sqrt{3}}\right] \quad (2.9)$$

to ensure that we can open edges in neighbouring rhombuses, see (2.2) and Step (iv) in the proof of Lemma 2.7 below. This completes the coarse graining set-up. We establish percolation in the mixed site-bond percolation model on \mathbb{Z}^2 , that is, the existence of an infinite chain of open sites and open bonds, and we relate it to the existence of an infinite connected component of open edges in Del_2 . This infinite connected component will connect with the complements of any finite boxes, and thus this connected component corresponds to an infinite connected component of edges where all sites carry the mark agreed for the boundary condition. To do this, we define $\text{CB}_{k:k+1,l}$ to be the straight line segment (also denoted the central band) between the centres of the rhombuses $\Delta_{k,l}$ and $\Delta_{k+1,l}$ and let

$$\text{H}_{k:k+1,l}(\zeta) = \{x \in \zeta : \text{Vor}_\zeta(x) \cap \text{CB}_{k:k+1,l} \neq \emptyset\} \quad (2.10)$$

be the subset of points of the configuration ζ , whose Voronoi cells intersect the line segment $\text{CB}_{k:k+1,l}$, see Figures 3 and 4, and where the Voronoi cell is defined as

$$\text{Vor}_\zeta(x) := \{y \in \mathbb{R}^2 : |x - y| \leq |\tilde{x} - y| \forall \tilde{x} \in \zeta\}.$$




Figure 4: The shaded area is the union of the Voronoi cells with centre $\in \text{H}_{k:k+1,l}(\zeta)$ (square version).

We need to consider the distribution of the points given by the marginal distribution

$$M_{\Lambda,\omega} = C_{\Lambda,\omega}(\cdot \times \mathcal{E})$$

of the Delaunay random-cluster measure on Ω . Note that (2.2) can be written as

$$C_{\Lambda,\omega}(d\zeta, dE) = M_{\Lambda,\omega}(d\zeta)\mu_{\zeta,\Lambda}^{(q)}(dE), \quad \mu_{\zeta,\Lambda}^{(q)}(dE) = \frac{q^{K(\zeta,E)}\mu_{\zeta,\Lambda}(dE)}{\int q^{K(\zeta,E)}\mu_{\zeta,\Lambda}(dE)}.$$

We define h_Λ to be the Radon-Nikodym density of $M_{\Lambda,\omega}$ with respect to $\Pi_{\Lambda,\omega}^z$, i.e., for $\zeta \in \Omega_{\Lambda,\omega}$,

$$h_\Lambda(\zeta) := \mathcal{Z}_\Lambda(\omega)^{-1} \int q^{K(\zeta,E)}\mu_{\zeta,\Lambda}(dE).$$

In the following lemma we derive a bound for the Papangelou conditional intensity of $M_{\Lambda,\zeta}$.

Lemma 2.3. *For any $\Lambda \Subset \mathbb{R}^2$ and any boundary condition $\omega \in \Omega$ and $M_{\Lambda,\zeta}$ -almost all $\zeta \in \Omega_{\Lambda,\omega}$ and a point $x_0 \in \Lambda$ with $x_0 \notin \zeta$,*

$$\frac{h_\Lambda(\zeta \cup \{x_0\})}{h_\Lambda(\zeta)} \geq q^{1-\alpha}, \tag{2.11}$$

where $\alpha \in (0, \infty)$ is given in Theorem 2.1.

Proof of Lemma 2.3. Recall the different edge drawing mechanisms $\mu_{x_0,\zeta}^{(-)}$, $\mu_{x_0,\zeta}^{(+)}$, and $\mu_{x_0,\zeta}^{(\text{ext})}$ on $E_{x_0,\zeta}^{(\text{ext})}$, $E_{x_0,\zeta}^{(+)}$, and $E_{x_0,\zeta}^{(-)}$, respectively, and the definition of the probability measure $\mu_{(\text{ext}),\zeta}^{(q)}$ in (2.4). It follows that

$$\begin{aligned} \frac{h_\Lambda(\zeta \cup \{x_0\})}{h_\Lambda(\zeta)} &= \frac{\int q^{K(\zeta,E_2)} \int q^{K(\zeta \cup \{x_0\}, E_1 \cup E_2) - K(\zeta, E_2)} \mu_{x_0,\zeta}^{(+)}(dE_1) \mu_{x_0,\zeta}^{(\text{ext})}(dE_2)}{\int q^{K(\zeta,E_3)} \int q^{K(\zeta, E_3 \cup E_4) - K(\zeta, E_3)} \mu_{x_0,\zeta}^{(-)}(dE_4) \mu_{x_0,\zeta}^{(\text{ext})}(dE_3)} \\ &\geq \iint q^{K(\zeta \cup \{x_0\}, E_1 \cup E_2) - K(\zeta, E_2)} \mu_{x_0,\zeta}^{(+)}(dE_1) \mu_{(\text{ext}),\zeta}^{(q)}(dE_2) \\ &\geq q^{\int (K(\zeta \cup \{x_0\}, E_1 \cup E_2) - K(\zeta, E_2))} \mu_{x_0,\zeta}^{(+)}(dE_1) \mu_{(\text{ext}),\zeta}^{(q)}(dE_2), \end{aligned}$$

since

$$K(\zeta, E_3 \cup E_4) - K(\zeta, E_3) \leq 0$$

due to the fact that adding any edge from $E_4 \subset E_{x_0,\zeta}^{(-)}$ will only fuse connected components of the remaining graph. The second inequality is just Jensen's inequality applied to the convex function $x \mapsto q^x$. Note that new edges from $E_{x_0,\zeta}^{(+)}$, made by the insertion of x_0 to the configuration ζ , are edges connecting x_0 to points in ζ and are open with respect to $\mu_{x_0,\zeta}^{(+)}$, and therefore

$$K(\zeta \cup \{x_0\}, E_1 \cup E_2) - K(\zeta, E_2) \geq -N_{x_0}^{(\text{cc})}(\zeta, E_2) + 1 \tag{2.12}$$

for any $E_1 \subset E_{x_0,\zeta}^{(+)}$ and any $E_2 \subset E_{x_0,\zeta}^{(\text{ext})}$, and we conclude with the statement using Theorem 2.1. \square

We establish an upper bound for the Papangelou conditional intensity for certain configurations below in Lemma 2.5.

An important component of our coarse graining method is to estimate the conditional probability that at least one point lies inside some $\Delta \subset \Lambda$ for Δ element of the partitioning (2.7) of Λ . For any $\omega' \in \Omega_{\Delta^c}$ with $\omega' \cap \Lambda^c = \omega$ we denote by $M_{\Lambda,\Delta,\omega'}$ the conditional distribution of the configuration in Δ given the configuration ω' in Δ^c relative to $M_{\Lambda,\omega}$. The details of the construction of the regular conditional probability distribution can be found in [Eye14] or [GH96]. Having a uniform lower bound for the quotient $h_\Lambda(\zeta \cup$

$\{x_0\}/h_\Lambda(\zeta)$ allows to exhibit some control over the distribution $M_{\Lambda, \nabla, \omega'}$ for any $\nabla \subset \Lambda$. In the following we write ∇ for any cell $\Delta_{k,l}^{i,j}, i, j = 0, \dots, 7$. We fix

$$\varepsilon = \frac{1 - \sqrt{p_c^{(\text{site})}(\mathbb{Z}^2)}}{4}, \tag{2.13}$$

where $p_c^{(\text{site})}(\mathbb{Z}^2) \in (0, 1)$ is the critical probability for site percolation on \mathbb{Z}^2 .

Lemma 2.4. *For*

$$zR^2 > z_0^*(\alpha^*, q, R) =: \frac{8 \cdot 64^2 \sqrt{3} q^{\alpha^* - 1}}{\varepsilon} \tag{2.14}$$

there exist $\ell = \ell(z) \in (0, \frac{R}{2\sqrt{3}}]$ such that for all pseudo-periodic boundary conditions $\omega \in \Omega$,

$$M_{\Lambda, \nabla, \omega'}(\#\{\zeta \cap \nabla\} \geq 1) > 1 - \frac{\varepsilon}{64}$$

for all cells $\nabla = \Delta_{k,l}^{i,j}, (k, l) \in \{-n, \dots, n\}^2, i, j = 0, \dots, 7$, of the partitioning (2.7) of Λ , and for any configuration $\omega' \in \Omega_{\nabla^c}$ with $\omega' \setminus \Lambda = \omega$.

Proof of Lemma 2.4. Fix some $\omega' \in \Omega_{\nabla^c}$ with $\omega' \setminus \Lambda = \omega$. Then the statement follows immediately from

$$\frac{M_{\Lambda, \nabla, \omega'}(\#\{\zeta \cap \nabla\} = 1)}{M_{\Lambda, \nabla, \omega'}(\#\{\zeta \cap \nabla\} = 0)} = z \int_{\nabla} \frac{h_\Lambda(\omega' \cup \{x\})}{h_\Lambda(\omega')} dx \geq zq^{1-\alpha} |\nabla|,$$

where the inequality follows from Lemma 2.3. It follows that

$$M_{\Lambda, \nabla, \omega'}(\#\{\zeta \cap \nabla\} = 0) \leq q^{\alpha-1} (z|\nabla|)^{-1},$$

and hence

$$M_{\Lambda, \nabla, \omega'}(\#\{\zeta \cap \nabla\} \geq 1) \geq 1 - q^{\alpha-1} (z|\nabla|)^{-1} = 1 - q^{\alpha-1} \left(\frac{2 \cdot 64}{z\sqrt{3}\ell^2} \right)$$

For any $zR^2 > z_0^*(\alpha^*, q, R)$ there exists $\ell = \ell(z) \in (0, \frac{R}{2\sqrt{3}}]$ such that

$$z\ell^2 > \frac{2 \cdot 64^2 q^{\alpha-1}}{\varepsilon\sqrt{3}}$$

and thus

$$M_{\Lambda, \nabla, \omega'}(\#\{\zeta \cap \nabla\} \geq 1) > 1 - \frac{\varepsilon}{64}$$

holds. □

Let $\Delta_{k,l}$ be an element of the partitioning (2.7) of Λ and define $F_{k,l}^{(\text{ext})}$ to be the event that each of the smaller boxes $\Delta_{k,l}^{i,j} \subset \Delta_{k,l}$ that are not in the centre region $\Delta_{k,l}^{(-)}$, contain at least one point. We call the elements in this event “well-behaved” configurations,

$$F_{k,l}^{(\text{ext})} = \bigcap_{\substack{i, j \in \{0, \dots, 7\}: \\ \Delta_{k,l}^{i,j} \not\subset \Delta_{k,l}^{(-)}}} \{\zeta \in \Omega_{\Lambda, \omega} : \#\{\zeta \cap \Delta_{k,l}^{i,j}\} \geq 1\}. \tag{2.15}$$

Lemma 2.5. *For any $\Lambda \Subset \mathbb{R}^2$ with partitioning (2.7), any boundary condition $\omega \in \Omega$ and well-behaved configuration $\zeta \in F_{k,l}^{(\text{ext})}$ and a point $x_0 \in \Delta_{k,l}^{(-)}$ with $x_0 \notin \zeta$ for any $k, l \in \{-n, \dots, n\}$,*

$$\frac{h_\Lambda(\zeta \cup \{x_0\})}{h_\Lambda(\zeta)} \leq q^\alpha, \tag{2.16}$$

where $\alpha \in (0, \infty)$ is given in Theorem 2.1.

Proof of Lemma 2.5. Then, adapting similar steps in the proof of Lemma 2.3, we see that

$$\begin{aligned} \frac{h_\Lambda(\zeta \cup \{x_0\})}{h_\Lambda(\zeta)} &= \frac{\int q^{K(\zeta, E_2)} \int q^{K(\zeta \cup \{x_0\}, E_1 \cup E_2) - K(\zeta, E_2)} \mu_{x_0, \zeta}^{(+)}(dE_1) \mu_{x_0, \zeta}^{(\text{ext})}(dE_2)}{\int q^{K(\zeta, E_3)} \int q^{K(\zeta, E_3 \cup E_4) - K(\zeta, E_3)} \mu_{x_0, \zeta}^{(-)}(dE_4) \mu_{x_0, \zeta}^{(\text{ext})}(dE_3)} \\ &\leq \frac{q \int q^{K(\zeta, E_2)} \mu_{x_0, \zeta}^{(\text{ext})}(dE_2)}{\int q^{K(\zeta, E_3)} \int q^{K(\zeta, E_3 \cup E_4) - K(\zeta, E_3)} \mu_{x_0, \zeta}^{(-)}(dE_3) \mu_{x_0, \zeta}^{(\text{ext})}(dE_4)} \\ &= q \left(\iint q^{K(\zeta, E_3 \cup E_4) - K(\zeta, E_4)} \mu_{x_0, \zeta}^{(-)}(dE_3) \mu_{(\text{ext}), \zeta}^{(q)}(dE_4) \right)^{-1}, \end{aligned}$$

where we used the inequality

$$K(\zeta \cup \{x_0\}, E_1 \cup E_2) - K(\zeta, E_2) \leq 1, \quad E_1 \subset E_{x_0, \zeta}^{(+)}, E_2 \subset E_{x_0, \zeta}^{(\text{ext})}, \quad (2.17)$$

as all new edges are connected to x_0 and thus can at most built one additional open component. We apply Jensen's inequality to the integral in the denominator to obtain the upper bound

$$\frac{h_\Lambda(\zeta \cup \{x_0\})}{h_\Lambda(\zeta)} \leq q \left(q^{\iint (K(\zeta, E_3 \cup E_4) - K(\zeta, E_4))} \mu_{x_0, \zeta}^{(-)}(dE_3) \mu_{(\text{ext}), \zeta}^{(q)}(dE_4) \right)^{-1}. \quad (2.18)$$

For all configurations $\zeta \in F_{k,l}^{(\text{ext})}$ and $x_0 \in \Delta_{k,l}^{(-)}$ with $x_0 \notin \zeta$ and all $x \in V_{x_0, \zeta}$ we have that $|x - x_0| < \frac{5}{8}\sqrt{3}\ell$ where the bound is the length of the longest diagonal of the rhombus $\Delta_{k,l}^{(-)}$ plus the length of the longer diagonal of one of the smaller rhombuses with side length $\ell/8$. This ensures that

$$V_{x_0, \zeta} \subset B_{\frac{5}{8}\sqrt{3}\ell}(x_0),$$

where $B_{\frac{5}{8}\sqrt{3}\ell}(x_0)$ is the ball of radius $\frac{5}{8}\sqrt{3}\ell$ around x_0 , and as $\ell \leq \frac{R}{2\sqrt{3}}$, the new edges are within the range R of the potential. Therefore, since $x, y \in V_{x_0, \zeta}$ for all $x, y \in E_{x_0, \zeta}^{(-)}$, it follows that adding edges in $E_{x_0, \zeta}^{(-)}$ can only fuse together two connected components (reducing the number of connected components by one) if they each intersect $V_{x_0, \zeta}$. Hence,

$$K(\zeta, E_3 \cup E_4) - K(\zeta, E_4) \geq -N_{x_0}^{(\text{cc})}(\zeta, E_4) + 1, \quad E_3 \subset E_{x_0, \zeta}^{(-)},$$

and thus with Theorem 2.1 we conclude with the statement. \square

We get a lower bound for the conditional probability, given well-behaved configurations $\zeta \in F_{k,l}^{(\text{ext})}$, that $\Delta_{k,l}^{(-)}$ contains no more than $m = m(z)$ points of the well-behaved configurations ζ .

Lemma 2.6. *Given the partitioning (2.7) of Λ with boundary condition $\omega \in \Omega$ and boundary condition $\zeta' \in \Omega$ for $\Delta_{k,l}^{(-)}$ with $\zeta' \cap \Lambda^c = \omega$ and $\zeta' \in F_{k,l}^{(\text{ext})}$ for any $k, l \in \{-n, \dots, n\}$, the estimate*

$$M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(\#\{\zeta \cap \Delta_{k,l}^{(-)} \leq \lfloor m(z) \rfloor\}) > 1 - \varepsilon$$

holds for $m(z) = 2\varepsilon^{-1}q^\alpha |\Delta_{k,l}^{(-)}|z$.

Proof of Lemma 2.6. Note that we can write $M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(d\zeta)$ as

$$M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(d\zeta) = \frac{1}{Z_{\Lambda, \Delta_{k,l}^{(-)}}(\zeta')} h_\Lambda(\zeta \cup \zeta') \Pi_{\Delta_{k,l}^{(-)}}^z(d\zeta),$$

where $Z_{\Lambda, \Delta_{k,l}^{(-)}}(\zeta')$ is the normalisation. Using the well-known fact

$$\int f(\zeta) \Pi_{\Delta_{k,l}^{(-)}}^z(d\zeta) = e^{-z|\Delta_{k,l}^{(-)}|} \sum_{n=0}^{\infty} \frac{z^n}{n!} \int_{\Delta_{k,l}^{(-)}} f(\{x_1, \dots, x_n\}) dx_1 \cdots dx_n$$

for any observable f of the underlying Poisson process and writing $N = \#\{\zeta \cap \Delta_{k,l}^{(-)}\}$ we obtain, setting $Z' = e^{-z|\Delta_{k,l}^{(-)}|} / Z_{\Lambda, \Delta_{k,l}^{(-)}}(\zeta')$ for brevity, the probability for any $n \in \mathbb{N}$,

$$\begin{aligned} M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(N = n + 1) &= \frac{z^{n+1}}{(n + 1)!} Z' \int_{(\Delta_{k,l}^{(-)})^{n+1}} h_{\Lambda}(\{x_1, \dots, x_{n+1}\} \cup \zeta') dx_1 \cdots dx_{n+1} \\ &= \frac{z^n}{n!} \left(\frac{z}{n + 1}\right) Z' \int_{(\Delta_{k,l}^{(-)})^n} \int_{\Delta_{k,l}^{(-)}} h_{\Lambda}(\{x_1, \dots, x_n\} \cup \zeta' \cup \{x\}) dx dx_1 \cdots dx_n \\ &= \frac{z^n}{n!} \left(\frac{z}{n + 1}\right) Z' \int_{(\Delta_{k,l}^{(-)})^n} h_{\Lambda}(\{y\} \cup \zeta') g_{\Delta_{k,l}^{(-)}, \zeta'}(y) dx dy \end{aligned}$$

where

$$g_{\Delta_{k,l}^{(-)}, \zeta'}(y) = \int_{\Delta_{k,l}^{(-)}} \frac{h_{\Lambda}(\{y\} \cup \zeta' \cup \{x\})}{h_{\Lambda}(\{y\} \cup \zeta')} dx, \quad y = \{x_1, \dots, x_n\}.$$

We obtain that

$$\begin{aligned} M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(N = n + 1) &= \left(\frac{z}{n + 1}\right) \int_{\Omega_{\Delta_{k,l}^{(-)}, \zeta'}} g_{\Delta_{k,l}^{(-)}, \zeta'}(\zeta) \mathbb{1}\{N(\zeta) = n\} M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(d\zeta) \\ &= \left(\frac{z}{n + 1}\right) \times M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(N = n) \int_{\Omega_{\Delta_{k,l}^{(-)}, \zeta'}} g_{\Delta_{k,l}^{(-)}, \zeta'}(\zeta) M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(d\zeta | N = n). \end{aligned}$$

By Lemma 2.5 we can bound the function $g_{\Delta_{k,l}^{(-)}, \zeta'}(\zeta)$ for point configurations $\zeta \cup \zeta' \in F_{k,l}^{(\text{ext})}$ from above by $q^\alpha |\Delta_{k,l}^{(-)}|$. Therefore,

$$\begin{aligned} \frac{M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(N = n + 1)}{M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(N = n)} &= \left(\frac{z}{n + 1}\right) \int_{\Omega_{\Delta_{k,l}^{(-)}, \zeta'}} g_{\Delta_{k,l}^{(-)}, \zeta'}(\zeta) M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(d\zeta | N = n) \\ &\leq \left(\frac{z}{n + 1}\right) q^\alpha |\Delta_{k,l}^{(-)}| \int_{\Omega_{\Delta_{k,l}^{(-)}, \zeta'}} M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(d\zeta | N = n) \leq \frac{q^\alpha |\Delta_{k,l}^{(-)}| z}{n + 1}. \end{aligned}$$

For all $n > m(z)$ we apply the previous step $n - \lfloor m(z) \rfloor$ times to obtain

$$\begin{aligned} M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(N = n) &\leq \frac{q^\alpha |\Delta_{k,l}^{(-)}| z}{n} M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(d\zeta | N = n - 1) \\ &\leq \frac{1}{n!} (q^\alpha |\Delta_{k,l}^{(-)}| z)^{n - \lfloor m(z) \rfloor} \lfloor m(z) \rfloor!. \end{aligned}$$

It follows that

$$\begin{aligned} M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(N > m(z)) &\leq \sum_{n=\lfloor m(z) \rfloor + 1}^{\infty} \frac{1}{n!} (q^\alpha |\Delta_{k,l}^{(-)}| z)^{n - \lfloor m(z) \rfloor} \lfloor m(z) \rfloor! \\ &\leq \sum_{n=\lfloor m(z) \rfloor + 1}^{\infty} \left(\frac{q^\alpha |\Delta_{k,l}^{(-)}| z}{\lfloor m(z) \rfloor}\right)^{n - \lfloor m(z) \rfloor}, \end{aligned}$$

and thus

$$M_{\Lambda, \Delta_{k,l}^{(-)}, \zeta'}(N > m(z)) \leq \sum_{n=\lfloor m(z) \rfloor + 1}^{\infty} \left(\frac{\varepsilon}{2}\right)^{n - \lfloor m(z) \rfloor} = \sum_{n=1}^{\infty} \left(\frac{\varepsilon}{2}\right)^n.$$

Since $\varepsilon < \frac{1}{2}$, the right hand side of the previous inequality is less than ε . We conclude with the statement. \square

Step 2: Random-cluster measure $\tilde{C}_{\Lambda, \omega}$.

We find a measure $\tilde{C}_{\Lambda, \omega}$ which is stochastically smaller than $C_{\Lambda, \omega}$. Here, $\tilde{C}_{\Lambda, \omega}$ is stochastically smaller than $C_{\Lambda, \omega}$, in symbols $C_{\Lambda, \omega} \succcurlyeq \tilde{C}_{\Lambda, \omega}$, if

$$\mathbb{E}_{C_{\Lambda, \omega}}[f] \geq \mathbb{E}_{\tilde{C}_{\Lambda, \omega}}[f]$$

for all functions $f: \Omega \times \mathcal{E} \rightarrow \mathbb{R}$ which are increasing in the second entry, see [GHM00, Gri94, GH96] and for an adaptation to the Delaunay graph [Eye14]. Then using coarse graining and comparison to mixed site-bond percolation on \mathbb{Z}^2 we establish percolation for $\tilde{C}_{\Lambda, \omega}$. Percolation for $\tilde{C}_{\Lambda, \omega}$ then implies percolation for the original random cluster measure $C_{\Lambda, \omega}$. We base the definition of the measure $\tilde{C}_{\Lambda, \omega}$ on a coarse graining method originally introduced in [Hag00] and later extended and adapted in [AE16].

For given $\zeta \in \Omega_{\Lambda, \omega}$ and $\ell \in (0, \frac{R}{2\sqrt{3}}]$ let

$$\text{Del}_2^*(\zeta) = \{\eta \in \text{Del}_2(\zeta) : \phi_\beta(\ell(\eta)) \geq g(\beta) := \log(1 + \beta/64\ell^4)\},$$

and let $\tilde{\mu}_{\zeta, \Lambda}$ be the distribution of the random set $\{\eta \in E_\zeta : v(\eta) = 1\}$ with

$$\mathbb{P}(v(\eta) = 1) = \tilde{p}_\ell(\eta) = \frac{1 - \exp\{-g(\beta)\}}{1 + (q - 1) \exp\{-g(\beta)\}} \mathbb{1}_{\text{Del}_2^*(\zeta)}(\eta). \tag{2.19}$$

The function g is increasing in β and is chosen according to the given potential and the coarse graining. Note that $\tilde{\mu}_{\zeta, \Lambda}$ depends on Λ only via the configuration $\zeta \in \Omega_{\Lambda, \omega}$. Note also the important fact that \tilde{p}_ℓ is increasing in β , although to reduce excessive notation, we don't explicitly write this. It is easy to show that $\mu_{\zeta, \Lambda} \succcurlyeq \tilde{\mu}_\zeta$ by noting that

$$\frac{p(\eta)}{q(1 - p(\eta))} = \frac{1 - \exp(-\phi_\beta(\eta))}{q \exp(-\phi_\beta(\eta))} \geq \frac{1 - \exp[-g(\beta)\mathbb{1}_{\text{Del}_2^*(\zeta)}(\eta)]}{q \exp[-g(\beta)\mathbb{1}_{\text{Del}_2^*(\zeta)}(\eta)]} = \frac{\tilde{p}_\ell(\eta)}{q(1 - \tilde{p}_\ell(\eta))}, \quad \eta \in \text{Del}_2(\zeta),$$

and using comparison inequalities in [Gri94, page 78]. Hence, $C_{\Lambda, \omega} \succcurlyeq \tilde{C}_{\Lambda, \omega}$.

As site percolation implies edge percolation we consider site percolation given by (2.19), that is, we open vertices in $\text{Del}_1(\zeta)$ independently of each other with probability \tilde{p}_ℓ . Formally this is defined as follows. We let $\tilde{\lambda}_{\zeta, \Lambda}$ be the distribution of the random mark vector $\tilde{\sigma} \in M_q^\zeta$ where $(\tilde{\sigma}_x)_{x \in \zeta}$ are Bernoulli random variables satisfying

$$\begin{aligned} \mathbb{P}(\tilde{\sigma}_x = 1) &= \tilde{p}_\ell \mathbb{1}_{\text{Del}_1^*(\zeta)}(x) \\ \mathbb{P}(\tilde{\sigma}_x = 0) &= 1 - \tilde{p}_\ell \mathbb{1}_{\text{Del}_1^*(\zeta)}(x), \end{aligned} \tag{2.20}$$

where \tilde{p}_ℓ is given in (2.19) and $\text{Del}_1^*(\zeta)$ is the set of points that build the edges of $\text{Del}_2^*(\zeta)$. Then the site percolation process is defined by the measure

$$\tilde{C}_{\Lambda, \omega}^{(\text{site})}(d\zeta) = M_{\Lambda, \omega}(d\zeta) \tilde{\lambda}_{\zeta, \Lambda}(d\zeta). \tag{2.21}$$

Note that for all $\eta \in \text{Del}_2^*(\zeta)$ and all $\ell \in (0, \frac{R}{2\sqrt{3}}]$,

$$\tilde{p}_\ell \geq \frac{1}{\frac{4R^4}{\beta} + 1} =: \tilde{p}. \tag{2.22}$$

Note that $\tilde{p} = \tilde{p}(\beta)$ is increasing in β with limit $\lim_{\beta \rightarrow \infty} \tilde{p}(\beta) = 1$ and $\tilde{p}(\beta) = 0$.

Step 3: Percolation for $\tilde{C}_{\Lambda,\omega}$.

We now establish percolation for the random-cluster measure $\tilde{C}_{\Lambda,\omega}$. The following lemma provides a uniform lower bound for the probability that a cell Δ in Λ is connected to the complement of Λ .

Lemma 2.7. *There is a $c > 0$ such that*

$$\tilde{C}_{\Lambda,\omega}(\{\Delta \longleftrightarrow \Lambda^c\}) \geq c > 0 \tag{2.23}$$

for any $\Delta = \Delta_{k,l} \subset \Lambda, (k, l) \in \{-n, \dots, n\}^2$, in the partitioning (2.7) of Λ , and any pseudo-periodic boundary condition $\omega \in \Omega_\Lambda^*$.

Proof of Lemma 2.7. The proof is split in several steps and employs the following idea. The site percolation measure $\tilde{C}_{\Lambda,\omega}^{(\text{site})}$ opens certain cells, whereas the corresponding bond percolation measure $\tilde{C}_{\Delta(\text{link}),\zeta'}^{(\text{site})}$ defined in Step (iii) below opens links (bonds) between two neighbouring cells. With comparison to mixed site-bond percolation in \mathbb{Z}^2 we obtain chains of good (open) cells and open links (bonds). This finally ensures percolation for $\tilde{C}_{\Lambda,\omega}$ and the uniform lower bound on the right hand side of (2.23).

Step (i) Probability that small cells have at least one point.

In the following we write $\Delta = \Delta_{k,l}$ and $\nabla = \Delta_{k,l}^{i,j}$ for any $k, l \in \{-n, \dots, n\}$ and for any $i, j \in \{0, 1, \dots, 7\}$. For all configurations $\zeta' \in \Omega_{\Delta^c}$ with $\zeta' \cap \Lambda^c = \omega$ we obtain with Lemma 2.4 the estimate

$$M_{\Lambda,\Delta,\zeta'}(F_{k,l}^{(\text{ext})}) \geq 1 - \sum_{i,j: \Delta_{k,l}^{i,j} \not\subset \Delta_{k,l}} M_{\Lambda,\Delta,\zeta'}(\#\{\zeta \cap \Delta_{k,l}^{i,j}\} = 0) \geq 1 - \frac{48\varepsilon}{64} = 1 - \frac{3\varepsilon}{4}. \tag{2.24}$$

Define the following two events, first the event

$$G_{k,l} = \{\zeta \in \Omega_{\Lambda,\omega} : \#\{\zeta \cap \Delta_{k,l}^{(-)}\} \leq m(z)\}$$

that there are at most $m(z)$ points in ζ in the centre $\Delta_{k,l}^{(-)}$ of $\Delta_{k,l}$, and the event that all smaller cells in that centre contain at least one point,

$$F_{k,l}^{(-)} = \bigcap_{i,j: \Delta_{k,l}^{i,j} \subset \Delta_{k,l}^{(-)}} \{\zeta \in \Omega_{\Lambda,\omega} : \#\{\zeta \cap \Delta_{k,l}^{i,j}\} \geq 1\}.$$

We have tacitly replaced $m(z)$ by $m(z) \vee 16$. Both events depend on point configurations in the centre region $\Delta_{k,l}^{(-)}$, and it suffices to estimate the probability of the intersection of these two events for any boundary condition outside of $\Delta_{k,l}$ and any point configuration ζ'' in $\Delta_{k,l} \setminus \Delta_{k,l}^{(-)}$. For all boundary conditions $\zeta' \in \Omega_{\Delta_{k,l}^c}$ with $\zeta' \cap \Lambda^c = \omega$,

$$M_{\Lambda,\Delta,\zeta'}(F_{k,l}^{(-)} \cap G_{k,l} \cap F_{k,l}^{(\text{ext})}) = \int \mathbb{1}_{F_{k,l}^{(\text{ext})}}(\zeta'' \cup \zeta') \left[\int \mathbb{1}_{F_{k,l}^{(-)}}(\zeta) \mathbb{1}_{G_{k,l}}(\zeta) M_{\Lambda,\Delta,\zeta'}(d\zeta | \zeta = \zeta'' \text{ on } \Delta \setminus \Delta_{k,l}^{(-)}) \right] M_{\Lambda,\Delta,\zeta'}(d\zeta'').$$

Using Lemma 2.6 it follows that

$$\begin{aligned} M_{\Lambda,\Delta_{k,l}^{(-)},\zeta'' \cup \zeta'}(F_{k,l}^{(-)} \cap G_{k,l}) &\geq 1 - M_{\Lambda,\Delta_{k,l}^{(-)},\zeta'' \cup \zeta'}((F_{k,l}^{(-)})^c) - M_{\Lambda,\Delta_{k,l}^{(-)},\zeta'' \cup \zeta'}((G_{k,l})^c) \\ &> 1 - \frac{16\varepsilon}{64} - \varepsilon = 1 - \frac{5\varepsilon}{4}, \end{aligned} \tag{2.25}$$

and hence we conclude with (2.24) that

$$\begin{aligned} M_{\Lambda,\Delta_{k,l},\zeta'}(F_{k,l}^{(-)} \cap G_{k,l} \cap F_{k,l}^{(\text{ext})}) &> \left(1 - \frac{5\varepsilon}{4}\right) \int \mathbb{1}_{F_{k,l}^{(\text{ext})}}(\zeta'' \cup \zeta') M_{\Lambda,\Delta,\zeta'}(d\zeta'') \\ &> \left(1 - \frac{5\varepsilon}{4}\right) \left(1 - \frac{3\varepsilon}{4}\right) > 1 - 2\varepsilon. \end{aligned} \tag{2.26}$$

Step (ii): Good cells and site percolation.

After these preparation steps we shall define when a cell $\Delta_{k,l}$ is good. A good cell at (k, l) will result in the site $(k, l) \in \mathbb{Z}^2$ being open. The next step is therefore to condition the marks of the points, that is, we pick $(k, l) \in \{-n, \dots, n\}^2$ and consider the event $C_{k,l}$ that each cell $\Delta_{k,l}^{i,j}$ has at least one point, $\Delta_{k,l}^{(-)}$ contains no more than $m(z)$ points and all points in $\Delta_{k,l}^{(-)} \cap \text{Del}_1^*(\zeta)$ are carrying mark 1,

$$C_{k,l} = \{\zeta \in \Omega_{\Lambda,\omega} : \zeta \in F_{k,l}^{(-)} \cap F_{k,l}^{(\text{ext})} \cap G_{k,l} \text{ and } \sigma_\zeta(x) = 1 \text{ for all } x \in \Delta_{k,l}^{(-)} \cap \text{Del}_1^*(\zeta)\}.$$

A cell $\Delta_{k,l}$ is declared to be “good” or “open” if $C_{k,l}$ occurs. Each vertex $x \in \text{Del}_1^*(\zeta)$ is open with probability \tilde{p}_ℓ (see (2.20)). It follows that

$$\begin{aligned} \tilde{C}_{\Lambda,\omega}^{(\text{site})}(C_{k,l}) &\geq \int M_{\Lambda,\Delta_{k,l},\zeta'}(d\zeta) \mathbb{1}_{F_{k,l}^{(-)}}(\zeta) \mathbb{1}_{F_{k,l}^{(\text{ext})}}(\zeta) \mathbb{1}_{G_{k,l}}(\zeta) \tilde{p}_\ell^{\#\{\text{Del}_1^*(\zeta) \cap \Delta_{k,l}^{(-)}\}} \\ &\geq \tilde{p}_\ell^{\lfloor m(z) \rfloor} M_{\Lambda,\Delta_{k,l},\zeta'}(F_{k,l}^{(-)} \cap G_{k,l} \cap F_{k,l}^{(\text{ext})}). \end{aligned}$$

Recall from Lemma 2.4 that for $z > z^*(\alpha^*, q, R)$ there is $\ell \in (0, \frac{R}{2\sqrt{3}}]$ such that the estimates. Recall that $|\Delta^{(-)}| = \frac{\sqrt{3}}{8} \ell^2 \leq \frac{R^2}{32\sqrt{3}}$ and

$$\lfloor m(z) \rfloor \leq \frac{2q^\alpha R^2 z}{32\sqrt{3}\varepsilon} \leq \lceil m(z) \rceil \leq \frac{2q^{\alpha^*} R^2 z}{32\sqrt{3}\varepsilon}.$$

For all $z > z_0^*(\alpha^*, q, R)$ choose $\beta_0 = \beta_0(q, R, z) > 0$ such that

$$(\tilde{p})^{\frac{q^{\alpha^*} R^2 z}{16\sqrt{3}\varepsilon}} \geq (1 - 2\varepsilon) \quad \text{for all } \beta \geq \beta_0 \vee q, \tag{2.27}$$

where α^* is the bound for α (2.5) for any $\beta > q$. Then, for all $z > z_0^*(\alpha^*, q, R)$ there is $\ell \in (0, \frac{R}{2\sqrt{3}}]$ and there is $\beta_0 = \beta_0(q, R, z)$ such that for all $\beta \geq \beta_0 \vee q$

$$\tilde{p}_\ell^{\lfloor m(z) \rfloor} \geq \tilde{p}^{\lfloor m(z) \rfloor} \geq (1 - 2\varepsilon). \tag{2.28}$$

Combining the above estimates, we conclude, for all $\beta \geq \beta_0 \vee q$, that

$$\tilde{C}_{\Delta_{k,l},\zeta'}^{(\text{site})}(C_{k,l}) \geq (1 - 2\varepsilon)^2 > 1 - 4\varepsilon > (p_c^{(\text{site})}(\mathbb{Z}^2))^{1/2}. \tag{2.29}$$

Step (iii) Neighbouring good cells and link percolation.

If $\zeta \in C_{k,l}$, we say that the cell $\Delta_{k,l}$ is a “good” cell. Two neighbouring cells $\Delta_{k,l}$ and $\Delta_{k+1,l}$ are said to be “linked” if the box $\Delta_{(\text{link})} := \Delta_{(\text{link})}^{k:k+1,l}$ defined in (2.8) has an intersection with $\text{Del}_1^*(\zeta)$ that contains only points of mark 1. More precisely, the event that $\Delta_{k,l}$ and $\Delta_{k+1,l}$ are linked, is

$$L_{k:k+1}^l = \{\zeta \in \Omega_{\Lambda,\omega} : \sigma_\zeta(x) = 1 \text{ for all } x \in \Delta_{(\text{link})}^{k:k+1,l} \cap \text{Del}_1^*(\zeta)\}.$$

We also define

$$F_{(\text{link})} = \left(F_{k,l}^{(-)} \cap F_{k,l}^{(\text{ext})} \right) \cap \left(F_{k+1,l}^{(-)} \cap F_{k+1,l}^{(\text{ext})} \right)$$

and

$$G_{(\text{link})} = \{\zeta \in \Omega : \#\{\zeta \cap \Delta_{(\text{link})}\} \leq m(z)\},$$

and let $\zeta' \in \Omega_{\Delta_{(\text{link})}^c}$ be the boundary condition outside $\Delta_{(\text{link})}$ such that $\zeta' \cap \Lambda^c = \omega$. The conditional probability that the cells $\Delta_{k,l}$ and $\Delta_{k+1,l}$ are linked, given they are both

“good” cells, is then given by

$$\begin{aligned} \tilde{C}_{\Delta_{(\text{link})}, \zeta'}^{(\text{site})}(L_{k,k+1}^l | C_{k,l} \cap C_{k+1,l}) &\geq \int \mathbb{1}_{G_{(\text{link})}}(\zeta) \tilde{p}^{\lfloor m(z) \rfloor} M_{\Lambda, \Delta_{(\text{link})}, \zeta'}(d\zeta | F_{(\text{link})}) \\ &\geq (1 - \varepsilon)(1 - 2\varepsilon) \geq (1 - 4\varepsilon) \geq (p_c^{(\text{site})}(\mathbb{Z}^2))^{1/2}, \end{aligned} \tag{2.30}$$

where the second inequality comes from an adaptation of Lemma 2.6, where $\Delta_{(\text{link})}$ takes the role of $\Delta_{k,l}^{(-)}$.

Step (iv): Mixed site-bond percolation and proof of (2.23).

Then by (2.29), (2.30) and the results of McDiarmid and Hammersley, in particular, (C.4), mixed site-bond percolation in \mathbb{Z}^2 occurs. There exists a chain of good cells joined by open links from $\Delta_{k,l} \subset \Lambda$ to Λ^c .

It remains to check that the preceding result implies $\{\Delta \leftrightarrow \Lambda^c\}$ in the Delaunay graph. We establish this by showing that we can connect a point $x \in \zeta$ whose Voronoi cell contains the centre of the cell $\Delta_{k,l}$ to a point y in the neighbouring cell $\Delta_{k+1,l}$ whose Voronoi cell contains the centre of $\Delta_{k+1,l}$. For this, recall the set $H_{k:k+1,l}(\zeta)$ defined in (2.10). We know by construction that all edges $\eta = \{u, v\} \in \text{Del}_2(\zeta)$ that have a non-empty intersection with $H_{k:k+1,l}(\zeta)$ satisfy $|u - v| < 2\sqrt{3}\frac{\ell}{8}$. This implies that $H_{k:k+1,l}(\zeta) \subset \text{Del}_1^*(\zeta)$, and therefore one can find an open chain of Delaunay edges intersecting the link in between cells and within cells. Let $x, y \in \zeta$ be such that $\text{Vor}_\zeta(x)$ and $\text{Vor}_\zeta(y)$ contain the centres of the boxes $\Delta_{k,l}$ and $\Delta_{k+1,l}$ respectively. Since $H_{k:k+1,l}(\zeta) \subset \text{Del}_1^*(\zeta)$, we can connect x and y in the graph $\text{Del}_2^*(\zeta)$ inside $\Delta_{k,l}^{(-)} \cup \Delta_{(\text{link})} \cup \Delta_{k+1,l}^{(-)}$. Hence, by (2.29) and (2.30), there exists $c > 0$ such that the following uniform lower bound holds

$$\tilde{C}_{\Lambda, \omega}^{(\text{site})}(\{\Delta \leftrightarrow \Lambda^c\}) > c > 0,$$

and the proof of Lemma 2.7 is finished. □

Step 4: Finish of the proof of Proposition 2.2.

The proof follows immediately from all previous steps as percolation for the percolation measure $\tilde{C}_{\Lambda, \omega}$ implies percolation in the Delaunay random cluster measure $C_{\Lambda, \omega}$ due to stochastic dominance,

$$C_{\Lambda, \omega} \succcurlyeq \tilde{C}_{\Lambda, \omega}. \tag{□}$$

2.3 Symmetry breaking of the mark distribution

To relate the influence of the boundary condition on the mark of a single point to the connectivity probabilities in the random-cluster model we follow [GH96]. For any $\Delta \subset \Lambda$, $s \in M_q$, $\zeta \in \Omega_{\Lambda, \omega}$ and (ζ, E) , with $E \subset E_\zeta$ we define

$$N_{\Delta, s}(\zeta) = \#\{\zeta^{(s)} \cap \Delta\}.$$

Then

$$N_{\Delta \leftrightarrow \Lambda^c}(\zeta, E) = \#\{x \in \zeta \cap \Delta : x \text{ belongs to a cluster connected to } \Lambda^c \text{ in } E \cap \text{Del}_2(\zeta)\}$$

is the number of points in $\zeta \cap \Delta$ connected to any point in Λ^c in the random graph $E \cap \text{Del}_2(\zeta)$. Because of the edge-drawing mechanism (edges are open when the two vertices share the same type), $\{\Delta \leftrightarrow \Lambda^c\} = \{N_{\Delta \leftrightarrow \Lambda^c} > 0\}$ is also the event that there exists a point in $\zeta \cap \Delta$ connected to infinity in $E \cap \text{Del}_2(\zeta)$.

The next Proposition is the key argument why percolation for the random cluster measures leads to a break of symmetry in the mark distribution.

Proposition 2.8. For any measurable $\Delta \subset \Lambda \in \mathbb{R}^2$,

$$\int (qN_{\Delta,1} - N_{\Delta}) d\gamma_{\Lambda,\omega} = (q-1) \int N_{\Delta \leftrightarrow \Lambda^c} dC_{\Lambda,\omega}.$$

Proof. This is proved in [Eye14, Lemma 2.17] following ideas in [GH96]. \square

The statement in Proposition 2.8 is well-known result for lattice systems, see [GHM00] for an overview. For continuum systems it appeared in [GH96] for the first time. It basically shows a break in the symmetry of the type distribution when percolation is established in the corresponding random cluster model. The aim is to find uniform (in Λ and ω) lower bounds for the left hand side away from zero to ensure the break of the symmetry.

3 Number of connected components

In this section we are going to prove the main technical tool for our phase transition proof, namely, Theorem 2.1. The proof of Theorem 2.1 is rather long, so we first outline the strategy. We want to bound the number of connected components in the graph $E_{x_0,\zeta}^{(\text{ext})}$ that intersect the boundary graph $G_{x_0,\zeta}$ under the edge drawing mechanism $\mu_{(\text{ext}),\zeta}^{(q)}$. We also define G_B the contraction of $G_{x_0,\zeta}$ to the ball $B := B_R(x_0)$ around x_0 as due to the finite range condition the point x_0 cannot be connected to any point further away than $R > 0$. More precisely, $G_B = (V_B, E_B)$ with $V_B = V_{x_0,\zeta} \cap B$ and $E_B = \{\eta_{x,y} \in \text{Del}_2(\zeta \cap B) : x, y \in V_B\}$. The pivotal point of the whole proof is to find an upper bound for the number of edges in the edge set E_B that have length greater than some fixed real number. This allows us to construct families consisting of edges in E_B , defined by edge length, to balance the unbounded number of points against the increased probability that they are connected. The shorter the edge length, the greater the possible number of edges in the subset, but also the greater the probability that these edges are open. It turns out that such an upper bound can be found in the scenario where there are no defects in the geometry. These defects which we give the logical name “kinks” are defined below in Section 3.1. An upper bound cannot be found if the geometry of the contracted graph contains kinks, so we devise a plan to discount them.

For $R > 0$ the following three cases depend on the configuration ζ and the point x_0 . (i) $V_B = \emptyset$ in which case there is no connection to any connected component of $E_{x_0,\zeta}^{(\text{ext})}$, (ii) $V_B \supset V_{x_0,\zeta}$, and (iii) $V_B \cap V_{x_0,\zeta} \neq \emptyset$ and $B_R(x_0)^c \cap V_{x_0,\zeta} \neq \emptyset$. In case (ii) we have that $E_B \subset E_{x_0,\zeta}^{(\text{ext})}$ but this does not hold in case (iii). This creates a problem when dealing with our edge drawing mechanism on $E_{x_0,\zeta}^{(\text{ext})}$. To overcome this, we introduce an edge drawing mechanism on the contracted graph and build a structure that will allow us to compare events between the two different probability spaces. All our techniques rely heavily on geometric properties of the Delaunay tessellation.

3.1 Notation and geometric facts

We introduce a polar coordinate system in \mathbb{R}^2 with x_0 being the pole, and we denote L the polar axis in horizontal direction. For any $z \in \mathbb{R}^2$, denote \hat{z} be the angular coordinate of z taken counter clockwise from the axis L . Given two points $x, y \in \mathbb{R}^2$, \overleftrightarrow{xy} denotes the unique straight line that intersects x and y in the plane, \overleftarrow{xy} denotes the half line that stops at y and \overline{xy} denotes the line segment between x and y only. Given two straight lines $\ell_1, \ell_2 \subset \mathbb{R}^2$ that intersect a point $z \in \mathbb{R}^2$, $\angle(\ell_1, \ell_2)$ denotes the angle between them. More precisely, it is the angle in order to rotate ℓ_1 onto ℓ_2 with z being the centre of rotation. Notice that $\angle(\ell_1, \ell_2) = \angle(\ell_2, \ell_1)$ only if $\angle(\ell_1, \ell_2) = \pi/2$, however, it holds that $\angle(\ell_1, \ell_2) + \angle(\ell_2, \ell_1) = \pi$. When we consider a triangle in the plane with vertices x, y, z ,

we refer to the interior angle at y as \widehat{xyz} . In this case, as we specify the interior angle, $\widehat{xyz} = \widehat{zyx}$.

Given a set of points $V = \{x_i \in \mathbb{R}^2 : 1 \leq i \leq n\}$ with $\hat{x}_1 < \dots < \hat{x}_n$, the graph

$$\Gamma = \left(V, \bigcup_{i=1}^{n-1} \eta_{x_i, x_{i+1}} \right) \tag{3.1}$$

is called a **spoked chain** with pole x_0 if $\eta_{x_0, x_i} \in \text{Del}_2(V \cup \{x_0\})$ for all $1 \leq i \leq n$. The polygon $P(\Gamma, x_0)$ created by adding the point x_0 and edges η_{x_0, x_1} and η_{x_0, x_n} to Γ is called the induced polygon of Γ , see Figure 5.

In order to quantify the number of connected components that intersect V_B , we analyse the shape of the contracted boundary graph G_B . For that we split B into four quadrants, $Q_i \subset \mathbb{R}^2, i = 1, 2, 3, 4$, where

$$Q_i = \{z \in B : \frac{\pi}{2}(i-1) \leq \hat{z} < \frac{\pi}{2}i\}.$$

This allows to bound the number of connected components in one quadrant, and then the final bound is just four times this bound because two points connected in a quadrant are also connected in the full disc. The reasons for doing this are twofold: it not only provides us a framework to define kinks, but also ensures that any two points that we consider will differ in angle by no more than $\pi/2$. This allows to find a lower bound for the probability that the two points belong to the same connected component, that is, a lower bound that neighbouring points are connected by an open edge. This lower bound is the crucial steps as it allows to balance a possible increase in the number of points as mentioned earlier.

Definition 3.1 (Kinks). Let $\Gamma = (V, E)$ be a spoked chain with pole $x_0 \notin V$. Suppose that $x_i, x_j, x_k \in V$ such that $\hat{x}_i < \hat{x}_j < \hat{x}_k$. We say that x_i, x_j and x_k form a **kink** in Γ if the following holds.

1. $\widehat{x_i x_j x_k} < \pi/2$,
2. $\widehat{x_{i'} x_{j'} x_{k'}} \geq \pi/2$ for all $x_{i'}, x_{j'}, x_{k'} \in V$ with $\hat{x}_{i'} < \hat{x}_{j'} < \hat{x}_{k'}$ and satisfying

$$\hat{x}_i \leq \hat{x}_{i'} < \hat{x}_{j'} < \hat{x}_{k'} < \hat{x}_k \text{ or } \hat{x}_i < \hat{x}_{i'} < \hat{x}_{j'} < \hat{x}_{k'} \leq \hat{x}_k.$$

Suppose that x_i, x_j and x_k form a kink in the spoked chain $\Gamma = (V, E)$. The kink is called **intruding** if the line segment $\overline{x_i x_k}$ lies outside of the induced Polygon $P(\Gamma, x_0)$ and **protruding** if it lies inside $P(\Gamma, x_0)$, see Figure 6.

Lemma 3.2. Let $\Gamma = (V, E)$ be a spoked chain with $V = \{x_1, \dots, x_n\}$ and $\hat{x}_1 < \dots < \hat{x}_n$ and pole $x_0 \notin V$. A kink in Γ is either intruding or protruding.

Proof of Lemma 3.2. Suppose the statement is false. Then there exist $1 \leq i < j < k \leq n$ such that $\hat{x}_i < \hat{x}_j < \hat{x}_k$ with x_0 being the pole such that x_i, x_j and x_k form a kink in Γ and $\overline{x_i x_k}$ lies neither inside nor outside of $P(\Gamma, x_0)$. Let $U \subset \mathbb{R}^2$ be the connected component of $\mathbb{R}^2 \setminus \overrightarrow{x_i x_k}$ that does not contain x_j . Consider first that $\overline{x_i x_k}$ lies inside of $P(\Gamma, x_0)$. It follows that edges of Γ cross $\overline{x_i x_k}$ between \hat{x}_i and \hat{x}_k and hence there exists $x_{j'} \in V \cap U$ with $\hat{x}_i < \hat{x}_{j'} < \hat{x}_k$. Without loss of generality, let $\hat{x}_j < \hat{x}_{j'} < \hat{x}_k$. Therefore, as $x_{j'} \in V \cap U$ with $x_j \notin U$ and $\widehat{x_i x_j x_k} < \pi/2$ one gets that $\widehat{x_i x_j x_{j'}} < \pi/2$ which contradicts property (2) in Definition 3.1 for the kink formed by x_i, x_j and x_k . The second alternative case follows analogously. \square




Figure 5: From top to bottom we have: (1.) A collection of points that neighbour x_0 in the Delaunay (Voronoi) tessellation. (2.) A spoked chain Γ with pole x_0 . (3.) The induced polygon $P(\Gamma, x_0)$.




Figure 6: (a) The points x_2, x_4 and x_5 form an intruding kink; (b) the points x_2, x_3 and x_5 form an extruding kink.

Lemma 3.3. Let $\Gamma = (V, E)$ be a spoked chain with $V = \{x_1, \dots, x_n\}$ and $\hat{x}_1 < \dots < \hat{x}_n$. If x_i, x_j and x_k form an intruding kink in Γ , then

$$\angle(\overrightarrow{\hat{x}_i x_{i+1}}, \overrightarrow{\hat{x}_{k-1} x_k}) < \pi/2.$$

Proof of Lemma 3.3. Let x_i, x_j and x_k form an intruding kink in Γ . Since the kink is intruding, we know that x_l lies in the interior of the triangle $\tau(x_0, x_i, x_k)$ for all $i < l < k$. Suppose the statement of the Lemma is false, that is,

$$\angle(\overrightarrow{\hat{x}_i x_{i+1}}, \overrightarrow{\hat{x}_{k-1} x_k}) \geq \pi/2.$$

This forces either x_{i+1} or x_{k-1} to be in the interior of the triangle $\tau(x_i, x_j, x_k)$. Without loss of generality, suppose, in fact, that x_{i+1} is in the interior of that triangle. Therefore, $\widehat{x_{i+1} x_j x_k} < \pi/2$ which, by Definition 3.1, contradicts the fact that the points x_i, x_j and x_k form a kink in Γ . \square




Figure 7: Lower bound for the angle $\widehat{x_0 x_{j-1} x_j}$.

3.2 Intermediary lemmas

In the following we assume that the points in the graph are ordered according to increasing angles and we define $G_B \cap Q_1$ be the of the sub-graph of G_B whose edges and points are contained in Q_1 . Then $G_B \cap Q_1$ is a spoked chain with pole x_0 .

Kinks of intruding and protruding nature may occur in $G_B \cap Q_1$, although their number is bounded from above, see Lemma 3.4 and Lemma 3.5 below. We shall separate $G_B \cap Q_1$ into a finite number of kink-less pieces, each of which a spoked chain, which are easier to work with.

Lemma 3.4. *The number of intruding kinks in $G_B \cap Q_1$ is bounded above by 2.*

Proof of Lemma 3.4. We show that the angle between two intruding kinks in a spoked chain is greater than $\pi/4$. Since $G_B \cap Q_1$ lies in the quadrant Q_1 , and is a spoked chain by definition, the statement follows immediately. Let $\Gamma = (V, E)$ be a spoked chain and order the elements of V such that $\hat{x}_1 < \dots < \hat{x}_n$. Suppose there is an intruding kink in Γ . By the definition of an intruding kink and Lemma 3.3, there exist $1 \leq i < n - 1$ and $i + 1 < j \leq n$, such that

$$\angle(\overrightarrow{\hat{x}_i x_{i+1}}, \overrightarrow{\hat{x}_{j-1} x_j}) < \pi/2, \tag{3.2}$$

and $\overline{x_i x_j}$ lies outside of the induced polygon $P(\Gamma, x_0)$. The straight lines $\overrightarrow{\hat{x}_i x_{i+1}}$ and $\overrightarrow{\hat{x}_{j-1} x_j}$ split the plane into four regions. Since the kink is intruding, the point x_0 must lie in the opposite region to that of the line segment $\overline{x_i x_j}$. Let L^* be the radial line of angle $(\hat{x}_{i+1} - \hat{x}_{j-1})/2$ and $z_1 \in \mathbb{R}^2$ be the point of the intersection of $\overrightarrow{\hat{x}_i x_{i+1}}$ and $\overrightarrow{\hat{x}_{j-1} x_j}$ and let z_2, z_3 be the points of intersection of L^* with $\overrightarrow{\hat{x}_i x_{i+1}}$ and $\overrightarrow{\hat{x}_{j-1} x_j}$ respectively, see Figure 7. Then,

$$\widehat{x_i z_1 x_j} + \widehat{x_i z_2 x_0} + \widehat{x_0 z_3 x_j} = 2\pi, \tag{3.3}$$

which implies, together with (3.2), that

$$\max\{\widehat{x_i z_2 x_0}, \widehat{x_0 z_3 x_j}\} \geq \frac{2\pi - \pi/2}{2} = \frac{3\pi}{4}. \tag{3.4}$$

Without loss of generality, let $\widehat{x_0 z_3 x_j} \geq 3\pi/4$. Because x_{j-1} lies on the line segment $\overline{z_3 x_j}$, it follows that

$$\widehat{x_0 x_{j-1} x_j} \geq \widehat{x_0 z_3 x_j} \geq 3\pi/4.$$

Suppose now that there is another intruding kink in Γ , formed by the points x_k, x_l and x_m for $j < k < l < m \leq n$. Then, by Lemma 3.3, we have that

$$\angle(\overrightarrow{\hat{x}_k x_{k+1}}, \overrightarrow{\hat{x}_{m-1} x_m}) < \pi/2. \tag{3.5}$$

Let t_{k+1} be the tangent to boundary $\partial B(\tau(x_0, x_k, x_{k+1}))$ at x_{k+1} , where

$$B(\tau(x_0, x_k, x_{k+1}))$$

is the ball with centre x_{k+1}^* whose boundary line intersects the points of the given triangle $\tau(x_0, x_k, x_{k+1})$, see Figure 8 for details. Then, by noting that

$$|V \cap B(\tau(x_0, x_{j-1}, x_j))| = 0, \tag{3.6}$$

which is a consequence of quadratic position, it follows that

$$\angle(t_{k+1}, \overrightarrow{\hat{x}_0 x_{k+1}}) \leq \angle(\overrightarrow{\hat{x}_k x_{k+1}}, \overrightarrow{\hat{x}_0 x_{k+1}}) \leq \angle(\overrightarrow{\hat{x}_k x_{k+1}}, \overrightarrow{\hat{x}_{m-1} x_m}) < \pi/2. \tag{3.7}$$

Here, the first inequality follows directly from the definition of the tangent line and (3.6) and the second inequality is a consequence of the fact that

$$\hat{x}_{k+1} < \hat{x}_{m-1} < \hat{x}_m.$$




Figure 8: The intruding kink formed by x_k, x_l and x_m .

For $1 \leq r < n$, let x_{r+1}^* denote the centre of the circumcircle of the triangle

$$\tau(x_0, x_r, x_{r+1}) \in \text{Del}_3(V \cup \{x_0\}),$$

whereas the third inequality follows from (3.5). Since the points $\{x_0, x_{k+1}^*, x_{k+1}\}$ form an isosceles, see Figure 8, we can conclude from (3.7) that

$$\hat{x}_{k+1} - \hat{x}_{k+1}^* = \pi/2 - \angle(t_{k+1}, \overrightarrow{x_0 x_{k+1}}) > 0. \tag{3.8}$$

Let y be the antipodal point to x_0 on the circumscribed ball of the triangle $\tau(x_0, x_{j-1}, x_j)$ in \mathbb{R}^2 . Since, $|x_0 - y|$ is equal to the diameter of that circle, it follows that $\widehat{x_0 x_j y} = \pi/2$, see Figure 9. The points x_0, x_{j-1}, x_j and y form a cyclic quadrilateral. Using

$$\widehat{x_0 x_{j-1} x_j} \geq \widehat{x_0 z_3 x_j} \geq 3\pi/4$$

from above, and the fact that opposite angles of a cyclic quadrilateral add up to π , we see that $\widehat{x_0 y x_j} \leq \pi/4$. Hence, by (3.8)

$$\hat{x}_{k+1} - \hat{x}_j > \hat{x}_{k+1}^* - \hat{x}_j \geq \hat{x}_j^* - \hat{x}_j = \widehat{y x_0 x_j} = \pi - \widehat{x_0 x_j y} - \widehat{x_0 y x_j} \geq \pi/4,$$

where the second inequality is due to a further property of the Delaunay structure, see Lemma D.1 in appendix D. This implies that the angle between intruding kinks must be greater than $\pi/4$. \square

Lemma 3.5. *There are no protruding kinks in $G_B \cap Q_1$.*

Proof of Lemma 3.5. We order the elements of $V_B = \{x_1, \dots, x_n\}$ such that

$$\hat{x}_1 < \dots < \hat{x}_n.$$

Suppose we have a protruding kink with pole $x_0 \notin V$, then we have for some

$$1 \leq i < j < k \leq n \quad \text{that} \quad \widehat{x_i x_j x_k} < \pi/2.$$

The pair $\{x_i, x_k\}$ does not form an edge of E_B , therefore, by the properties of the Delaunay tessellation, x_j lies inside the circumcircle $\partial B(\tau(x_0, x_i, x_k))$ of the triangle $\tau(x_0, x_i, x_k)$. The line segment $\overline{x_i x_k}$ is a chord which splits the ball $B(\tau(x_0, x_i, x_k))$ into two regions. Since we have a protruding kink, $\overline{x_i x_k}$ lies inside the induced polygon $P(G_B \cap Q_1; x_0)$, and so x_j does not lie in the same region as x_0 . The angle $\widehat{x_i x_j x_k} = \pi/2$ once the point x_j lies on the boundary of the ball $B(\tau(x_0, x_i, x_k))$ for the case $\widehat{x_i x_0 x_k} = \pi$, and due to the fact that $\widehat{x_i x_j x_k} < \pi/2$ we get that $\widehat{x_i x_0 x_k} \geq \pi/2$, and hence, there are no protruding kinks in $G_B \cap Q_1$. \square




Figure 9: Lower bound for the angle between kinks of type 2.

3.3 Edge drawing

Before we can finish the proof of Theorem 2.1 in Section 3.4 below we need two more results to gain some control over the edge drawing mechanism in $E_{x_0, \zeta}^{(\text{ext})}$. Denote by $\tilde{\nu}_\zeta$ the edge drawing mechanism with probability

$$\tilde{p}_2(\eta_{xy}) = \begin{cases} \frac{\mathbb{1}\{|x-y| \leq R\}}{\frac{q}{\beta}|x-y|^4 + 1} & \text{if } \eta_{xy} \in \text{Del}_2(\zeta) \cap (E_{\mathbb{R}^2} \setminus E_{\Lambda^c}), \\ \mathbb{1}_{\text{Del}_2(\zeta)}(\eta_{xy}) & \text{if } \eta_{xy} \in E_{\Lambda^c}, \end{cases} \quad (3.9)$$

and denote $\tilde{\nu}_\zeta^{(\text{ext})}$ the corresponding edge drawing mechanism on $E_{x_0, \zeta}^{(\text{ext})}$.

Lemma 3.6. For all $q \geq 1$, $\Lambda \in \mathbb{R}^2$ and all $\zeta \in \Omega_{\Lambda, \omega}$,

$$\mu_{\Lambda, \zeta}^{(q)} \succcurlyeq \tilde{\nu}_\zeta.$$

Proof of Lemma 3.6. It suffices to show that for all edges $\eta \in \text{Del}_2(\zeta)$ with $\eta \notin E_{\Lambda^c}$,

$$\frac{p(\eta)}{q(1-p(\eta))} \geq \frac{\tilde{p}_2(\eta)}{(1-\tilde{p}_2(\eta))}. \quad (3.10)$$

Recall from (2.1) that

$$p(\eta) = \frac{1}{\frac{1}{\beta} \ell(\eta)^4 + 1}.$$

Thus, if $\ell(\eta) > R$, then $\tilde{p}(\eta) = 0$ and the inequality (3.10) trivially holds. Suppose that $\ell(\eta) < R$, then, in fact, we also have

$$\frac{p(\eta)}{q(1-p(\eta))} = \frac{\tilde{p}_2(\eta)}{(1-\tilde{p}_2(\eta))}.$$

Henceforth, (3.10) holds for all $\eta \in \text{Del}_2(\zeta)$. □

Note that E_B is not necessarily a subset of $E_{x_0, \zeta}^{(\text{ext})}$, in fact, they belong to different Delaunay tessellations

$$E_B \subset \text{Del}_2(V_B \cup \{x_0\}) \text{ and } E_{x_0, \zeta}^{(\text{ext})} \subset \text{Del}_2(\zeta \cup \{x_0\}).$$

We therefore introduce another edge drawing mechanism, but this time on E_B . Let μ^* denote the distribution of the random edge configurations $\{\eta \in E_B : v(\eta) = 1\}$, where $((v(\eta))_{\eta \in E_B})$ are independent Bernoulli random variables with probability

$$\mathbb{P}(v(\eta) = 1) = p^*(\eta) = \frac{\mathbb{1}\{\ell(\eta) \leq \frac{2}{\pi} \wedge R\} \mathbb{1}\{|\hat{x} - \hat{y}| \leq \frac{\pi}{2}\}}{\frac{q}{\beta} (\frac{\pi}{2} \ell(\eta)^4) + 1} \mathbb{1}_{E_B}(\eta), \quad \text{for } \eta = \eta_{xy}. \quad (3.11)$$

We now compare the probability that two points are connected with respect to $\tilde{\nu}_\zeta$ and with respect to μ^* .

Lemma 3.7. *Pick $\zeta \in \Omega_{\Lambda, \omega}$. Let $\eta_{xy} \in E_B$ and let $x \leftrightarrow y$ denote the event that x and y lie in the same connected component of (ζ, E) , where E is a \tilde{p}_2 -thinning of the edge set $E_{x_0, \zeta}^{(\text{ext})}$. Then,*

$$\tilde{\nu}_\zeta^{(\text{ext})}(x \leftrightarrow y) \geq p^*(\eta_{xy}). \quad (3.12)$$

Proof of Lemma 3.7. By the definition of p^* , (3.12) follows for $x, y \in V_B$ with

$$|x - y| > \frac{2}{\pi} \wedge R \quad \text{or with} \quad |\hat{x} - \hat{y}| > \frac{\pi}{2}.$$

Therefore, we assume that $|x - y| \leq \frac{2}{\pi} \wedge R$ and $|\hat{x} - \hat{y}| \leq \frac{\pi}{2}$.

Case I: If $\eta_{xy} \in E_{x_0, \zeta}^{(\text{ext})}$, we get

$$\tilde{\nu}_\zeta^{(\text{ext})}(x \leftrightarrow y) \geq \tilde{p}_2(\eta_{xy}) = \frac{1}{\frac{q}{\beta} \ell(\eta_{xy})^4 + 1} \geq \frac{1}{\frac{q}{\beta} (\frac{\pi}{2} \ell(\eta_{xy}))^4 + 1} = p^*(\eta_{xy}).$$

Case II: If $\eta_{xy} \notin E_{x_0, \zeta}^{(\text{ext})}$, the proof is no longer straightforward and will take some care. Since $\eta_{xy} \notin E_{x_0, \zeta}^{(\text{ext})}$, and $x, y \in V_B$ there exists $z \in \zeta \cap V_B^c$, such that $\eta_{zx_0} \in \text{Del}_2(\zeta \cup \{x_0\})$. This implies that $z \in V_{x_0, \zeta} \setminus V_B$ and $\hat{x} < \hat{z} < \hat{y}$. We now check whether $\eta_{xz}, \eta_{zy} \in E_{x_0, \zeta}^{(\text{ext})}$. If they are not, we find more points $z \in V_{x_0, \zeta} \setminus V_B$ with $\hat{x} < \hat{z} < \hat{y}$. Therefore, there exists a (finite) sequence $z_1, \dots, z_n \in V_{x_0, \zeta} \setminus V_B$ with $\hat{x} < \hat{z}_1 < \dots < \hat{z}_n < \hat{y}$ such that

$$\eta_{xz_1}, \eta_{z_1 z_2}, \dots, \eta_{z_n y} \in E_{x_0, \zeta}^{(\text{ext})}.$$

The event that each of these edges is open implies the event that x and y belong to the same connected component of open edges, hence

$$\tilde{v}_\zeta^{(\text{ext})}(x \leftrightarrow y) \geq \tilde{p}_2(\eta_{xz_1})\tilde{p}_2(\eta_{z_1z_2}) \cdots \tilde{p}_2(\eta_{z_ny}). \tag{3.13}$$

For any two points $x_1, x_2 \in \zeta$ with $\hat{x}_1 < \hat{x}_2$, define $C_{x_1, x_2}^{x_0}$ to be the arc on the circumcircle $\partial B(\tau(x_1, x_2, x_0))$ of the triangle $\tau(x_1, x_2, x_0)$ between x_1 and x_2 , and define U_{x_1, x_2} to be the subset of \mathbb{R}^2 bounded by this arc $C_{x_1, x_2}^{x_0}$ and $\overline{x_1x_2}$, that is, the convex hull of $C_{x_1, x_2}^{x_0}$. Let $n = \#\{z \in V_{x_0, \zeta} : \hat{x} < \hat{z} < \hat{y}\}$. We claim that

$$L(C_{x, z_1}^{x_0}) + \cdots + L(C_{z_n, y}^{x_0}) \leq L(C_{x, y}^{x_0}), \quad n \in \mathbb{N}, \tag{3.14}$$

where $L(C)$ denotes the length of the arc C . We will prove the claim (3.14) below after we finish the proof of the statement in the lemma. By our assumption that $|x - y| \leq \frac{2}{\pi} \wedge R$ and $|\hat{x} - \hat{y}| \leq \frac{\pi}{2}$, it follows that $L(C_{x, y}^{x_0}) \leq 1$. To see this note that with r being the radius of the circumcircle and with $\theta = \angle x_0xy = |\hat{x} - \hat{y}|$,

$$L(C_{x, y}^{x_0}) = 2r\theta = 2\theta \frac{|x - y||x_0 - x||x_0 - y|}{4 \text{area}(\tau(x, x_0, y))} = \frac{|x - y|\theta}{\sin(\theta)} \leq 1,$$

where we used that $\sin(\theta)/\theta > 2/\pi$ for $\theta \in (0, \pi/2)$. With our claim (3.14) we obtain

$$L(C_{x, z_1}^{x_0}) + \cdots + L(C_{z_n, y}^{x_0}) \leq 1.$$

Obviously, this shows that

$$|x - z_1| + |z_1 - z_2| + \cdots + |z_{n-1} - z_n| + |z_n - y| \leq 1. \tag{3.15}$$

Now choose $\beta > 0$ such that $q/\beta < 1$. Then for $a, b \in \mathbb{R}$ with $0 \leq a \leq b \leq 1$, we have the following simple fact

$$\left(\frac{1}{\frac{q}{\beta}a^4 + 1}\right)\left(\frac{1}{\frac{q}{\beta}b^4 + 1}\right) = \frac{1}{\frac{q}{\beta}(\frac{q}{\beta}a^4b^4 + a^4 + b^4) + 1} \geq \frac{1}{\frac{q}{\beta}(a + b)^4 + 1}, \tag{3.16}$$

where the inequality follows because of $q/\beta < 1$ and the given constraints on a and b . Hence, using (3.15), we obtain

$$\begin{aligned} \tilde{p}_2(\eta_{xz_1})\tilde{p}_2(\eta_{z_1z_2}) \cdots \tilde{p}_2(\eta_{z_ny}) &\geq \left(\frac{1}{\frac{q}{\beta}|x - z_1|^4 + 1}\right) \cdots \left(\frac{1}{\frac{q}{\beta}|z_n - y|^4 + 1}\right) \\ &\geq \frac{1}{\frac{q}{\beta}(|x - z_1| + |z_1 - z_2| + \cdots + |z_{n-1} - z_n| + |z_n - y|)^4 + 1} \\ &\geq \frac{1}{\frac{q}{\beta}L(C_{x, y}^{x_0})^4 + 1} \geq \frac{1}{\frac{q}{\beta}(\frac{\pi}{2}|x - y|)^4 + 1} = p^*(\eta_{xy}), \end{aligned} \tag{3.17}$$

where the second inequality results from repeated use of relation (3.16) with $a = |z_i - z_{i+1}|$ and $b = |z_j - z_{j+1}|$. We conclude with the statement in the lemma.

We are left to verify the claim (3.14): Suppose there exists $z \in V_{x_0, \zeta} \setminus V_B$ such that $\eta_{xz}, \eta_{zy} \in E_{x_0, \zeta}^{(\text{ext})}$. Since $z \notin B$, it must be true that $z \in U_{xy}$. Therefore, by a direct application of Theorem D.2, we have

$$L(C_{x, z}^{x_0}) + L(C_{z, y}^{x_0}) \leq L(C_{x, y}^{x_0}),$$

and the claim follows for $n = 1$. We shall proceed by induction with respect to $n \in \mathbb{N}$. Assume the claim holds for $n = k - 1$. There exist $z_1, \dots, z_k \in V_{x_0, \zeta} \setminus V_B$ such that $\hat{x}_1 < \cdots < \hat{x}_k$ and $\eta_{x, z_1}, \dots, \eta_{z_k y} \in E_{x_0, \zeta}^{(\text{ext})}$. Let

$$i = \operatorname{argmax}_{1 \leq j \leq k} |z_j - \overline{xy}|.$$

It follows that $z_i \in U_{z_{i-1}, z_{i+1}}$, where, for convenience, we write $z_0 = x$ and $z_{k+1} = y$. By Theorem D.2 again,

$$L(C_{z_{i-1}, z_i}^{x_0}) + L(C_{z_i, z_{i+1}}^{x_0}) \leq L(C_{z_{i-1}, z_{i+1}}^{x_0}). \tag{3.18}$$

By changing the notation $z'_j = z_j$ for $1 \leq j < i$ and $z'_j = z_{j+1}$ for $i \leq j \leq k - 1$, it follows from the previous inequality that

$$L(C_{x, z_1}^{x_0}) + \dots + L(C_{z_k, y}^{x_0}) \leq L(C_{x, z'_1}^{x_0}) + \dots + L(C_{z'_{k-1}, y}^{x_0}),$$

and hence, by our assumption for $n = k - 1$, we conclude with the statement of the claim (3.14). □




Figure 10: The sectors S_i of a spoked chain in Q_1 .

Lemma 3.8. *Let $\delta > 0$ and $\Gamma = (V, E)$ be a spoked chain with $V \subset Q_1$. If Γ does not contain a kink, then the number of edges in E with length greater than 2δ is at most $6(\frac{R}{\delta})^2$.*

Proof of Lemma 3.8. Let $V = \{x_1, \dots, x_n\}$ with $\hat{x}_1 < \dots < \hat{x}_n$ be given. For $1 \leq i < n$, let $D_i \subset \mathbb{R}^2$ be the disc of radius $|x_i - x_{i+1}|/2$ centred at x_i . Let $S_i \subset \mathbb{R}^2$ be the sector of D_i

with interior angle $\pi/2$ and line of symmetry $\overline{x_i x_{i+1}}$. We claim the following,

$$S_i \cap S_j = \emptyset \text{ for } i \neq j, 1 \leq i, j \leq n, \tag{3.19}$$

and

$$\bigcup_{i=1}^{n-1} S_i \subset \mathcal{U}_{\frac{R}{\sqrt{2}}}(Q_1), \tag{3.20}$$

where $\mathcal{U}_{\frac{R}{\sqrt{2}}}(Q_1) = \{x \in \mathbb{R}^2 : \text{dist}(x, Q_1) \leq \frac{R}{\sqrt{2}}\}$ is the $R/\sqrt{2}$ neighbourhood of the sector Q_1 . Assume our claim (3.20) is true, the sum of the areas of the sectors S_i must not exceed the area of $\mathcal{U}_{\frac{R}{\sqrt{2}}}(Q_1)$ which is less than $\frac{3}{2}\pi R^2$, see Figure 10. Now each edge $\eta \in E$ of length greater than 2δ contributes a sector of area greater than $\pi/4\delta^2$, therefore, the maximum number of such edges in Γ is simply

$$\frac{(3/2)\pi R^2}{\pi/4\delta^2} = 6\left(\frac{R}{\delta}\right)^2,$$

which gives the result. We are left to prove our claim (3.20) above. Pick $x_i \in V$ and let ℓ_1 be the image of the line $\overleftarrow{x_i x_{i+1}}$ under a rotation with an angle $\pi/2$, centred at point x_{i+1} . There are exactly two connected components of $\mathbb{R}^2 \setminus \ell_1$. Let U denote the component that contains x_i . Now suppose $x_k \in U$ for some $i + 1 < k \leq n$. This implies that $\widehat{x_i x_{i+1} x_k} < \pi/2$. Then, by Definition 3.1, this contradicts the fact that Γ does not contain a kink. Therefore, $x_k \in U^c$ for all $i + 1 < k \leq n$. Let ℓ_2 and ℓ_3 be the images of the half line $\overleftarrow{x_i x_{i+1}}$ under rotations, centred at x_{i+1} , of angles $\pi/4$ and $-\pi/4$ respectively, see Figure 11.




Figure 11: The point x'_k is the first time after x_{i+1} that the chain enters U .

Again, there are two connected components of $\mathbb{R}^2 \setminus (\ell_2 \cup \ell_3)$. Let \tilde{U} denote the one that contains x_i . Now (3.19) follows by noticing that $S_i \subset \tilde{U}$ and $S_k \subset \tilde{U}^c$ for all $i + 1 < k \leq n$. Claim (3.20) follows easily $S_i \subset D_i$ for all $1 \leq i < n$ and that the maximal radius for D_i is half the maximal edge length, which is $\sqrt{2}R$. \square

3.4 Final proof of Theorem 2.1

Recall that we split $B = B_R(x_0) \subset \mathbb{R}^2$ into four quadrants, $Q_i \subset \mathbb{R}^2, i = 1, 2, 3, 4$. Now $G_B \cap Q_1$ contains all vertices and edges of G_B that lie wholly in Q_1 . By construction, $G_B \cap Q_1$ is a spoked chain. It follows from Lemma 3.4 and Lemma 3.5 that there are at most 2 intruding kinks in the spoked chain $G_B \cap Q_1$ and no single protruding kink. For each intruding kink x_i, x_j, x_k , we remove the edge $\eta_{x_j x_{j+1}}$ from $G_B \cap Q_1$. Since removing an edge anywhere except from the end of the spoked chain will result in leaving two spoked chains, we are left with at most 3 spoked chains in Q_1 . Importantly, none of these contain an intruding or protruding kink. Let denote $\Gamma = (V^\Gamma, E^\Gamma)$ one of these kink-less spoked chains in Q_1 . We denote (compare with Theorem 2.1) $N_\Gamma^{(cc)}(\zeta, E)$ to be the number of connected components (clusters) of (ζ, E) that intersect V^Γ . We endeavour to bound the expectation of $N_{x_0}^{(cc)}(\zeta, \cdot)$ with respect to the edge drawing mechanism $\mu_{(ext), \zeta}^{(q)}$ on $E_{x_0, \zeta}^{(ext)}$ given in (2.4). To conclude the Theorem, we shall use

$$\int N_{x_0}^{(cc)}(\zeta, E) \mu_{(ext), \zeta}^{(q)}(dE) \leq 12 \int N_\Gamma^{(cc)}(\zeta, E) \mu_{(ext), \zeta}^{(q)}(dE),$$

where the factor 12 is considering at most three kinkless spoked chains in each of the four quadrants. Order the elements in $V_\Gamma = \{x_1, \dots, x_n\}$ such that $\hat{x}_1 < \dots < \hat{x}_n$. Recall that $\{x \leftrightarrow y\}$ denotes the event that x and y belong to the same cluster of (ζ, E) and notice that

$$\begin{aligned} \int N_\Gamma^{(cc)}(\zeta, E) \mu_{(ext), \zeta}^{(q)}(dE) &\leq 1 + \sum_{j=1}^{n-1} (1 - \mu_{(ext), \zeta}^{(q)}(\{x_j \leftrightarrow x_{j+1}\})) \\ &\leq 1 + \sum_{j=1}^{n-1} (1 - \tilde{\nu}_\zeta^{(ext)}(\{x_j \leftrightarrow x_{j+1}\})) \\ &\leq 1 + \sum_{j=1}^{n-1} (1 - p^*(\eta_{x_j x_{j+1}})) \\ &\leq 1 + \sum_{\eta \in E^\Gamma} (1 - p^*(\eta)). \end{aligned} \tag{3.21}$$

We partition the edge set E^Γ of the spoked chain Γ into subsets of edges according to their lengths. Let

$$\begin{aligned} E_1 &= \{\eta_{xy} \in E^\Gamma : |x - y| > \frac{2}{\pi} \wedge R\}, \\ E_i &= \left\{ \eta_{xy} \in E^\Gamma : \frac{2}{i} \wedge R < |x - y| \leq \frac{2}{i-1} \wedge R \right\}, \quad i \geq 2, i \in \mathbb{N}. \end{aligned}$$

By recalling that

$$p^*(\eta_{xy}) = \frac{\mathbb{1}\{\ell(\eta) \leq \frac{2}{\pi} \wedge R\} \mathbb{1}\{|\hat{x} - \hat{y}| \leq \frac{\pi}{2}\}}{\frac{q}{\beta} \left(\frac{\pi}{2} \ell(\eta)\right)^4 + 1} \mathbb{1}_{E_B}(\eta)$$

from (3.11), we see that $1 - p^*(\eta) = 1$ for all $\eta \in E_1$. However, since Γ is contained in Q_1 , and henceforth $|\hat{x} - \hat{y}| < \frac{\pi}{2}$, we have

$$1 - p^*(\eta_{xy}) = \frac{1}{\frac{q}{\beta} \left(\frac{\pi}{2}|x - y|\right)^4 + 1}$$

for all $\eta_{xy} \in E_i, i \geq 2$. Let $r := 1 \wedge \frac{R\pi}{2}$. Then, considering $\frac{2r}{\pi i} < |x - y| \leq \frac{2r}{\pi(i-1)}$ for all $\eta_{xy} \in E_i$ and noticing that $\bigcup_{i=1}^{\infty} E_i = E^\Gamma$ and $E_i \cap E_j = \emptyset$ for $i \neq j$, it follows readily that

$$\begin{aligned} \sum_{\eta \in E^\Gamma} (1 - p^*(\eta)) &= \sum_{i=1}^{\infty} \sum_{\eta \in E_i} (1 - p^*(\eta)) \leq \sum_{\eta \in E_1} 1 + \sum_{i=2}^{\infty} \sum_{\eta \in E_i} \left(\frac{1}{\frac{\beta}{q} \left(\frac{i-1}{r}\right)^4 + 1} \right) \\ &\leq 6R^2\pi^2r^{-2} + \sum_{i=2}^{\infty} (6R^2\pi^2i^2r^{-2}) \left(\frac{1}{\frac{\beta}{q} \left(\frac{i-1}{r}\right)^4 + 1} \right) \\ &\leq 6R^2\pi^2r^{-2} \left(1 + \frac{qr^2}{\beta} \sum_{i=2}^{\infty} \frac{i^2}{(i-1)^4} \right), \end{aligned}$$

where the second inequality comes from an application of Lemma 3.8. We use that

$$\sum_{i=2}^{\infty} \frac{i^2}{(i-1)^4} \leq \sum_{i=2}^{\infty} \frac{4}{(i-1)^2} = 4 \sum_{i=1}^{\infty} \frac{1}{i^2} = \frac{2}{3}\pi^2. \tag{3.22}$$

Combining all our previous steps we obtain from (3.21) that

$$\int N_\Gamma^{(cc)}(\zeta, E) \mu_{(ext),\zeta}^{(q)}(dE) \leq 1 + 6R^2\pi^2r^{-2} \left(1 + \frac{2q\pi^2r^2}{3\beta} \right)$$

We finish the proof of Theorem 2.1 by setting $\alpha = \alpha(R, q, \beta) = 1 + 6R^2\pi^2r^{-2} \left(1 + \frac{2q\pi^2r^2}{3\beta} \right)$. Note that for given β and q the function $\alpha(R, q, \beta)$ grows quadratic in the finite range radius R (in both cases $R\pi/2 > 1$ and $R\pi/2 < 1$). Furthermore,

$$\lim_{\beta \rightarrow \infty} \alpha(R, q, \beta) = 1 + 6R^2\pi^2r^{-2}. \quad \square$$

4 Proofs

This section delivers the remaining open proofs of our results. We first establish the existence of Gibbs measures. In Section 4.2 we finally finish the proof of Theorem 1.4.

4.1 Existence of Gibbs measures

To show the existence of Gibbs measures (Proposition 1.2) for our Delaunay Potts model we follow [DDG12]. The potential ϕ_β depends solely on the individual Delaunay hyperedges in $\text{Del}_2(\zeta)$, of a marked configuration ζ . Every marked hyperedge $\eta \in \text{Del}_2(\zeta)$ has the so-called finite horizon $\overline{B}(\eta, \zeta)$, where $B(\eta, \zeta)$ is the open ball with $\partial B(\eta, \zeta) \cap \zeta = \eta$ that contains no points of ζ . Thus ϕ_β satisfies the range condition (R) in [DDG12], see [DDG12, Proposition 4.1 & 4.3], with finite horizon being the ball $\overline{B}(\eta, \zeta)$. The finite-horizon property of a general hyperedge potential $\varphi: \Omega \times \Omega \rightarrow \mathbb{R}$ says that for each pair (η, ζ) with $\eta \in \text{Del}_2(\zeta)$ there exists some $\Delta \Subset \mathbb{R}^2$ such that for the pair $(\eta, \tilde{\zeta})$ with $\eta \in \text{Del}_2(\tilde{\zeta})$ we have that $\varphi(\eta, \zeta) = \varphi(\eta, \tilde{\zeta})$ when $\tilde{\zeta} = \zeta$ on $\Delta \equiv \overline{B}(\eta, \zeta)$.

The second requirement for existence of Gibbs measures is the stability condition (S). A hyperedge potential is called stable if there is a lower bound for the Hamiltonian for any $\Lambda \Subset \mathbb{R}^2$, and, as $\phi_\beta(\ell) \geq 0$ for all $\ell \geq 0$, the stability condition (S) is satisfied. The third condition to be checked is a partial complementary upper bound for the

Hamiltonian in any $\Lambda \in \mathbb{R}^2$. This is a bit more involved, and we shall first define appropriate configurations, the so-called *pseudo-periodic marked configurations*. We consider the partition of \mathbb{R}^2 as given in Appendix A. Note that in Appendix A we have introduced a length scale $\ell > 0$ which is not necessary for the existence proof as we can put $\ell = 1$. We let $B_r(0)$ be an open ball around the origin of radius $r \leq \rho_0 \ell$, where we choose $\rho_0 \in (0, 1/2)$ sufficiently small such that $B_r(0) \subset \Delta_{0,0}$. Note that

$$B := \{\zeta \in \Omega_{\Delta_{0,0}} : \zeta = \{x\} \text{ for some } x \in B(0, r)\}$$

is a measurable set of $\Omega_{\Delta_{0,0}} \setminus \{\emptyset\}$. Then

$$G := \{\omega \in \Omega : \theta_{Mz}(\omega_{\Delta_{k,l}}) \in B \text{ for all } z = (k, l) \in \mathbb{Z}^2\}$$

is a set of pseudo-periodic configurations, see (A.2) for details. These configurations are not marked yet. The reason is that when a point is shifted its mark remains unchanged. Thus we define the set of pseudo-periodic marked configurations as

$$\{\omega = (\omega^{(1)}, \dots, \omega^{(q)}) : \omega^{(i)} \in G \text{ for all } i \in M_q\}.$$

The required control of the Hamiltonian from above will be achieved by the following properties. As our hyperedge potential depends only on the single hyperedge the so-called uniform confinement (see [DDG12]) is trivially satisfied. In addition, we need the uniform summability, that is,

$$c_r := \sup_{\zeta \in \Gamma^r} \sum_{\eta \in \text{Del}_2(\zeta) : \eta \cap \Delta \neq \emptyset} \frac{\phi_\beta(\ell(\eta))(1 - \delta_\sigma(\eta))}{\#\hat{\eta}} < \infty,$$

where $\hat{\eta} = \{(k, l) \in \mathbb{Z}^2 : \eta \cap \Delta_{k,l} \neq \emptyset\}$ and where $\Delta = \Delta_{0,0}$. The length $\ell(\eta)$ of any $\eta \in \text{Del}_2(\zeta) \cap \Delta$ when ζ is any pseudo-periodic configuration satisfies

$$\ell(1 - 2\rho_0) \leq \ell(\eta) \leq \ell(1 + 2\rho_0).$$

There are at most six edges from the centre ball in $\Delta = \Delta_{0,0}$ and each Delaunay edge touches exactly two cells and thus $\hat{\eta} = 2$. We obtain an upper bound for each edge by considering the shortest possible length for each edge, that is,

$$c_r = 3 \log \left(\frac{(\ell(1 - 2\rho_0))^4 + \beta}{(\ell(1 - 2\rho_0))^4} \right) < \infty.$$

We need furthermore the so-called weak non-rigidity, that is

$$\Pi_{\Delta_{0,0}}(\Gamma^r) = qe^{-|\Delta_{0,0}|} |\Delta_{0,0}| z > 0.$$

Using [DDG12, Theorem 3.3] and [DDG12, Corollary 3.4] we obtain all the statements in Proposition 1.2. \square

4.2 Breaking of the symmetry of the mark distribution

In this section we complete the proof of Theorem 1.4 by analysing the Gibbs distributions $\gamma_{\Lambda, \omega}$ in the limit $\Lambda \uparrow \mathbb{R}^2$. We pick a boundary condition $\omega \in \Omega$, and we let $\omega = (\omega \setminus \Lambda_n, \emptyset, \dots, \emptyset) \in \omega \in \Omega$ be the monochromatic boundary condition. We write γ_n for $\gamma_{\Lambda_n, \omega}$ and let P_n be the probability measure on Ω relative to which the marked configurations in distinct rhombuses $\Lambda_n + (2n + 1)M(k, l)$, $(k, l) \in \mathbb{Z}^2$, are independent with identical distribution γ_n . As we are dealing with a cell structure for the partition of \mathbb{R}^2 , we confine ourself first to lattice shifts when we employ spatial averaging. Thus,

$$\bar{P}_n = \frac{1}{2n + 1} \sum_{(k,l) \in \{-n, \dots, n\}^2} P_n \circ \theta_{M(k,l)}^{-1}.$$

By the periodicity of P_n the measure \bar{P}_n is \mathbb{Z}^2 -shift-invariant. The proof in [DDG12, Chapter 5] shows that $(\bar{P}_n)_{n \geq 1}$ has a subsequence which converges with respect to the topology of local convergence to some $\hat{P} \in \mathcal{M}_1(\Omega)$. As outlined in [DDG12] it is difficult to show that \hat{P} is concentrated on pseudo periodic configurations. As \hat{P} is non-degenerate the proof in [DDG12, Chapter 5] shows that $P = \hat{P}(\cdot | \{\emptyset\}^c)$ is a Gibbs measure with $P(\{\emptyset\}) = 0$. In order to obtain an \mathbb{R}^2 -shift-invariant Gibbs measure one needs to apply another averaging,

$$P^{(1)} = \int_{\Delta_{0,0}} P \circ \theta_{Mx}^{-1} dx.$$

Applying Propositions 2.8 and 2.2, we see that for $\Delta = \Delta_{0,0}$,

$$\begin{aligned} \int (qN_{\Delta,1} - N_{\Delta}) d\bar{P}_n &\geq \frac{(q-1)}{2n+1} \sum_{(k,l) \in \{-n, \dots, n\}^2} \int N_{\Delta_{k,l} \leftrightarrow \infty} dC_{\Lambda_n, \omega} \\ &\geq (q-1)\varepsilon. \end{aligned}$$

Thus

$$\int (qN_{\Delta,1} - N_{\Delta}) dP^{(1)} > 0,$$

and we observe the following break of symmetry in the expected density of particles of type 1 and of any other type, that is,

$$\rho_1(P^{(1)}) > \rho_2(P^{(1)}) = \dots = \rho_q(P^{(1)}),$$

where $\rho_s(P^{(1)}) = 1/|\Delta| \mathbb{E}_{P^{(1)}}[N_{\Delta,s}]$, $s \in M_q$. We conclude with our statement as in [GH96] by showing that the matrix

$$(\rho_s(P^{(t)}))_{s,t \in M_q}$$

is regular, where $P^{(t)}$ is obtained from $P^{(1)}$ by swapping the role of 1 and t .

Appendix

A Pseudo-periodic configurations

We define pseudo-periodic configurations as in [DDG12]. We first obtain a partition of \mathbb{R}^2 into rhombuses. Pick a length scale $\ell > 0$ and consider the matrix

$$M = (M_1 \quad M_2) = \begin{pmatrix} \ell & \ell/2 \\ 0 & \sqrt{3}/2\ell \end{pmatrix}.$$

Note that $|M_i| = \ell$, $i = 1, 2$, and $\angle(M_1, M_2) = \pi/3$. For each $(k, l) \in \mathbb{Z}^2$ we define the cell

$$\Delta_{k,l} = \{Mx \in \mathbb{R}^2 : x - (k, l) \in [-1/2, 1/2)^2\} \tag{A.1}$$

with area $|\Delta_{k,l}| = \frac{\sqrt{3}}{2}\ell^2$. For example, $\Delta_{0,0}$ is the rhombus with corners

$$(3\ell/4, \sqrt{3}\ell/4), (\ell/4, -\sqrt{3}\ell/4), (-3\ell/4, -\sqrt{3}\ell/4), (-\ell/4, \sqrt{3}\ell/4),$$

and horizontal side length of ℓ . These cells constitute a periodic partition of \mathbb{R}^2 into rhombuses. Let

$$\Gamma = \{\omega \in \Omega : \theta_{Mz}(\omega_{\Delta_{k,l}}) \in B \text{ for all } z = (k, l) \in \mathbb{Z}^2, B \text{ measurable set of } \Omega_{\Delta_{0,0}} \setminus \{\emptyset\}\} \tag{A.2}$$

be the set of all configurations whose restriction to an arbitrary cell, when shifted back to $\Delta_{0,0}$, belongs to the measurable set B for all measurable sets B of $\Omega_{\Delta_{0,0}} \setminus \{\emptyset\}$. Elements of Γ are called **pseudo-periodic** configurations. We define marked pseudo-periodic configurations in an analogous way.

B Topology of local convergence

We write $\mathcal{M}_1^\theta(\Omega)$ (resp. $\mathcal{M}_1^\theta(\Omega)$) for the set of all shift-invariant probability measures on (Ω, \mathcal{F}) (resp. (Ω, \mathcal{F})). A measurable function $f: \Omega \rightarrow \mathbb{R}$ is called local and tame if

$$f(\omega) = f(\omega_\Lambda) \quad \text{and} \quad |f(\omega)| \leq aN_\Lambda(\omega) + b$$

for all $\omega \in \Omega$ and some $\Lambda \Subset \mathbb{R}^2$ and suitable constants $a, b \geq 0$. Let \mathcal{L} be the set of all local and tame functions. The topology of local convergence, or \mathcal{L} -topology, on $\mathcal{M}_1^\theta(\Omega)$ is then defined as the weak* topology induced by \mathcal{L} , i.e., as the smallest topology for which the mappings $P \mapsto \int f dP$ with $f \in \mathcal{L}$ are continuous.

C Mixed site-bond percolation

Given a graph $G = (V, E)$, let \mathbb{P}_p be the probability measure on configurations of open and closed vertices of G . Each vertex of G is open with probability p and closed with probability $1 - p$. Similarly, let $\tilde{\mathbb{P}}_p$ be the probability measure on configurations of open and closed edges of G . Each edge of G is open with probability p and closed with probability $1 - p$. For $x_0 \in V$ and a subset of vertices $X \subset V$, let

$$\sigma(p, x_0, X, G) = \mathbb{P}_p(\exists \text{ a path } x_0 = v_0, e_1, v_1, \dots, e_n, v_n \text{ with } v_n \in X \text{ and all vertices are open}),$$

$$\beta(p, x_0, X, G) = \tilde{\mathbb{P}}_p(\exists \text{ a path } x_0 = v_0, e_1, v_1, \dots, e_n, v_n \text{ with } v_n \in X \text{ and all edges are open}).$$

It is known since [Kes82] that site percolation implies bond percolation, that is, for any $0 \leq p \leq 1$,

$$\sigma(p, x_0, X, G) \leq \beta(p, x_0, X, G). \quad (\text{C.1})$$

In mixed site-bond percolation, both edges and vertices may be open or closed, possibly with different probabilities. Each edge or bond is open independently of anything else with probability p' and each vertex is open independently of anything else with probability p . The edges and vertices that are not open, along with the edges to or from these vertices, are closed. We shall consider paths of open vertices and open edges. For $x_0 \in V$ and a subset of vertices $X \subset V$, let

$$\gamma(p, p', x_0, X, G) = \mathbb{P}_{pp'}(\exists \text{ a path } x_0 = v_0, e_1, v_1, \dots, e_n, v_n \text{ with } v_n \in X \text{ and all vertices and all edges are open}).$$

Let G' be the reduced graph where each edge and vertex of G is removed independently with probability $1 - p'$ and $1 - p$ respectively. By taking the expectation on both sides of inequality (C.1), on G' , with respect to \mathbb{P}_δ and $\tilde{\mathbb{P}}_\delta$, we arrive at the mixed site-bond percolation result of Hammersley, a generalisation of the work of McDiarmid, see [Ham80]. That is, for $\delta, p, p' \in [0, 1]$ one gets that

$$\gamma(\delta p, p', x_0, X, G) \leq \gamma(p, \delta p', x_0, X, G). \quad (\text{C.2})$$

By setting $\delta = p$ and $p' = 1$ in (C.2), and noticing that $\gamma(p^2, 1, x_0, X, G) = \sigma(p^2, x_0, X, G)$, we arrive at

$$\sigma(p^2, x_0, X, G) \leq \gamma(p, p, x_0, X, G), \quad (\text{C.3})$$

and hence

$$\theta_{(\text{mixed})}(p, p) \geq \theta_{(\text{site})}(p^2), \quad (\text{C.4})$$

where $\theta_{(\text{mixed})}(p, p')$ is the mixed site-bond percolation probability with parameters p and p' , and $\theta_{(\text{site})}(p)$ is the vertex percolation probability with parameter p , defined as $\theta_{(\text{site})}(p) := \mathbb{P}_p(|C| = \infty)$, where C is an open cluster, which is a connected component of open vertices.

D Geometrical lemmas

Lemma D.1. Let $\Gamma = (V, E)$ be a spoked chain (see definition in (3.1)) with $V = \{x_1, \dots, x_n\}$ and $\hat{x}_1 < \dots < \hat{x}_n$. For $1 < k \leq n$, let x_k^* and x_{k+1}^* be the centres of the circumscribing circles of the triangles $\tau(x_0, x_{k-1}, x_k)$ and $\tau(x_0, x_k, x_{k+1})$ respectively. Then $\hat{x}_{k+1}^* \geq \hat{x}_k^*$.

Proof. The points x_{k+1}^* and x_k^* both lie on the bisector of the line segment $\overline{x_0 x_k}$. Suppose $\hat{x}_k^* > \hat{x}_{k+1}^*$, then the radius of circumcircle $\partial B(\tau(x_0, x_k, x_{k+1}))$ is greater than the radius of $B(\tau(x_0, x_{k-1}, x_k))$ and hence $\hat{x}_{k+1}^* \geq \hat{x}_k^*$. Now suppose $\hat{x}_k^* \leq \hat{x}_{k+1}^*$. If $\hat{x}_{k+1}^* < \hat{x}_k^*$, then x_{k+1} lies in the interior of the circumcircle $\partial B(\tau(x_0, x_{k-1}, x_k))$ which contradicts properties of the Delaunay tessellation. Therefore, $\hat{x}_{k+1}^* \geq \hat{x}_k^*$. \square

Let $a \in \mathbb{R}^2$ be the pole in a polar coordinate system where \hat{x} denotes the angular coordinate of $x \in \mathbb{R}^2$. For $x, y \in \mathbb{R}^2$ with $\hat{x} < \hat{y}$, let $\partial B(\tau(a, x, y))$ be the unique circum-circle to a, x, y . Let C_{xy}^a be the arc opposite the vertex a which is the intersection of the circle-line through the points x and y with centre at a and the sector given by the three points x, a, y . For any arc C , let $L(C)$ denote its length.

Theorem D.2. Suppose $a \in \mathbb{R}^2$ is the pole. Let $b, c \in \mathbb{R}^2$ with $0 < \hat{b} < \hat{c} < \pi$. Let U be the convex hull of C_{bc}^a . Then, for all $z \in U$,

$$L(C_{bz}^a) + L(C_{zc}^a) \leq L(C_{bc}^a). \tag{D.1}$$

Proof. Let $r > 0$ denote the radius of the circumcircle $\partial B(\tau(a, b, c))$ and define for $z \in U$,

$$\begin{aligned} M &:= |b - c|; \quad h_1 := |b - z|; \quad h_2 := |z - c|; \quad t := |z - a|; \quad s_1 := |b - a|; \quad s_2 := |c - a|, \\ \theta_1 &:= \hat{z} - \hat{b}; \quad \theta_2 := \hat{c} - \hat{z}; \quad \theta := \theta_1 + \theta_2. \end{aligned}$$

Then, $L(C_{bz}^a) = 2\theta \text{radius}(B(\tau(a, b, z)))$ with $\text{radius}(B(\tau(a, b, z))) = h_1/2 \sin(\theta_1)$. Thus the following holds:

$$L(C_{bz}^a) = h_1 \frac{\theta_1}{\sin(\theta_1)}, \quad L(C_{zc}^a) = h_2 \frac{\theta_2}{\sin(\theta_2)}, \quad L(C_{bc}^a) = M \frac{\theta}{\sin(\theta)}.$$

The strategy of the proof is to first show that $L(C_{bz}^a) + L(C_{zc}^a) = L(C_{bc}^a)$ for $z \in C_{bc}^a$ and $L(C_{bz}^a) + L(C_{zc}^a) \leq L(C_{bc}^a)$ for $z \in \overline{bc}$. We then define $L(C_{bz}^a) + L(C_{zc}^a)$ as a function of θ_1, s_1, t and r , and show that it is convex with respect to t . Noting that $z \in U$ is uniquely determined by t and θ_1 , we conclude with the result for all $z \in U$.

Let $z \in C_{bc}^a$. Then $B(\tau(a, b, c)) = B(\tau(a, b, z)) = B(\tau(a, z, c))$. Therefore, $C_{bz}^a \cup C_{zc}^a = C_{bc}^a$ and thus

$$L(C_{bz}^a) + L(C_{zc}^a) = L(C_{bc}^a). \tag{D.2}$$

Now let $z \in \partial U \cap \overline{bc}$. Then, $h_1 + h_2 = M$ and

$$\begin{aligned} L(C_{bz}^a) + L(C_{zc}^a) &= h_1 \frac{\theta_1}{\sin(\theta_1)} + h_2 \frac{\theta_2}{\sin(\theta_2)} = h_1 \frac{\theta_1}{\sin(\theta_1)} + (M - h_1) \frac{\theta_2}{\sin(\theta_2)} \\ &\leq h_1 \frac{\theta}{\sin(\theta)} + (M - h_1) \frac{\theta}{\sin(\theta)} = M \frac{\theta}{\sin(\theta)} = L(C_{bc}^a), \end{aligned} \tag{D.3}$$

where the inequality holds because $\theta \geq \max\{\theta_1, \theta_2\} > 0$ and $g(x) := \frac{x}{\sin(x)}$ is an increasing function on the interval $[0, \pi]$. To write $L(C_{bz}^a) + L(C_{zc}^a)$ as a function of θ_1, s_1 and t , note that by the cosine rule of triangles,

$$h_1^2 = t^2 - 2s_1 t \cos(\theta_1) + s_1^2 \quad \text{and} \quad h_2^2 = t^2 - 2s_2 t \cos(\theta_2) + s_2^2,$$

and thus

$$L(C_{bz}^a) = (t^2 - 2s_1t \cos(\theta_1) + s_1^2)^{1/2} \frac{\theta_1}{\sin(\theta_1)} =: f_1(\theta_1, s_1, t).$$

Furthermore, s_2 is a function of s_1 and θ_1 since

$$M^2 = s_1^2 + s_2^2 - 2s_1s_2 \cos\left(\sin^{-1}\left(\frac{M}{2r}\right)\right),$$

and θ_2 is a function of θ_1 ,

$$\theta_2 = \theta - \theta_1 = \sin^{-1}\left(\frac{M}{2r}\right) - \theta_1.$$

We obtain from these relations the expression

$$L(C_{zc}^a) = (t^2 - 2s_2(s_1, \theta_1)t \cos(\theta_2) + s_2(s_1, \theta_1)^2)^{1/2} \frac{\theta_2(\theta_1)}{\sin(\theta_2(\theta_1))} =: f_2(\theta_1, s_1, t). \quad (D.4)$$

We will show that $f(\theta_1, s_1, t) = f_1(\theta_1, s_1, t) + f_2(\theta_1, s_1, t)$ is convex with respect to t . We obtain

$$\begin{aligned} \frac{d^2}{dt^2} f_1(\theta_1, s_1, t) &= \\ &= \frac{\theta_1}{\sin(\theta_1)} \left(\frac{(t^2 - 2s_1t \cos(\theta_1) + s_1^2)^3/2}{(t^2 - 2s_1t \cos(\theta_1) + s_1^2)^2} - \frac{(t^2 - 2s_1t \cos(\theta_1) + s_1^2)^{1/2}(t - s_1 \cos(\theta_1))^2}{(t^2 - 2s_1t \cos(\theta_1) + s_1^2)^2} \right). \end{aligned}$$

The function $\frac{x}{\sin(x)}$ is positive for $0 \leq x \leq \pi$. The denominator in the bracket is just h_1^2 and thus positive. The numerator in the bracket reads as

$$\begin{aligned} &t^2 - 2s_1t \cos(\theta_1) + s_1^2)^3/2 - (t^2 - 2s_1t \cos(\theta_1) + s_1^2)^{1/2}(t - s_1 \cos(\theta_1))^2 \\ &= (t^2 - 2s_1t \cos(\theta_1) - s_1^2)^{1/2}(t^2 - 2s_1t \cos(\theta_1) + s_1^2 - (t - s_1 \cos(\theta_1))^2) \\ &= (t^2 - 2s_1t \cos(\theta_1) - s_1^2)^{1/2}s_1^2(1 - \cos^2(\theta_1)) \geq 0, \end{aligned}$$

since $(t^2 - 2s_1t \cos(\theta_1) - s_1^2)^{1/2} = h_1 \geq 0$. Therefore, the function f_1 is convex with respect to t . Similarly, show that f_2 is convex with respect to t to see that the function f is convex with respect to t . Pick $0 \leq \theta_1 \leq \sin^{-1}\left(\frac{M}{2r}\right)$. There exist $0 < t_{\min}(\theta_1) < t_{\max}(\theta_1) < 2r$ such that $t_{\min}(\theta_1) \leq |z| \leq t_{\max}(\theta_1)$ for all $z \in U$ with $\hat{z} - \hat{b} = \theta_1$. We have shown (see (D.2) and (D.3)) that

$$f(\theta_1, s_1, t_{\min}(\theta_1)) \leq L(C_{bc}^a) \quad \text{and} \quad f(\theta_1, s_1, t_{\max}(\theta_1)) = L(C_{bc}^a).$$

Therefore, by the convexity of f , for all $t \in [t_{\min}(\theta_1), t_{\max}(\theta_1)]$,

$$\begin{aligned} f(\theta_1, s_1, t) &\leq \frac{t - t_{\min}(\theta_1)}{t_{\max}(\theta_1) - t_{\min}(\theta_1)} f(\theta_1, s_1, t_{\min}(\theta_1)) + \frac{t_{\max}(\theta_1) - t}{t_{\max}(\theta_1) - t_{\min}(\theta_1)} f(\theta_1, s_1, t_{\max}(\theta_1)) \\ &\leq f(\theta_1, s_1, t_{\max}(\theta_1)) = L(C_{bc}^a). \quad \square \end{aligned}$$

References

- [AE16] S. Adams and M. Evers, Phase Transitions in Delaunay Potts Models, *Journal of Statistical Physics* **162**, Issue 1, 162–185, DOI 10.1007/s10955-015-1393-8 (2016). MR-3439335
- [BBD99] E. Bertin, J.M. Billiot and R. Drouilhet, Existence of Delaunay pairwise Gibbs point process with superstable component, *Journal of Statistical Physics* **95**, 719–744 (1999). MR-1700922
- [BBD02] E. Bertin, J.M. Billiot and R. Drouilhet, Continuum Percolation in The Gabriel Graph, *Advances in Applied Probability*, Vol. **34**, No. 4, 689–701 (2002). MR-1938937

Phase transition on the Delaunay graph

- [BBD04] E. Bertin, J.M. Billiot and R. Drouilhet, Phase Transition in the Nearest-Neighbor Continuum Potts Model, *Journal of Statistical Physics* **114**, Nos. 1/2, 79–100 (2004). MR-2032125
- [BKL84] J. Bricomont, K. Kuroda and J.L. Lebowitz, The structure of Gibbs states and coexistence for non-symmetric continuum Widom-Rowlinson models, *Prob. Theory Relat. Fields* **67**, 121–138 (1984). MR-0758069
- [CCK95] J.T. Chayes, L. Chayes and R. Kotecký, The Analysis of the Widom-Rowlinson Model by Stochastic Geometric Methods, *Commun. Math. Phys.* **172**, 551–569 (1995). MR-1354260
- [Der08] D. Dereudre, Gibbs Delaunay tessellations with geometric hard core conditions, *Journal of Statistical Physics* **131**, 127–151 (2008). MR-2394701
- [DDG12] D. Dereudre, R. Drouilhet and H.O. Georgii, Existence of Gibbsian point processes with geometry-dependent interactions, *Probab. Theory Relat. Fields* **153**, 643–670 (2012). MR-2948688
- [DG09] D. Dereudre and H.O. Georgii, Variational characterisation of Gibbs measures with Delaunay triangle interaction, *Electronic Journal of Probability* **14**, 2438–2462 (2009). MR-2563247
- [DL11] D. Dereudre and F. Lavancier, Practical Simulation and estimation for Gibbs Delaunay-Voronoi tessellations with geometric hard-core interaction, *Computational Statistics and Data Analysis* **55**, 498–519 (2011). MR-2736572
- [Eye14] M. Evers, *On Delaunay Random-Cluster Models*, PhD thesis University of Warwick (2014), available at <http://wrap.warwick.ac.uk/67144/>.
- [Geo94] H.-O. Georgii, Large deviations and the equivalence of ensembles for Gibbsian particle systems with superstable interaction, *Prob. Theory Relat. Fields* **99**, 171–195 (1994). MR-1278881
- [GH96] H.O. Georgii and O. Häggström, Phase transition in continuum Potts models, *Commun. Math. Phys.* **181**, 507–528 (1996). MR-1414841
- [GHM00] H.O. Georgii, O. Häggström and C. Maes, The random geometry of equilibrium phases, In: C. Domb and J.L. Lebowitz (eds.) *Phase Transitions and Critical Phenomena*, Vol. **18**, Academic Press, London, pp. 1–142 (2000). MR-2014387
- [GK97] H.O. Georgii and T. Kühneth, Stochastic comparison of point random fields, *Journal of Applied Probability* **34**, 868–881 (1997). MR-1484021
- [Gri94] G. Grimmett, Potts-models and random-cluster processes with many-body interactions, *Journal in Statistical Physics* **75**, 67–121 (1994). MR-1273054
- [Hag00] O. Häggström, Markov random fields and percolation on general graphs, *Adv. Appl. Prob.* **32**, 39–66 (2000). MR-1765172
- [Ham80] J.M. Hammersley, A generalization of McDiarmid’s theorem for mixed Bernoulli percolation, *Mathematical proceedings of the Cambridge mathematical society*, **88**, 167–170 (1980). MR-0569643
- [Kes82] H. Kesten, *Percolation theory for mathematicians*, Birkhäuser, Boston (1982). MR-0692943
- [LMP99] J.L. Lebowitz, A. Mazel and E. Presutti, Liquid-vapor phase transitions for systems with finite range interactions, *Journal in Statistical Physics* **94**, 955–1025 (1999). MR-1694123
- [LL72] J.L. Lebowitz and E.H. Lieb, Phase transition in a continuum classical system with finite interactions, *Phys. Lett.* **39A**, 98–100 (1972).
- [Lis94] D. Lischinski, Incremental Delaunay Triangulation, in *Graphic gems IV*, 47–59, Academic Press (1994).
- [MSS15] A. Mazel, Y. Suhov and I. Stuhl, A classical WR model with q particle types, *Journal in Statistical Physics* **159**, 1040–1086 (2015). MR-3345410
- [MR09] F. Merkl and S.W.W. Rolles, Spontaneous breaking of continuous rotational symmetry in two dimensions, *Electron. J. Probab.* **14**, 1705–1726 (2009). MR-2535010

Phase transition on the Delaunay graph

- [Mø94] J. Møller, *Lectures in Random Voronoi Tessellations*, Springer Lecture Notes in Statistics **87**, Springer (1994). MR-1295245
- [Rue70] D. Ruelle, Superstable interactions in classical statistical mechanics, *Commun. Math. Phys.* **18**, 127–159 (1970). MR-0266565
- [Rue71] D. Ruelle, Existence of a phase transition in a continuous classical system, *Phys. Rev. Lett.* **27**, 1040–1041 (1971).
- [WR70] B. Widom and J. Rowlinson, New model for the study of liquid-vapour transitions, *J. Chem. Phys.* **52**, 1670–1684 (1970).

Acknowledgments. S.A. thanks D. Dereudre for helpful discussions and exchange of ideas visiting Lille.

Electronic Journal of Probability

Electronic Communications in Probability

Advantages of publishing in EJP-ECP

- Very high standards
- Free for authors, free for readers
- Quick publication (no backlog)
- Secure publication (LOCKSS¹)
- Easy interface (EJMS²)

Economical model of EJP-ECP

- Non profit, sponsored by IMS³, BS⁴, ProjectEuclid⁵
- Purely electronic

Help keep the journal free and vigorous

- Donate to the IMS open access fund⁶ (click here to donate!)
- Submit your best articles to EJP-ECP
- Choose EJP-ECP over for-profit journals

¹LOCKSS: Lots of Copies Keep Stuff Safe <http://www.lockss.org/>

²EJMS: Electronic Journal Management System <http://www.vtex.lt/en/ejms.html>

³IMS: Institute of Mathematical Statistics <http://www.imstat.org/>

⁴BS: Bernoulli Society <http://www.bernoulli-society.org/>

⁵Project Euclid: <https://projecteuclid.org/>

⁶IMS Open Access Fund: <http://www.imstat.org/publications/open.htm>