

Int. J. Aquat. Biol. (2013) 1(4): 185-187  
 E-ISSN: 2322-5270; P-ISSN: 2383-0956  
 Journal homepage: www.ij-aquaticbiology.com  
 © 2013 Iranian Society of Ichthyology

## Short Communication

# Length-weight relationship and condition factor in giant harpiosquillid Mantis shrimp, *Harpiosquilla raphidea* (Crustacea: Stomatopoda) in Banten Bay waters, Indonesia

Mugi Mulyono \*<sup>1</sup>, Mufti Petala Patria<sup>1</sup>, Abi Abinawanto<sup>1</sup>, Ridwan Affandi<sup>2</sup>

<sup>1</sup>Department of Biology, Faculty of Mathematics and Science, University of Indonesia, Depok, Indonesia.

<sup>2</sup>Departement of Aquatic Resources Management, Faculty of Fisheries and Marine Sciences, Bogor Agricultural University, Bogor, Indonesia.

**Abstract:** Length-weight relationship of *Harpiosquilla raphidea* from Banten Bay waters was studied from 146 males and 186 females ranging from 38 to 354 and from 37 to 348 mm, respectively. The following equations were obtained  $\ln W = 5.164 + 2.478 \ln X$  (for males),  $\ln W = 5.333 + 2.596 \ln X$  (for females) and  $\ln W = 5.264 + 2.549 \ln X$ ,  $\ln X$  (pooled). The 'b' values (2.5497) found not near to 3 indicating allometric growth being matched up with the cube law. The ANCOVA indicated not much variation between the 'b' values for males and females. In general, the growth rate of the mantis shrimp *H. raphidea* almost the same or not much different from other mantis shrimp species of the order Stomatopoda.

### Article history:

Received 3 June 2013

Accepted 30 July 2013

Available online 20 August 2013

### Keywords:

Mantis shrimps

*Harpiosquilla raphidea*

Stomatopoda

Length-weight relationship

## Introduction

Stomatopods are a group of crustaceans coming under the class Malacostraca and commonly known as 'mantis shrimp'. The spearer mantis shrimp, *Harpiosquilla raphidea* lives on muddy bottoms in coastal waters around Indonesia. In a mudflat develop in the Banten Bay of Province Banten, this species is exploited commercially, mainly by small bottom-trawlers and gill net due to its economic value. Little is known about this group and distribution of the adults.

Weight of the fish is a function of length. Information on length-weight relationship is essential for studies on growth and sexual maturity of animals. Our literature review shows that except few works by Antony et al. (2004) in *Harpiosquilla harpax*; Sukumaran (1987); James and Thirumilu (1993); Tanuja (1996) in *Oratosquilla nepa* and Lyla et al. (1993) in *Harpiosquilla melanoura*, the works concerned with the assessment of weight-length relationship in mantis shrimps from the Indian

waters are scanty. Moreover no work has so far been carried out on the length weight relationship of *Harpiosquilla raphidea*. Thus, the present study aimed to explain the mathematical relationship between two variables namely the length and the weight. The present study provides detailed information on the biology of *H. raphidea* along Banten Bay waters.

## Materials and methods

Around 146 males ranging from 38 to 354 mm and 186 females ranging from 37 to 348 mm were collected from the Banten Bay waters from December 2011 to January 2012. The length was measured from the tip of the rostrum to the end of telson using vernier calipers. The weight was measured to the nearest 0.01 g using an electronic balance. The length-weight relationship was calculated separately for each sex as well as the pooled. The logarithmic equation for this relationship is  $\ln(W) = \ln a + b \ln(L)$  i.e.,  $Y = a +$

\* Corresponding author: Mugi Mulyono  
 E-mail address: mulyonomugi@gmail.com




Figure 1. Logarithmic relationship between the length and weight in male *Harpiosquilla raphidea*.




Figure 2. Logarithmic relationship between the length and weight in female *Harpiosquilla raphidea*.

bX, according to Wardiatno and Mashar (2010), where 'W' is weight (in gram), 'a' and 'b' value are the constant and the regression coefficient, which were estimated using the least-squares regression analysis. The condition factor (K) determines the effects of seasonal and habitat difference in the robustness and general well-being of the species being calculated using the equation  $K = 100W/L^3$  (Gayaniilo and Pauly, 1997), where: L = Length (cm); W = Weight (g); K = Condition factor ( $g/cm^3$ ). The linear equation was fitted separately as well as pooled for males and females of *H. harpax*. Analysis of covariance (ANCOVA) was employed to test the significance of difference between regression coefficients (b) of both sexes (Welcome, 2001).

## Results

The log values corresponding to the length and weight of males and females are plotted in the Figures 1 and 2. There was linear relationship between the weight and length. The following relationships were found:

Males:  $\text{Ln } W = 5.164 + 2.478 \text{ Ln } L$

Females:  $\text{Ln } W = 5.333 + 2.596 \text{ Ln } L$




Figure 1. Logarithmic relationship between the length and weight in male *Harpiosquilla raphidea*.

There were differences in the condition factors for males and females, i.e. the K values for females and males were 1.21 and 1.19, respectively. The K value for the combined sexes was 1.20. The results of covariance analysis on the length-weight equation revealed significant difference between the regressions of males and females at 5% level ( $F = 2.8134$ ;  $P > 0.05$ ). Also, the length-weight relationship of the pooled was found: Pooled:  $\text{Ln } W = 5.265 + 2.549 \text{ Ln } L$ .

## Discussion

The 'b' values of both males and females were less than 3, which is in agreement with Antony et al. (2004) in *Harpiosquilla harpax* of Parangipettai waters of India and Sukumaran (1987) who also described an allometric pattern of growth in *O. nepa* of South Kanara coast.

Rocket et al. (1984) investigated the length-weight relationship in *Squilla empusa* in the northwestern Gulf of Mexico, where they concluded that the species shows an isometric growth pattern as the 'b' values (2.9574 for males and 2.9362 for females) did not deviated much from the cube value 3. Giovanardi and Manfrin (1984), Tanuja (1996) and Lyla et al. (1998) also presented separate equations for males and females in different species of stomatopods.

James and Thirumilu (1993) also recorded a similar kind of observation in *O. nepa* off the Madras coast and found a similar equation for males and females (b value, 2.9661).

The mean condition factor obtained from this study showed significant difference between sexes, thus, indicating sex dependent between female and male.

## Acknowledgement

Authors are thankful to Ai Setiadi, Nurlaela, Sigit and Mbak Riri for their assistance in the laboratorial works. We are indebted to Prof. M. Kasim Moosa due to his help for the identification of the shrimps.

Indonesia with a highlight of its reproductive aspects. *Journal of Tropical and Conservation*, 7: 63-73.  
 Welcome R.L. (2001). *Inland fisheries, ecology and management*. London: Fishing News Book, Blackwell Science. 358 pp.

## References

- Antony P.J., Mani E.P., Khan S.A. (2004). Length-weight relationships in Mantis shrimp *Harpisquilla harpax* (de Haan) (Crustacea: Stomatopoda). *Journal of Aquatic Biology*, 19: 39-42.
- Gayanilo F.C., Pauly D. (1997). *FAO-ICLARM Stock Assessment Tools (FiSAT)*. FAO Computerized Information Series (fisheries) No. 8, Rome, 262 pp.
- Giovanardi O., Piccinetti-Manfrin P. (1984). Summary of biological parameters of *Squilla mantis* L. in the Adriatic Sea. *FAO Fisheries and Aquaculture Report*, 290: 131- 134.
- James U.B., Thirumilu P. (1993). Population dynamics of *Oratosquilla nepa* in the trawling grounds off Madras. *Journal of the Marine Biological Association of India*, 35: 135-110.
- Lyla P.S., Panchatcharam K., Khan S.A. (1998). Age, growth and length-weight relationship in the stomatopod *Harpisquilla melanoura* (Manning). In: *Proc. Symp. Advances and Priorities in Fish. Tech.*, K.K. Balachandran, T.S.G. Iyer, P. Madhavan, J. Joseph P.A. Perigreen M.R. Raghunath and M.D. Varghese (Ed.). *Society of Fisheries Technologists (India)*. Cochin, 44-47.
- Rocket M.D., Standard G.W., Chittenden M.E. (1984). Bathymetric distribution, spawning periodicity, sex ratios, and size compositions of mantis shrimp *Squilla empusa* in the northwestern Gulf of Mexico. *Fishery Bulletin*, 82: 418-426.
- Sukumaran K.K. (1987). Study on the fishery and biology of the mantis shrimp *Oratosquilla nepa* (Latreille) of South Kanara coast during 1979-1983. *Indian Journal of Fisheries*, 34: 292-305.
- Tanuja R. (1996). Some aspects of biology and utilization of the mantis shrimp *Oratosquilla nepa* from Cochin waters. Ph.D. Thesis. Cochin University of Science and Technology, India.
- Wardiatno Y., Mashar A. (2010). Biological information on the mantis shrimp, *Harpisquilla raphidea* (Fabricius 1798) (Stomatopoda, Crustacea) in