

Washington University in St. Louis Washington University Open Scholarship

Mathematics Faculty Publications

Mathematics and Statistics

8-24-2016

Determinantal Representations of Semihyperbolic Polynomials

Greg Knese

Washington University in St. Louis, geknes@wustl.edu

Follow this and additional works at: https://openscholarship.wustl.edu/math_facpubs

 Part of the [Mathematics Commons](#)

Recommended Citation

Knese, Greg, "Determinantal Representations of Semihyperbolic Polynomials" (2016). *Mathematics Faculty Publications*. 40.
https://openscholarship.wustl.edu/math_facpubs/40

This Article is brought to you for free and open access by the Mathematics and Statistics at Washington University Open Scholarship. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

Determinantal Representations of Semihyperbolic Polynomials

GREG KNESE

ABSTRACT. We prove a generalization of the Hermitian version of the Helton–Vinnikov determinantal representation for hyperbolic polynomials to the class of *semihyperbolic* polynomials, a strictly larger class, as shown by an example. We also prove that certain hyperbolic polynomials affine in two out of four variables divide a determinantal polynomial. The proofs are based on work related to polynomials with no zeros on the bidisk and tridisk.

1. Introduction

A homogeneous polynomial $P \in \mathbb{R}[x_0, x_1, \dots, x_n]$ is *hyperbolic* of degree d with respect to $e \in \mathbb{R}^{n+1}$ if $P(e) \neq 0$ and if for all $x \in \mathbb{R}^{n+1}$ the one-variable polynomial $t \mapsto P(x - te)$ has only real zeros. This concept was originally studied by Gårding for its relation to PDE (see [7; 15]), but it—and the related concept of stable polynomials—has since become important to convex optimization, combinatorics, probability, and analysis. See the papers and surveys [31; 14; 13; 34; 29; 15; 26].

A deep result in the area is a determinantal representation for trivariate hyperbolic polynomials due to Helton and Vinnikov [16; 32], who solved a 1958 conjecture of Lax [17] (see [18]) and, as is mentioned in [15], can be used to develop the full Gårding theory of hyperbolicity.

THEOREM A. *Let $p \in \mathbb{R}[x_0, x_1, x_2]$ be hyperbolic of degree d with respect to e_2 and monic in x_2 . Then, there exist $d \times d$ real symmetric matrices A_0 and A_1 such that*

$$p(x_0, x_1, x_2) = \det(x_0 A_0 + x_1 A_1 + x_2 I).$$

If we relax the problem to finding self-adjoint matrices instead of real symmetric matrices, then proofs more amenable to computations are possible (see [11; 30; 33]). The resulting theorem is just as useful for most purposes.

THEOREM A*. *Let $p \in \mathbb{R}[x_0, x_1, x_2]$ be hyperbolic of degree d with respect to e_2 and monic in x_2 . Then, there exist $d \times d$ self-adjoint matrices A_0 and A_1 such that*

$$p(x_0, x_1, x_2) = \det(x_0 A_0 + x_1 A_1 + x_2 I).$$

Received December 17, 2014. Revision received May 30, 2016.
 This research was supported by NSF grant DMS-1363239.

Our immediate goal is to prove a generalization of this result based on a result of Geronimo et al. [8] and an extension to four variables based on a result of Bickel and Knese [2], whereas our larger goal is to advertise the close connection between determinantal representations of hyperbolic polynomials and sums of squares decompositions for multivariable Schur stable polynomials. See [12; 10; 22; 23] for background on the latter topic.

Our main result establishes a determinantal representation with the assumption of hyperbolicity weakened. We shall call a homogeneous polynomial $P \in \mathbb{R}[x_0, x_1, \dots, x_n]$ a *semihyperbolic polynomial* with respect to the direction $e \in \mathbb{R}^{n+1} \setminus \{0\}$ if for every $x \in \mathbb{R}^{n+1}$ the univariate polynomial $t \mapsto P(x - te)$ is either identically zero or only has real roots. The key distinction between hyperbolic and semihyperbolic polynomials is that we do not assume that $P(e) \neq 0$. Some references actually confuse the two, whereas Renegar [31] is the only reference we have found that emphasizes the distinction. We elaborate on our motivations in Section 6. We do need to allow for $t \mapsto P(x - te)$ to be identically zero because, for instance, if $P(e) = 0$ and $x = 0$, then $P(-te) \equiv 0$. We give an example of a semihyperbolic polynomial that is not hyperbolic in any direction in Section 3.

Here is our main theorem.

THEOREM 1. *Let $p \in \mathbb{R}[x_0, x_1, x_2]$ of degree d be semihyperbolic with respect to $e_2 = (0, 0, 1)$. Then, there exist $d \times d$ self-adjoint matrices A_0, A_1 , and A_2 with A_2 positive semidefinite and a constant $c \in \mathbb{R}$ such that*

$$p(x) = c \det(x_0 A_0 + x_1 A_1 + x_2 A_2).$$

Assuming that p has no factors depending on x_0, x_1 alone, the data can be chosen to additionally satisfy:

-
- $\text{rank } A_1 = \text{deg}_1 p, \quad \text{rank } A_2 = \text{deg}_2 p,$
- $A_1 = B_+ - B_-$ with B_{\pm} both positive semidefinite where $\text{rank } B_-$ equals the number of roots of $p(1, t, i)$ in the upper half plane and $\text{rank } B_+ + \text{rank } B_- = \text{rank } A_1,$
- and $B_- + B_+ + A_2 = I.$

See Section 2 for the proof. We can recover Theorem A* when $p(e_2) \neq 0$ since p will then have degree d in x_2 and then A_2 will be positive definite. We can then factor $A_2^{1/2}$ from the right and left of $\sum_{j=0}^2 x_j A_j$ in order to get a determinantal representation of the form given in Theorem A*, namely with $A_2 = I.$

There is nothing special about the vector e_2 ; a linear change of variables could be used to establish a determinantal representation for other semihyperbolic polynomials. The assumption of no factors depending on only x_0, x_1 is there to avoid certain annoyances that such trivial factors introduce. For instance, $p(x_0, x_1, x_2) = x_1$ is certainly semihyperbolic in the direction e_2 , but then $A_2 = A_0 = 0$, and the signature of the 1×1 matrix A_1 does not really provide any useful information.

It follows that a trivariate semihyperbolic polynomial p can be lifted to a four-variable polynomial

$$P(x_0, x_1, y_1, x_2) = c \det(x_0 A_0 + x_1 B_+ + y_1 B_- + x_2 A_2),$$

which is hyperbolic in the direction $(0, 1, 1, 1)$ and such that $P(x_0, x_1, -x_1, x_2) = p(x_0, x_1, x_2)$. So, we are projecting a hyperbolic polynomial (possessing a definite determinantal representation) of four variables to a set where it is not necessarily hyperbolic. It also follows that a trivariate semihyperbolic polynomial is a limit of hyperbolic polynomials. Indeed, writing p as in Theorem 1, define, for $\varepsilon > 0$,

$$p_\varepsilon(z) = c \det(x_0 A_0 + x_1 A_1 + x_2 (A_2 + \varepsilon I)). \tag{1.1}$$

Then, $p_\varepsilon \rightarrow p$ as $\varepsilon \searrow 0$. We do not know if semihyperbolic polynomials in more than three variables are the limits of hyperbolic polynomials.

Theorem 1 has a curious asymmetry in its treatment of x_0 and x_1 . This is partly due to idiosyncrasies of our proof, but we also think there are some subtleties to resolve. To be specific, we could certainly break up A_0 into a difference of positive semidefinite matrices according to its signature, but we have been unable to connect the signature of the A_0 we construct with geometric properties of p . We have no reason to believe this cannot be done, especially because this issue does not arise in the hyperbolic case. Indeed, we can take $A_2 = I$, and the number of zeros of $t \mapsto p(tx_0, tx_1, i)$ in \mathbb{C}_+ equals the number of negative eigenvalues of $x_0 A_0 + x_1 A_1$. Similarly, the number of zeros of $t \mapsto p(tx_0, tx_1, -i)$ equals the number of positive eigenvalues of $x_0 A_0 + x_1 A_1$. Thus, the signature of $x_0 A_0 + x_1 A_1$ can be derived from properties of p in the hyperbolic case. Notice that we evaluate p on the complex line $(0, t, i)$ to determine the signature of A_1 , whereas in the theorem we evaluate on the line $(1, t, i)$, which actually seems less natural. The example in Section 3 shows this is actually necessary: using the line $(1, t, i)$, we get a correct count of the negative eigenvalues of A_1 , whereas using the line $(0, t, i)$, we get an incorrect count. The details are recorded in Section 3.

As a nice corollary, we can quickly recover the following variant of Theorem A*. The original proof, while not difficult, requires transforming a real stable polynomial to a hyperbolic polynomial through a linear transformation. Our signature count of A_1 in Theorem 1 makes the proof go smoothly.

COROLLARY 1 (See Theorem 6.6 of [3]). *If $p \in \mathbb{R}[x_0, x_1, x_2]$ is homogeneous of degree d and hyperbolic with respect to all vectors in the cone $\{(0, v_1, v_2) : v_1, v_2 > 0\}$, then there exist $d \times d$ self-adjoint matrices A_0, A_1 , and A_2 and a constant $c \in \mathbb{R}$ such that A_1 and A_2 are positive semidefinite, $A_1 + A_2 = I$, and*

$$p(x) = c \det(x_0 A_0 + x_1 A_1 + x_2 A_2).$$

Since [3] uses Theorem A to prove this result, all of the matrices can be taken to be real, but our proof does not yield this. For p as in the corollary, $p(1, x_1, x_2)$ is known as a *real stable* polynomial. This formula was used in the recent paper regarding the Kadison–Singer problem [26]. See Section 4 for the very short proof of the corollary.

The key tool for the proof of Theorem 1 is a determinantal representation proven by Geronimo, Iliev, and Knesé [8] for certain polynomials on the bidisk $\mathbb{D}^2 = \mathbb{D} \times \mathbb{D}$ (here \mathbb{D} is the open unit disk in the complex plane \mathbb{C}). Define $D(z) = z_1 D_1 + z_2 D_2$ where the D_1 and D_2 are $(n + m) \times (n + m)$ matrices given by

$$D_1 = \begin{pmatrix} I_n & 0 \\ 0 & 0 \end{pmatrix}, \quad D_2 = \begin{pmatrix} 0 & 0 \\ 0 & I_m \end{pmatrix}.$$

For $n = n_1 + n_2$, define

$$P_+ = \begin{pmatrix} I_{n_1} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad P_- = \begin{pmatrix} 0 & 0 & 0 \\ 0 & I_{n_2} & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

where the blocks correspond to the orthogonal decomposition $\mathbb{C}^{n+m} = \mathbb{C}^{n_1} \oplus \mathbb{C}^{n_2} \oplus \mathbb{C}^m$. Let $\mathbb{E} = \{z \in \mathbb{C} : |z| > 1\}$ and $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$.

THEOREM B. *Suppose $p \in \mathbb{C}[z_1, z_2]$ has bidegree (n, m) , no zeros in $(\mathbb{T} \times \mathbb{D}) \cup (\mathbb{T} \times \mathbb{E})$, and no factors depending on z_1 alone. Let n_2 be the number of zeros of $p(z_1, 0)$ in \mathbb{D} . Then, there exists an $(n + m) \times (n + m)$ unitary U and a constant $c \in \mathbb{C}$ such that*

$$p(z_1, z_2) = c \det((z_1 P_- + P_+ + D_2) - U(P_- + z_1 P_+ + z_2 D_2)).$$

This is referred to as a determinantal representation for “generalized distinguished varieties” in [8] since it generalizes a determinantal representation for the “distinguished varieties” of Agler and McCarthy [1], which correspond to the case $n_2 = 0$. Polynomials defining distinguished varieties are essentially a Cayley transform of real stable polynomials, and distinguished varieties have their own motivation in terms of operator theory as shown in [1]. Theorem B is based on first proving a sums-of-squares decomposition for polynomials $p \in \mathbb{C}[z_1, z_2]$ with no zeros in $\mathbb{T} \times \mathbb{D}$ (“a face of the bidisk”) and no factors in common with $\tilde{p}(z) = z_1^n z_2^m \overline{p(1/\bar{z}_1, 1/\bar{z}_2)}$. Namely,

$$|p(z)|^2 - |\tilde{p}(z)|^2 = (1 - |z_1|^2)(|E_1(z)|^2 - |E_2(z)|^2) + (1 - |z_2|^2)|F(z)|^2,$$

where $E_1 \in \mathbb{C}^{n_1}[z]$, $E_2 \in \mathbb{C}^{n_2}[z]$, $F \in \mathbb{C}^m[z]$, and $n = n_1 + n_2$ where n_2 is the number of zeros of $p(z_1, 0)$ in \mathbb{D} . This formula generalizes a sums-of-squares formula of Cole and Wermer [5] related to Andô’s inequality from operator theory (see also [9] and [20]). It would be interesting to characterize which polynomials possess such a sums-of-squares formula where $|F(z)|^2$ is also given by a difference of squares $|F_1(z)|^2 - |F_2(z)|^2$, and—going further—it would be interesting to see what sort of determinantal representation for real homogeneous polynomials comes out of the corresponding development from Theorem B to Theorem 1 presented here.

Beyond trivariate polynomials, there are many results on the existence or nonexistence of determinantal representations. See [33; 19; 27; 28; 4; 25] for recent results and convenient summaries of the state of the art. Vinnikov [33] conjectures that hyperbolic polynomials always divide a hyperbolic polynomial

that has a determinantal representation but with additional requirements placed on the set where the determinantal polynomial is positive. Our next theorem offers a step in the right direction for this conjecture albeit in a special situation. A polynomial p is *affine* with respect to a variable x_j if it has degree one in that variable.

THEOREM 2. *Let $p \in \mathbb{R}[x_0, x_1, x_2, x_3]$ be hyperbolic of degree d with respect to the cone $\{(0, v_1, v_2, v_3) : v_1, v_2, v_3 > 0\}$. Assume that p is affine in x_2 and x_3 and of degree n in x_1 . Then, there exist $k \leq 2n + 4$ and $k \times k$ self-adjoint matrices A_0, A_1, A_2, A_3 such that p divides*

$$\det\left(\sum_{j=0}^3 x_j A_j\right),$$

A_1, A_2, A_3 are positive semi-definite, and $A_1 + A_2 + A_3 = I$.

See Section 5. Theorem 2 seems to be one of the few higher-dimensional situations where we get a determinantal representation from simple hypotheses. The recent article of Kummer [24] proves the interesting result that a hyperbolic polynomial in n variables with no real singularities divides a determinantal polynomial. This article also obtains bounds on the sizes of the matrices involved under the assumption that some power of the polynomial has a determinantal representation. Theorem 2 requires no assumptions of smoothness and obtains general bounds on the sizes of the matrices involved, but Kummer’s result has the advantage that it works in n variables and does not assume degree restrictions.

The key tool for Theorem 2 is the following sums-of-squares decomposition from Bickel and Knese [2].

THEOREM C (Theorem 1.12 of [2]). *Let $p \in \mathbb{C}[z_1, z_2, z_3]$ have multidegree $(n, 1, 1)$ and no zeros on $\overline{\mathbb{D}}^3$. Then, there exist column-vector-valued polynomials $E_1 \in \mathbb{C}^n[z_1, z_2, z_3], E_2, E_3 \in \mathbb{C}^2[z_1, z_2, z_3]$ such that, for $z = (z_1, z_2, z_3)$ and $w = (w_1, w_2, w_3)$,*

$$p(z)\overline{p(w)} - \tilde{p}(z)\overline{\tilde{p}(w)} = \sum_{j=1}^3 (1 - z_j \bar{w}_j) E_j(w)^* E_j(z)$$

where $\tilde{p}(z) = z_1^n z_2 z_3 \overline{p(1/\bar{z}_1, 1/\bar{z}_2, 1/\bar{z}_3)}$.

2. Proof of Theorem 1 from Theorem B

Let $\mathbb{C}_+ = \{z \in \mathbb{C} : \Im z > 0\}$ and $\mathbb{C}_- = \{z \in \mathbb{C} : \Im z < 0\}$.

Assume that $P \in \mathbb{R}[x_0, x_1, x_2]$ is homogeneous of degree d and, for every $x \in \mathbb{R}^3$,

$$t \mapsto P(x - te_2)$$

is either identically zero or only has real zeros. We will assume that P has no factors depending only on x_0, x_1 , which can easily be incorporated into our final determinantal representation by appending diagonal blocks to our matrices.

Consider

$$q(z_1, z_2) = P(1, z_1, z_2),$$

which has no zeros in $(\mathbb{R} \times \mathbb{C}_+) \cup (\mathbb{R} \times \mathbb{C}_-)$. To see this, take $z = (a_1, a_2 + ib_2) \in (\mathbb{R} \times \mathbb{C}_+) \cup (\mathbb{R} \times \mathbb{C}_-)$ with $q(z) = 0$. Then, $P((1, a_1, a_2) + te_2)$ has the imaginary root $t = ib_2$, which would imply that $t \mapsto P(1, a_1, a_2 + t)$ is identically zero. This means that $x_1 - a_1x_0$ divides P , which we have ruled out.

Now, define

$$p(z_1, z_2) = q\left(i \frac{1+z_1}{1-z_1}, i \frac{1+z_2}{1-z_2}\right) \left(\frac{1-z_1}{2i}\right)^n \left(\frac{1-z_2}{2i}\right)^m$$

where q has degree n in x_1 and degree m in x_2 . Setting $x_0 = 1$ in $P(x_0, x_1, x_2)$ cannot lower the degree in x_1 or x_2 , so $n = \deg_1 P$ and $m = \deg_2 P$. Recall that

$$z \mapsto i \frac{1+z}{1-z}$$

is a conformal map of the unit disk onto the upper half-plane sending \mathbb{T} to $\mathbb{R} \cup \{\infty\}$ where $1 \mapsto \infty$. Thus, p has no zeros in $(\mathbb{T} \setminus \{1\}) \times \mathbb{D}$ as well as $(\mathbb{T} \setminus \{1\}) \times \mathbb{E}$ where $\mathbb{E} = \{z \in \mathbb{C} : |z| > 1\}$. We cannot have $p(1, z_2) = 0$ unless $p(z_1, z_2)$ has $z_1 - 1$ as a factor. This follows by Hurwitz’s theorem since the polynomials $z_2 \mapsto p(z_1, z_2)$ will have no zeros in $\mathbb{C} \setminus \mathbb{T}$ for $z_1 \in \mathbb{T}$ with $z_1 \rightarrow 1$, and then $p(1, z_2)$ will either have the same property or will be identically zero. However, such factors cannot exist since they imply that q has degree less than n in x_1 . In any case, we can safely divide out factors of p that depend only on z_1 since these can easily be incorporated into our final determinantal representation. Having done this, p satisfies the hypotheses of Theorem B, and we may write

$$p(z_1, z_2) = c \det((z_1 P_- + P_+ + D_2) - U(P_- + z_1 P_+ + z_2 D_2))$$

for a unitary U . Notice that n_2 is the number of roots of $z_1 \mapsto p(z_1, 0)$ in \mathbb{D} , which is the same as the number of roots of $z_1 \mapsto q(z_1, i) = P(1, z_1, i)$ in \mathbb{C}_+ .

We convert back to q via $z \mapsto \frac{z-i}{z+i}$. So,

$$\begin{aligned} q(z_1, z_2) &= p\left(\frac{z_1-i}{z_1+i}, \frac{z_2-i}{z_2+i}\right) (z_1+i)^n (z_2+i)^m \\ &= p\left(\frac{z_1-i}{z_1+i}, \frac{z_2-i}{z_2+i}\right) \det((z_1+i)D_1 + (z_2+i)D_2) \\ &= c \det((z_1-i)P_- + (z_1+i)P_+ + (z_2+i)D_2 \\ &\quad - U((z_1+i)P_- + (z_1-i)P_+ + (z_2-i)D_2)) \\ &= c \det((I-U)D(z) - i(I+U)(P_- - P_+ - D_2)) \\ &= \pm c \det((I-U)(-z_1 P_- + z_1 P_+ + z_2 D_2) + i(I+U)). \end{aligned} \tag{2.1}$$

The last line comes from multiplying on the right by $\det(-P_- + P_+ + D_2)$. Letting $M(z) = -z_1 P_- + z_1 P_+ + z_2 D_2$, we now form the spectral decomposition $U = V \begin{pmatrix} u & 0 \\ 0 & I \end{pmatrix} V^*$; V is a unitary, u is a $k \times k$ diagonal unitary with no 1s on the diagonal,

and k is the rank of $U - I$. Factoring V and V^* out from the left and right of (2.1) leaves

$$\begin{aligned} q(z) &= \pm c \det \left(\begin{pmatrix} I - u & 0 \\ 0 & 0 \end{pmatrix} V^* M(z) V + i \begin{pmatrix} I + u & 0 \\ 0 & 2I \end{pmatrix} \right) \\ &= \pm c \det(I - u) \det \left(\begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix} V^* M(z) V + \begin{pmatrix} a & 0 \\ 0 & 2iI \end{pmatrix} \right) \\ &= \pm c \det(I - u) \det \begin{pmatrix} (V^* M(z) V)_{kk} + a & * \\ 0 & 2iI \end{pmatrix} \\ &= C \det((V^* M(z) V)_{kk} + a) \end{aligned}$$

where $a = i(I + u)(I - u)^{-1}$ is a diagonal matrix with real entries, $(V^* M(z) V)_{kk}$ is the upper $k \times k$ block of $V^* M(z) M$, and C is a constant. Now, $V^* M(z) V = -z_1 V^* P_- V + z_1 V^* P_+ V + z_2 V^* D_2 V$, and if we set $A_0 = a$, $A_1 = (-V^* P_- V + V^* P_+ V)_{kk}$, and $A_2 = (V^* D_2 V)_{kk}$, then we have a determinantal representation for q :

$$q(z) = C \det(A_0 + z_1 A_1 + z_2 A_2).$$

Notice that A_0, A_1, A_2 are evidently self-adjoint with A_2 positive semidefinite, and since $\deg q = d$, we have $d \leq k$. Once we show that $k = d$, we can homogenize to get the determinantal representation for P . It helps to first establish some of the additional details listed in Theorem 1.

It is a general fact that for matrices A, B , the degree of $\det(tA + B)$ is at most $\text{rank } A$ (we leave this as an exercise). So, $\deg_j q \leq \text{rank } A_j$ for $j = 1, 2$. On the other hand, by construction $\text{rank } A_1 \leq \text{rank}(-P_- + P_+) = \deg_1 q$ and $\text{rank } A_2 \leq \text{rank } D_2 = \deg_2 q$, yielding $\deg_j q = \text{rank } A_j$ for $j = 1, 2$. Next, setting $B_{\pm} = (V^* P_{\pm} V)_{kk}$, we have $A_1 = B_+ - B_-$. Since $\text{rank } A_1 = n_1 + n_2$, $\text{rank } B_+ \leq n_1$, and $\text{rank } B_- \leq n_2$, we must have equality in both inequalities. This also shows that the ranges of B_+, B_- have trivial intersection by considering dimensions. Since $P_+ + P_- + D_2 = I$, we must have $B_+ + B_- + A_2 = I$.

In order to show $k = d$, it suffices to show that $Q(t) := tA_1 + A_2$ is nonsingular for some t . For then, there would be a t_0 such that

$$t \mapsto q(t(t_0, 1))$$

has degree k , and since q has degree d , we would have $k \leq d$ and thus $k = d$.

Note that $Q(t) = I + (t - 1)B_+ - (t + 1)B_-$. By the spectral theorem

$$B_+ = \sum_{j=1}^{n_1} v_j v_j v_j^* \quad \text{and} \quad B_- = \sum_{j=1}^{n_2} \mu_j w_j w_j^*$$

where $V = \{v_1, \dots, v_{n_1}\}$ and $W = \{w_1, \dots, w_{n_2}\}$ form orthonormal sets of eigenvectors corresponding to the positive eigenvalues $\{v_1, \dots, v_{n_1}\}$ and $\{\mu_1, \dots, \mu_{n_2}\}$ of B_+ and B_- , respectively. Let $Y = \{y_1, \dots, y_{k-n}\}$ be an orthonormal basis for the complement of V and W . Then, $\mathcal{B} = V \cup W \cup Y$ is a basis for \mathbb{C}^k . Let \mathcal{C} be a basis dual to \mathcal{B} . (Two bases $\{b_1, \dots, b_N\}$ and $\{c_1, \dots, c_N\}$ are dual if $b_j^* c_k = \delta_{jk}$.)

The matrix for $Q(t)$ obtained by using \mathcal{C} as a basis for the domain and \mathcal{B} for the range is of the form

$$\begin{pmatrix} I + (t - 1)d_+ & 0 & 0 \\ 0 & I - (t + 1)d_- & 0 \\ 0 & 0 & I \end{pmatrix}$$

for diagonal matrices d_+, d_- containing the eigenvalues $\nu_1, \dots, \nu_{n_1}, \mu_1, \dots, \mu_{n_2}$ on the diagonal. The determinant of this vanishes for only finitely many t , and so $Q(t_0)$ is certainly nonsingular for some t_0 . Thus, $k = d$, and we homogenize q at degree d to see that

$$P(x) = C \det(x_0 A_0 + x_1 A_1 + x_2 A_2).$$

This concludes the proof of Theorem 1.

3. Example

Renegar [31] has an example of a polynomial that is semihyperbolic but not hyperbolic in any direction (see Section 2 of that paper); however, we have constructed an example that is more illustrative for our purposes.

Let

$$p(x_0, x_1, x_2) = 2x_0^2 x_1 - (x_0^2 + 3x_1^2)x_2.$$

Then, $t \mapsto p(x - te_2)$ clearly has only real roots for $x \in \mathbb{R}^3$ since this one-variable polynomial has degree 1 and real coefficients. Let

$$A_0 = \frac{i}{3} \begin{pmatrix} 0 & -3 & -\sqrt{3} \\ 3 & 0 & \sqrt{3} \\ \sqrt{3} & -\sqrt{3} & 0 \end{pmatrix},$$

$$A_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad A_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

We see that

$$p(x) = 3 \det(x_0 A_0 + x_1 A_1 + x_2 A_2).$$

As remarked in the Introduction, we can lift to

$$P(x_0, x_1, y_1, x_2) = 3x_1 y_1 x_2 - (x_2 + x_1 + 3y_1)x_0^2,$$

which is hyperbolic in the direction $(0, 1, 1, 1)$ and such that $P(x_0, x_1, -x_1, x_2) = p(x_0, x_1, x_2)$. We now explain why p is not hyperbolic in any direction.

We first show that $\{x : p(x) \neq 0\}$ consists of the two connected components $\mathcal{P}_+ = \{x : p(x) > 0\}$ and $\mathcal{P}_- = \{x : p(x) < 0\}$. I thank the referee for the following simplified explanation. The hypersurface $\{x : p(x) = 0\}$ is the graph of the continuous function $(x_0, x_1) \mapsto 2x_0^2 x_1 / (x_0^2 + 3x_1^2)$. Thus, $\{x : p(x) \neq 0\}$ is divided into exactly two components, the part above the graph and the part below.

Next, neither component $\mathcal{P}_+, \mathcal{P}_-$ is convex. For instance, $(-1, 0, -1), (1, 0, -1) \in \mathcal{P}_+$, but $(0, 0, -1) \notin \mathcal{P}_+$. We can similarly show that \mathcal{P}_- is not convex. This implies that p is not hyperbolic in any direction since it is a fundamental

result of Gårding that if p is hyperbolic in some direction e , then the connected component of $\{x : p(x) \neq 0\}$ containing e is convex.

This brings up a potential paradox. Since p is a limit of hyperbolic polynomials p_ε as in equation (1.1), how is it possible that the connected components of $\{x : p(x) \neq 0\}$ are nonconvex in our example? An answer is that a convex component of $\{x : p_\varepsilon(x) \neq 0\}$ could shrink to an isolated point (in projective space) as $\varepsilon \searrow 0$. This is something we have seen graphically using our example.

Finally, in connection with our discussion after Theorem 1 regarding the signatures of A_0 and A_1 , let us point out that

$$p(1, t, i) = 2t - (1 + 3t^2)i$$

has one zero in \mathbb{C}_+ , which agrees with the number of negative eigenvalues of A_1 . On the other hand, $p(0, t, i) = -3t^2i$ has no zeros in \mathbb{C}_+ . Notice also that $p(t, 1, i) = (2 - i)t^2 - 3i$ has one zero in \mathbb{C}_+ . This matches the number of negative eigenvalues of A_0 , which is what we would like to have more generally in order for Theorem 1 to have a more symmetric statement.

4. Proof of Corollary 1

Notice that $t \mapsto p(x - te_2)$ is either identically zero or only has real roots by Hurwitz's theorem since this polynomial can be obtained as the limit as $a \searrow 0$ of

$$t \mapsto p(x - t(ae_1 + e_2)).$$

Any factors depending only on x_0, x_1 can easily be dealt with separately, so we may assume that there are no such factors. So, p satisfies the hypotheses of Theorem 1. Also, $t \mapsto p(1, t, i)$ can have no zeros in the upper half-plane for if it had such a zero $z = x + iy$ where $y > 0$, then

$$t \mapsto p((1, x, 0) + t(0, y, 1))$$

would have the nonreal zero $t = i$, contradicting hyperbolicity in the direction $(0, y, 1)$. This shows that $\text{rank } B_- = 0$ in Theorem 1 and therefore A_1 is positive semidefinite, as desired.

5. Proof of Theorem 2 from Theorem C

We largely follow the scheme of [21]. Let $P \in \mathbb{R}[x_0, x_1, x_2, x_3]$ be homogeneous of degree d , of degree 1 in x_2, x_3 , and of degree n in x_1 . Assume that P is hyperbolic with respect to the cone $\{(0, v_1, v_2, v_3) : v_1, v_2, v_3 > 0\}$. Then, for $x = (x_1, x_2, x_3)$,

$$q(x) = P(1, x)$$

has no zeros in $\mathbb{C}_+^3 \cup \mathbb{C}_-^3$, and $\overline{q(\bar{x})} = q(x)$. Switching to the tridisk, we see that

$$f(z) = q\left(i \frac{1+z_1}{1-z_1}, i \frac{1+z_2}{1-z_2}, i \frac{1+z_3}{1-z_3}\right) \left(\frac{1-z_1}{2i}\right)^n \left(\frac{1-z_2}{2i}\right) \left(\frac{1-z_3}{2i}\right)$$

has no zeros in $\mathbb{D}^3 \cup \mathbb{E}^3$. Note that we may as well assume that f is irreducible since otherwise f would have a factor depending on one or two variables alone, in which case there is no issue with having a determinantal representation.

Let $1/\bar{z} = (1/\bar{z}_1, 1/\bar{z}_2, 1/\bar{z}_3)$ for $z \in \mathbb{C}^3$ and define

$$\begin{aligned} \tilde{f}(z) &= z_1^n z_2 z_3 \overline{f(1/\bar{z})}, \\ \frac{\partial \tilde{f}}{\partial z_j} &= \frac{z_1^n z_2 z_3}{z_j} \overline{\frac{\partial f}{\partial z_j}(1/\bar{z})} \quad \text{for } j = 1, 2, 3. \end{aligned}$$

Since q has real coefficients, we can show that $\tilde{f} = f$ and

$$\begin{aligned} nf &= z_1 \frac{\partial f}{\partial z_1} + \frac{\partial \tilde{f}}{\partial z_1}, \\ f &= z_j \frac{\partial f}{\partial z_j} + \frac{\partial \tilde{f}}{\partial z_j} \quad \text{for } j = 2, 3 \end{aligned}$$

after some simple computations. Thus, $(n + 2)f = p + \tilde{p}$ where

$$p(z) = \sum_{j=1}^3 \frac{\partial \tilde{f}}{\partial z_j}, \quad \tilde{p}(z) = \sum_{j=1}^3 z_j \frac{\partial f}{\partial z_j}.$$

Let $f_t(z) = f(tz)$ for $0 < t < 1$. Then, f_t has no zeros in $\overline{\mathbb{D}^3}$, and if we set $\tilde{f}_t(z) = t^{n+2} f(z/t)$, then $|f_t| = |\tilde{f}_t|$ on \mathbb{T}^3 (since $\tilde{f} = f$), and so \tilde{f}_t/f_t is analytic and bounded by 1 in modulus for $z \in \mathbb{D}^3$ by the maximum principle. Now, for $z \in \overline{\mathbb{D}^3}$,

$$\begin{aligned} 0 &\leq \lim_{t \nearrow 1} \frac{|f(tz)|^2 - |t^{n+2} f(z/t)|^2}{1 - t^2} (n + 2) \\ &= (n + 2)^2 |f(z)|^2 - 2 \operatorname{Re}(\tilde{p}(z)(n + 2)\overline{f(z)}) \\ &= |p(z)|^2 - |\tilde{p}(z)|^2 \quad \text{since } (n + 2)f = p + \tilde{p} \end{aligned}$$

with some computations omitted (see [21] for more details). This shows that if p vanishes in \mathbb{D}^3 , then so does \tilde{p} , and so does f , which by assumption does not happen. Hence, p has no zeros in \mathbb{D}^3 .

Note that if p and \tilde{p} had a common factor, then this would be a factor of f , which we have already ruled out; we point out that p and \tilde{p} cannot be multiples of one another since \tilde{p} vanishes at the origin. The *conclusion* of Theorem C holds for such a p , but since we have only stated it for polynomials with no zeros on $\overline{\mathbb{D}^3}$ (as opposed to \mathbb{D}^3), we must explain how to address the case at hand. The main point is that for $0 < t < 1$, $p_t(z) = p(tz)$ will satisfy the hypotheses of Theorem C, and therefore there exist vector polynomials E_1^t, E_2^t, E_3^t corresponding to p_t as in Theorem C. Then,

$$\sup_{\mathbb{T}^3} |p(z)|^2 \geq (1 - |z_j|^2) |E_j^t(z)|^2$$

shows that the vector polynomials E_j^t are locally bounded in \mathbb{D}^3 , and hence we can choose subsequences of $t \nearrow 1$ such that $E_1^t \in \mathbb{C}^{2n}[z]$, $E_2^t, E_3^t \in \mathbb{C}^2[z]$ converge to vector polynomials $E_1 \in \mathbb{C}^{2n}[z]$, $E_2, E_3 \in \mathbb{C}^2[z]$, and hence we will get

a sums-of-squares decomposition as in Theorem C. Note the polynomials in E_1, E_2, E_3 necessarily have degree at most $(n - 1, 1, 1), (n, 0, 1), (n, 1, 0)$ (this is proven in [23] for instance), and they will be nontrivial since p and \bar{p} have no factors in common. On the zero set Z_f of $f, p = -\bar{p}$, and therefore

$$0 = \sum_{j=1}^3 (1 - z_j \bar{w}_j) E_j(w)^* E_j(z) \tag{5.1}$$

for $z, w \in Z_f$. This equation ensures that the map

$$\begin{pmatrix} z_1 E_1(z) \\ z_2 E_2(z) \\ z_3 E_3(z) \end{pmatrix} \mapsto \begin{pmatrix} E_1(z) \\ E_2(z) \\ E_3(z) \end{pmatrix} \tag{5.2}$$

defined initially for vectors of the above form with $z \in Z_f$, extends linearly to a well-defined $(2n + 4) \times (2n + 4)$ unitary U . Some details: If a combination of vectors from the left side of (5.2) sums to zero, then (5.1) shows the corresponding combination on the right sums to zero. So, we get a well-defined linear map from the span of the left side of (5.2) to the span of the right side. Now, (5.1) shows this map is an isometry. Since we are in finite dimensions, it can be extended to a unitary.

Note that E_1, E_2, E_3 cannot vanish identically in Z_f without vanishing in all of \mathbb{C}^3 since the degrees are lower and f is irreducible. Let P_j for $j = 1, 2, 3$ be the projection onto the j th component in the orthogonal decomposition of $\mathbb{C}^{2n+4} = \mathbb{C}^{2n} \oplus \mathbb{C}^2 \oplus \mathbb{C}^2$, and let $M(z) = \sum_{j=1}^3 z_j P_j$. By (5.2), for $z \in Z_f$,

$$(I - UM(z)) \begin{pmatrix} E_1(z) \\ E_2(z) \\ E_3(z) \end{pmatrix} = 0,$$

and therefore $\det(I - UM(z)) = 0$ for $z \in Z_f \setminus \{z : E_1, E_2, E_3 = 0\}$. Basic results in algebraic geometry (such as in Chap. 4, Sect. 4 of [6]) can be used to establish that this implies that $\det(I - UM(z))$ vanishes for $z \in Z_f$ (i.e., $Z_f \setminus \{z : E_1, E_2, E_3 = 0\}$ is Zariski dense in Z_f) since f is irreducible and none of E_1, E_2, E_3 vanishes identically on Z_f .

Therefore, f divides $\det(I - UM(z))$. Write

$$f(z)g(z) = \det(I - UM(z))$$

for some polynomial g of degree at most $(n, 1, 1)$. As with Section 2, we convert back to q . There is some repetition in what follows, but since the situations are slightly different, we include the details. Now,

$$\begin{aligned} q(z)r(z) &= \det\left(\left(\sum_{j=1}^3 (z_j + i) P_j\right) - U\left(\sum_{j=1}^3 (z_j - i) P_j\right)\right) \\ &= \det((I - U)M(z) + i(I + U)) \end{aligned} \tag{5.3}$$

for

$$r(z) = (z_1 + i)^n (z_2 + i)(z_3 + i)g\left(\frac{z_1 - i}{z_1 + i}, \frac{z_2 - i}{z_2 + i}, \frac{z_3 - i}{z_3 + i}\right).$$

Let $U = V\begin{pmatrix} u & 0 \\ 0 & I \end{pmatrix}V^*$ be the spectral decomposition of U where u is $k \times k$ diagonal with unimodular entries, none of which equals 1. Here k is the rank of $I - U$. As in Section 2, the determinant (5.3) can be converted to

$$q(z)r(z) = (\text{const}) \det((V^*M(z)V)_{kk} + a) \tag{5.4}$$

where again $(V^*M(z)V)_{kk}$ refers to taking the upper $k \times k$ block of the given matrix, and $a = i(I + u)(I - u)^{-1}$. Finally, if we homogenize (5.4) at degree k —note this is at most $2n + 4$ —then

$$P(x)R(x) = (\text{const}) \det\left(x_0a + \sum_{j=1}^3 x_j A_j\right)$$

with $A_j = (V^*P_jV)_{kk}$ and $A_1 + A_2 + A_3 = (V^*IV)_{kk} = I$, and where $R(x) = x_0^{k-d}r((1/x_0)(x_1, x_2, x_3))$. This concludes the proof.

6. Concluding Questions and Remarks

We think it is worth discussing or rehashing some of the motivations and lingering questions of this paper in more detail.

Semihyperbolic polynomials have perhaps been overlooked because they lack one of the key features of hyperbolic polynomials. Specifically, if p is hyperbolic in the direction e , then the connected component of $\{x : p(x) \neq 0\}$ containing e is convex (see [31]). No such result holds for semihyperbolic polynomials (see Renegar [31], Section 2, or Section 3 of this paper). This convexity property ties hyperbolic polynomials to optimization and is “the cornerstone of hyperbolic programming” [31]. This begs the question why study *semihyperbolic* polynomials, which may lack this property?

First, we think it is a good general principle in mathematics to understand the degenerate versions of objects of interest. Notice that the (local uniform) limit of a sequence of homogeneous polynomials of degree d that are semihyperbolic with respect to a specific direction e is either semihyperbolic or identically zero. This follows from Hurwitz’s theorem applied to each polynomial $t \mapsto P(x - te)$. Hyperbolic polynomials do not share this property. Somewhat related is the following question mentioned in the Introduction.

QUESTION 1. Every trivariate semihyperbolic polynomial is a limit of hyperbolic polynomials. Is this true in the case of more variables?

Second, our main theorem, Theorem 1, shows that trivariate semihyperbolic polynomials possess determinantal representations just as in the hyperbolic case. We think this in itself provides a good justification for the study of semihyperbolic polynomials. This is a good point to formally state a question from the Introduction.

QUESTION 2. In Theorem 1, can the signature and rank of A_0 be determined directly from properties of p ?

Our own personal motivations for studying semihyperbolic polynomials came from the natural connection we presented between semihyperbolic polynomials and two-variable polynomials with no zeros on $\mathbb{T} \times \mathbb{D}$. These latter polynomials appeared in [8] essentially because of the realization that some of the theory of polynomials with no zeros in \mathbb{D}^2 could be pushed further to the situation of no zeros in $\mathbb{T} \times \mathbb{D}$. It was realized later that this initially unnatural condition is closely related to hyperbolicity and indeed is essentially equivalent to semihyperbolicity.

Finally, we wish to rehash our larger goal of the paper of connecting sums of squares formulas to determinantal representations. This description will be somewhat imprecise. The approach of this paper shows that if $p(z_1, z_2, \dots, z_n)$ has no zeros in \mathbb{D}^n and possesses a Hermitian sums-of-squares formula

$$|p|^2 - |\tilde{p}|^2 = \sum_{j=1}^n (1 - |z_j|^2) SOS_j \tag{6.1}$$

(here each SOS_j term is a sum of squared moduli of polynomials), then $p + \tilde{p}$ divides a unitary determinantal polynomial

$$\det(I - UD(z)).$$

Here U is a unitary matrix, and $D(z)$ is a diagonal matrix with coordinate functions on the diagonal. We can then convert $p + \tilde{p}$ via the Cayley transform and homogenization to a hyperbolic polynomial (hyperbolic with respect to all vectors with positive entries) and through some linear algebra get a self-adjoint determinantal polynomial. We can reverse engineer some of this: take a hyperbolic polynomial P (again hyperbolic with respect to vectors with positive entries) and convert to a polynomial q satisfying $q = \tilde{q}$. If q can be written as $p + \tilde{p}$ where p satisfies (6.1), then P divides a determinantal representation. If $n > 2$, not every $p \in \mathbb{C}[z_1, \dots, z_n]$ with no zeros in \mathbb{D}^n satisfies an equation of the form (6.1); such polynomials are called *Agler denominators*. A polynomial is an Agler denominator if and only if \tilde{p}/p satisfies a multivariable von Neumann inequality (see [23]).

We have also presented a modification to hyperbolicity/semihyperbolicity with respect to a specific direction. In n variables, this would entail, after various conversions, to understanding polynomials satisfying (6.1) where SOS_1, \dots, SOS_{n-1} are replaced with differences of squares. With the exception of our work in [8], this is relatively uncharted territory.

ACKNOWLEDGMENTS. I would like to sincerely thank my coauthors J. Geronimo and P. Iliev [8], my coauthor K. Bickel [2], and the anonymous referee. I would like to also thank Victor Vinnikov for introducing me to hyperbolic polynomials.

References

- [1] J. Agler and J. E. McCarthy, *Distinguished varieties*, Acta Math. 194 (2005), no. 2, 133–153, doi:[doi:10.1007/BF02393219](https://doi.org/10.1007/BF02393219).
- [2] K. Bickel and G. Knese, *Inner functions on the bidisk and associated Hilbert spaces*, J. Funct. Anal. 265 (2013), no. 11, 2753–2790, doi:[doi:10.1016/j.jfa.2013.08.002](https://doi.org/10.1016/j.jfa.2013.08.002).
- [3] J. Borcea and P. Brändén, *Multivariate Pólya–Schur classification problems in the Weyl algebra*, Proc. Lond. Math. Soc. (3) 101 (2010), no. 1, 73–104, doi:[10.1112/plms/pdp049](https://doi.org/10.1112/plms/pdp049).
- [4] P. Brändén, *Obstructions to determinantal representability*, Adv. Math. 226 (2011), no. 2, 1202–1212, doi:[10.1016/j.aim.2010.08.003](https://doi.org/10.1016/j.aim.2010.08.003).
- [5] B. J. Cole and J. Wermer, *Ando’s theorem and sums of squares*, Indiana Univ. Math. J. 48 (1999), no. 3, 767–791, doi:[10.1512/iumj.1999.48.1716](https://doi.org/10.1512/iumj.1999.48.1716).
- [6] D. Cox, J. Little, and D. O’Shea, *Ideals, varieties, and algorithms: an introduction to computational algebraic geometry and commutative algebra*, Undergrad. Texts Math., 3rd edition, Springer, New York, 2007.
- [7] L. Gårding, *An inequality for hyperbolic polynomials*, J. Math. Mech. 8 (1959), 957–965.
- [8] J. S. Geronimo, P. Iliev, and G. Knese, *Polynomials with no zeros on a face of the bidisk*, J. Funct. Anal. 270 (2016), no. 9, 3505–3558, doi:[10.1016/j.jfa.2016.02.002](https://doi.org/10.1016/j.jfa.2016.02.002).
- [9] J. S. Geronimo and H. J. Woerdeman, *Positive extensions, Fejér–Riesz factorization and autoregressive filters in two variables*, Ann. of Math. (2) 160 (2004), no. 3, 839–906, doi:[10.4007/annals.2004.160.839](https://doi.org/10.4007/annals.2004.160.839).
- [10] A. Grinshpan, D. S. Kaliuzhnyi-Verbovetskyi, V. Vinnikov, and H. J. Woerdeman, *Classes of tuples of commuting contractions satisfying the multivariable von Neumann inequality*, J. Funct. Anal. 256 (2009), no. 9, 3035–3054, doi:[10.1016/j.jfa.2008.09.012](https://doi.org/10.1016/j.jfa.2008.09.012).
- [11] ———, *Stable and real-zero polynomials in two variables*, Multidimens. Syst. Signal Process. 27 (2016), no. 1, 1–26, doi:[10.1007/s11045-014-0286-3](https://doi.org/10.1007/s11045-014-0286-3).
- [12] A. Grinshpan, D. S. Kaliuzhnyi-Verbovetskyi, and H. J. Woerdeman, *Norm-constrained determinantal representations of multivariable polynomials*, Complex Anal. Oper. Theory 7 (2013), no. 3, 635–654, doi:[10.1007/s11785-012-0262-6](https://doi.org/10.1007/s11785-012-0262-6).
- [13] L. Gurvits, *Combinatorics hidden in hyperbolic polynomials and related topics*, 2004, [arXiv:math/0402088v1](https://arxiv.org/abs/math/0402088v1).
- [14] ———, *Van der Waerden/Schrijver-Valiant like conjectures and stable (aka hyperbolic) homogeneous polynomials: one theorem for all*, Electron. J. Combin. 15 (2008), no. 1, Research Paper 66, 26. With a corrigendum.
- [15] F. R. Harvey and H. B. Lawson Jr., *Gårding’s theory of hyperbolic polynomials*, Comm. Pure Appl. Math. 66 (2013), no. 7, 1102–1128.
- [16] J. W. Helton and V. Vinnikov, *Linear matrix inequality representation of sets*, Comm. Pure Appl. Math. 60 (2007), no. 5, 654–674, doi:[10.1002/cpa.20155](https://doi.org/10.1002/cpa.20155).
- [17] P. D. Lax, *Differential equations, difference equations and matrix theory*, Comm. Pure Appl. Math. 11 (1958), 175–194.
- [18] A. S. Lewis, P. A. Parrilo, and M. V. Ramana, *The Lax conjecture is true*, Proc. Amer. Math. Soc. 133 (2005), no. 9, 2495–2499 (electronic), doi:[doi:10.1090/S0002-9939-05-07752-X](https://doi.org/10.1090/S0002-9939-05-07752-X).
- [19] D. Kerner and V. Vinnikov, *Determinantal representations of singular hypersurfaces in \mathbb{P}^n* , Adv. Math. 231 (2012), no. 3–4, 1619–1654, doi:[10.1016/j.aim.2012.06.014](https://doi.org/10.1016/j.aim.2012.06.014).
- [20] ———, *Polynomials with no zeros on the bidisk*, Anal. PDE 3 (2010), no. 2, 109–149, doi:[10.2140/apde.2010.3.109](https://doi.org/10.2140/apde.2010.3.109).

- [21] G. Knese, *Polynomials defining distinguished varieties*, Trans. Amer. Math. Soc. 362 (2010), no. 11, 5635–5655, doi:[10.2140/apde.2010.3.109](https://doi.org/10.2140/apde.2010.3.109).
- [22] ———, *Schur–Agler class rational inner functions on the tridisk*, Proc. Amer. Math. Soc. 139 (2011), no. 11, 4063–4072, doi:[10.1090/S0002-9939-2011-10975-4](https://doi.org/10.1090/S0002-9939-2011-10975-4).
- [23] ———, *Rational inner functions in the Schur–Agler class of the polydisk*, Publ. Mat. 55 (2011), no. 2, 343–357, doi:[10.1090/S0002-9939-2011-10975-4](https://doi.org/10.1090/S0002-9939-2011-10975-4).
- [24] M. Kummer, *Determinantal representations and Bézoutions*, Math. Z. (2016), doi:[10.1007/s00209-016-1715-9](https://doi.org/10.1007/s00209-016-1715-9).
- [25] M. Kummer, D. Plaumann, and C. Vinzant, *Hyperbolic polynomials, interlacers, and sums of squares*, Math. Program., Ser. B 153 (2015), no. 1, 223–245, doi:[10.1007/s10107-013-0736-y](https://doi.org/10.1007/s10107-013-0736-y).
- [26] A. W. Marcus, D. A. Spielman, and N. Srivastava, *Interlacing families II: mixed characteristic polynomials and the Kadison–Singer problem*, Ann. of Math. (2) 182 (2015), no. 1, 327–350, doi:[10.4007/annals.2015.182.1.8](https://doi.org/10.4007/annals.2015.182.1.8).
- [27] T. Netzer and A. Thom, *Polynomials with and without determinantal representations*, Linear Algebra Appl. 437 (2012), no. 7, 1579–1595, doi:[10.1016/j.laa.2012.04.043](https://doi.org/10.1016/j.laa.2012.04.043).
- [28] T. Netzer, D. Plaumann, and A. Thom, *Determinantal representations and the Hermite matrix*, Michigan Math. J. 62 (2013), no. 2, 407–420, doi:[10.1307/mmj/1370870379](https://doi.org/10.1307/mmj/1370870379).
- [29] R. Pemantle, *Hyperbolicity and stable polynomials in combinatorics and probability*, Current developments in mathematics, 2011, pp. 57–123, Int. Press, Somerville, MA, 2012.
- [30] D. Plaumann and C. Vinzant, *Determinantal representations of hyperbolic plane curves: an elementary approach*, J. Symbolic Comput. 57 (2013), 48–60, doi:[10.1016/j.jsc.2013.05.004](https://doi.org/10.1016/j.jsc.2013.05.004).
- [31] J. Renegar, *Hyperbolic programs, and their derivative relaxations*, Found. Comput. Math. 6 (2006), no. 1, 59–79, doi:[10.1007/s10208-004-0136-z](https://doi.org/10.1007/s10208-004-0136-z).
- [32] V. Vinnikov, *Selfadjoint determinantal representations of real plane curves*, Math. Ann. 296 (1993), no. 3, 453–479, doi:[10.1007/BF01445115](https://doi.org/10.1007/BF01445115).
- [33] ———, *LMI representations of convex semialgebraic sets and determinantal representations of algebraic hypersurfaces: past, present, and future*, Mathematical methods in systems, optimization, and control, Oper. Theory Adv. Appl., 222, pp. 325–349, Birkhäuser/Springer Basel AG, Basel, 2012, doi:[10.1007/978-3-0348-0411-0_23](https://doi.org/10.1007/978-3-0348-0411-0_23).
- [34] D. G. Wagner, *Multivariate stable polynomials: theory and applications*, Bull. Amer. Math. Soc. (N.S.) 48 (2011), no. 1, 53–84, doi:[10.1090/S0273-0979-2010-01321-5](https://doi.org/10.1090/S0273-0979-2010-01321-5).

Department of Mathematics
Washington University in St. Louis
St. Louis, MO 63130
USA

geknese@wustl.edu