

Washington University in St. Louis
Washington University Open Scholarship

Mathematics Faculty Publications

Mathematics and Statistics

5-1-2013

On the Isomorphism Question for Complete Pick Multiplier Algebras

Matt Kerr

Washington University in St Louis

John E. McCarthy

Washington University in St Louis

Orr Moshe Shalit

Ben-Gurion University of the Negev

Follow this and additional works at: https://openscholarship.wustl.edu/math_facpubs

Recommended Citation

Kerr, Matt; McCarthy, John E.; and Shalit, Orr Moshe, "On the Isomorphism Question for Complete Pick Multiplier Algebras" (2013). *Mathematics Faculty Publications*. 13.

https://openscholarship.wustl.edu/math_facpubs/13

This Article is brought to you for free and open access by the Mathematics and Statistics at Washington University Open Scholarship. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of Washington University Open Scholarship. For more information, please contact digital@wumail.wustl.edu.

ON THE ISOMORPHISM QUESTION FOR COMPLETE PICK MULTIPLIER ALGEBRAS

MATT KERR, JOHN E. MCCARTHY, AND ORR MOSHE SHALIT

ABSTRACT. Every multiplier algebra of an irreducible complete Pick kernel arises as the restriction algebra $\mathcal{M}_V = \{f|_V : f \in \mathcal{M}_d\}$, where d is some integer or ∞ , \mathcal{M}_d is the multiplier algebra of the Drury-Arveson space H_d^2 , and V is a subvariety of the unit ball. For finite d it is known that, under mild assumptions, every isomorphism between two such algebras \mathcal{M}_V and \mathcal{M}_W is induced by a biholomorphism between W and V . In this paper we consider the converse, and obtain positive results in two directions. The first deals with the case where V is the proper image of a finite Riemann surface. The second deals with the case where V is a disjoint union of varieties.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

The general theme of this note is the following question.

(Q) Given two specific function algebras \mathcal{M}_V and \mathcal{M}_W associated with two analytic varieties V and W , if there is a biholomorphic equivalence $h : V \rightarrow W$, does composition with h induce an isomorphism from \mathcal{M}_W to \mathcal{M}_V ?

This type of question — and its converse, whether isomorphism of the algebras implies biholomorphic equivalence of the varieties — can be asked in many contexts, *e.g.* in algebraic geometry it is well known that two affine algebraic varieties are isomorphic if and only if the algebras of polynomial functions on the varieties are isomorphic [15, Cor. I.3.7]. Coming back to complex analysis, a theorem of L. Bers [9] says that two domains V and W in the plane are conformally equivalent if and only if the corresponding algebras of holomorphic functions $O(V)$ and $O(W)$ are isomorphic. Note that in the setting of Bers's theorem,

Date: February 13, 2012.

1991 *Mathematics Subject Classification.* 30H50, 47B32.

Partially supported by National Science Foundation Grant DMS 1068974.

Partially supported by National Science Foundation Grant DMS 0966845.

Partially supported by Israel Science Foundation Grant no. 474/12 and by the European Union Seventh Framework Programme (*FP7/2007-2013*) under grant agreement no. 321749.

(Q) is trivial and it is the converse that is the interesting part. In our setting, (Q) turns out to be harder than its converse (though the converse is by no means trivial).

Here we shall consider multiplier algebras of a certain Hilbert space associated with a variety V in the unit ball \mathbb{B}_d .

Let H_d^2 be the reproducing kernel Hilbert space on \mathbb{B}_d , the unit ball of \mathbb{C}^d , with kernel

$$k_d(z, w) = \frac{1}{1 - \langle z, w \rangle}.$$

This space was first introduced by S. Drury [12], and later studied in [18, 8, 2]. Let \mathcal{M}_d be the multiplier algebra $\text{Mult}(H_d^2)$ of H_d^2 . Here d is some integer or ∞ .

We shall use the word *variety* to mean a joint zero set in the ball of H_d^2 functions. That is, V is a variety in our sense if there is some set $E \subseteq H_d^2$ such that

$$V = \{z \in \mathbb{B}_d : \forall f \in E, f(z) = 0\}.$$

By [3, Theorem 9.27] this is the same as being a joint zero set of functions in \mathcal{M}_d . We shall denote by \overline{V} the closure of V in $\overline{\mathbb{B}_d}$.

Following [10, Section 5], let us say that a variety V is *irreducible* if for any regular point $\lambda \in V$, and any non-empty ball B centered at λ and contained in \mathbb{B}_d , the intersection of the zero sets of all multipliers vanishing on $V \cap B$ is exactly V . Note that if V cannot be written as the proper union of two analytic varieties *in the classical sense* then V is irreducible according to our definition. Indeed, this follows from the fact that in this case the set of regular points of V is a connected submanifold (see [14, Section III.C]), thus any analytic function vanishing on a neighbourhood in V of a regular point vanishes everywhere on V .

If V is a variety we define

$$\mathcal{F}_V = \overline{\text{span}}\{k_d(\cdot, \lambda) : \lambda \in V\}.$$

The Hilbert space \mathcal{F}_V is naturally a reproducing kernel Hilbert space of functions on the variety V . We define

$$\mathcal{M}_V = \{f|_V : f \in \mathcal{M}_d\}.$$

By [10, Proposition 2.6], \mathcal{M}_V is equal to $\text{Mult}(\mathcal{F}_V)$, and is completely isometrically isomorphic to the quotient of \mathcal{M}_d by the ideal

$$J_V = \{f \in \mathcal{M}_d : f|_V = 0\}.$$

By universality of H_d^2 , every multiplier algebra of an irreducible complete Pick kernel is of the form \mathcal{M}_V for some d and some variety $V \subseteq \mathbb{B}_d$ (see [1]).

Question (Q) in the setting just described was first considered by K. Dav-idson, C. Ramsey and O. Shalit in [11] for the case where V and W are homogeneous varieties. Under additional technical assumptions it was shown that the answer to question (Q) is *yes*. The additional technical assumptions were later removed by M. Hartz [16]. In [10] question (Q) and its converse were treated in greater generality, and it was shown that, when $d < \infty$, the converse direction holds true for varieties which are finite unions of irreducible varieties plus a discrete variety. Moreover, it was shown that the answer to (Q) is positive in some very special cases, and that it is negative in some rather pathological cases. The goal of the present work is to obtain a positive answer to (Q) for a wide class of “nice” varieties.

In Theorem 4.2 and Corollary 4.11, we answer question (Q) in the affirmative in the case that V is a finite Riemann surface with a finite number of pairs of points identified (under the assumption of sufficient regularity on the boundary). This generalizes work of D. Alpay, M. Putinar and V. Vinnikov [6], who proved the result when V was the unit disk, and N. Arcozzi, R. Rochberg and E. Sawyer [7, Subsection 2.3.6], who proved the result for V a planar domain. As a corollary we obtain an extension theorem: *every bounded holomorphic function on V extends to a function in \mathcal{M}_d* (see Corollary 4.12). We also prove that the converse to (Q) holds for H^∞ of finite Riemann surfaces in Corollary 4.9. Our work requires an understanding of Hardy spaces on finite Riemann surfaces — we give background information on this in Section 3.

In Section 6, we discuss varieties with a finite number of disjoint components, and find sufficient conditions for an isomorphism on each separate component to give an isomorphism on the whole variety. For this we require some results on the maximal ideal spaces of \mathcal{M}_V and of its norm closed analogue \mathcal{A}_V , results which are obtained in Section 5.

2. ISOMORPHISM BETWEEN \mathcal{M}_V AND \mathcal{M}_W

In this section, we shall assume that $d < \infty$.

Let us denote by $\Delta(\mathcal{M}_V)$ the *character space* of \mathcal{M}_V , that is, the space of all nonzero multiplicative linear functionals on \mathcal{M}_V . All characters are unital and contractive, hence completely contractive (since every contractive functional of an operator space is a complete contraction [17, Proposition 3.8]). Note that, since the algebras are semi-simple, every homomorphism $\alpha : \mathcal{M}_V \rightarrow \mathcal{M}_W$ is automatically norm continuous. Thus, if $\alpha : \mathcal{M}_V \rightarrow \mathcal{M}_W$ is a unital homomorphism, then

there is an induced continuous map

$$\alpha^* : \Delta(\mathcal{M}_W) \rightarrow \Delta(\mathcal{M}_V)$$

defined by $\alpha^*(\rho) = \rho \circ \alpha$ for all $\rho \in \Delta(\mathcal{M}_W)$.

Let us denote by z_1, \dots, z_d the coordinate functions in \mathcal{M}_V . Then (z_1, \dots, z_d) is a row contraction and since all characters are completely contractive, $(\rho(z_1), \dots, \rho(z_d))$ is in $\overline{\mathbb{B}_d}$ for every $\rho \in \Delta(\mathcal{M}_V)$. This gives rise to a continuous map

$$\pi : \Delta(\mathcal{M}_V) \rightarrow \overline{\mathbb{B}_d}.$$

For every $\lambda \in \mathbb{B}_d$ we call $\pi^{-1}(\lambda)$ *the fiber over λ* . In Section 3 of [10] it is shown that $\pi(\Delta(\mathcal{M}_V)) \cap \mathbb{B}_d = V$, that the fiber over every $\lambda \in V$ is a singleton, and that π identifies the space of WOT-continuous characters with V . If V extends analytically across $\partial\mathbb{B}_d$, by Corollary 5.4 below $\pi(\Delta(\mathcal{M}_V)) = \overline{V}$. It is easy to see that $\pi(\Delta(\mathcal{M}_V)) \supseteq \overline{V}$ always holds.

Question 2.1. *Is $\pi(\Delta(\mathcal{M}_V)) = \overline{V}$ for all V ?*

Theorem 2.2. *Let $V, W \subseteq \mathbb{B}_d$ be varieties, $d < \infty$. Suppose that V and W are each a union of finitely many varieties, each of which is either irreducible or discrete. Let α be an isomorphism between \mathcal{M}_V and \mathcal{M}_W . Then α induces a holomorphic map $h : \mathbb{B}_d \rightarrow \mathbb{C}^d$ that restricts to a biholomorphism from W onto V . The isomorphism α is implemented by composition with h . Moreover, if $[h_1, \dots, h_d] = h$ are the components of h , then $h_i \in \mathcal{M}_d$ for all $i = 1, \dots, d$.*

PROOF: This theorem and some variants are treated in detail in Section 5 of [10], but we would like to repeat some of the arguments. Consider the coordinate functions $z_i := z_i|_V$ in \mathcal{M}_V . For all i , $\alpha(z_i) \in \mathcal{M}_W$, and is therefore the restriction to W of multiplier $h_i \in \mathcal{M}_d$:

$$\alpha(z_i) = h_i|_V.$$

This gives rise to a holomorphic function $h = [h_1, \dots, h_d] : \mathbb{B}_d \rightarrow \mathbb{C}^d$ with components in \mathcal{M}_d .

Fix $\lambda \in W$, and let ρ_λ be the evaluation functional at λ on \mathcal{M}_W . Then $\alpha^*(\rho_\lambda)$ is a character in $\Delta(\mathcal{M}_V)$. We now compute

$$[\alpha^*(\rho_\lambda)](z_i) = \rho_\lambda(\alpha(z_i)) = \alpha(z_i)(\lambda).$$

So $\alpha^*(\rho_\lambda)$ is in the fiber over $(\alpha(z_1)(\lambda), \dots, \alpha(z_d)(\lambda)) = h(\lambda)$.

In [10, Section 5] it is proved that $h(\lambda) \in V$, and therefore that $\alpha^*(\rho_\lambda)$ is also an evaluation functional. Thus, for every $f \in \mathcal{M}_V$ and every $\lambda \in W$,

$$\begin{aligned} \alpha(f)(\lambda) &= \rho_\lambda(\alpha(f)) = \alpha^*(\rho_\lambda)(f) \\ &= \rho_{h(\lambda)}(f) = (f \circ h)(\lambda). \end{aligned}$$

Therefore $\alpha(f) = f \circ h$.

Finally, since the same reasoning holds for α^{-1} , h is a biholomorphism. \square

In general, the function h appearing in the above theorem extends to a homeomorphism between the character spaces, but not to a homeomorphism between \overline{V} and \overline{W} . Indeed, suppose that V and W are both interpolating sequences. Then \mathcal{M}_V and \mathcal{M}_W are isomorphic (both are isomorphic to ℓ^∞), and consequently V and W are biholomorphic, but \overline{V} and \overline{W} need not be homeomorphic.

Let \mathcal{A}_V be the norm closure of the polynomials in \mathcal{M}_V .

Question 2.3. *When is $\mathcal{A}_V = \mathcal{M}_V \cap C(\overline{V})$?*

We will see in Section 5 that always $\mathcal{A}_V \subseteq \mathcal{M}_V \cap C(\overline{V})$.

Theorem 2.4. *Let $V, W \subseteq \mathbb{B}_d$ be varieties, $d < \infty$. Let α be an isomorphism between \mathcal{M}_V and \mathcal{M}_W , and let h be as in Theorem 2.2. Then α maps \mathcal{A}_V into \mathcal{A}_W , (respectively $\mathcal{M}_V \cap C(\overline{V})$ into $\mathcal{M}_W \cap C(\overline{W})$), if and only if $h_i|_W \in \mathcal{A}_W$ (respectively, $\mathcal{M}_W \cap C(\overline{W})$) for all $i = 1, \dots, d$. Moreover, α is an isomorphism from $\mathcal{M}_V \cap C(\overline{V})$ onto $\mathcal{M}_W \cap C(\overline{W})$ if and only if $h|_W$ extends to be a homeomorphism from \overline{W} to \overline{V} .*

PROOF: We use the notation of the proof of the above theorem. If α takes \mathcal{A}_V into \mathcal{A}_W then for all i , $h_i|_W = \alpha(z_i) \in \mathcal{A}_W$ (similarly for $\mathcal{M}_V \cap C(\overline{V})$ and $\mathcal{M}_W \cap C(\overline{W})$).

Conversely, if $h_i|_W = \alpha(z_i) \in \mathcal{A}_W$ for all i , then, since α is continuous, we have that $\alpha(\mathcal{A}_V) \subseteq \mathcal{A}_W$. If $h_i|_W \in \mathcal{M}_W \cap C(\overline{W})$, then $f \circ h$ is in $\mathcal{M}_V \cap C(\overline{V})$ for every f that is in $\mathcal{M}_V \cap C(\overline{V})$, so $\alpha(\mathcal{M}_V \cap C(\overline{V})) \subseteq \mathcal{M}_W \cap C(\overline{W})$.

If $\alpha : \alpha(\mathcal{M}_V \cap C(\overline{V})) \rightarrow \mathcal{M}_W \cap C(\overline{W})$ is an isomorphism, then h must be continuous, and one-to-one, on \overline{W} , and so must h^{-1} , therefore h is a homeomorphism. \square

3. BACKGROUND ON FINITE RIEMANN SURFACES

A connected *finite Riemann surface* Σ is a connected open proper subset of some compact Riemann surface such that the boundary $\partial\Sigma$ is also the boundary of the closure and is the union of finitely many disjoint simple closed analytic curves. A general finite Riemann surface is a finite disjoint union of connected ones. Figure 1 shows a picture of a connected finite Riemann surface with three boundary circles. We




FIGURE 1. Finite Riemann Surface with two handles and three boundary components

shall identify Σ with its image in the double, $\hat{\Sigma}$, the compact Riemann surface obtained by attaching another copy of Σ along $\partial\Sigma$.

Let $O(\Sigma)$ denote the algebra of all holomorphic functions on Σ , and $A(\Sigma)$ the holomorphic functions that extend continuously to $\partial\Sigma$. Since Σ is not compact, there is a plentiful supply of holomorphic functions; indeed, by a result of J. Wermer [20], the real parts of functions in $A(\Sigma)$ form a finite codimensional subspace of the space of all continuous real-valued functions on $\partial\Sigma$. This means that in many important ways function theory on Σ is like function theory on the unit disk, subject to a finite number of extra conditions. See for example the paper [5].

Let us fix some base-point a in Σ , and let ω be the harmonic measure with respect to a , *i.e.* the measure on $\partial\Sigma$ with the property that

$$\int_{\partial\Sigma} u(\zeta) d\omega(\zeta) = u(a)$$

for every function u that is harmonic on Σ and continuous on $\bar{\Sigma}$.

We shall define $H^2(\Sigma)$ to be the closure in $L^2(\omega)$ of $A(\Sigma)$. Its multiplier algebra is $H^\infty(\Sigma)$, the bounded analytic functions on Σ . Note that the norm in $H^2(\Sigma)$ depends on the choice of base-point a , but the norm in $H^\infty(\Sigma)$ does not — it is the supremum of the modulus on Σ [5].

J.D. Fay [13, Prop 6.15] gave an explicit formula for the Szegő kernel for $H^2(\Sigma)$, *i.e.* for the function $j(\zeta, \eta)$ that is conjugate-holomorphic in η , as a function of ζ is holomorphic and lies in $H^2(\Sigma)$ for each fixed η , and satisfies

$$(3.1) \quad \int_{\partial\Sigma} f(z) \overline{j(\zeta, \eta)} d\omega(\zeta) = f(\eta) \quad \forall f \in H^2(\Sigma).$$

We need to know how j behaves qualitatively near the boundary. Fix some point η in $\partial\Sigma$, and let us introduce local coordinates. Let U be a small neighborhood of η in $\hat{\Sigma}$, and V a neighborhood of an arc of the unit circle in the complex plane, and $\phi : V \rightarrow U$ a biholomorphic map that sends $V \cap \mathbb{D}$ to $U \cap \Sigma$, and $V \cap \mathbb{D}^c$ to $U \cap \Sigma^c$. Let us write $\zeta = \phi(z)$ and $\eta = \phi(w)$. Let $J(z, w) = j(\phi(z), \phi(w))$.

Proposition 3.2. *With notation as above, let $w_0 \in V \cap \partial\mathbb{D}$. Then there is a neighborhood V_0 of w_0 that is contained in V and so that on $V_0 \times (V_0 \cap \partial\mathbb{D})$ we have*

$$(3.3) \quad J(z, w) = \frac{G(z, w)}{w - z}$$

where G is a holomorphic function that is non-vanishing on $V \times (V_0 \cap \partial\mathbb{D})$.

Proposition 3.2 is a corollary of Fay's theorem, which gives an explicit formula for j in terms of the theta function θ and the prime form E of Σ :

Theorem 3.4 (Fay). *The reproducing kernel $j(\zeta, \eta)$ for $H^2(\Sigma)$ is given by:*

$$(3.5) \quad j(\zeta, \eta) = \frac{\theta(\eta^* - \zeta - e)\theta(a^* - a - e)E(\zeta, a^*)E(\eta^*, a)}{\theta(a^* - \zeta - e)\theta(\eta^* - a - e)E(\eta^*, \zeta)E(a, a^*)}.$$

Moreover, if $U = U^*$ is a small symmetric neighborhood of some point in $\partial\Sigma$, then on $U \times U$, the function $k(\zeta, \eta) := j(\zeta, \eta^*)$ is meromorphic, non-vanishing, and is holomorphic except for simple poles where $\eta = \zeta$.

Here e is a certain even half-period, E is the prime form, which in local coordinates has the expansion

$$E(z, w) = \overline{E(z^*, w^*)} = \frac{(z - w) + O(z - w)^2}{\sqrt{dzdw}},$$

θ is the theta function for $\hat{\Sigma}$, and we use ζ^* to mean the involution on $\hat{\Sigma}$ that swaps Σ and $\hat{\Sigma} \setminus \bar{\Sigma}$. For details see Fay's monograph [13].

A Toeplitz operator on $H^2(\Sigma)$ is the operator of multiplication by a continuous function f on $\partial\Sigma$ followed by projection from $L^2(\sigma)$ onto $H^2(\Sigma)$; we denote this operator by T_f . The C^* -algebra generated by all the Toeplitz operators is called the Toeplitz algebra and is denoted \mathcal{T} . Let us denote the algebra of compact operators on $H^2(\Sigma)$ by \mathcal{K} . We need to know that \mathcal{T} is an extension of $\mathcal{T} \cap \mathcal{K}$ by the continuous functions on $\partial\Sigma$. This result is well-known, but we do not know of a convenient reference, so we include a proof for completeness.

Proposition 3.6. *The quotient $\mathcal{T}/(\mathcal{K} \cap \mathcal{T})$ is isometrically isomorphic to $C(\partial\Sigma)$ via the map $f \leftrightarrow T_f + \mathcal{K} \cap \mathcal{T}$.*

PROOF: If Σ is the disjoint union of $\Sigma_1, \dots, \Sigma_k$, then every Toeplitz operator on $H^2(\Sigma)$ decomposes into a direct sum of Toeplitz operators on $H^2(\Sigma_i)$, $i = 1, \dots, k$. Thus, we may assume that Σ is connected.

For an inner function B in $A(\Sigma)$ (inner means that the modulus is 1 on $\partial\Sigma$), the codimension of $BH^2(\Sigma)$ in $H^2(\Sigma)$ is equal to the number of zeroes of B in Σ , and hence finite. It follows that for such a B , the operator T_B is essentially normal, because the kernel of $T_B T_{B^*} - T_{B^*} T_B$ is $BH^2(\Sigma)$.

The Toeplitz algebra \mathcal{T} is irreducible. Indeed, any operator in the commutant of $A(\Sigma)$ must be (multiplication by) a function in $H^\infty(\Sigma)$, so the commutant of \mathcal{T} is contained in the intersection of $H^\infty(\Sigma)$ with its complex conjugate, and since Σ is connected, this reduces to the scalars. Therefore $\mathcal{K} \subseteq \mathcal{T}$.

A result of E. L. Stout [19] is that every connected finite Riemann surface can be embedded in \mathbb{C}^3 with a triple B_1, B_2, B_3 of inner functions in $A(\Sigma)$. We may therefore assume that $\Sigma \subseteq \mathbb{D}^3$ and that B_i is equal to the coordinate function z_i . Then B_1, B_2, B_3 generate $A(\Sigma)$, and these functions and their conjugates generate $C(\Sigma)$. Since $T_g T_f = T_{gf}$ for every $f \in A(\Sigma)$ and $g \in C(\partial\Sigma)$, it follows that $T_{B_1}, T_{B_2}, T_{B_3}$ generate the C*-algebra \mathcal{T} . It follows from $T_B T_{B^*} - T_{B^*} T_B \in \mathcal{K}$ that \mathcal{T}/\mathcal{K} is commutative. The spectrum of this commutative C*-algebra can be naturally identified with $\partial\Sigma$, thus the result follows. \square

4. BIHOLOMORPHIC MAPS INDUCE ISOMORPHISMS

In [6], Alpay, Putinar and Vinnikov showed that if $h : \mathbb{D} \rightarrow W$ is a biholomorphic unramified C^2 map that is transversal at the boundary, then there is an isomorphism of multiplier algebras from $\mathcal{M}_{\mathbb{D}} = H^\infty(\mathbb{D})$ to \mathcal{M}_W . Arcozzi, Rochberg and Sawyer extended this result by allowing \mathbb{D} to be replaced by a finitely connected planar domain [7]. We extend one step further, to the case of a finite Riemann surface. Note that we do not require d to be finite.

In [4], the following definition was introduced and studied:

Definition 4.1. *A holomap is a proper holomorphic map h from a Riemann surface Σ into a bounded open set $U \subseteq \mathbb{C}^n$ such that there is a finite subset E of Σ with the property that h is non-singular and injective on $\Sigma \setminus E$.*

If h is a holomap from Σ to W and X is a space of holomorphic functions on Σ , then by X_h we mean the functions in X that are of

the form $F \circ h$ for some holomorphic function F on W (thus we get $H_h^p(\Sigma)$, $A_h(\Sigma)$, etc).

Theorem 4.2. *Let Σ be a finite Riemann surface, and W a variety in \mathbb{B}_d . Let $h : \Sigma \rightarrow \mathbb{B}_d$ be a holomap that maps Σ onto W , is C^2 up to $\partial\Sigma$, and is one-to-one on $\partial\Sigma$. Assume that h is transversal, i.e.*

$$(4.3) \quad \langle Dh(\zeta), h(\zeta) \rangle \neq 0, \quad \forall \zeta \in \partial\Sigma.$$

Then the map

$$\alpha : f \mapsto f \circ h$$

is an isomorphism from \mathcal{F}_W onto $H_h^2(\Sigma)$. Consequently, the mapping $\phi \mapsto \phi \circ h$ implements an isomorphism of \mathcal{M}_W onto $H_h^\infty(\Sigma)$.

Remark 4.4. The calculation of Dh , the derivative of h , depends on the choice of local coordinate, and another choice will differ by a non-zero multiplicative factor (the derivative of the change of variable map). This will not, however, affect whether or not $\langle Dh(\zeta), h(\zeta) \rangle$ is non-zero.

Remark 4.5. One can think of W as the space Σ with a finite number of pairs of points identified, or limiting cases of this where one gets cusps. See [4] for a detailed description.

PROOF: The main idea of the proof goes back to [6]. To show that α given by the formula $f \mapsto f \circ h$ is a well defined bounded and invertible map from \mathcal{F}_W onto $H_h^2(\Sigma)$, we will formally compute α^* and $R := \alpha\alpha^*$, and show that R is an injective Fredholm operator. Being positive and Fredholm, injectivity implies invertibility, and the first claim in the theorem follows. A straightforward computation then shows that the asserted isomorphism between \mathcal{M}_W and $H_h^\infty(\Sigma)$ is the similarity induced by α .

For f in $H^2(\Sigma)$ (not necessarily in $H_h^2(\Sigma)$) let us calculate $\alpha^*(f)$. This is the function on W whose value at a point w is given by

$$\langle \alpha^*f, k_d(\cdot, w) \rangle_{H_d^2} = \langle f(\zeta), k_d(h(\zeta), w) \rangle_{H^2(\Sigma)}$$

Let $j(\zeta, \eta) = j_\eta(\zeta)$ be the reproducing kernel for $H^2(\Sigma)$, defined by (3.1). We have that

$$\begin{aligned} \alpha^*j_\eta(w) &= \langle j_\eta(\zeta), k_d(h(\zeta), w) \rangle \\ &= \frac{1}{1 - \langle w, h(\eta) \rangle} \\ &= k_d(w, h(\eta)). \end{aligned}$$

Let $R = \alpha\alpha^*$. We have shown

$$(4.6) \quad Rf(\zeta) = \int_{\partial\Sigma} f(\eta) \frac{1}{1 - \langle h(\zeta), h(\eta) \rangle} d\omega(\eta).$$

Let us write

$$L(\zeta, \eta) = \frac{1}{j(\zeta, \eta)} \frac{1}{1 - \langle h(\zeta), h(\eta) \rangle}.$$

For $\eta \in \partial\Sigma$, let $L(\eta, \eta) = \lim_{\zeta \rightarrow \eta} L(\zeta, \eta)$. Break R up as

$$\begin{aligned} R_1 f(\zeta) &= \int_{\partial\Sigma} f(\eta) j(\zeta, \eta) L(\eta, \eta) d\omega(\eta) \\ R_2 f(\zeta) &= \int_{\partial\Sigma} f(\eta) j(\zeta, \eta) [L(\zeta, \eta) - L(\eta, \eta)] d\omega(\eta). \end{aligned}$$

Fix some point η in $\partial\Sigma$, and let us introduce local coordinates. Let U be a small neighborhood of η in $\hat{\Sigma}$, and V a neighborhood of an arc of the unit circle in the complex plane, and $\phi : V \rightarrow U$ a biholomorphic map that sends $V \cap \mathbb{D}$ to $U \cap \Sigma$, and $V \cap \mathbb{D}^c$ to $U \cap \Sigma^c$. Let us write $\zeta = \phi(z)$ and $\eta = \phi(w)$. Let $H(z) = h(\phi(z))$ and $J(z, w) = j(\phi(z), \phi(w))$.

By Proposition 3.2, since h is C^1 , we have

$$\begin{aligned} L(\eta, \eta) &= \lim_{z \rightarrow w} \frac{1}{J(z, w)} \frac{1}{1 - \langle H(z), H(w) \rangle} \\ &= \lim_{z \rightarrow w} \frac{w - z}{G(w, w)} \frac{1}{\langle H(w), H(w) \rangle - \langle H(z), H(w) \rangle} \\ &= \lim_{z \rightarrow w} \frac{w - z}{G(w, w)} \frac{1}{\langle H(w) - H(z), H(w) \rangle} \\ &= \frac{1}{G(w, w) \langle DH(w), H(w) \rangle} \end{aligned}$$

Therefore R_1 is a Toeplitz operator with a continuous non-vanishing symbol, and so is Fredholm by Proposition 3.6.

We claim that R_2 is Hilbert-Schmidt, hence compact, so R is a positive Fredholm operator. Indeed, R_2 is an integral operator, and will be Hilbert-Schmidt provided the function

$$M(\zeta, \eta) := j(\zeta, \eta) [L(\zeta, \eta) - L(\eta, \eta)]$$

is continuous on $\partial\Sigma \times \partial\Sigma$. The function M is clearly continuous when ζ stays away from η , so we just need to check the limit exists as $(\zeta, \eta) \in \partial\Sigma \times \partial\Sigma$ tends to the diagonal of $\partial\Sigma \times \partial\Sigma$.

Let us use local coordinates as before. Then we wish to calculate, for points z, w and u in the unit circle,

$$\begin{aligned} &\lim_{(z, w) \rightarrow (u, u)} \left[\frac{1}{1 - \langle H(z), H(w) \rangle} - \frac{J(z, w)}{G(w, w) \langle DH(w), H(w) \rangle} \right] \\ &= \lim_{(z, w) \rightarrow (u, u)} \left[\frac{1}{\langle H(w) - H(z), H(w) \rangle} - \frac{G(z, w)/G(w, w)}{(w - z) \langle DH(w), H(w) \rangle} \right] \end{aligned}$$

Using the assumption that h (and therefore H) is C^2 , we have that

$$H(z) = H(w) + (z - w)DH(w) + (z - w)^2\phi(z, w)$$

where ϕ is continuous. Moreover, we have

$$G(z, w)/G(w, w) = 1 + (z - w)\psi(w) + O(z - w)^2,$$

where

$$\psi(w) = \frac{1}{G(w, w)} \frac{\partial}{\partial z} G(z, w)|_{(w, w)}$$

is continuous.

So the above limit becomes

$$\begin{aligned} & \lim_{(z, w) \rightarrow (u, u)} \frac{1}{w - z} \left[\frac{1}{\langle DH(w) - (w - z)\phi(z, w), H(w) \rangle} \right. \\ & \quad \left. - \frac{1 + (w - z)\psi(w) + O(w - z)^2}{\langle DH(w), H(w) \rangle} \right] \\ &= \lim_{(z, w) \rightarrow (u, u)} \left[\frac{\langle \psi(w)DH(w) + \phi(z, w), H(w) \rangle}{\langle DH(u), H(u) \rangle^2} \right] \\ &= \left[\frac{\langle \psi(u)DH(u) + \phi(u, u), H(u) \rangle}{\langle DH(u), H(u) \rangle^2} \right]. \end{aligned}$$

So M , *a priori* defined only for $\zeta \neq \eta$, agrees with a continuous function on all of $\partial\Sigma \times \partial\Sigma$, as required.

Finally, suppose f is in the kernel of R . By expanding $\frac{1}{1 - \langle h(\zeta), h(\eta) \rangle}$ in powers of $h(\zeta)$ and using (4.6), we see that for all $\zeta \in \Sigma$,

$$\sum_{\gamma} \frac{|\gamma|!}{\gamma!} h(\zeta)^\gamma \langle f, h^\gamma \rangle = [\alpha\alpha^*f](\zeta) = 0.$$

Taking the inner product of the function $\alpha\alpha^*f$ with f we find

$$\langle \alpha\alpha^*f, f \rangle = \sum_{\gamma} \frac{|\gamma|!}{\gamma!} \langle f, h^\gamma \rangle \langle h^\gamma, f \rangle = 0,$$

therefore $\langle f, h^\gamma \rangle = 0$ for all γ . As polynomials in h span $H_h^2(\Sigma)$, this means $f \perp H_h^2(\Sigma)$. It follows that R is invertible, thus α is an isomorphism from \mathcal{F}_W onto $H_h^2(\Sigma)$, as required. \square

Example 4.7. Let (a, b) be a vector in C^2 of length one, with neither coordinate zero, and consider the map $h : \mathbb{D} \rightarrow \mathbb{B}_2$ given by $h(z) = (az^2, bz^3)$. This is a holomap that is transversal. The algebra H_h^∞ consists of all bounded holomorphic functions on the disk whose derivatives vanish at the origin.

Corollary 4.8. *Let W be as in Theorem 4.2. Then $\mathcal{A}_W = \mathcal{M}_W \cap C(\overline{W})$.*

PROOF: On the one hand, $A_h(\Sigma) = H_h^\infty(\Sigma) \cap C(\overline{\Sigma})$, and this algebra is mapped by composition with h isomorphically onto $\mathcal{M}_W \cap C(\overline{W})$ (as in Theorem 2.4). On the other hand, $A_h(\Sigma)$ is equal to the norm closure of $\{\text{polynomials}\} \circ h$ in H_h^∞ , and this algebra is mapped by composition with h isomorphically onto \mathcal{A}_W . The result follows. \square

Corollary 4.9. *Let Σ_1 and Σ_2 be finite Riemann surfaces. Then $H^\infty(\Sigma_1)$ and $H^\infty(\Sigma_2)$ are isomorphic if and only if Σ_1 and Σ_2 are biholomorphic.*

PROOF: Using Stout's result (mentioned in the proof of Proposition 3.6), and replacing each inner function B_i with $\frac{1}{\sqrt{3}}B_i$, one can embed any finite Riemann surface Σ in \mathbb{B}_3 with a map h . By the Schwarz reflection principle, each B_i continues analytically across $\partial\Sigma$, and maps some neighborhood U of $\overline{\Sigma}$ in $\hat{\Sigma}$ into \mathbb{C} in such a way that $B_i(U \setminus \overline{\Sigma}) \subset \mathbb{C} \setminus \overline{\mathbb{D}}$. Therefore B_i cannot have a vanishing derivative anywhere on $\partial\Sigma$.

Let $\zeta \in \partial\Sigma$, and let z be a local coordinate around ζ as in the proof of Theorem 4.2 so that $z(\zeta) = 1$. Since $|B_i(z)|^2$ is increasing as one crosses $\partial\Sigma$, we have $\frac{\partial}{\partial r}|B_i(z)|^2|_\zeta > 0$. But

$$\frac{\partial}{\partial r}|B_i(z)|^2 = \frac{z}{r}B_i'\overline{B_i} + \frac{\overline{z}}{r}B_i\overline{B_i'} = 2\Re\frac{z}{r}B_i'\overline{B_i}.$$

It follows that $\Re B_i'\overline{B_i} > 0$ for all i , therefore

$$\langle Dh(\zeta), h(\zeta) \rangle = \sum_{i=1}^3 B_i'(\zeta)\overline{B_i(\zeta)} \neq 0,$$

and h is transversal.

So we can assume that each of Σ_1 and Σ_2 are embedded transversally in \mathbb{B}_3 by maps that are C^∞ up to the boundary. By Theorem 4.2, the multiplier algebras \mathcal{M}_{Σ_1} and \mathcal{M}_{Σ_2} are isomorphic to $H^\infty(\Sigma_1)$ and $H^\infty(\Sigma_2)$, respectively. Therefore by Theorem 2.2, if $H^\infty(\Sigma_1)$ and $H^\infty(\Sigma_2)$ are isomorphic, then Σ_1 and Σ_2 are biholomorphic.

The converse is immediate. \square

We can now answer question (Q) in the affirmative for a wide class of one dimensional varieties when the biholomorphism is sufficiently regular.

Definition 4.10. *Let $g : W \rightarrow V$ be a holomorphic map that is C^1 up to ∂W and that maps ∂W onto ∂V . We say that g is transversal on*

∂W if for every $w \in \partial W$ and every analytic curve $\alpha : \mathbb{D} \rightarrow W$ which extends to a C^1 homeomorphism from $\mathbb{D} \cup \{1\}$ onto $\alpha(\mathbb{D}) \cup \{w\}$,

$$\langle D(g \circ \alpha)(1), g \circ \alpha(1) \rangle \neq 0.$$

Corollary 4.11. *Let Σ, W and h be as in Theorem 4.2. Let V be a variety in \mathbb{B}_d , and let $g : W \rightarrow V$ be biholomorphism that extends to be C^2 , one-to-one and transversal on ∂W . Then the mapping $\phi \mapsto \phi \circ g$ implements an isomorphism of \mathcal{M}_V onto \mathcal{M}_W .*

PROOF: By Theorem 4.2, \mathcal{M}_W is isomorphic to $H_h^\infty(\Sigma)$ via the composition map. Since g is transversal on ∂W , it follows that $g \circ h$ is transversal, so Theorem 4.2 implies that \mathcal{M}_V is isomorphic to $H_{g \circ h}^\infty(\Sigma)$ via the composition map. But $H_{g \circ h}^\infty(\Sigma) = H_h^\infty(\Sigma)$, therefore the mapping $\phi \mapsto \phi \circ (g \circ h) \circ h^{-1} = \phi \circ g$ is an isomorphism from \mathcal{M}_V to \mathcal{M}_W . \square

To conclude this section, we want to emphasize the following application to the extension problem of bounded holomorphic functions, which is a corollary of Theorem 4.2 (this type of result was also observed, in lesser generality, in [6, 7]).

Corollary 4.12. *Let W be as in Theorem 4.2. Then $\mathcal{M}_W = H^\infty(W)$, and every bounded analytic function on W extends to a multiplier in \mathcal{M}_d , and in particular to a bounded holomorphic function on the ball. Moreover, the extension operator is bounded.*

5. THE CHARACTER SPACE OF \mathcal{A}_V

In this and the next section, we assume that $d < \infty$. Denote by \mathcal{A}_d the norm closure of the polynomials in \mathcal{M}_d .

Lemma 5.1. *For every $\epsilon > 0$*

$$O((1 + \epsilon)\mathbb{B}_d) \subseteq \mathcal{A}_d.$$

Proof. Since $(M_{z_1}, \dots, M_{z_d})$ is a row contraction, this follows from the fact that the power series for every $f \in O((1 + \epsilon)\mathbb{B}_d)$ converges absolutely at every point in $(1 + \epsilon)\mathbb{B}_d$. \square

Recall the notation \mathcal{A}_V for the closure of the polynomials in \mathcal{M}_V .

Lemma 5.2. $\mathcal{A}_V \subseteq \mathcal{M}_V \cap C(\bar{V})$.

PROOF: Every $f \in \mathcal{A}_V$ is the norm limit of a sequence $\{p_n\}$ of polynomials. Since the operator norm dominates the sup norm, the sequence $\{p_n\}$ converges uniformly on V . It follows that the sequence $\{p_n\}$ converges uniformly on \bar{V} . Hence f can be extended continuously to all of \bar{V} . \square

We write $\Delta(\mathcal{A}_V)$ for the space of multiplicative linear functionals of the algebra \mathcal{A}_V . Since functions in \mathcal{A}_V are continuous on \overline{V} , it is clear that every $\lambda \in \overline{V}$ gives rise to an evaluation functional ρ_λ given by

$$\rho_\lambda(f) = f(\lambda) , f \in \mathcal{A}_V.$$

Theorem 5.3. *Assume that V is a variety in \mathbb{B}_d such that there exists a non-trivial ideal $I_V \subseteq \mathcal{A}_d$ for which*

$$\overline{V} = \{z \in \overline{\mathbb{B}_d} : \forall f \in I_V, f(z) = 0\}.$$

Then $\Delta(\mathcal{A}_V)$ can be identified homeomorphically with \overline{V} in the following way: to each $\rho \in \Delta(\mathcal{A}_V)$ there is some $\lambda \in \overline{V}$ such that $\rho = \rho_\lambda$, and every $\lambda \in \overline{V}$ gives rise to an evaluation functional.

Proof. Let $\lambda \in \overline{V}$. It is clear that λ gives rise to an evaluation character ρ_λ of $\mathcal{M}_V \cap C(\overline{V})$. By Lemma 5.2, ρ_λ restricts to a character on \mathcal{A}_V .

Conversely, let $\rho \in \Delta(\mathcal{A}_V)$. Let $\lambda = (\lambda_1, \dots, \lambda_d)$, where for all i we put $\lambda_i = \rho(z_i)$. As ρ is completely contractive, $\lambda \in \overline{\mathbb{B}_d}$. It is clear that the restriction of ρ to $\mathbb{C}[z]$ must be equal to the evaluation functional ρ_λ for this λ . We will show that ρ must agree with ρ_λ on \mathcal{A}_V , and that $\lambda \in \overline{V}$.

Since $(z_1|_V, \dots, z_d|_V)$ is a row contraction that generates \mathcal{A}_V , and since \mathcal{A}_d is the universal operator algebra generated by a row contraction [8], it follows that the restriction map is a homomorphism $q : \mathcal{A}_d \rightarrow \mathcal{A}_V$. Therefore ρ lifts to a character $\hat{\rho} = \rho \circ q$ of \mathcal{A}_d , which, as is well known, must be evaluation at the point λ .

To show that λ must be in \overline{V} , assume for contradiction that $\lambda \notin \overline{V}$. Then there exists some $f \in I_V$ such that $f|_{\overline{V}} \equiv 0$ while $f(\lambda) \neq 0$. But then we have

$$0 = \rho(f|_V) = \hat{\rho}(f) = f(\lambda) \neq 0.$$

It follows that λ has to be in \overline{V} and that $\rho = \rho_\lambda$.

Finally, the mapping $\overline{V} \ni \lambda \mapsto \rho_\lambda \in \Delta(\mathcal{A}_V)$ is bijective and clearly weak-* continuous, hence a homeomorphism. \square

Corollary 5.4. *Assume that V is a variety in \mathbb{B}_d such that there exists a non-trivial ideal $I_V \subseteq \mathcal{A}_d$ for which*

$$\overline{V} = \{z \in \overline{\mathbb{B}_d} : \forall f \in I_V, f(z) = 0\}.$$

If $\pi : \Delta(\mathcal{M}_V) \rightarrow \overline{\mathbb{B}_d}$ is the natural projection given by $\pi(\rho) = (\rho(z_1), \dots, \rho(z_d))$, then $\pi(\Delta(\mathcal{M}_V)) = \overline{V}$.

6. UNIONS OF STRONGLY DISJOINT VARIETIES

In this section we will consider varieties in \mathbb{B}_d which decompose in a very controllable way, to obtain some positive results towards question (Q).

Definition 6.1. *Let V_1, \dots, V_k be varieties in \mathbb{B}_d . We will say that V_1, \dots, V_k are strongly disjoint if for all $i = 1, \dots, k$, there exists a function $\phi_i \in \mathcal{M}_d$ such that $\phi_i \equiv 1$ on V_i and $\phi_i \equiv 0$ on V_j for $j \neq i$.*

An easy example of strongly disjoint varieties is given in the following lemma.

Lemma 6.2. *Suppose that $V = V_1 \cup \dots \cup V_k$ is such that for every i there is some $\epsilon > 0$ and a variety (in the usual sense) \tilde{V}_i in $(1 + \epsilon)\mathbb{B}_d$, such that $V_i = \tilde{V}_i \cap \mathbb{B}_d$. Assume further that $\tilde{V}_i \cap \tilde{V}_j = \emptyset$ for all $i \neq j$. Then V_1, \dots, V_k are strongly disjoint.*

Proof. Define ϕ_i by $\phi_i \equiv 1$ on \tilde{V}_i and $\phi_i \equiv 0$ on \tilde{V}_j for $j \neq i$. Then ϕ_i is holomorphic on \tilde{V} . By Cartan's extension Theorem [14, Theorem VIII.A.18] ϕ_i extends to a holomorphic function on $(1 + \epsilon)\mathbb{B}_d$. By Lemma 5.1, the restriction of ϕ_i to the unit ball is in $\mathcal{A}_d \subset \mathcal{M}_d$. \square

Here is a stronger result.

Lemma 6.3. *Let $V = V_1 \cup \dots \cup V_k$, where V_1, \dots, V_k are varieties in \mathbb{B}_d such that $\overline{V}_i \cap \overline{V}_j = \emptyset$ for all $i \neq j$. Assume further that there exist non-trivial, finitely generated ideals I_1, \dots, I_k in \mathcal{A}_d for which*

$$\overline{V}_i = \{z \in \overline{\mathbb{B}_d} : \forall f \in I_i, f(z) = 0\}.$$

Then V_1, \dots, V_k are strongly disjoint.

PROOF: If $k = 1$ there is nothing to prove, so let us consider $k = 2$ first. For each V_i , there are finitely many functions $\psi_{i,n}$ such that \overline{V}_i is the common zero set. If $\overline{V}_1 \cap \overline{V}_2$ is empty, then the functions $\psi_{i,n}$ can't all lie in a maximal ideal (as this would have to be in a fiber over some point in $\overline{V}_1 \cap \overline{V}_2$, by Theorem 5.3), so the algebra they generate is all of \mathcal{A}_d . Therefore one can find $g_{i,n} \in \mathcal{A}_d$ such that

$$1 = \sum_n g_{1,n} \psi_{1,n} + \sum_n g_{2,n} \psi_{2,n}.$$

Define $\phi_1 = \sum_n g_{2,n} \psi_{2,n}$ and $\phi_2 = \sum_n g_{1,n} \psi_{1,n}$. Since $\psi_{i,n}$ vanishes on V_i , the functions ϕ_1, ϕ_2 satisfy the definition of strong disjointness.

Suppose now that $k > 2$. By the case $k = 2$ which we just proved, for each $i \neq j$, there exist functions ψ_{ij} that are 1 on V_i and 0 on V_j . Then the functions $\phi_i = \prod_{j \neq i} \psi_{ij}$ satisfy the definition of strong disjointness. \square

Theorem 6.4. *Let $V = V_1 \cup \dots \cup V_k$, where V_1, \dots, V_k are strongly disjoint varieties in \mathbb{B}_d . Then $f \in \mathcal{M}_V$ if and only if for all $i = 1, \dots, k$, the restriction $f|_{V_i}$ is in \mathcal{M}_{V_i} .*

Proof. Since $\mathcal{M}_W = \mathcal{M}_d|_W$ for every variety W , the “only if” implication is obvious. Assume therefore that for all $i = 1, \dots, k$, $f|_{V_i} \in \mathcal{M}_{V_i}$. For all i , let $f_i \in \mathcal{M}_d$ satisfy $f_i|_{V_i} = f|_{V_i}$. Let $\phi_1, \dots, \phi_k \in \mathcal{A}_d$ be as in the definition of strong disjointness. Then $F = f_1\phi_1 + \dots + f_k\phi_k \in \mathcal{M}_d$, and the restriction of F to V equals f . Thus $f \in \mathcal{M}_V$. \square

Theorem 6.5. *Let $V = V_1 \cup \dots \cup V_k$ and $W = W_1 \cup \dots \cup W_k$ be two varieties such that the V_1, \dots, V_k and W_1, \dots, W_k are strongly disjoint. If \mathcal{M}_{V_i} is isomorphic to \mathcal{M}_{W_i} for all $i = 1, \dots, k$, then \mathcal{M}_V is isomorphic to \mathcal{M}_W .*

Proof. By Theorem 2.2, for all $i = 1, \dots, k$ there is a biholomorphism $H_i : W_i \rightarrow V_i$ with \mathcal{M}_d components, such that the isomorphism $\mathcal{M}_{V_i} \cong \mathcal{M}_{W_i}$ is implemented by $f \mapsto f \circ H_i$. Let h be defined on W by $h|_{W_i} = H_i$. Since for all $f \in \mathcal{M}_V$ we have that $f \circ h|_{W_i} = f|_{V_i} \circ H_i$, we conclude from Theorem 6.4 that $f \mapsto f \circ h$ implements an isomorphism of \mathcal{M}_V onto \mathcal{M}_W . \square

Remark 6.6. It now follows from Theorem 2.2 that the function h defined in the proof above can be extended to a function with components in \mathcal{M}_d .

REFERENCES

- [1] J. Agler and J.E. McCarthy. Complete Nevanlinna-Pick kernels. *J. Funct. Anal.*, 175(1):111–124, 2000.
- [2] J. Agler and J.E. McCarthy. Nevanlinna-Pick kernels and localization. In A. Gheondea, R.N. Gologan, and D. Timotin, editors, *Proceedings of 17th International Conference on Operator Theory at Timisoara, 1998*, pages 1–20. Theta Foundation, Bucharest, 2000.
- [3] J. Agler and J.E. McCarthy. *Pick Interpolation and Hilbert Function Spaces*. American Mathematical Society, Providence, 2002.
- [4] J. Agler and J.E. McCarthy. Hyperbolic algebraic and analytic curves. *Indiana Math. J.*, 56(6):2899–2933, 2007.
- [5] P. Ahern and D. Sarason. The H^p spaces of a class of function algebras. *Acta Math.*, 117:123–163, 1967.
- [6] D. Alpay, M. Putinar, and V. Vinnikov. A Hilbert space approach to bounded analytic extension in the ball. *Commun. Pure Appl. Anal.*, 2(2):139–145, 2003.
- [7] N. Arcozzi, R. Rochberg, and E. Sawyer. Carleson measures for the Drury-Arveson Hardy space and other Besov-Sobolev spaces on complex balls. *Adv. Math.*, 218(4):1107–1180, 2008.
- [8] W.B. Arveson. Subalgebras of C^* -algebras III: Multivariable operator theory. *Acta Math.*, 181:159–228, 1998.

- [9] L. Bers. On rings of analytic functions. *Bull. Amer. Math. Soc.*, 54:311–315, 1948.
- [10] K.R. Davidson, C. Ramsey, and O.M. Shalit. Operator algebras for analytic varieties. *Trans. Amer. Math. Soc.* To appear.
- [11] K.R. Davidson, C. Ramsey, and O.M. Shalit. The isomorphism problem for some universal operator algebras. *Adv. Math.*, 228:167–218, 2011.
- [12] S.W. Drury. Remarks on von Neumann’s inequality. In R.C. Blei and S.J. Sidney, editors, *Banach spaces, Harmonic analysis, and Probability theory*, volume 995 of *Lecture notes in Math.*, pages 14–32. Springer Verlag, Berlin, 1983.
- [13] J.D. Fay. *Theta functions on Riemann Surfaces*. Number 352 in LNMS. Springer, New York, 1973.
- [14] R. Gunning and H. Rossi. *Analytic functions of several complex variables*. Prentice-Hall, Englewood Cliffs, 1965.
- [15] R. Hartshorne. *Algebraic Geometry*. Springer, New York, 1977.
- [16] M. Hartz. Topological isomorphisms for some universal operator algebras. *J. Funct. Anal.*, 263(11):3564–3587, 2012.
- [17] V.I. Paulsen. *Completely bounded maps and operator algebras*. Cambridge University Press, Cambridge, 2002.
- [18] G. Popescu. Von Neumann inequality for $(B(\mathcal{H})^n)_1$. *Math. Scand.*, 68:292–304, 1991.
- [19] E.L. Stout. On some algebras of functions on finite open Riemann surfaces. *Math. Z.*, 92:366–379, 1966.
- [20] J. Wermer. Analytic disks in maximal ideal spaces. *Amer. J. Math.*, 86:161–170, 1964.

WASHINGTON UNIVERSITY, ST. LOUIS, MO 63130, USA
E-mail address: `mattkerr@math.wustl.edu`

WASHINGTON UNIVERSITY, ST. LOUIS, MO 63130, USA
E-mail address: `mccarthy@wustl.edu`

BEN-GURION UNIVERSITY OF THE NEGEV, BEER-SHEVA, ISRAEL
E-mail address: `oshalit@math.bgu.ac.il`