

QR CODE



Observed Risk Factors and the Prevalence of Potential Oral Malignant Lesions among a North Indian Population

VASUDA BHAGAT^{*1}, MEHTAB SINGH BHAGAT²

INTRODUCTION: The menace of oral cancer has become a public health concern and its prevalence is increasing with every passing day, making it the sixth most common cancer across the globe.

AIM: To evaluate the prevalence and risk factors for potential oral malignant lesions in a middle aged north Indian population.

MATERIALS AND METHODS: Adopting a Cross-sectional study design, this study examined a total of 944 participants in the Delhi-NCR region and males formed a majority of the population (569, 60.3%). Data was collected using a pre-tested and pre-validated questionnaire. Descriptive statistics was applied and the chi-squared test (for qualitative data) and independent samples t-test (for quantitative data) was applied to find out statistical differences (Using SPSS 21.0), if any.

RESULTS: It was reported that 201 (21.29%) of the study population suffered from oral malignant lesions, and the most prevalent lesion was OSMF [115, 12.18%] followed by leukoplakia [49, 5.19%]. The prevalence of oral malignant lesions among males was found to be higher (18.97%) as compared to females. A significant difference was found between the oral lesions and tobacco ever and never chewers ($p < 0.001$).

CONCLUSION: The results indicate a need to focus tobacco cessation programmes in the population of Delhi-NCR and reinforce it from time to time to reduce the burden of such Oral Malignant Lesions.

KEYWORDS: Leukoplakia, OSMF, Oral Malignant Lesions, Tobacco

A
B
S
T
R
A
C
T

INTRODUCTION

Although similar to human skin, the oral mucosa faces exposure to different kinds of complex environments due the habits and dietary patterns of an individual.¹ This exposure to various kinds of substances have the potential to harm the integrity of the oral mucosa which can cause a change in colour, shape, size and consistency leading to various disorders and/or Oral Mucosal Lesions (OMLs) in an individual.

Although a majority of OMLs are benign in nature and require no specialized treatment, a few lesions might be of concern as they may have certain pathological changes that can transform them into potentially malignant disorder.²

The various reasons for the occurrence of such lesions ranges from infections (bacterial, viral, fungal) to local trauma and/or irritation (e.g. traumatic keratosis, irritational fibroma, burns), along with systemic diseases or excessive consumption of tobacco, betel quid and also alcohol.³ Such lesions have the capacity to significantly impact one's quality of life and increase the burden of these diseases in a society/nation.⁴ Treatment of such lesions add an extra financial burden on the individual in addition to regular visits to the treatment centre wasting one's precious time and effort.

The use of tobacco products has been considered as a prime etiological factor for the formation of OMLs and its easy availability and cheap pricing leads to its increased consumption across the globe. Use of tobacco products among individuals can be due to a variety of reasons which includes use for recreational purposes, peer pressure, to relieve stress, etc. In Asians population, the cause for a steep rise of such OMLs is due to cigarette smoking, excess alcohol intake, and areca quid chewing.⁵ Various others factors documented by various researchers across of the globe are Body Mass Index and diabetes coupled with various dietary preferences (low vegetable inclusion, lesser fruit incorporation).⁶

The prevalence of oral mucosal diseases varies from 10.8%–81.3% in different populations⁷, with malignant tumors, representing only a minor percentage among these lesions.

Keeping in mind the harmful effects of various etiological factors in formation of OMLs and its capability to increase the burden of disease significantly in a society, the present study was undertaken with an aim to record and evaluate the Prevalence and risk factors for Potential oral malignant disorders in a



© Vasuda Bhagat et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY-NC 4.0, which permits unrestricted use, distribution and reproduction in any medium, provided the use is not commercial and the original author and source are cited.

middle aged North Indian population.

MATERIALS AND METHODS

This study adopted a cross-sectional designed and data was collected in the region of Delhi-NCR through a pre-tested and pre-validated proforma (adopted & modified from Kumar S et al.⁸) to record data of various patients visiting various oral health screening/ treatment camps. The camps were organized during the months of March 2018- February 2019 (12 months duration) after an ethical clearance was obtained from the institutional review board. It was kept in mind that the participants (aged >18 years) were duly explained about the study objectives and a written consent was obtained from them after they were assured of the confidentiality of the data obtained. Also they were informed that their data recorded could be used for research purposes, however, their individual identities shall remain masked.

The sample size was selected through the prevalence of OMLs obtained in the pilot study done to check for the feasibility of the study. It was observed that approximately 29.4% of the total screened population (125 in number) were having some kind of OML present in them. This prevalence rate was amenable to statistical analysis and using statistical deductions done by a certified statistician, the sample was found to be 783. However, we aimed to include >800 people in the study to account for loss of people/data.

The inclusion criteria was above 18 years as they could legally give consent on participation in the study. Exclusion criteria included participants who did not give their consent to be a part of the study. The part of the proforma first contained data regarding socio-demographic profile, socio economic status (based on Kuppaswamy scale), age and gender of the participants. A total of 944 participants were included in the study and the data obtained from them was analysed.

The second sub-section focused on the oral hygiene, the preventive measures taken to maintain it and recorded the adverse oral habits with its detailed noting of the quantified amount along with its quantity consumed. was registered. The third subsection emphasised primarily on the adverse oral habits present in the study population which majorly included data on tobacco chewing, areca nut use, and alcohol consumption. The participants were divided into ever chewers and never chewers with respect to smokeless tobacco.

The examinations were carried out by two observers who had two recoding clerks by their side and all of them were duly standardized prior to the study. Descriptive statistics was applied and the chi-squared test (for qualitative data) and independent samples t-test (for quantitative data) was applied to find out statistical differences (Using SPSS Version 21.0), if any among the study population.

RESULTS

Of the total 944 participants in the study, males formed a majority of the population (569, 60.3%) and a Greater portion belonged to the middle class (35.5%) or lower class (34.6%) divisional percentage based on their socioeconomic status. 70.6% of the population were using a toothbrush to clean their teeth. (Table 1)

It was documented that 21.29% of the Northern Indian Population which suffered from oral malignant lesions, with the prevalence rates of OSMF being 115 (12.18%) followed by leukoplakia in 5.19%, Lichen Planus in 2.54% and lowest being erythroplakia seen in 1.37 of the study subjects (Table 2).

CHARACTERISTICS	CATEGORIES	NUMBER (%)
GENDER	MALE	569 (60.3%)
	FEMALE	379 (39.7%)
SOCIO ECONOMIC POSITION	UPPER INCOME GROUP	376 (29.8%)
	MIDDLE INCOME GROUP	448 (35.5%)
	LOWER INCOME GROUP	436 (34.6%)
BRUSHING METHODOLOGY	TOOTHBRUSH	890 (70.6%)
	FINGER	156 (12.3%)
	OTHER DENTRIFICES	214 (16.9%)
BRUSHING FREQUENCY	ONCE A DAY	887 (70.3%)
	TWICE A DAY	235 (18.6%)
	THRICE AND MORE A DAY	138 (10.9%)

Table 1. Socio demographic and oral hygiene inclination and maintenance by the subjects.

PREVALENCE	NUMBER %
OSMF	115 12.18%
LEUKOPLAKIA	49 5.19%
LICHEN PLANUS	24 2.54%
ERYTHROPLAKIA	13 1.37%
TOTAL	201 21.29%

Table 2. Prevalence of oral probable malignant disorders in the northern Indian Population

The prevalence of oral malignant lesions among males was found to be higher(18.97%) as compared to their female counterparts (18.1.91%), but the difference was found to be non-significant. It was observed that in both groups, most the study population brushed once a day and this²⁰²

was found to be statistical significant. Tobacco consumption was another factor that was found to have a significant statistical significance ($p < 0.001$) among those found with and without oral mucosal lesions. Although the results were non significant, it was observed that almost all people used a toothbrush to brush their teeth and only 02(0.2%) people reported using other methods of cleaning their teeth (Table No 3).

FACTORS	CATEGORIES	PRESENCE OF OML (n,%)	ABSENCE OF OML (n,%)	p VALUE
GENDER	Male	183(18.97%)	386 (40.88%)	0.24
	Female	18 (1.91%)	361 (38.24%)	
		201	743	
SOCIOECONOMIC STATUS	Upperclass	44 (21.9%)	286 (38.5%)	0.375
	Middle class	212(30.8%)	212(28.5%)	
	Lower class	245(47.3%)	245(33.0%)	
CLEANING METHOD	Tooth brush	193(96.1%)	739(99.5%)	0.481
	Finger	07(3.48%)	03(0.40%)	
	Others	01(0.41%)	01(0.1%)	
BRUSHING FREQUENCY	Once	188(93.53%)	701 (94.3%)	0.01*
	Twice	13(6.47%)	40(5.38%)	
	Thrice	0(0%)	02(.03%)	
ALCOHOL CONSUMPTION	Ever consumer	147(71.3%)	312(42%)	0.456
	Never consumer	54(28.87%)	431(58%)	
TOBACCO	Ever consumer	121(60.2%)	216(27.05%)	<0.001*
	Never consumer	80(39.8%)	542(72.94%)	

Table 3. Interrelations of sociodemographic traits, oral hygiene maintenance and adverse oral Habits with oral probable malignant disorders

DISCUSSION

The present study, with an aim to record and evaluate the prevalence and risk factors for potential oral malignant disorders in a middle aged North Indian population found varying results with 21.29% of the total screened population presenting with some kind of potential oral malignant lesion. This is in agreement with Faraz SA et al (27.84%)⁸, higher as compared to Cebeci ARI et al.⁹ and lower as compared to Priya MK et

al. (42.4%)¹⁰ and Chung et al.¹¹ (12.7%). This variance could be attributed to factors including but not limited to genetic predisposition, geographic factors, consumption of tobacco, habits etc.

Most of the population screened comprised of males 569 (60.3%) and this result was supported by Kamble KA et al. [416 (70.8%)].¹² We postulate that females,

with their busy schedules and aiming to maintain their work-home balance find very little time to attend such camps, or go out of their homes/offices and get their check-up done.

The results of the present study that from the total prevalence of total oral malignant lesions was 21.29%, out of which the most prevalent lesion was OSMF [115, 12.18%], and this percentage is higher as compared Kamble KA et al (5.96%).¹² The second most lesion found to be prevalent was leukoplakia [49, 5.19%], and this percentage is in partial agreement to Mathew AL et al. (1.59%).¹³ Such variances in the prevalence of oral mucosal lesions is seen across the globe and it has been stated in the literature that males are at a higher risk of developing such potential oral malignant disorders as compared to their female counterparts (Nair et al.).¹⁴ This is attributed to the increased consumption of tobacco related products among males.

The study has certain limitations one of which is that the diagnosis of a person suffering from any potential oral malignant lesions was made on presumptive tests provided in textbooks and solely on the clinical manifestations examination. No Biopsy or confirmatory tests were conducted among the examined subjects. Another limitation is the societal pressure (i.e. social desirability bias) faced by the women as they unable to admit their adverse oral habit, leading to underreporting of data.

CONCLUSION

The results of the present study document a high prevalence of Oral Malignant disorders among the population of Delhi-NCR, thus warranting a need to provide tobacco cessation counselling from time to time so that the burden of such disorders is significantly reduced in the general population.

REFERENCES

1. Al-Gburi SM, Mudhir SH. The prevalence of the oral mucosal lesions among adult patients in Abu Ghraib city (Iraq), *J Res Med Dent Sci* 2018;6(5):145-8.
2. Toum SE, Cassia A, Bouchi N, Kassab I. Prevalence and Distribution of Oral Mucosal Lesions by Sex and Age Categories: A Retrospective Study of Patients Attending Lebanese School of Dentistry *Int J Dent*. 2018; 2018: 4030134.

3. Sridharan G. Epidemiology, control and prevention of tobacco induced oral mucosal lesions in India. *Indian J Cancer* 2014; 51:80-5.
4. Triantos D. Intra-oral findings and general health conditions among institutionalized and non-institutionalized elderly in Greece. *J Oral Pathol Med* 2005; 34: 577-82.
5. Hernandez BY, Zhu X, Goodman MT, Gatewood R, Mendiola P, Quinata K, et al. Betel nut chewing, oral premalignant lesions, and the oral microbiome. *PLoS One*. 2017; 12(2): e0172196.
6. Jain A, Sharma B. The Prevalence and Risk Factors for Potential Oral Malignant Disorders in a Middle Aged North Indian Population. *Int Healthc Res J*. 2019;3(3):123-7. <https://doi.org/10.26440/IHRJ/0303.06247>
7. Kansky AA, Didanovic V, Dovsak T, Brzak BL, Pelivan I, Terlevic D. Epidemiology of oral mucosal lesions in Slovenia. *Radiol Oncol* 2018; 52(3): 263-6.
8. Faraz SA, Huda I, Nabi AT, Fraz AA, Fatima A, Singh N. Prevalence of oral mucosal lesions/conditions in 35-44 years age group of adults of Patna city, Bihar, India. *J Dent Specialities*. 2019;7(1):23-7. <http://doi.org/10.18231/j.jds.2019.005>
9. Cebeci ARI, Gulsahi A, Kamburoglu K, Orhan BK, Oztas B. Prevalence and distribution of oral mucosal lesions in an adult Turkish population. *Med Oral Pathol Oral Buccal* 2009;14(6):E 272-7.
10. Krishna Priya M, Srinivas P, Devaki T. Evaluation of the prevalence of oral mucosal lesions in a population of eastern coast of South India. *J Int Soc Prevent Communit Dent* 2018;8:396-401.
11. Chung CH, Yang YH, Wang TY, Shieh TY, Warnakulasuriya S. Oral precancerous disorders associated with areca quid chewing, smoking, and alcohol drinking in southern Taiwan. *J Oral Pathol Med* 2005;34:460-6.
12. Kamble KA, Guddad SS, Nayak AG, Suragimath A, Sanade AR. Prevalence of Oral Mucosal Lesions in Western Maharashtra: A Prospective Study. *J Indian Acad Oral Med Radiol* 2017;29:282-7.
13. Mathew AL, Pai KM, Sholapurkar AA, Vengal M. The prevalence of oral mucosal lesions in patients visiting a dental school in Southern India. *Indian J Dent Res* 2008;19:99-103
14. Nair DR, Pruthy R, Pawar U, Chaturvedi P. Oral cancer: Premalignant conditions and screening - an update. *J Can Res Ther* 2012;8, Suppl S2:57-66.

Source of support: Nil, **Conflict of interest:** None declared

Cite this article as:

Bhagat V, Bhagat MS. Observed Risk Factors and the Prevalence of Potential Oral Malignant Lesions among a North Indian Population. *Int Healthc Res J.* 2019;3(6):201-205. <https://doi.org/10.26440/IHRJ/0306.09274>

AUTHOR AFFILIATIONS:

1. PG Student (Department of Oral Medicine and Radiology), DJ College of Dental Sciences and Research, Modinagar [(ORCID ID: <https://orcid.org/0000-0001-8344-4771>), (Corresponding Author)]
2. MBBS, Medical Officer, Government of Jammu and Kashmir, India

Contact corresponding author at: [vasudhabhagat1991\[at\]gmail\[dot\]com](mailto:vasudhabhagat1991[at]gmail[dot]com)