

Original Article

Proximate composition and antimicrobial effect of *Ocimum gratissimum* on broiler gut microfloraRuth T. S. Ofongo¹ and Elijah I. Ohimain*²¹Department of Animal Science, Faculty of Agricultural Technology, Niger Delta University, Wilberforce Island, Bayelsa State, Nigeria²Agricultural/Veterinary Microbiology Research Unit, Department of Biological Sciences, Faculty of Science Niger Delta University, Wilberforce Island, Bayelsa State, Nigeria.***Corresponding Author**

Elijah I. Ohimain

Agricultural/Veterinary Microbiology

Research Unit,

Department of Biological Sciences,

Faculty of Science Niger Delta University,

Wilberforce Island, Bayelsa State, Nigeria.

E-mail: ohimain@yahoo.com**Keywords:**

Botanicals,

E. coli,

Ethnoveterinary medicine,

Herb,

Poultry infections,

Lactobacillus,*Salmonella***Abstract**

The poultry industry is challenged by microbial infections, but the use of antibiotic growth promoters is restricted in many countries. This study is designed to assess the use of *Ocimum gratissimum* for the control of pathogenic gut microflora (*E. coli* and *Salmonella* sp). One hundred day old (ANAK 2000) commercial broilers were purchased and used for the experiment. The birds were brooded for seven days before random distribution to their respective treatments and replicates. Aqueous *Ocimum gratissimum* (0.11g/l) was administered to one set of 50 birds for 7 days (ocimum treatment), while the second sets of 50 birds which were not given the *Ocimum* extracts served as the control. Result show that the herb had a crude protein content of 4.56%, carbohydrate 13.40% and dry matter of 78.63%, but low in crude fibre (1.06%) and ash (1.21%). Prior to the administration of *Ocimum* extract the population of *Salmonella* was highest at the ileum (2.05 log cfu/g) followed by the crop (1.79 log cfu/g) and least at the caecum (1.77 log cfu/g). *E. coli* was highest at the ileum (2.05 log cfu/g), followed by the caecum (1.93 log cfu/g) and least at the crop (1.83 log cfu/g). *Lactobacillus* followed the pattern of *E. coli* but with different population, being highest at the ileum (1.95 log cfu/g) flowed by the caecum (1.90 log cfu/g) and least at the crop (1.79 log cfu/g). One week after the administration of *Ocimum* extracts to the chickens, the population of microbes in the gut decreased significantly ($P < 0.05$) in relation to the control, suggesting the efficacy of *Ocimum* at reducing the population of enteric bacteria.

1. Introduction

Chicken meat is consumed by most people irrespective of race, tribe and religion. But the poultry enterprise is challenged by infection and high cost of feeds. Due to the problem of antibiotic resistance and possible carry over effect on consumers, the use of antibiotic growth promoter (AGP) is being restricted in many countries. Hence, alternative feedstock and drugs are increasing being sought after. Recently, there is increased research interest and awareness on the ethnoveterinary medicines, which involved livestock diseases generally[1-4] and poultry in particular[5,6]. For instance, the leaves of Christmas bush, *Alchornea cordifolia* have been used to control infections in human[7,8] and for the replacement of soya bean meal in the diet of New Zealand white rabbits[9]. Timibitei et al[10] demonstrated the use of *Alchornea cordifolia* in the modulation of gonads, serum enzymes, testosterone, estrogen and blood corpuscles of rabbits.

One of the herb/spice that is increasingly being used is alternative medicine in Asia, south America and Africa, particularly in India[11,12] and Nigeria[13,14]. *Ocimum gratissimum* Linn. Literature appears to be in conflict concerning the family of the plant. While some reported that *Ocimum gratissimum* belong to the Lamiaceae[13-18], other authors classified the plant as belonging to Labiata family [11,12,19-21]. The plant is variously called basil, clove basil, sweet basil, teabush, scent leaf and fever plant [13]. The plant has many medicinal uses including the treatment of upper headaches, neuralgia, abdominal pains, convulsion, tooth ache, regulation of menstruation, cure for prolapsed rectum, rheumatism and paralysis, seminal weakness, asthma, anaemia, pile, diabetes, bleeding, strokes[13-16,22]. It has been well documented that *O. gratissimum* exhibit antimicrobial properties against many infectious diseases including

upper respiratory tract infections, cough and pneumonia, ear disease, conjunctivitis, tooth infection, skin disease (eczema, scabies, dermatitis), bronchitis, urogenital infections including sexually transmitted diseases, and enteric diseases (diarrhea, typhoid fever and dysentery [13,19,21]. The plant has been reported to have broad antimicrobial properties against bacteria and fungi including *Escherichia coli*, *Pseudomonas* species, *Klebsilla pneumonia*, *Shigella sonnei* and *Staphylococcus aureus* [12], *E. coli*, *Salmonella typhi*, *Shigella dysenteriae* and *Staphylococcus aureus* [20], *E. coli*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Klebsiella pneumonia*, *proteus mirabilis* and *Candida albicans* [15], *Listeria monocytogenes*[23,24]. *O. gratissimum* is also effective against several species of fungi including *Trichophyton rubrum*, *T. mentagraphytes*, *Penicillium islandi* and *Candida albicans*[24]. The plant has been considered for the management of HIV/AIDS infections [12]. A detailed review of the medicinal, ethopharmacological, chemical and microbial properties of *O. gratissimum* reported by Prabhu et al[11].

The antimicrobial and other medicinal properties of *O. gratissimum* is linked to the presence of phytochemicals such as essential oils, flavonoids, steroids, phenols, alkaloids, tannins, mucilages, gums, resins, glucosides and glycosides, terpenes, saponins and phytate [23,25]. Other authors linked the medicinal properties of *O. gratissimum* to the chemicals found in essential oil including eugenol, methyl eugenol, linalool, methyl cinnamate, Camphor, thymol, cis-and trans-ocimene, terpenes, germacrene, limonene, camphene, citral, thujene and myrene[14-18,22,26]. To the best of our knowledge, the use of *O. gratissimum* in the control of poultry infectious has not been documented. Hence, this study is focused on the demonstration of the use of crude extract of *O.*

gratissimum for the control of pathogenic enteric bacteria (*Salmonella* and *E. coli*) and assesses their effects on beneficial lactobacillus species in the gastrointestinal tract (GIT) of broilers.

2. Materials and Methods

2.1 Source of Experimental Birds and preparation of brood house

One hundred day old (ANAK 2000) commercial broilers were purchased at CHI farm, Ibadan, Nigeria and transported to Niger Delta University Teaching and Research Farm where the experiment was carried out. Vittalyte was administered to the birds due to stress resulting from transportation. The brooder house and its environment was cleaned with detergent and disinfectant (Z-germicide) two weeks prior to the arrival of the birds. Electric bulb (200Watts) was used at the brooding stage as a source of heat and light. The feeders and drinkers were properly washed prior to brooding. The birds were brooded for seven days before random distribution to their respective treatments and replicates. The birds were fed with a commercial broiler starter diet for two weeks.

2.2 Source of *Ocimum gratissimum* and its preparation methods

Ocimum gratissimum used in this experiment were obtained from Swali market in Yenagoa Local Government Area of Bayelsa State, Nigeria. The *Ocimum gratissimum* was detached from the stalk and washed with clean water. *Ocimum gratissimum* leaves were allowed to drop excess water before shredding it. Five hundred grams (500g) was blended with an electric blender using 1.5 liters of distilled water. The aqueous *Ocimum gratissimum* solution was filtered with cheese cloth, after adding 3.0 liters of distilled water to the solution. The solution divided into 5 equal portions before IT was administered to the birds (*Ocimum* treatment). The procedure was repeated in 3 days interval. Another set of fifty birds were not given the aqueous *Ocimum gratissimum* solution (Control). The administration of aqueous *Ocimum gratissimum* was done on day seven.

2.3 Digesta Collection

Two birds per replicate were slaughtered prior to administration of aqueous *Ocimum gratissimum* and digesta from the gastrointestinal tracts i.e. ileum, caecum and crop were collected into sterile container for microbial analysis. Digesta was collected again seven days after (day 14) administration of aqueous *Ocimum gratissimum* for a second set of microbial analysis.

2.4 Enumeration of microorganisms from the gastrointestinal tract of birds

The populations of microorganisms in the different samples were enumerated using serial dilution pour plate method of Pepper and Gerba [27](2005). About 1g of the sample was serially diluted in sterile distilled/deionized water and aliquots of the dilutions were aseptically plated into growth media; MRS Agar supplement with cycloheximide to enumerate total lactobacillus species. The medium were anaerobically incubated at 30°C for 7 days. For the isolation of *E. coli* EMB Agar were employed and it was incubated aerobically at 30°C for 24 hours. *Salmonella*-Shigella Agar was used to enumerate total *Salmonella* population. The medium was incubated aerobically at 30°C for 24 hours, however, presence of black colonies indicates salmonella species. After incubation, the colonies that grew on the medium were counted and expressed as colony forming units (cfu)/g of the samples.

2.5 Proximate composition of *Ocimum gratissimum*

The proximate parameters determined for the plant including protein, lipid, ash, fiber, moisture etc. the guide provided by AOAC (1977) was used for the analysis[28].

2.6 Statistical analysis

Log transformations were carried out on total bacteria count (log CFU) before subjecting the results obtained to general linear model analysis using SPSS version 16. Mean separation was carried out using Duncan Multiple Range test.

3. Results and Discussion

The proximate composition of *O. gratissimum* used in the experiment is presented in Table 1. The herb had a crude protein content of 4.56%, carbohydrate 13.40% and dry matter of 78.63%, but low in crude fibre (1.06%) and ash (1.21%). Several authors have analyzed the composition of *O. gratissimum* and have obtained similar results. Belewu et al[29] in 2009 reported the composition of *O. gratissimum* to be 93.33% dry matter, 21.37% of moisture, 20.78% crude protein, 11.75% fat, 14.99% crude fiber and 3.58% ash. Mlitan et al[30] in 2014 reported the proximate composition of *O. gratissimum* as 9.10 – 9.80% protein 10.40 – 10.60% fat 10.80 – 11.16% moisture, 13.1 – 14.50% ash and 50.35 – 55.20% carbohydrate. Aluko et al[31] in 2012 reported the proximate composition of *O. canum* leaves to be 12% ash, 17% crude fibre and 7% crude fat. Emeka and Chimaobi [32] in 2012 reported 4.28 – 5.56% ash, 6.21% fat, 4.659.68 – 11.30% fibre, 30.35 – 34.05% moisture, 5.02 – 6.77% protein and 78.22 – 87.23% carbohydrate. Fagbohun et al[33] in 2012 reported the proximate composition of *O. gratissimum* leaves as follows; 5.11 – 7.77% ash, 5.04 – 6.54% moisture, 14.6 – 19.30% crude protein, 6.80 – 7.57% fat, 9.61 – 12.66% crude fibre and 50.08 – 56.16% carbohydrate. Adewole [34] in 2014 presented the proximate composition of *O. gratissimum* as follows; 10.30% moisture, 2.45% ash, 2.18% fat, 16.51% protein, 9.07% crude fibre and 58.89% carbohydrate. Efiog [35] in 2014 reported 4.43% protein, 2.7% fat, 4.2% crude fibre and 1.15% ash, which is comparable to our findings in this study. Idris et al [36] in 2007 reported 82% moisture, 3.33% protein, 8.50% lipid, 9.52% fibre and 64.98% carbohydrate with calorific value of 343.08 kcal/100g. Oboh et al [37] in 2009 reported the proximate composition of *O. gratissimum* as follows; 81.35% moisture, 1.2% protein and 0.57% ash. In addition, Abdurahman et al [38] in 2012 reported the presence of several minerals in the plant including 0.62mg/g calcium, 0.21mg/g fluoride, 1.6mg/g chromium, 14.83mg/g iron, 0.47mg/g manganese and 2.14mg/g zinc. The differences in the proximate composition of *O. gratissimum* as reported by different authors may be due to the differences in the preparation and extraction of the plant components, which affected the moisture and ash contents.

Table 1 Proximate Composition of *Ocimum gratissimum*

Parameters	Proximate composition (g/100g)
Crude fibre	1.06
Dry matter content	78.63
Crude protein	4.56
Ash content	1.21
Moisture content	21.37
Carbohydrate	13.40
Percentage tannin	3.25

Prior to the administration of *Ocimum* extract the population diversity at the crop, ileum and caecum of the chicks were different (Table 2). The population of *Salmonella* was highest at the ileum (2.05 log cfu/g) followed by the crop (1.79 log cfu/g) and least at the caecum (1.77 log cfu/g). *E. coli* was highest at the ileum (2.05 log cfu/g), followed by the caecum (1.93 log cfu/g) and least at the crop (1.83 log cfu/g). *Lactobacillus* followed the pattern of *E. coli* but with different population, being highest at the ileum (1.95 log cfu/g) followed by the caecum (1.90 log cfu/g) and least at the crop (1.79 log cfu/g). In a previous study Ohimain and Ofongo[39] in 2013 reported different populations of microbes in the crop, ileum and caecum of broilers. The population of *E. coli* was 7.05 log cfu/g at the crop, 7.02 log cfu/g at the ileum and 6.98 log cfu/g at the caecum. *Lactobacillus* has 6.63 log cfu/g at the crop, 6.66 log cfu/g at the ileum and 6.73 log cfu/g at the caecum.

Table 2: Microbial population from the gastrointestinal tracts of the birds before administration of aqueous *Ocimum gratissimum*

Bacterial Species	<i>Salmonella</i>	<i>Escherichia coli</i>	<i>Lactobacillus</i>
CROP	1.79b	1.83a	1.79a
ILEUM	2.05c	2.05c	1.95c
CAECUM	1.77a	1.93b	1.90b

Along the columns, means with different alphabets are significantly different according to the Duncan statistics ($p < 0.05$)

Generally, the population of microbes in the GIT increases with age until a stable population is reached. One week after the administration of *Ocimum* extracts to the chickens, the population of microbes in the GIT decreased significantly ($P < 0.05$) in relation to the control (Table 3), suggesting the efficacy of *Ocimum* at reducing the population of enteric bacteria. Lower population of *Salmonella*, *E. coli* and *Lactobacillus* was recorded at the crop, ileum and caecum

($P < 0.05$). The result therefore shows that the antibiotic property of *O. gratissimum* does not discriminate between pathogenic (*Salmonella* and *E. coli*) and beneficial microbes (*Lactobacillus*) hence its use has to be applied with caution. Many authors have reported the efficacy of *Ocimum* against *Salmonella* [22,40-42], *lactobacillus* [43,44] and *E. coli* infections [45-51].

Table 3: Microbial population in the gastrointestinal tracts of broilers after administration of aqueous *Ocimum gratissimum* extracts

Bacterial Species	<i>Ocimum</i> Treatment, Log cfu/g	Control, Log cfu/g	SEM	P.VALUE
CROP				
<i>Salmonella</i>	1.85 ^b	2.20 ^a	0.106	0.016**
<i>Escherichia coli</i>	1.93 ^b	2.06 ^a	0.156	0.050*
<i>Lactobacillus</i>	1.95 ^b	2.14 ^a	0.075	0.030**
ILEUM				
<i>Salmonella</i>	2.04 ^b	2.27 ^a	0.113	0.041**
<i>Escherichia coli</i>	1.93 ^b	2.09 ^a	0.120	0.000***
<i>Lactobacillus</i>	1.88 ^b	2.08 ^a	0.096	0.010***
CEACUM				
<i>Salmonella</i>	1.82 ^b	1.95 ^a	0.139	0.030**
<i>Escherichia coli</i>	1.97 ^b	2.17 ^a	0.122	0.000***
<i>Lactobacillus</i>	1.67 ^b	1.95 ^a	0.125	0.020***

Along the rows, means with different alphabets are significantly different according to the Duncan statistics ($p < 0.05$)

4. Conclusion

This study tested the effects of *Ocimum gratissimum* on the gut microflora of broiler chickens. The administration of aqueous extracts of *Ocimum gratissimum* WAS found to be able to control the population of *E. coli* and salmonella, but also negatively affected beneficial microbes like lactobacillus. Hence, we conclude by recommending the use of the herb for the control of microbial infections caused by detrimental gut microbes, while enhancing the probiotic effects of lactobacillus.

Acknowledgement

This publication was based on the undergraduate research work carried out by Charles A.A.R. Itekesi at the Niger Delta University under the supervision of the authors. The authors also wish to thank Sylvester C. Izah for the editorial works.

References

- [1] Bhardwaj AK, Lone PA, Dar MM, Parray JA, Shan KW. Ethnoveterinary medicinal uses of plants of district Bandipora of Jammu and Kashmir, India. *International Journal of Traditional and Natural Medicines* 2013; 2(3): 164 – 178.
- [2] Njoroge GN, Bussmann RW, Herbal usage and informant consensus in ethnoveterinary management of cattle diseases among the Kikuyus (Central Kenya). *Journal of Ethnopharmacology* 2007; 108(3):332-9.
- [3] Maroyi A. Use of traditional veterinary medicine in Nhema communal area of the midlands province, Zimbabwe. *Afr. J. Tradit. Complement. Altern. Med.*, 2012; 9(3): 315 – 322.
- [4] Shicai S, Andreas W, Vernooy R. The importance of ethnoveterinary treatments for pig illness in poor, ethnic minority communities: a case study of Nu people in Yunnan, China. *Intern. J. Appl. Res. Vet. Med.*, 2010; 8(1): 53 – 59.
- [5] Moreki JC, Poroga B, Dikeme R, Seabo D. Ethnoveterinary medicine and health management in poultry in southern and western districts, Botswana. *Livestock Research for Rural Development*, 2010; 22(22).
- [6] Matekaire T, Bwakura TM. Ethnoveterinary medicine: a potential alternative to orthodox animal health delivery in Zimbabwe. *Intern. J. Appl. Res. Vet. Med.*, 2004; 2(4): 269 – 273.
- [7] Kigigha L.T. and Atuzie M.N. Assessment of traditional medicinal application of *Alchornea cordifolia*. *African Journal of Biotechnology*, 2012; 11(8): 2083 – 2086.
- [8] George NJ, Obot IB, Ikot AN, Akpan AE, Obi-Egbedi NO. Phytochemical and antitmicrobial properties of leaves of

Alchornea cordifolia. *E-Journal of Chemistry*, 2010; 7(3): 1071 – 1079.

- [9] Alikwe, P.C.N., Ohimain, E. I., Kester, A. E. Performance evaluation of New Zealand white rabbits fed *Alchornea cordifolia* leaf meal as replacement for soya bean meal. *American Journal of Agriculture and Forestry*. 2014; 2 (2): 51-54
- [10] Timibitei, K. O., Alikwe, P. C. N., Ohimain, E. I. and Wekhe, S.N. Evaluating the impact of *Alchornea cordifolia* (christmas bush) root bark, seeds and pod husks on the gonads, serum level of testosterone, estrogen, serum enzymes and blood corpuscles of rabbits. *International Journal of Current Research in Life Sciences*. 2014; 3 (5): 46-50.
- [11] Prabhu KS, Lobo R, Shirwaik AA, Shirwaikar A. *Ocimum gratissimum*: a review of its chemical, pharmacological and ethnomedicinal properties. *The Open Complementary Medicine Journal*, 2009; 1: 1 – 15.
- [12] Rajalakshmi G, Komathi S, Savetha S. Antibacterial potential of *Ocimum gratissimum* leaf extracts. *Int. J. Pharm. Sci. Rev. Res.*, 2013; 19(2): 70 – 71.
- [13] Mann A. Phytochemical constituents and antimicrobial and grain protectant activities of clove basil (*Ocimum gratissimum* L.) grown in Nigeria. *International Journal of Plant Research* 2012; 2(1): 51 – 58.
- [14] Matasyoh LG, Matasyoh JC, Wachira FN, Kinyua MG, Muigai AWT, Mukiyama TK. Variation in the antimicrobial activity of essential oils of *Ocimum gratissimum* L. from different populations of Kenya. *African Crop Science Conference Proceedings*, 2007; 8: 1745 – 1750.
- [15] Matasyoh LG, Matasyoh JC, Wachira FN, Kinyua MG, Muigai AWT, Mukiyama TK. Antimicrobial activity of essential oils of *Ocimum gratissimum* L. from different populations of Kenya. *Afr. J. Trad. CAM*, 2008; 5(2): 187 – 193.
- [16] Verma RS, Bisht PS, Padalia RC, Saikia D, Chauhan A. Chemical composition and antibacterial activity of essential oil from two *Ocimum* spp grown in sub-tropical India during spring-summer cropping season. *Journal of Traditional Medicines*, 2011; 6(5): 211 – 217.
- [17] Sastry KP, Kumar RR, Kumar AN, Sneha G, Elizabeth M. Morpho-chemical description and antimicrobial activity of different *Ocimum* species. *J. Plant Develop.*, 2012; 19: 53 – 64.
- [18] Benitez NP, Leon EMM, Stashenko EE. Eugenol and methyl eugenol chemotypes of essential oil of species *Ocimum gratissimum* L. and *Ocimum campechianum* mill. From Colombia. *Journal of Chromatographic Science*, 2009; 47: 800 – 803.

- [19] Adebolu T.T, Oladimeji SA. Antimicrobial activity of leaf extracts of *Ocimum gratissimum* on selected diarrhea causing bacteria in southwestern Nigeria. *African Journal of Biotechnology*, 2005; 4(7): 682 – 684.
- [20] Mabekoje OO, Bello OO, Egberongbe HO. Antimicrobial efficacy of *Ocimum gratissimum* and *Vernonia amygdalina* on gastrointestinal bacteria. *Canadian Journal of Pure and Applied Sciences*, 2013; 7(2): 2341 – 2345.
- [21] Nakamura CV, Ueda-Nakamura T, Bando E, Melo AFN, Cortez DAG, Filho BPD. Antibacterial activity of *Ocimum gratissimum* L. essential oil. Mem. Inst. Oswaldo Cruz, Rio de Janeiro 1999; 94(5): 675 – 678.
- [22] Matasyoh LG, Matasyoh JC, Wachira FN, Kinyua MG, Muigai AWT, Mukiyama TK. Chemical composition and antimicrobial activity of the essential oil of *Ocimum gratissimum* L. growing in Eastern Kenya. *African Journal of Botany*, 2013; 1(4): 050 – 054.
- [23] Mbata T, Saikia A. Antibacterial activity and phytochemical screening of crude ethanolic extracts of leaves of *Ocimum gratissimum* L on *Listeria monocytogenes*. *The Internet Journal of Microbiology*, 2007; 4(2).
- [24] Mbata T, Saikia A. Antibacterial activity of essential oil from *Ocimum gratissimum* on *Listeria monocytogenes*. *Internal Journal of Food Safety*, 2007; 5(7): 15-19.
- [25] Ishiwu CN, Umenwanne CP, Obiegbuna JE, Uchegbu NN. *In vitro* assessment of anti bacterial effect of extracts of *Ocimum gratissimum* and *Carica papaya* leaves. *International Journal of Applied Science and Technology*, 2014; 4(1): 171-177.
- [26] Kpoviessi BGHK, Ladekan EY, Kpoviessi DSS, Gbaguidi F, Yehouenou B, Quetin-Leclercq J, Figueredo G, Moudachirou M, Accrombessi GC. Chemical variation of essential oil constituents of *Ocimum gratissimum* L. from Benin, and impact on antimicrobial properties and toxicity against *Artemia salina* leach. *Chemistry and Biodiversity*, 2012; 9: 139 – 149.
- [27] Association of Official Analytical Chemist (AOAC) (1977). Analytical food analysis. British England. 332 pp.
- [28] Pepper, I.L. and Gerba, C.P. (2005). Environmental microbiology. A laboratory manual. Second edition. Elsevier academic press.
- [29] Belewu MA, Olatunde OA, Giwa TA. Underutilized medicinal plants and spices: chemical composition and phytochemical properties. *Journal of Medicinal Plants Research*, 2009; 3(12): 1099 – 1103.
- [30] Mlitan AM, Sasi MS, Alkherraz AM. 2014. Proximate and minor mineral content in some selected basil leaves of *Ocimum gratissimum* L, in Libya. *International Journal of Chemical Engineering and Applications*, 5(6): 502 – 505.
- [31] Aluko BT, Oloyede OI, Afolayan AJ. Phytochemical and nutrient compositions of the leaves of *Ocimum canus* Sims. *African Journal of Biotechnology*, 2012; 11(63): 12697 – 12701.
- [32] Emeka NG, Chimaobi A. Chemical composition and variability among some *ocimum gratissimum* accessions. *Int. J. Med. Arom. Plants*, 2012; 2(3): 460 – 467.
- [33] Fagbohun ED, Lawal OU, Ore ME. The proximate, mineral and phytochemical analysis of the leave of *Ocimum gratissimum* L., *Melanthera scandens* A. and *Leea guineensis* L. and their medicinal value. *International Journal of Applied Biology and Pharmaceutical Technology*, 2012; 3(1): 5- 21.
- [34] Adewole E. Proximate and phytochemical constituents of *Ocimum gratissimum*. *Journal of Physical and Chemical Sciences*, 2014; 1(1): 1 – 3.
- [35] Efiang EE. Phytochemical, proximate, vitamins and mineral composition of *Ocimum gratissimum* leaves. *Journal of Physical and Chemical Sciences*, 2014; 1(4): 1-5.
- [36] Idris S, Iyaka YA, Ndamitso MM, Paiko YB. Nutritional composition of the leaves and stems of *Ocimum gratissimum*. *Journal of Emerging Trends in Engineering and Applied Sciences*, 2011; 2(5): 801 – 805.
- [37] Oboh FOJ, Masodje HI, Enabulele SA. Nutritional and antimicrobial properties of *Ocimum gratissimum* leaves. *Journal of Biological Sciences*, 2009; 9(4): 377 – 380.
- [38] Abdurahman FI, Tijjani MA, Osuji UO. 2012. Proximate content and chemical composition of *Ocimum viridis* leaf and *Ocimum gratissimum* leaf. *International Research Journal of Pharmacy*, 2012; 3(4): 153 – 156.
- [39] Ohimain, E. I. and Ofongo, R. T. S. Effect of enzyme supplemented diet on gut microflora, digesta pH and performance of broiler chickens. *Journal of Microbiology, Biotechnology & Food Science*. 2013; 3 (2): 127 – 131
- [40] Alo MN, Anyim C, Igwe JC, Elom M, Uchenna DS. Antibacterial activity of water, ethanol and methanol extracts of *Ocimum gratissimum*, *Vernonia amygdalina* and *Aframomum melegueta*. *Advances In Applied Science Research*, 2012; 3(2): 844 – 848.
- [41] Olamide SO, Agu GC. The assessment of the antimicrobial activities of *Ocimum gratissimum* (wild basil) and *Vernonia amygdalina* (bitter leaf) on some enteric pathogen causing dysentery or diarrhea in patients. *The International Journal of Engineering and Science*, 2013; 2(9): 83 – 96.
- [42] Bankole HA, Anjorin AA, Kazeem MI, Ogbeche ME, Agbafor U. Antibacterial activity of *Ocimum gratissimum* and *Gongronrma latifolium* on *Staphylococcus aureus* and *Salmonella typhi*. *Ast African Journal of Science and Technology*, 2012; 2(1): 114-128.
- [43] Ijeh I.I, Omodamiro OD, Nwanna IJ. Antimicrobial effects of aqueous and ethanolic fractions of two species, *Ocimum gratissimum* and *Xylopiya aethiopicia*. *African Journal of Botechnology*, 2005; 4(9): 953-956.
- [44] Olusola OO, Oyadeyi OS, Omojola AB, Olugbemi TS. 2014. Antimicrobial activity of *Ocimum gratissimum* extract on suya (an intermediate moisture meat in Nigeria). *African Journal of Food, Agriculture, Nutrition and Development*, 2014; 14(6): 9361-9375.
- [45] Mbajiuka CS, Obeagu EI. The antibacterial activity of leaf extracts of *Ocimum gratissimum* and *sida acuta*. *International Journal of Microbiological Research*, 2014; 5(2): 124 – 129.
- [46] Ladipo MK, Doherty VF, Kanife UC. Phytochemical screening and antibacterial investigation of the extract of *Ocimum gratissimum* (scent leaf) on selected Enterobacteriaceae. *Publication, Agriculture and Technology*, 2010; 6(2): 75 – 84.
- [47] Ighodaro O, Macdonald AS, Oludare AO. Phytotoxic and antimicrobial activities of flavonoids in *Ocimum gratissimum*. *Life Science Journal*, 2010; 7(3): 45- 48.
- [48] Omodamiro OD, Jimoh MA. Antioxidant and antibacterial activities of *Ocimum gratissimum*. *American Journal of Phytomedicine and Clinical Therapeutics*, 2015; 3(1): 010 – 019.
- [49] Nwinyi OC, Chinedu NS, Ajani OO, Ikpo CO, Ogunniran KO. Antibacterial effects of extracts of *Ocimum gratissimum* and *Piper guineense* on *Escherichia coli* and *Staphylococcus aureus*. *African Journal of Food Science*, 2009; 3(1): 22 – 25.
- [50] Nweze EI, Eze EE. Justification for the use of *Ocimum gratissimum* L. in herbal medicine and its interaction with disc antibiotics, *BMC Complementary and Alternative Medicine*, 2009; 9:37; doi:10.1186/1472-6882-9-37.
- [51] Lawal IO, Borokini TI, Oyeleye A, Willams OA, Olayemi JO. Evaluation of extract of *ficus exasperata* vahl root bark for antimicrobial activities against some strains of clinical isolates of bacteria and fungi. *International Journal of Mordern Botany*, 2012; 2(1): 6-12.