



---

Articles by College of Natural and Applied Sciences Faculty

---

9-1-1988

## **Depression of aerobic metabolism and intracellular pH by hypercapnia in land snails, *Otala lactea***

M. Christopher Barnhart

Brian R. McMahon

Follow this and additional works at: <https://bearworks.missouristate.edu/articles-cnas>

---

### **Recommended Citation**

Barnhart, M. Christopher, and Brian R. McMahon. "Depression of aerobic metabolism and intracellular pH by hypercapnia in land snails, *Otala lactea*." *Journal of Experimental Biology* 138, no. 1 (1988): 289-299.

This article or document was made available through BearWorks, the institutional repository of Missouri State University. The work contained in it may be protected by copyright and require permission of the copyright holder for reuse or redistribution.

For more information, please contact [BearWorks@library.missouristate.edu](mailto: BearWorks@library.missouristate.edu).

## DEPRESSION OF AEROBIC METABOLISM AND INTRACELLULAR pH BY HYPERCAPNIA IN LAND SNAILS, *OTALA LACTEA*

BY M. CHRISTOPHER BARNHART\* AND BRIAN R. McMAHON  
*Department of Biology, University of Calgary, Calgary, Alberta T2N 1N4,  
Canada*

*Accepted 7 April 1988*

### Summary

The pulmonate land snail *Otala lactea* undergoes simultaneous hypercapnia, hypoxia, extracellular acidosis and metabolic depression during dormancy. We tested the effects of ambient hypercapnia and hypoxia on oxygen consumption ( $\dot{V}_{O_2}$ ) and on extracellular and intracellular pH of active (i.e. non-dormant) individuals. Active snails reduced  $\dot{V}_{O_2}$  by 50% within 1 h when exposed to 65 mmHg (1 mmHg = 133.3 Pa) ambient  $P_{CO_2}$ , and by 63% in 98 mmHg. These levels of  $CO_2$  are within the range that occurs naturally in the lung and blood during dormancy.  $\dot{V}_{O_2}$  of hypercapnic snails remained below that of controls for the duration of exposure (up to 9 h) and returned to control levels within 1 h when  $CO_2$  was removed. Both pHe and whole-body pH<sub>i</sub> (measured using [<sup>14</sup>C]DMO) fell with increasing haemolymph  $P_{CO_2}$  by approximately  $0.7 \log P_{CO_2}$ . Critical ( $\dot{V}_{O_2}$ -limiting) ambient  $P_{O_2}$  of active snails was 90 mmHg in the absence of  $CO_2$  and dropped to 50 mmHg when  $\dot{V}_{O_2}$  was reduced 45% by exposure to  $CO_2$ . Estimated critical  $P_{O_2}$  at the lower  $\dot{V}_{O_2}$  typical of dormancy is well below the typical lung  $P_{O_2}$  of dormant *Otala*, suggesting that  $P_{O_2}$  in the lung does not normally limit oxygen consumption during dormancy. These results support the hypothesis that hypercapnia or resulting respiratory acidosis depresses metabolic rate during dormancy, and argue against a limitation of  $\dot{V}_{O_2}$  by hypoxia.

### Introduction

Studies of the physiology and biochemistry of metabolic arrest suggest that hypoxia and changes of intracellular pH are possible factors governing reversible transitions between activity and dormancy in many animals (Hochachka, 1988; Busa & Nuccitelli, 1984; Hochachka & Guppy, 1987). Pulmonate land snails provide an interesting opportunity to investigate both these mechanisms. Dormancy in *Otala lactea* (Helicidae) is characterized by reduced aerobic metabolism

\* Present address: Physiological Research Laboratory, Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA 92093, USA.

Key words: Pulmonata, Helicidae, dormancy, metabolic arrest, intracellular pH, hypercapnia, hypoxia.

and by simultaneous hypoxia, hypercapnia and respiratory acidosis brought about by hypoventilation of the lung. Mean  $\dot{V}_{O_2}$  declines to about 15 % of that of resting, non-dormant individuals (Barnhart & McMahon, 1987).  $P_{O_2}$  in the lung often falls below 10 mmHg, compared with 137 mmHg in active snails, whereas  $P_{CO_2}$  reaches 100 mmHg in dormancy, compared with 12 mmHg in active snails. Mean haemolymph pH falls 0.4 units as a result of hypercapnia and up to 0.5 units further during prolonged dormancy as  $[HCO_3^-]$  decreases (Barnhart, 1986a,b).

Cycles of  $CO_2$  accumulation and release coincide with fluctuations of  $\dot{V}_{O_2}$  in dormant *Otala*, and  $\dot{V}_{O_2}$  is inversely related to whole-body  $CO_2$  content (Barnhart & McMahon, 1987). Both the broad correlation between hypercapnia and dormancy and the short-term correlation between body  $CO_2$  content and  $\dot{V}_{O_2}$  suggest that  $CO_2$  accumulation and/or resultant acidosis might be factors in the control of metabolic rate during dormancy. However, low  $P_{O_2}$  values in the lung and haemolymph suggest that  $\dot{V}_{O_2}$  might be oxygen-limited during dormancy. To test these possibilities, the present study examines the effects of acute experimental hypercapnia and hypoxia on the oxygen consumption of active *Otala*. The effect of hypercapnia on whole-body intracellular pH (pHi) was also measured because intracellular acidification is a possible mechanism by which  $CO_2$  could affect metabolism.

## Materials and methods

### *Subjects*

Adult *Otala* were collected at Playa del Rey in Los Angeles County, California, and maintained in the laboratory as described previously (Barnhart & McMahon, 1987). The individuals used had a mean whole mass of  $8.44 \pm 0.84$  g (S.D.).

### *Respirometry*

$O_2$  uptake and  $CO_2$  release of individuals were measured over successive 1-h intervals by closed-chamber respirometry, using 60 ml syringes as chambers (Barnhart, 1986b). For measurements in the absence of  $CO_2$ , an open vial holding a folded wick and 1 ml of 5 % KOH was placed in the chamber to absorb respiratory  $CO_2$ . For measurements in the presence of  $CO_2$  the KOH was replaced with distilled water. Temperature was 20–23°C.

After a brief period (2–7 days) of dormancy, snails were placed on wet paper towelling for 2 h to induce arousal and permit hydration, and then placed in the respirometry chambers for 1 h of acclimation. To measure gas exchange, the chambers were flushed with mixtures of air and  $CO_2$  or  $N_2$  prepared by Wösthoff gas-mixing pumps, closed for 1 h, and then placed in an infusion pump to expel the gas successively through a desiccant tube, an Applied Electrochemistry S3A electrochemical  $O_2$  analyser and a CD3A infrared  $CO_2$  analyser. Oxygen and  $CO_2$  fractions were measured to the nearest 0.01 % volume. The analysers were calibrated using nitrogen, dry  $CO_2$ -free air and a 5 %  $CO_2$  standard analysed with

a Scholander device. Change in oxygen fraction in the respirometers was typically 1–2 % of volume. Gas exchange was calculated according to Vleck (1987).

#### *Effects of $P_{CO_2}$ and $P_{O_2}$ on $\dot{V}_{O_2}$*

The effect of ambient  $P_{CO_2}$  on  $\dot{V}_{O_2}$  was examined in four experiments which ranged in duration from 5 to 14 h. In each experiment, 24 individuals were distributed randomly among a control (zero  $CO_2$ ) and several  $CO_2$  treatment groups.  $\dot{V}_{O_2}$  of all individuals was first measured in the absence of  $CO_2$ . The treatment groups were subsequently exposed to ambient  $P_{CO_2}$  between 13 and 98 mmHg.

The relationship between  $P_{O_2}$  and  $\dot{V}_{O_2}$  was measured in two experiments, one in the absence of  $CO_2$  and the other at 66 mmHg  $P_{CO_2}$ .  $\dot{V}_{O_2}$  of each individual was determined initially in air and afterwards at one of six levels of  $P_{O_2}$  between 14 and 160 mmHg.

#### *Effects of hypercapnia on pH*

The effects of hypercapnia on pH of venous haemolymph (pHe) and of intracellular fluids (pHi) were measured in individuals which had previously been catheterized to permit marker injection and serial sampling of haemolymph. pHi was estimated with the labelled weak acid distribution technique (Waddell & Butler, 1959), using 5,5-dimethyl[ $^{14}C$ ]oxazolidine-2,4-dione (DMO) as the pH indicator and polyethylene glycol[1,2- $^3H$ ] (PEG) as the extracellular fluid volume marker (isotopes from New England Nuclear). Catheters consisted of 21 gauge hypodermic infusion needles with attached tubing (Deseret Company). The needle was lightly coated with cyanoacrylate glue just behind the tip, then inserted through a hole in the shell into the visceral sinus. If no leakage occurred around the insertion, the needle was sealed to the shell and supported with epoxy paste. A 1 cm length of tubing was left attached to the needle, and was reversibly closed by a crimp secured by a band of larger tubing. Animals were placed on moist paper towelling and used within 1–2 days of preparation.

The catheterized snails were placed in plastic bags for exposure to gas mixtures. The bags were flushed continuously with humidified mixtures of  $CO_2$  and air to maintain 0, 46 or 98 mmHg  $P_{CO_2}$ . After waiting 3 h to permit equilibration with ambient  $CO_2$ , each snail was injected with 0.15  $\mu Ci$  of DMO and 0.20  $\mu Ci$  of PEG in 30  $\mu l$  of 200 mmol  $l^{-1}$  NaCl, followed by 100  $\mu l$  of snail haemolymph to flush the catheter. Haemolymph samples (about 60  $\mu l$  each) were collected at hourly intervals thereafter for 6 h. The catheter dead-space volume was discarded. The next 40  $\mu l$  was transferred anaerobically to a Radiometer capillary pH electrode for determination of pHe. Subsamples (10  $\mu l$ ) were analysed for isotope activities (see below).

Haemolymph  $P_{CO_2}$  was estimated from pH using the *in vitro* relationship between pH and  $P_{CO_2}$  (Astrup method). Haemolymph from each individual was equilibrated at 46 and 98 mmHg  $P_{CO_2}$  in a microtonometer and pH was measured. The results were plotted, assuming a linear relationship between pH and  $\log P_{CO_2}$

(Barnhart, 1986a). *In vivo* pH was compared with these plots to estimate *in vivo*  $P_{CO_2}$ .

Isotope activities in haemolymph (disints  $\text{min}^{-1} \text{ml}^{-1}$ ) were measured using a dual-channel liquid scintillation counter with external standard (Tracor Mark III). Haemolymph was digested in NCS tissue solubilizer and counted in OCS (Amersham), with correction for counting efficiencies in the calculation of activity. Extracellular volume was calculated from PEG dilution, extrapolating the exponential decline of haemolymph PEG activity to the time of injection. Extracellular water content (ECW, g) was calculated from extracellular volume assuming haemolymph water content to be  $1 \text{ g ml}^{-1}$ . Total body water (TBW, g) was measured as the difference between wet and dry body mass, and intracellular water (ICW, g) as the difference between TBW and ECW.  $\text{pHi}$  was calculated for the total intracellular space using the equation:

$$\text{pHi} = \text{pK}_{\text{DMO}} + \log \frac{[\text{DMO}]_i}{[\text{DMO}]_e} (1 + 10^{\text{pHe} - \text{pK}_{\text{DMO}}}) - 1 \quad (1)$$

(Waddell & Butler, 1959), assuming  $\text{pK}_{\text{DMO}} = 6.27$  (Boron & Roos, 1976).  $[\text{DMO}]_i$  was calculated as follows:

$$[\text{DMO}]_i (\text{disints min}^{-1} \text{g}^{-1}) = \frac{\text{DMO}_{\text{inj}} (\text{disints min}^{-1}) - \{[\text{DMO}]_e (\text{disints min}^{-1} \text{g}^{-1}) \times \text{ECW}(\text{g})\}}{\text{ICW}(\text{g})}, \quad (2)$$

where  $\text{DMO}_{\text{inj}} (\text{disints min}^{-1})$  is the total DMO injected.

## Results

### *Effect of $CO_2$ on $\dot{V}_{O_2}$*

$\dot{V}_{O_2}$  decreased with time in snails respiring in the absence of  $CO_2$  (see Fig. 2). Typically,  $\dot{V}_{O_2}$  declined gradually over 6–8 h to about 70% of the initial level, then remained stable for the remainder of the measurement periods (up to 14 h).

$\dot{V}_{O_2}$  was rapidly depressed by ambient hypercapnia, and the reduction of  $\dot{V}_{O_2}$  was greater at higher  $P_{CO_2}$ . The relationship between  $\dot{V}_{O_2}$  and  $P_{CO_2}$  appears to be nonlinear, with maximum slope at roughly 40–60 mmHg  $P_{CO_2}$  (Fig. 1). The reduction of  $\dot{V}_{O_2}$  relative to the control group was maximal within the first hour of exposure (Fig. 2), although equilibration with altered ambient  $P_{CO_2}$  requires more than 2 h (Fig. 3, see below). When  $CO_2$  was removed,  $\dot{V}_{O_2}$  increased within 1 h to a level similar to or somewhat higher than that of the control group (Fig. 2).

The behaviour of each animal was noted at hourly intervals. All individuals remained at least partly extended from the shell during respirometry. Crawling was infrequent in all groups (2% of observations). Over time, an increasing proportion of the snails partly retracted into the shell; this occurred sooner in low  $CO_2$  than in high  $CO_2$ . In the experiment illustrated in Fig. 2, half or more of the snails in 0 mmHg  $P_{CO_2}$  were partly retracted after 2 h, whereas this occurred only after 7 and 11 h in 46 and 98 mmHg  $P_{CO_2}$ , respectively.




Fig. 1. Effect of carbon dioxide on oxygen consumption of active *Otala lactea*. Ordinate shows the rate of oxygen consumption during hypercapnia as a fraction of that in the absence of  $CO_2$ . Oxygen consumption of each individual was measured over two consecutive 1-h periods: first without  $CO_2$ , then with  $CO_2$  present. Horizontal bars, boxes and vertical bars indicate, respectively, mean, 95% confidence interval of mean, standard deviation and range. Numbers beside boxes indicate sample size. Points show individual values where sample size was less than 10.

#### *Equilibration with ambient $P_{CO_2}$*

The time course of change of internal  $P_{CO_2}$  during ambient hypercapnia can be inferred from the respiratory exchange ratio,  $R$  (Fig. 3).  $R$  was low or negative immediately following an increase of ambient  $P_{CO_2}$ , which may be attributed to the accumulation of  $CO_2$  in the body fluids as  $P_{CO_2}$  increased.  $R$  increased rapidly and became positive within 2 h, which shows that  $P_{CO_2}$  of the body fluids rose rapidly and exceeded ambient  $P_{CO_2}$  within this period.  $R$  eventually rose beyond the control level (Fig. 3). The overshoot was greater at higher  $P_{CO_2}$ , and may have been due to an increase of haemolymph  $[HCO_3^-]$  during hypercapnia (Burton, 1976; Barnhart, 1986a). Addition of  $HCO_3^-$  to the body fluids shifts the  $CO_2$ -carbonic acid-bicarbonate equilibrium and should therefore increase  $CO_2$  loss and elevate  $R$ .

#### *Effects of hypercapnia on $pH_e$ and $pH_i$*

Mixing of the isotopes was judged to be complete within 3 h of injection; after this period the mean  $[DMO]_e$  was stable, although PEG levels declined exponentially, as indicated by the linearity of plots of the logarithm of activity ( $\text{disintegrations min}^{-1}$ ) vs time from 3 to 6 h after injection. These plots were extrapolated to time zero to estimate extracellular fluid volumes for each individual. Mean extracellular water as a percentage of total body water was  $51.1 \pm 6.74\%$  (s.d.,  $N = 15$ ).




Fig. 2. Time course of change in rate of oxygen consumption in the presence and absence of  $\text{CO}_2$ . Points and bars indicate means and 95 % confidence limits of means,  $N=8$ .  $\dot{V}_{\text{O}_2}$  of each individual was expressed as a fraction of an initial control measurement in the absence of  $\text{CO}_2$ . Dotted lines indicate the period of  $\text{CO}_2$  exposure.  $P_{\text{CO}_2} = 0$  mmHg (●);  $P_{\text{CO}_2} = 46$  mmHg (▲);  $P_{\text{CO}_2} = 98$  mmHg (■).

Between 3 and 6 h after injection the mean values of  $[\text{DMO}]_e$  and  $\text{pHe}$  in the three groups of snails did not change significantly. Therefore, the  $[\text{DMO}]_e$  and  $\text{pHe}$  results were averaged over this period for each individual and used to calculate a single estimate of  $\text{pHi}$  (from equations 1, 2) and  $P_{\text{CO}_2}$  (from the Astrup line) for each (Fig. 4). Both  $\text{pHe}$  and  $\text{pHi}$  declined similarly with increase of haemolymph  $P_{\text{CO}_2}$ . Linear regressions of  $\text{pHe}$  and  $\text{pHi}$  on the logarithm of haemolymph  $P_{\text{CO}_2}$  yielded the following equations:

$$\text{pHi} = 8.446 - 0.667 \log P_{\text{CO}_2} \quad (r^2 = 0.81), \quad (3)$$

$$\text{pHe} = 8.631 - 0.700 \log P_{\text{CO}_2} \quad (r^2 = 0.93). \quad (4)$$

The regression coefficients for  $\text{pHe}$  and  $\text{pHi}$  did not differ significantly. Mean  $\text{pHi}$  was lower than  $\text{pHe}$  by 0.12 units; this difference was significant (analysis of covariance, critical  $P = 0.05$ ).

#### *Effect of hypoxia on $\dot{V}_{\text{O}_2}$*

$\dot{V}_{\text{O}_2}$  was independent of ambient  $P_{\text{O}_2}$  above about 90 mmHg in the absence of  $\text{CO}_2$ , and independent above about 50 mmHg when  $\dot{V}_{\text{O}_2}$  was lowered 45 % by the presence of  $\text{CO}_2$  (Fig. 5). Below these levels,  $\dot{V}_{\text{O}_2}$  varied with  $P_{\text{O}_2}$ . The relationship between ambient  $P_{\text{O}_2}$  and  $P_{\text{O}_2}$ -limited  $\dot{V}_{\text{O}_2}$  is approximately described by the following regression equation (see Fig. 5):

$$\dot{V}_{\text{O}_2} (\mu\text{l g}^{-1} \text{h}^{-1}) = 1.05 P_{\text{O}_2} (\text{mmHg}). \quad (5)$$




Fig. 3. Time course of change in the respiratory exchange ratio ( $R = \dot{V}_{CO_2}/\dot{V}_{O_2}$ ) during  $CO_2$  exposure. Points and bars indicate means and 95% confidence limits of means ( $N=6$ ). Note that the vertical scale differs above and below zero.  $P_{CO_2} = 33$  mmHg (●);  $P_{CO_2} = 66$  mmHg (▲);  $P_{CO_2} = 98$  mmHg (■).

The y-intercept did not differ significantly from zero (5.1 mmHg;  $P = 0.2$ ); consequently, the regression was forced through zero. This equation was used to predict the limit of  $\dot{V}_{O_2}$  as a function of  $P_{O_2}$  (see Discussion).

## Discussion

### *Effect of hypercapnia on $\dot{V}_{O_2}$*

Previous work has shown that  $CO_2$  retention is correlated with lowered  $\dot{V}_{O_2}$  in dormant *Otala* (Barnhart & McMahon, 1987). The present results show that the oxygen consumption of active *Otala* is rapidly and reversibly depressed by imposed hypercapnia. Both results appear to be consistent with the hypothesis that hypercapnia resulting from hypoventilation reduces metabolic rate during dormancy. However,  $\dot{V}_{O_2}$  of hypercapnic active snails was not depressed to the same degree as during dormancy. The highest ambient  $P_{CO_2}$  tested lowered mean  $\dot{V}_{O_2}$  by only 63% (Fig. 1) compared with the 85% reduction during dormancy (Barnhart & McMahon, 1987). In absolute terms, the mean  $\dot{V}_{O_2}$  in 98 mmHg ambient  $P_{CO_2}$  ( $37 \mu l g^{-1} h^{-1}$ ) was more than double the mean dormant  $\dot{V}_{O_2}$  and six times the minimum dormant  $\dot{V}_{O_2}$  observed previously (13.7 and  $5.6 \mu l g^{-1} h^{-1}$ , respectively; Barnhart & McMahon, 1987). Thus, it is still uncertain whether the



effects of hypercapnia limit  $\dot{V}_{O_2}$  during true dormancy. Moreover, hypercapnia did not hasten withdrawal into the shell or induce epiphragm formation, events which normally accompany entry into dormancy.




Fig. 4. Relationship between extracellular (i.e. haemolymph) pH (pHe, ●), whole-body intracellular pH (pHi, ○), and haemolymph  $P_{CO_2}$  in active *Otala lactea* after 6–9 h of ambient hypercapnia. pHe was measured directly, pHi was estimated using DMO, and  $P_{CO_2}$  was estimated from pHe using the Astrup method. Points represent individual animals; data for each individual are based on averages of 3–4 measurements of pHe and  $[DMO]_e$  (see Results).




Fig. 5. Effect of  $P_{O_2}$  on  $\dot{V}_{O_2}$ . Symbols and bars indicate means and 95% confidence limits of means ( $N = 4$  for each). Open and closed circles are animals respiring in 0 and 66 mmHg  $P_{CO_2}$ , respectively.  $\dot{V}_{O_2}$  is expressed per gram whole body mass (including shell). The line is the least-squares linear regression of  $P_{O_2}$ -dependent  $\dot{V}_{O_2}$  on  $P_{O_2}$ . The regression was forced through zero (see Results).

*Hypercapnia and O<sub>2</sub> delivery*

A possible mechanism for depression of  $\dot{V}_{O_2}$  by CO<sub>2</sub> in *Otala* is through interference with O<sub>2</sub> delivery to the tissues, either by depressing circulation (heart rate decreases in hypercapnic individuals) or by the reversed Bohr effect of haemocyanin (Barnhart, 1986b) which might reduce P<sub>O<sub>2</sub></sub> at the tissues. Either effect might conceivably cause oxygen limitation of  $\dot{V}_{O_2}$  during hypercapnia. However, if hypercapnia inhibited oxygen transport, oxygen-limited  $\dot{V}_{O_2}$  should be lower at similar P<sub>O<sub>2</sub></sub> during hypercapnia due to reduced conductance for oxygen (Herreid, 1980). This is not the case (Fig. 5), and the results therefore argue that hypercapnia does not interfere with oxygen transport.

*Hypercapnia and acid-base balance*

Change of intracellular pH is one of several possible mechanisms by which hypercapnia could initiate metabolic effects (Walsh *et al.* 1988). The present results show that intracellular acidosis must be considered as a potential mechanism for the observed effect of CO<sub>2</sub> on  $\dot{V}_{O_2}$  in *Otala*. The whole-body pH<sub>i</sub> measurement leaves open the question of the exact values of pH<sub>i</sub> in different tissues. These values were not determined in the present study because of uncertainty regarding the fate of the extracellular volume marker (PEG) as it left the haemolymph. The similar extracellular and intracellular acidosis during hypercapnia is curious because of the presumably higher buffering capacity of the intracellular space (Burton, 1976). Similar results were, however, obtained in the bivalve *Mytilus* (Lindinger *et al.* 1984).

Although intracellular pH has not been measured in dormant *Otala*, Rees & Hand (1987) report that pH<sub>e</sub> and whole-body pH<sub>i</sub> decline by similar increments during dormancy in the land pulmonate *Oreohelix*. It appears that pH<sub>i</sub> may not only fall but also fluctuate over a substantial range during dormancy due to changes in P<sub>CO<sub>2</sub></sub>. CO<sub>2</sub> release of dormant *Otala* is periodic (Barnhart & McMahon, 1987) and haemolymph P<sub>CO<sub>2</sub></sub> ranges between 25 and 100 mmHg (Barnhart, 1986b). Comparison of this range with Fig. 4 suggests that pH<sub>i</sub> fluctuates over about 0.4 units due to changes of P<sub>CO<sub>2</sub></sub> during dormancy. These changes of pH may occur rapidly, particularly during CO<sub>2</sub> release. Measurement of pH<sub>i</sub> during dormancy using DMO thus presents special difficulties. It should be noted in this regard that dormant pulmonates are sensitive to disturbances associated with handling (Machin, 1975; Herreid & Rokitka, 1976; Herreid, 1977). The manipulations required by the present application of the DMO method for measuring pH<sub>i</sub> induce hyperventilation, release of CO<sub>2</sub> and increase of  $\dot{V}_{O_2}$  in dormant *Otala* (M. C. Barnhart, unpublished results).

*Effects of acidosis on metabolism*

Although the biochemical mechanisms that reduce the rate of energy utilization during dormancy and other hypometabolic states are not yet understood, considerable evidence indicates that glycolysis is suppressed (Hochachka &

Guppy, 1987). Intracellular acidosis appears to suppress glycolysis in hibernating mammals (see review by Malan, 1986) and in the brine shrimp, *Artemia*, during metabolic arrest (Busa & Nuccitelli, 1984; Carpenter & Hand, 1986; Hand & Carpenter, 1986). Among molluscs, acid pH reduces phosphofructokinase activity in the adductor muscle of *Mytilus* (Ebberink, 1982), and acidosis has been implicated in reduction of phosphofructokinase activity and glycolysis during anoxia in the whelk *Busycon* (Ellington, 1983). Pyruvate kinase from *Otala* shows decreased affinity for phosphoenolpyruvate and ADP, increased inhibition by alanine and ATP, and decreased activation by fructose 1,6-diphosphate *in vitro* when pH is lowered from 7.0 to 6.5 (J. Fields, personal communication).

Other mechanisms which have been proposed to suppress glycolysis during hypometabolism involve neither CO<sub>2</sub> nor change in pH. These include phosphorylation of regulatory enzymes, dissociation of enzyme complexes to produce less active soluble forms, and decreased levels of fructose 2,6-bisphosphate, an activator of phosphofructokinase (Storey, 1988). Recent studies show that fructose 2,6-bisphosphate levels fall by 86–93% in *Otala* during 4 days of dormancy and subcellular binding of glycolytic enzymes decreases (K. Storey, personal communication). Thus, evidence suggests that both pH-dependent and pH-independent mechanisms may suppress glycolysis in *Otala* during dormancy.

#### *Hypoxia and $\dot{V}_{O_2}$*

P<sub>O<sub>2</sub></sub> in the lung and haemolymph of dormant *Otala* is on average only about one-third that of active snails, and is often less than 10 mmHg owing to hypoventilation (Barnhart, 1986b). Might  $\dot{V}_{O_2}$  be oxygen-limited during dormancy? One approach to this question is to compare  $\dot{V}_{O_2}$  in dormancy with the limit of  $\dot{V}_{O_2}$  predicted at the low levels of P<sub>O<sub>2</sub></sub> typical of dormancy. As a first approximation, lung P<sub>O<sub>2</sub></sub> can be substituted for ambient P<sub>O<sub>2</sub></sub> in equation 5 [lung and ambient P<sub>O<sub>2</sub></sub> do not differ greatly in active *Otala* (Barnhart, 1986b)]. The mean of the lowest 10% of lung P<sub>O<sub>2</sub></sub> values observed in dormant *Otala* was 4.5 mmHg (data of Barnhart, 1986b). The predicted limit of  $\dot{V}_{O_2}$  at this P<sub>O<sub>2</sub></sub> is about 5  $\mu\text{l g}^{-1} \text{h}^{-1}$  (from equation 5). This value is similar to the minimum sustained  $\dot{V}_{O_2}$ , observed during continuous respirometry ( $5.6 \pm 0.57 \mu\text{mol g}^{-1} \text{h}^{-1}$ , s.e.m.; Barnhart & McMahon, 1987). Thus it appears that lung P<sub>O<sub>2</sub></sub> sometimes approaches a critical level. However, the data do not indicate that P<sub>O<sub>2</sub></sub> normally limits  $\dot{V}_{O_2}$  during dormancy. Mean lung P<sub>O<sub>2</sub></sub> during dormancy is 38 mmHg (Barnhart, 1986a), which corresponds to a  $\dot{V}_{O_2}$  limit of 40  $\mu\text{l g}^{-1} \text{h}^{-1}$ , much higher than the mean  $\dot{V}_{O_2}$  during dormancy (14  $\mu\text{l g}^{-1} \text{h}^{-1}$ ; Barnhart & McMahon, 1987).

In conclusion, the effects of acute hypercapnia and hypoxia on oxygen consumption of active *Otala lactea* indicate that hypercapnia, but not hypoxia, explains much of the reduction of aerobic metabolism during dormancy. However,  $\dot{V}_{O_2}$  of acutely hypercapnic active snails is not depressed to the lowest levels typical of dormancy, suggesting that other factors also contribute to metabolic depression.

This work was supported by a postdoctoral fellowship from the Alberta Heritage Foundation for Medical Research, and by NSERC grant A5762.

### References

- BARNHART, M. C. (1986a). Control of acid-base status in active and dormant land snails, *Otala lactea* (Pulmonata, Helicidae). *J. comp. Physiol. B* **156**, 347–354.
- BARNHART, M. C. (1986b). Respiratory gas tensions and gas exchange in active and dormant land snails, *Otala lactea*. *Physiol. Zool.* **59**, 733–745.
- BARNHART, M. C. & MCMAHON, B. R. (1987). Discontinuous carbon dioxide release and metabolic depression in dormant land snails. *J. exp. Biol.* **128**, 123–138.
- BORON, W. F. & ROOS, A. (1976). Comparison of microelectrode, DMO, and methylamine methods for measuring intracellular pH. *Am. J. Physiol.* **231**, 799–809.
- BURTON, R. F. (1976). Calcium metabolism and acid-base balance in *Helix pomatia*. In *Perspectives in Experimental Biology*, vol. 1 (ed. P. S. Davies), pp. 7–16. Oxford: Pergamon Press.
- BUSA, W. B. & NUCCITELLI, R. (1984). Metabolic regulation via intracellular pH. *Am. J. Physiol.* **246**, R409–R438.
- CARPENTER, J. F. & HAND, S. C. (1986). Arrestment of carbohydrate metabolism during anaerobic dormancy and aerobic acidosis in *Artemia* embryos: Determination of pH sensitive control point. *J. comp. Physiol. B* **156**, 451–460.
- EBBERINK, R. H. M. (1982). Control of adductor muscle phosphofructokinase activity in the sea mussel *Mytilus edulis* during anaerobiosis. *Molec. Physiol.* **2**, 345–355.
- ELLINGTON, W. R. (1983). Phosphorus nuclear magnetic resonance studies of energy metabolism in molluscan tissues: Effect of anoxia and ischemia on the intracellular pH and high energy phosphates in the ventricle of the whelk, *Busycon contrarium*. *J. comp. Physiol.* **153**, 159–166.
- HAND, S. C. & CARPENTER, J. F. (1986). pH-induced metabolic transitions in *Artemia* embryos mediated by a novel hysteretic trehalase. *Science* **232**, 1535–1537.
- HERREID, C. F. (1977). Metabolism of land snails (*Otala lactea*) during dormancy, arousal, and activity. *Comp. Biochem. Physiol.* **56A**, 211–215.
- HERREID, C. F., III (1980). Hypoxia in invertebrates. *Comp. Physiol. Biochem.* **67A**, 311–320.
- HERREID, C. F. & ROKITKA, M. A. (1976). Environmental stimuli for arousal from dormancy in the land snail, *Otala lactea*. *Physiol. Zool.* **49**, 181–190.
- HOCHACHKA, P. W. (1988). Metabolic suppression and oxygen availability. *Can. J. Zool.* **66**, 152–158.
- HOCHACHKA, P. W. & GUPPY, M. (1987). *Metabolic Arrest and the Control of Biological Time*. Cambridge, MA: Harvard University Press. 227pp.
- LINDINGER, M. I., LAUREN, D. J. & McDONALD, D. G. (1984). Acid-base balance in the sea mussel, *Mytilus edulis*. III. Effects of environmental hypercapnia on intra- and extracellular acid-base balance. *Mar. Biol. Letts* **5**, 371–381.
- MACHIN, J. (1975). Water relationships. In *The Pulmonates* (ed. V. Fretter & J. Peake), pp. 105–163. New York: Academic Press.
- MALAN, A. (1986). pH as a control factor in hibernation. In *Living in the Cold. Physiological and Biochemical Adaptations* (ed. H. C. Heller, X. J. Musacchia & L. C. H. Wang), pp. 61–70. New York: Elsevier.
- REES, B. & HAND, S. C. (1987). Acidosis and urea accumulation during dormancy in the mountain snail, *Oreohelix*. *Physiologist* **30**, 222.
- STOREY, K. (1988). Suspended animation: the molecular basis of metabolic depression. *Can. J. Zool.* **66**, 124–132.
- VLECK, D. (1987). Measurement of O<sub>2</sub> consumption, CO<sub>2</sub> production, and water vapor production in a closed system. *J. appl. Physiol.* **62**, 2103–2106.
- WADDELL, W. J. & BUTLER, T. C. (1959). Calculation of intracellular pH from the distribution of 5,5-dimethyl-2,4-oxazolinedione (DMO). Application to skeletal muscle of the dog. *J. clin. Invest.* **38**, 720–729.
- WALSH, P. J., MOMMSEN, T. P., MOON, T. W. & PERRY, S. F. (1988). Effects of acid-base variables on *in vitro* hepatic metabolism in rainbow trout. *J. exp. Biol.* **135**, 231–241.