



JOURNAL OF BIOMEDICINE AND TRANSLATIONAL RESEARCH

Copyright©2016 by Faculty of Medicine Diponegoro University and Indonesian Doctor Association, Central Java Region

Polymorphisms of *TLR4* Asp299Gly and *TNF-α* -308G/A in Leptospirosis

Nur Farhanah¹, Muhammad Hussein Gasem¹, Sultana MH Faradz²

1. Division of Tropical Medicine and Infectious Diseases, Faculty of Medicine Diponegoro University/Dr. Kariadi Hospital Semarang, Indonesia
2. Center for Biomedical Research (CEBIOR), Faculty of Medicine, Diponegoro University, Semarang, Indonesia

Article Info

History:

Received 1 July 2016

Accepted 20 July 2016

Available 30 June 2016

ABSTRACT

Background: *TLR4* Asp299Gly and *TNF-α* -308G/A polymorphisms have been shown to be associated with increased susceptibility and severity of infection. *TLR4* Asp299Gly polymorphism could affect the host's ability to respond to *leptospira sp.* *TNF-α* -308G/A polymorphism, is associated with the high producer of *TNF-α*.

Methods: Total of 36 leptospirosis patients (IgM anti leptospira and MAT positive) and healthy individual with equal number were included. The polymorphisms were determined by polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) using site specific restriction enzyme.

Results: Distribution of homozygous wild-type *TLR4* Asp299Gly polymorphism was higher in both of groups (94.5: 97.2%) and homozygous mutant allele was absent. There was not significantly difference of *TLR4* Asp299Gly in leptospirosis patients and healthy group ($p=1.00$; OR 0.5; 95%CI, 0.04-5.6) and between mild and severe leptospirosis ($p=0.54$; OR 1.54; 95% CI, 1.20-1.98). The presence of homozygous wild-type *TNF-α* -308G/A polymorphism was higher between leptospirosis patients and healthy group (100: 94.5%) and homozygous mutant allele was not found in both of the groups. No significantly different of *TNF-α* -308G/A polymorphism between leptospirosis patient and healthy group ($p=0.49$).

Conclusions: In this study, the polymorphisms of *TLR4* Asp299Gly and *TNF-α* -308G/A are not associated with the susceptibility and severity of leptospirosis.

Keywords: Leptospirosis, *TLR4* Asp299Gly polymorphism, *TNF-α* -308G/A polymorphism.

INTRODUCTION

Leptospirosis is a zoonotic disease caused by *Leptospira interrogans*. Leptospirosis is an endemic

health problem in South-East Asia region^{1,2}. The clinical manifestations are varying widely, ranging from mild leptospirosis (anicteric) to severe leptospirosis (icteric). Severe leptospirosis known as Weil's disease is characterized by icteric, acute renal failure and bleeding tendency^{3,4}. Severity and

*Corresponding Author:

Nur Farhanah

Phone: +62 812 2524 318

Email: nurfarhanahams@gmail.com

mortality rates of leptospirosis influence by several factors: types of serovars, and host factors including genetic, age, immunity status, comorbidity, occupation or activity. Genetic factor is one of the main important in susceptibility and severity of infections^{1,3}.

The pathogenesis of leptospirosis is not fully understood. It seems mimic the pathogenesis of sepsis caused by gram-negative bacteria infection⁵. *Leptospira sp* has immunogenic lipopolysaccharide wall cell (LPS) a potent activator of the natural immune system (innate/nonspecific immunity), which will be responded by *TLR4* (*Toll Like Receptor-4*). *TLR4* is a key receptor for gram negative bacteria, mycobacteria, fungi and parasites of malaria. *TLR4* will stimulate macrophages to produce proinflammatory cytokines such as tumor necrosis factor- α (TNF- α), interleukin-1 (IL-1) and interleukin-6 (IL-6)⁶⁻⁸.

Polymorphism of *TLR4* with the common frequency >5% is *TLR4* Asp299Gly change from asparagine (Asp) at codon GAT to glycine (Gly) to GGT codon at 299 of amino acid. The presence of these polymorphisms alter the function of *TLR4* receptor, associated with hyporesponsive to LPS and increased the susceptibility of gram negative infection, sepsis, septic shock, tuberculosis and malaria⁹⁻¹³.

The major cytokines in the acute inflammatory response of gram negative bacteria and microbes is TNF- α ¹⁴. There are several type of TNF- α polymorphisms that have been studied. TNF- α -308G/A is associated with increased of transcriptional activity of TNF- α ¹⁵. Several studies showed that the TNF- α -308G/A associated with the incidence of infection in old age, respiratory failure, septic shock in community acquired pneumonia, and severe sepsis in burn trauma. There was a significant difference between mutant-type allele A/A of TNF- α -308G/A polymorphism with septic shock¹⁵⁻¹⁷. The study about dengue in Semarang showed that there was correlation of TNF- α polymorphism between dengue cases and control¹⁸.

The objective of this study were to know whether polymorphisms of *TLR4* Asp299Gly gene and TNF- α -308G/A related to susceptibility and severity of leptospirosis.

MATERIALS AND METHODS

Study Design

The subjects were consecutively derived from an observational cohort study conducted in Dr. Kariadi Hospital and Center for Biomedical Research (CEBIOR) Semarang from 2010 to 2012.

Study Subjects

Inclusion criteria for patients were individual aged 14 years and older who match clinical criteria of leptospirosis (WHO-SEARO, Chennai 2010) with positive of rapid diagnostic test of IgM anti leptospira and confirmed by MAT (microagglutination test) and agreeing to become a study subject by signing the consent form. Inclusion criteria for control were healthy volunteers not suffering from leptospirosis and signing the consent form. Thirty six controls and 36 leptospirosis patients were selected; 24 patients (72.2%) were classified as severe leptospirosis, while 12 patients (27.8%) were mild.

DNA extraction

DNA extraction used the salting out method as described by previous investigators with slight modification¹⁹.

Analysis of gene polymorphisms

Analysis of TNF- α gene polymorphism was performed by amplification of the TNF- α -308 gene in the promoter region at position -308, using a PCR cyclor (Applied Biosystems). The total volume of each sample mixture (20ul) contained 2ul DNA, 0.2 ul of forward primer (5'AGG CAA TAG GTT TTG AGG GCC AT 3') and reverse primer (5' TCC TCC CTG CTC CGA TTC CG 3') respectively, 0.4 ul dNTP, 0.4 ul MgCl₂ and 0.2 ul of the Taq enzyme.

The polymerase chain reaction (PCR) profile was consisted of 5 minutes at 95°C for initial denaturation, followed by final extension for 7 minutes at 72°C. The PCR product of 107 bp was then cut by restriction fragment length polymorphism (RFLP), by incubation at 55°C for 18 hours (overnight) with the *NcoI* restriction enzyme. The PCR product was visualized by 2% agarose gel electrophoresis. In individuals homozygous for the -308 (A/A) allele, the PCR product was not cut, resulting in a band on the gel of 107 bp. In contrast, homozygous G/G yielded fragments of 87bp and 20 bp, while heterozygous G/A yielded fragments of 107 bp, 87 bp and 20 bp.

Analysis of *TLR-4* Asp299Gly gene polymorphism was using a PCR. The total volume of each sample mixture (20ul) contained 2ul DNA, 0.5 ul of forward primer (5' GAT TAG CAT ACT TAG ACT ACT ACC TCC ATG 3') and reverse primer (5' TCC TCC CTG CTC CGA TTC CG 3'), respectively, 1.2 ul dNTP, 2.5 ul MgCl₂ and 0.3 ul of the Taq enzyme. The polymerase chain reaction (PCR) cycle was consisted of 5 minutes at 95°C for initial denaturation, followed by final extension for 7 minutes at 72°C. The PCR product of 107 bp was then cut by restriction fragment length polymorphism (RFLP), by incubation at 55°C for 18 hours (overnight) with the *NcoI* restriction enzyme. We used 100 bp for marker ladder. The PCR product was visualized using 2% agarose

gel. In individuals homozygous for the *TLR4* Asp299Gly (A/A) allele, the PCR product was not cut, resulting in a band on the gel 195bp. In contrast, homozygous G/G yielded fragments of 176 bp and 19 bp, while heterozygous A/G yielded fragments of 195 bp, 176 bp and 19 bp.

Data Analysis

The data were analysed by Mann-Whitney U test, Hardy-Weinberg Equilibrium for allele frequency and Fischer-Exact test. All analyses were performed by means of the SPSS program version 15 for Windows, at a significance level of 5%.

Ethical Clearance

The study was approved by Ethics Committee of Medical Research and Health Faculty of Medicine, Diponegoro University/Dr Kariadi Hospital, Semarang, with ethical clearance No 102/EC/FK/RSDK/2010.

RESULTS

Thirty six leptospirosis patients were included in this study, consisted of male 67.2% and the mean of age was 47.25 (SD 12.52) years. The distribution of homozygous wild-type *TLR4* Asp299Gly polymorphism was high in leptospirosis patients and control groups (94.5 vs 97.2%) and homozygous mutant allele was absent in both studied groups. The heterozygote allele of *TLR4* Asp299Gly was 5.5% while in control was 2.8%. The comparison between leptospirosis patients and control showed that there was not significantly difference in patients than control group ($\rho=1.00$; OR 0.5; 95%CI, 0.04-5.6). Frequency for G allele in each group was 2.8 : 1.4 % for leptospirosis group and control. There was no significantly difference of *TLR4* Asp299Gly polymorphism between mild and severe leptospirosis ($\rho=0.5$; OR 1.5; 95%CI 1.20-1.98).




Figure 1.A. PCR-RFLP results for *TLR4* Asp299Gly in Leptospirosis

Lane A1 marker 100 bp ladder, lane A2 blanks. Lanes A3, 5-7, 9-17 homozygous A/A Lanes A4,8 heterozygous A/G. PCR product digested using *NcoI* restriction enzyme.

Figure 1.B. PCR-RFLP for *TLR4* Asp299Gly in healthy control

Lane B1 marker 100 bp ladder, lane B2 blanks.

Lanes B3-5, 7-13 homozygous A/A. Lane B6 heterozygous A/G. Lane B11: blanks. The presence of *TNF- α* -308G/A polymorphism between leptospirosis patients and control group were as follow: homozygous wild-type (100 vs 94.5%), heterozygous allele (0.0 vs 5.5%) and homozygous mutant allele were not found in both of the group. There was not significantly different of *TNF- α* -308G/A polymorphism between leptospirosis patient and control ($\rho=0.49$).




Figure 2.A. PCR-RFLP *TNF-α* -308G/A polymorphism in leptospirosis.

Lane C1: marker 100 bp ladder, lane C2: blanks
Lanes C3-11: homozygote allele G/G. PCR product digested using *NcoI* restriction enzyme.

Figure 2.B. PCR-RFLP *TNF-α* -308G/A polymorphism in healthy control.

Lane D1: marker 100 bp ladder, lane D2: blanks
Lanes D3-5: homozygote allele G/G. Lane D6: heterozygote A/G. Lanes E1-E6: PCR product undigested by enzyme *NcoI*.

DISCUSSION

In this study, we found that there were no significant differences between *TLR4* Asp299Gly polymorphism of leptospirosis patients and healthy control, or between mild and severe leptospirosis. It was contrast with previous studies that the polymorphism *TLR4* Asp299Gly gene was associated with susceptibility and severity of gram negative infections^{5,10}. Other studies showed similar result that there were no significant differences of *TLR4* Asp299Gly between DHF/DSS and control group, or severity of meningococcal infection and outcome of gram negative sepsis²⁰⁻²³. Metaanalysis studies in Caucasian and in China found that there were not strong correlation between sepsis and *TLR4* Asp299Gly polymorphism²⁴⁻²⁷.

Race, ethnic and geography seem to affect the frequency of *TLR4* Asp299Gly polymorphism. In African, frequency of *TLR4* Asp299Gly haplotype was higher than Caucasian that had co-segregation of *TLR4* Asp299Gly/Thr399Ile, while in Asia was less. Environmental pressure when human migration to Europe, Africa, Asia and New World affected the frequency of *TLR4* Asp299Gly²⁸.

Distribution of *TNF-α* -308 G/A in leptospirosis patients was dominant allele homozygote G/G. This study found that there were no significant difference in *TNF-α* -308G/A between leptospirosis and control groups. *TNF-α* -308G/A causes increased production of *TNF-α* that could affect susceptibility and outcome of infection^{29,30}. The A allele causes increased production of *TNF-α* in a higher level than allele G³¹. Study of sepsis and severe sepsis in the Chinese population obtained that allele A was higher but was not influence on mortality. *TNF-α* serum level was high in leptospirosis but did not correlated with *TNF-α* -308G/A³².

A meta-analysis study showed that *TNF-α* -308G/A was associated with the incidence of sepsis not in death, and Asian had a strong contribution compared with other races³³. In Caucasian study showed that allele homozygotes A/A *TNF-α* -308 G/A effected on susceptibility and poor outcome and duration of ventilator used in critical illness.³⁴ Other studies showed that polymorphism of *TNF-α* -308G/A was not correlated with the outcome of sepsis gram negative and malaria³⁵⁻³⁶.

Study in Aceh revealed that no significant association between genotype distributions and allele frequency of SNP *TNF-α* -308G/A and spontaneous preterm birth in the Acehnese ethnic group³⁷. Study in endemic area in Indonesia showed that *TNF-α* -308G/A was not associated with typhoid fever³⁸. Ethnic affected the variation of allele and haplotypes of *TNF-α* -308G/A in three ethnic populations of Sulawesi. Some haplotypes might be restricted to Asian or Asian-derived populations³⁹.

This study was not consistent with the law of Hardy - Weinberg Equilibrium might be due to the small sample size and homogeneous population of Javanese.

CONCLUSION

TLR4 Asp299Gly and *TNF- α* - 308G/A genes polymorphism were not associated with susceptibility and severity of leptospirosis.

SUGGESTION

Further research is needed with a heterogeneous population including such ethnics and a large number of cases and the controls. Sequencing of *TLR4* and *TNF- α* genes are needed to determine the possibility of other polymorphisms.

Competing interest: The authors have declared that no competing interests exist.

ACKNOWLEDGEMENTS

We thank the personnel of CEBIOR. We also gratefully acknowledge the contribution of residents of Department of Medicine Dr Kariadi Hospital-Diponegoro University who helped the study. This study was supported by personal grant by Nur Farhanah (NF).

NF was recipient of fellowship of DIKTI-Beasiswa Unggulan for study in Magister of Biomedics (concentration in Genetic Counseling), Diponegoro University.

REFERENCES

- World Health Organization/International Leptospirosis Society. Human Leptospirosis Guidance for Diagnosis, Surveillance and Control. Geneva.2003.
- World Health Organization-Regional Officer for South-East Asia. Informal Expert Consultation on Surveillance, Diagnosis and Risk Reduction of Leptospirosis. 2009.
- Levett PN. Leptospirosis. Clin Microbiol Rev.2001; 14(2): 296-326.
- Speelman P. Leptospirosis. Fauci AS (ed). Harrison's Principle of Internal Medicine16 ed. Vol 1. Mc Graw Hill Companies Inc. New York. 2005:988-91
- Elves A.P, Maciel DAA, Eliana A.G, et al. High Serum Nitric Oxide Levels in Patients with Severe Leptospirosis. Acta Trop. 2006; 100(3): 256-60.
- Abbas AK, Pober JS, Conlin A. Innate Immunity. Cellular and Molecular Immunology. 5 ed. WB Saunders Company. Philadelphia. 2007 (5) : 22-6.
- Maat S. Toll-Like Receptor. Airlangga University Press. Surabaya. 2009 (1) : 1-23
- Akira Y, Takeda K. Role of Adapters in Toll-Like Receptor Signalling. Biochem Soc Trans. 2003; 31(3): 637-42.
- Ferwerda B, Verheijen K, Kullberg BJ, et al. Functional Consequences of Toll-like Receptor 4 Polymorphisms. Mol Mer. 2008; 14(5-6): 346-52.
- Lorenz E, Frees KL, Schwartz DA. Relevance of Mutations in the TLR4 Receptor in Patients With Gram-Negative Septic Shock. Arch Intern Med 2002; 162(13) : 591-601.
- Najmi KG, Sharma SK, Mehra NK. Human Toll-like receptor 4 polymorphisms TLR4 Asp299Gly and Thr399Ile influence susceptibility and severity of pulmonary tuberculosis in the Asian Indian population. Tissue Antigens. 2010; 76(2): 102-9.
- Zakeril S , Mehrizi A, Djadid ND. Genetic variation of TLR-4, TLR-9 and TIRAP gene in Iranian malaria patients. Malar J. 2011; 10:77
- Arbour NC, Schutte BC, Zabner J, et al. TLR Mutations are Associated with Endotoxin Hyporesponsiveness in Humans. Nat Genet .2000; 25: 187-91.
- Bratawijaya KG, Rengganis I. Sitokin. Immunologi Dasar. 9 ed. Jakarta: Balai Penerbit FK UI. 2010; 217-55.
- Waterer GW, Quasney M, Cantor RM. Septic Shock and Respiratory Failure in Community-acquired Pneumonia Have Different TNF Polymorphism Associations. J Respir Crit Care Med. 2001; 13: 163.
- Barber RC, Riverachavez FA. TLR4 and TNF- α polymorphism are associated with an increased risk for severe sepsis following burn injury. J Med 2004; 41: 808-13.
- Cipprino C, Caruso C, Lio D, et al. TNF a-308G/A polymorphism of TNF-alpha influences immunological parameters in old subjects affected by infectious diseases. Eur J Immunogenet. 2005; 32: 13-8.
- Ngestingisih D, Faradz SMH. Polimorfisme alel TNF- α pada penderita DHF di Semarang. M Med Indonesia. 2006; 41: 46-50.
- Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. Nucleic Acid Res.1988; 16(3) : 1215
- Djamiatun K, Ferwerda B, Netea MG, et al. Toll-like receptor 4 polymorphism in Dengue virus –infected Children. Am J Trop Med Hyg. 2011; 85(2): 352-4.
- RC Read, J Pullin S, Gregory, et al. A functional polymorphism of toll-like receptor 4 is not associated with likelihood or severity of meningococcal disease. J Infect Dis; 184(5): 640-2.

22. Jessen KM , Lindboe S, Petersen AL, et al. Common TNF- α , IL-1 β , PAI-1, uPA, CD14 and TLR4 polymorphisms are not associated with disease severity or outcome from Gram negative sepsis. *BMC Infect Dis* .2007; 7(108): 1-7.
 23. Schroder NW, Schuman RR. Single nucleotide polymorphisms of Toll-like receptors and susceptibility to infectious disease. *Lancet Infect Dis*. 2005; 5: 156-64.
 24. Zhu L, Li X, Miao C. Lack of association between TLR4 Asp299Gly and Thr399Ile polymorphisms and sepsis susceptibility: A meta-analysis. *Gene*. 2012; 601(2): 213-8.
 25. Shan XO, Wu Y, Ding JY, et al. Gene polymorphisms of Toll-like receptors in Chinese han children with sepsis in Wenzhou. *PubMed*. 2010; 48(1): 15-8.
 26. Song Z, Yin J, Yao C, et al. Variants in the Toll-interacting protein gene are associated with susceptibility to sepsis in the Chinese Han population. *Crit Care med*. 2011; 15(1): 1-10.
 27. Hang J, Zhou W, Zhang H, et al. TLR4 Asp299Gly and Thr399Ile polymorphisms are rare in the Chinese population. *J Endotoxin Res*. 2004; 10(4): 238-40.
 28. Ferwerda B, Mc Call MB, Alonso S et al. TLR4 polymorphisms, infectious Diseases, and evolutionary Pressure during Migration of Modern Humans. *Proc natl Acad Sci U S A*. 2007; 104(42): 16645-50.
 29. Sallakci N, Ackurin G, Koksoy S, et al. TNF-alpha G-308A polymorphism is associated with rheumatic fever and correlates with increased TNF-alpha production. *J Autoimmun*. 2005; 25(2): 150-4.
 30. Wong HR. Genetic and genomic in pediatric septic shock. *Crit Care Med*. 2012; 40 :1618-26
 31. Tuglular BP, Berthoux P. Berthoux F. Polymorphism of tumor necrosis factor a gene at position -308 and TNF α microsatellite in primary IgA nephropathy. *Nephrol Dial Transplant* 2003; 18: 724-31.
 32. Song Z, Yin J, Yao C, et al. Genetic variation in the TNF gene is associated with susceptibility to severe sepsis, but not with mortality. *PLoS One*. 2012; 7(9): e46113.
 33. Teuffel O, Beyene J, Sung L. Association between tumor necrosis factor-alpha promoter-308 A/G polymorphism and susceptibility to sepsis and sepsis mortality : a systemic review and meta-analysis. *Crit Care Med*. 2010; 38(1): 276-82.
 34. Watanabe E, Zehnbaue B, Oda S, et al. Tumor necrosis factor -308 polymorphism (rs1800629) is associated with mortality and ventilator duration in 1057 Caucasian patients. *Cytokine*. 2012; 60(2): 249-56.
 35. Jessen KM , Lindboe S, Petersen AL, et al. Common TNF- α , IL-1 β , PAI-1, uPA, CD14 and TLR4 polymorphisms are not associated with disease severity or outcome from Gram negative sepsis. *BMC Infect Dis* .2007; 7(108): 1-7.
 36. Mergani A, Khamis AH, Haboor AB, et al. Lack of association between - 308 tumor necrosis factor polymorphism and susceptibility to cerebral malaria among Central Sudanese children . *Int J Genet Mol Bio*. 2010; 2(5): 67-71.
 37. Andalas M, Hakimi M, Nurdiati DS, Astuti I, Imrn I, Harapan H. Association of _308G/A TNF-a gene polymorphism and spontaneous preterm birth in Acehnese ethnic group, Indonesia: This polymorphism is not associated with preterm birth. *Egypt J Med Hum Genet* (2015), <http://dx.doi.org/10.1016/j.ejmhg.2015.05.001>
 38. Ali S, Vallard AM, Kremer D, de Visser AW, Marina CA, Widjaja S, et al. Polymorphism in proinflammatory genes and susceptibility to typhoid fever and paratyphoid fever. *J Interferon Cytokine Res*. 2007; 27(4): 271-4
 39. Lamsis F, Flannery GR, White NG, Muratore R, Kaelan C, Mitchell RJ et al. Alleles and haplotypes of tumour necrosis factors (TNF) alpha and beta genes in three etnik populations of Sulawesi Indonesia. *Hum Biol* 2002; 74(3): 381-96.
-