

## Philadelphia College of Osteopathic Medicine DigitalCommons@PCOM

---

### Scholarly Posters

---

2012

# Autophagy and apoptotic genes implicated in Alzheimer's disease are modulated following infection of neuronal cells with *Chlamydia pneumoniae*

Denah M. Appelt

*Philadelphia College of Osteopathic Medicine, DenahA@pcom.edu*

Ian Kohler

*Philadelphia College of Osteopathic Medicine*

Annette K. Slutter

*Philadelphia College of Osteopathic Medicine, AnnetteKS5@gmail.com*

Juliana Zoga

*Philadelphia College of Osteopathic Medicine, JulianaZo@pcom.edu*

Susan T. Hingley

*Philadelphia College of Osteopathic Medicine, susanh@pcom.edu*

Follow this and additional works at: <http://digitalcommons.pcom.edu/posters>



Part of the [Bacterial Infections and Mycoses Commons](#), [Medical Biochemistry Commons](#), and the [Nervous System Diseases Commons](#)

---

### Recommended Citation

Appelt, Denah M.; Kohler, Ian; Slutter, Annette K.; Zoga, Juliana; Hingley, Susan T.; and Balin, Brian J., "Autophagy and apoptotic genes implicated in Alzheimer's disease are modulated following infection of neuronal cells with *Chlamydia pneumoniae*" (2012). *Scholarly Posters*. Book 10.

<http://digitalcommons.pcom.edu/posters/10>

This Book is brought to you for free and open access by DigitalCommons@PCOM. It has been accepted for inclusion in Scholarly Posters by an authorized administrator of DigitalCommons@PCOM. For more information, please contact [library@pcom.edu](mailto:library@pcom.edu).

---

**Authors**

Denah M. Appelt, Ian Kohler, Annette K. Slutter, Juliana Zoga, Susan T. Hingley, and Brian J. Balin

# Autophagy and apoptotic genes implicated in Alzheimer's disease are modulated following infection of neuronal cells with *Chlamydia pneumoniae*

Denah Appelt, Ian Kohler, Annette Slutter, Juliana Zoga, Susan Hingley, Brian Balin  
Philadelphia College of Osteopathic Medicine, Center for Chronic Disorders of Aging, Philadelphia, PA, USA

## Abstract

**Background:** The focus of the current studies was to determine the relationship between the molecular mechanisms interconnecting autophagy and apoptosis following *Chlamydia pneumoniae* infection in neuronal cells. Dysfunctions in apoptosis and autophagy have been implicated in the neurodegeneration associated with Alzheimer's disease (AD). Autophagy in AD pathogenesis has been shown to play a role in amyloid processing through the endosomal-lysosomal system. Apoptosis may contribute to the neuronal cell loss observed in AD; however, there is limited evidence of the apoptotic process proceeding to terminal completion. Although A $\beta$ 1-42 has been shown to induce apoptosis in neurons and may be an early factor in AD, our previous investigations demonstrated that neurons infected with *Chlamydia pneumoniae* are resistant to apoptosis, and that A $\beta$ 1-42 is induced following this infection. Thus, these studies address infection as an initiator/trigger or inhibitor for the processes of autophagy and apoptosis observed in Alzheimer's disease.

**Methods:** SKNMC neuronal cells obtained from ATCC were infected with the AR39 strain of *Chlamydia pneumoniae* at an MOI=1 for 24, 48, and 72hrs and were analyzed using Real-time PCR arrays from SABiosciences specific for autophagy and apoptosis genetic markers.

**Results:** Some major genes associated with apoptosis such as BID, DAPK1, TP53, TP73 were down regulated by 72hrs post-infection. Genes associated with the regulation of autophagic vacuole formation such as ATG3, ATG4B, ATG4C, ATG9A, ATG9B, ATG12, IRGM, and BECN1 were up-regulated within 72hrs post-infection. With regards to genes involved with co-regulation of autophagy and apoptosis, BNIP3 was significantly up-regulated within 48-72hrs post-infection. Of the genes linking autophagosomes to lysosomes, FAM176A was up-regulated throughout 24-72hrs post-infection.

**Conclusions:** Modulation of autophagy and apoptosis genes occurs in neuronal cells at 24, 48, and 72hrs post-infection with *Chlamydia pneumoniae*. These genetic changes lead to dysfunction in these basic cellular processes; dysfunction in these processes has been shown to contribute to the neuropathology of late-onset Alzheimer's disease. This work will allow future studies to further focus on the apoptotic and autophagic pathways to better understand how a pathogen such as *Chlamydia pneumoniae* plays a role in the development of late-onset Alzheimer's disease.

## Introduction

Neurodegeneration has been well documented in the CNS of Alzheimer individuals and a continuum of abnormalities has been identified within the AD brain. Strong evidence suggests that abnormalities of the autophagy and apoptosis pathways are contributing factors in the pathogenesis of Alzheimer's disease. Our laboratory has focused on studying the role of infection with *C. pneumoniae* as a causative agent in late onset AD. In separate studies, polymerase chain reaction detected *C. pneumoniae* DNA in 80 to 90% of postmortem sporadic AD brain samples [1,2], but only 5-11% of postmortem brain samples from age-matched, non-AD, control individuals. Furthermore, a murine model has been developed in which mice infected with *C. pneumoniae* demonstrate deposits of amyloid in brain areas typically affected in AD [3].

Recently, we have demonstrated that *C. pneumoniae* is capable of inhibiting apoptosis in neuronal cells thereby prolonging the viability of the infected neuronal cell [4]. Other laboratories have demonstrated that chlamydiae-infected host cells were resistant to proapoptotic stimuli such as TNF $\alpha$ , Fas antibody, staurosporine, and UV-light [5]. *C. pneumoniae* infection has also been shown to modulate pro-apoptotic cytoplasmic proteins, such as caspase-3 and cytochrome c, as well as the anti-apoptotic mitochondrial protein Bcl-2 and the anti-apoptotic nuclear protein NF-KB[6].

In some instances, cells may activate the autophagy pathway instead of, or before, initiating apoptosis. Induction of the autophagic process may lead to modulation of the apoptotic process. Autophagy is associated with the endosomal-lysosomal system, in which contents of an autophagosome are degraded as a result of the autophagosome fusing with the lysosome[7,8]. An increase in the number of autophagic vacuoles (AVs) was identified in neurons from AD brains implicating autophagy as a pathological process in AD [9]. The endosomal pathway is linked to the lysosomal system because early endosomes fuse with late endosomes or lysosomes. Neurons from AD brains have been found to exhibit enlarged early endosomes. This is significant in AD because early endosomes take in proteins such as apolipoprotein E and APP, and it has been determined that A $\beta$  is formed in early endosomes [10].





Typically, autophagy and apoptosis are common pathways by which infected cells attempt to rid themselves of the infectious agent and cells incapable of eliminating the infectious agent, undergo cell death. Intriguingly, these pathways appear to be altered in *C. pneumoniae* infected neuronal cells. These studies are an attempt to identify the defects in the regulation of genes associated with apoptosis and autophagy in neuronal cells as observed in AD following an infection by *C. pneumoniae*

## Material and Methods

The SKNMC (ATCC) human neuroblastoma cell line were infected with ATCC's AR39 strain of *Chlamydia pneumoniae* at an MOI = 1 for 24, 48, and 72 hrs. Cells were immunolabeled with FITC-chlamydia antibody 61C75 (Fitzgerald) for verification of infection. The Human Autophagy RT<sup>2</sup>-Profiler<sup>TM</sup> PCR Array from SABiosciences was used to analyze the expression of a panel of genes involved in autophagy and apoptosis. All data was derived from experiments performed in triplicate.


## Results

RT-PCR of SKNMC neuronal cells infected with *C. pneumoniae* for 24, 48, and 72hrs



Verification of *C. pneumoniae* infection for 24, 48, and 72hrs by immunocytochemistry.

Verification of *C. pneumoniae* infection in neuronal cells used in RT-PCR gene arrays



## Conclusions

Modulation of genes associated with the autophagy and apoptosis pathways is observed in neuronal cells after 24, 48, and 72hrs of infection with *C. pneumoniae*. These genetic changes lead to dysfunction in the basic cellular processes of autophagy and apoptosis; consequently dysfunction in these processes has been shown to contribute to the neuropathology of late onset Alzheimer's disease. This work will allow future studies to further focus on the analysis of genes involved in the apoptotic and autophagic pathways to better understand how a pathogen such as *C. pneumoniae* plays a role in the development of AD.

## Funding

This work was funded by the Center for Chronic Disorders of Aging at the Philadelphia College of Osteopathic Medicine, the Adolph and Rose Levis Foundation for Alzheimer's Disease Research, Health Research/CURE Program, PA Department of Health and the undergraduate science education program grant awarded to Washington and Jefferson from the Howard Hughes Medical Institute and W&J College's 2008 Undergraduate Science Education Program Grant #5200632

	Function	Gene Symbol	Gene Name
Autophagy Machinery Components			
•Autophagic Vacuole Formation			
	AMBRA1	Autophagy/beclin-1 regulator 1	
	ATG4A	ATG4 autophagy related 4 homolog A (S. cerevisiae)	
	ATG4C	ATG4 autophagy related 4 homolog C (S. cerevisiae)	
	ATG9A	ATG9 autophagy related 9 homolog A (S. cerevisiae)	
	BECN1	Beclin 1, autophagy related	
	MAP1LC3B	Microtubule-associated protein 1 light chain 3 beta	
	ATG4D	ATG4 autophagy related 4 homolog D (S. cerevisiae)	
•Genes Linking Autophagosome to Lysosome			
	DRAM	Damage-regulated autophagy modulator	
	FAM176A	Family with sequence similarity 176, member A	
• Protein Transport			
	ATG10	ATG10 autophagy related 10 homolog (S. cerevisiae)	
	ATG16L1	ATG16 autophagy related 16-like 1 (S. cerevisiae)	
	RAB24	RAB24, member RAS oncogene family	
Co-Regulators of Autophagy and Apoptosis			
	BAK1	BCL2-antagonist/killer 1	
	BNIP3	BCL2/adenovirus E1B 19kDa interacting protein 3	
	CASP3	Caspase 3, apoptosis-related cysteine peptidase	
	FAS	Fas (TNF receptor superfamily, member 6)	
	NFKB1	Nuclear factor of kappa light polypeptide gene enhancer in B-cells 1	
Autophagy Induction by Intracellular Pathogens			
	IFNA2	Interferon, alpha 2	
	IFNA4	Interferon, alpha 4	
	IFNG	Interferon, gamma	
Autophagy in Response to Other Intracellular Signals			
	ARSA	Arylsulfatase A	
	EIF4G1	Eukaryotic translation initiation factor 4 gamma, 1	
	ESR1	Estrogen receptor 1	
	PIK3C3	Phosphoinositide-3-kinase, class 3	
	PIK3R4	Phosphoinositide-3-kinase, regulatory subunit 4	
	PRKAA2	Protein kinase, AMP-activated, alpha 2 catalytic subunit	
	TMEM74	Transmembrane protein 74	
	TMEM77	Transmembrane protein 77	
	ULK2	Unc-51-like kinase 2 (C. elegans)	
	UVRAG	UV radiation resistance associated gene	

## References

- [1] B. J. Balin, H. C. Gerard, E. J. Arking, D. M. Appelt, P. J. Branigan, J. T. Abrams, J. A. Whitum-Hudson, and A. P. Hudson. (1998). Identification and localization of *Chlamydia pneumoniae* in the Alzheimer's brain. *Med. Microbiol. Immunol.* 187, 23-42.
- [2] H. C. Gerard, U. Dreses-Werringloer, K. S. Wildt, S. Deka, C. Oszust, B. J. Balin, W. H. Frey 2nd, E. Z. Bordojo, J. A. Whitum-Hudson, and A. P. Hudson. (2006). Chlamydia (*Chlamydia pneumoniae*) in the Alzheimer's brain. *FEMS Immunol. Med. Microbiol.* 48, 355-66.
- [3] C. S. Little, C. J. Hammond, A. MacIntyre, B. J. Balin, and D. M. Appelt. (2004). *Chlamydia pneumoniae* induces Alzheimer-like amyloid plaques in brains of BALB/c mice. *Neurobiol. Aging* 25, 419-29.
- [4] D. M. Appelt, M. R. Roupas, D. S. Way, M. G. Bell, E. V. Albert, C. J. Hammond, and B. J. Balin. (2008). Inhibition of apoptosis in neuronal cells infected with Chlamydia (*Chlamydia pneumoniae*). *BMC Neurosci.* 9, 13.
- [5] Fischer SF, Schwarz C, Vier J, Häcker G. (2001). Characterization of antiapoptotic activities of *Chlamydia pneumoniae* in human cells. *Infection and Immunity* 69 (11), 7121-9.
- [6] S. F. Fischer and G. Hacker. (2003). Characterization of antiapoptotic activities of *Chlamydia pneumoniae* in infected cells. *Ann. N. Y. Acad. Sci.* 1010, 565-7.
- [7] S. F. Fischer, J. Vier, S. Kirschnek, A. Klos, S. Hess, S. Ying, and G. Hacker. (2004). Chlamydia inhibit host cell apoptosis by degradation of proapoptotic BH3-only proteins. *J. Exp. Med.* 200, 905-16.
- [8] Funderburk, S. F., Marcellino, B. K., & Yue, Z. (2010). Cell "Self-eating" (autophagy) mechanism in Alzheimer's disease. *Mount Sinai Journal of Medicine* 77(1), 59-68.
- [9] S. Lee, Y. Sato, and R. A. Nixon. (2011). Lysosomal Proteolysis Inhibition Selectively Disrupts Axonal Transport of Degradative Organelles and Causes an Alzheimer's-Like Axonal Dystrophy. *J. Neurosci.* 31, 7817-30.
- [10] Nixon, R. A., Cataldo, A. M., & Matthews, P. M. (2000). The endosomal-lysosomal system of neurons in Alzheimer's disease pathogenesis. *Neurochemical Research* 25, 1161-1172.