

Philadelphia College of Osteopathic Medicine DigitalCommons@PCOM

Scholarly Posters

2013

Analysis of autophagy and inflammasome regulation in neuronal cells and monocytes infected with *Chlamydia pneumoniae*: Implications for Alzheimer's disease

Brian J. Balin

Philadelphia College of Osteopathic Medicine, brianba@pcom.edu

Christine J. Hammond

Philadelphia College of Osteopathic Medicine, christineh@pcom.edu

Juliana Zoga

Philadelphia College of Osteopathic Medicine

Ahmad B. Cader

Philadelphia College of Osteopathic Medicine, ahmadca@pcom.edu

Annette K. Slutter

Philadelphia College of Osteopathic Medicine, AnnetteKSS@gmail.com

Follow up on additional works at: <http://digitalcommons.pcom.edu/posters>



Part of the [Medical Immunology Commons](#), [Nervous System Diseases Commons](#), and the [Neurology Commons](#)

Recommended Citation

Balin, Brian J.; Hammond, Christine J.; Zoga, Juliana; Cader, Ahmad B.; Slutter, Annette K.; Anzman, Jonathan M.; Kohler, Ian; Hingley, Susan T.; and Appelt, Denah M., "Analysis of autophagy and inflammasome regulation in neuronal cells and monocytes infected with *Chlamydia pneumoniae*: Implications for Alzheimer's disease" (2013). *Scholarly Posters*. Book 3.

<http://digitalcommons.pcom.edu/posters/3>

This Book is brought to you for free and open access by DigitalCommons@PCOM. It has been accepted for inclusion in Scholarly Posters by an authorized administrator of DigitalCommons@PCOM. For more information, please contact library@pcom.edu.

Authors

Brian J. Balin, Christine J. Hammond, Juliana Zoga, Ahmad B. Cader, Annette K. Slutter, Jonathan M. Anzman, Ian Kohler, Susan T. Hingley, and Denah M. Appelt

B. Balin, C. Hammond, J. Zoga, A. Cader, A. Slutter, J. Anzmann, I. Kohler, S. Hingley, D. Appelt
 Center for Chronic Disorders of Aging, Philadelphia College of Osteopathic Medicine,
 Philadelphia, PA, USA

Abstract

Objectives: Our laboratory has been studying the role of infection with the obligate intracellular bacterium *Chlamydia pneumoniae* in sporadic late-onset Alzheimer disease (LOAD). This infection may be a trigger for the pathology observed in LOAD as a function of initiating changes in gene regulation following entry of the organism into the brain. As such, we are analyzing how this infection can promote changes in autophagy and inflammasome gene regulation as both have been shown to be altered in LOAD.

Methods: Human SKNMC neuronal cells and THP1 monocytes were infected in vitro for 24-72 hrs with a laboratory strain of *Chlamydia pneumoniae* followed by RNA extraction, cDNA synthesis and analysis using Real-Time PCR microarrays for autophagy and inflammasome genes.

Results: Gene expression for autophagy and inflammasome pathways was altered dramatically following infection. Genes encoding for co-regulation of autophagy, apoptosis, and the cell cycle that were significantly changed included: BCL2L1, FAS, PIK3CG, APP, and TP53. In addition, ATG3, and GABARAP, genes encoding for protein transport & ubiquitination and autophagic vacuole formation were significantly deregulated. Of the inflammasome genes, 4 NOD-like receptor genes were significantly up-regulated. IL-1beta, AIM2, CCL2, and CCL7 genes were all dramatically up-regulated in monocytes during the 72 hrs of infection.

Conclusions: Our data suggest that *Chlamydia pneumoniae*-infected human SKNMC neuronal cells and THP1 monocytes exhibit specific changes in gene regulation for both autophagy and inflammasome pathways. These gene changes appear to correlate with pathologic changes previously reported in AD and further support the contention that infection with *Chlamydia pneumoniae* plays a role in LOAD pathogenesis.

Introduction

Neurodegeneration has been well documented in the CNS of Alzheimer individuals. Strong evidence suggests that abnormalities of autophagy and apoptosis pathways as well as activation of inflammasomes are contributing factors in Alzheimer's disease (AD) pathogenesis. Our laboratory has focused on infection with *Chlamydia pneumoniae* (Cpn) as a risk factor/co-susceptible agent in LOAD. Cpn is an obligate, intracellular, parasitic bacterium. Cpn is transmitted from person to person via respiration. Once inhaled, Cpn may enter the brain along 2 pathways, directly through olfaction and/or blood-borne in monocytes. In studies of AD brain tissues, we have identified Cpn in areas of neuropathology by PCR and immunohistochemistry. Cpn was detected in 80 to 90% of post-mortem LOAD brain samples, but only in 5-11% of brains from age-matched non-AD controls (Balin et al., 1998; Gerard et al., 2006; Hammond et al., 2010). Infected glia, perivascular macrophages, monocytes, and neurons have been observed in the AD brain.

Infection may result in early neuroinflammation and neuronal damage in specific vulnerable regions of the brain (Balin et al., 1998). In analyzing cellular changes following infection, we have demonstrated that Cpn can inhibit apoptosis in neuronal cells thereby prolonging the viability of the infected cells (Appelt et al., 2008). Other laboratories have demonstrated that *chlamydiae*-infected host cells are resistant to proapoptotic stimuli such as TNF α , Fas antibody, staurosporine, and UV-light (Fischer et al., 2004). Further, as Cpn is an **intracellular bacterium**, we have started investigating autophagy and inflammasome activation of the host cell as these mechanisms are commonly employed by eukaryotic cells to eradicate intracellular organisms.

Autophagy and apoptosis are common pathways by which infected cells attempt to rid themselves of an infectious agent and cells incapable of eliminating the infectious agent undergo cell death. Autophagy is associated with the endosomal-lysosomal system. The endosomal-lysosomal system is linked to the lysosomal system as early endosomes fuse with late endosomes or lysosomes. Contents of an autophagosome are degraded as a result of fusing with lysosomes (Fischer et al., 2004; Funderburk et al., 2010). An increase in the number of autophagic vacuoles (AVs) has been identified in neurons from AD brains implicating autophagy as a pathological process in AD (Lee et al., 2011). Neurons from AD brains do have enlarged early endosomes which is significant because early endosomes take in proteins such as apolipoprotein E and APP, and A β has been demonstrated to be formed in early endosomes (Nixon et al., 2011).

An additional prominent feature in AD is neuroinflammation (Akiyama et al., 2000). Recent evidence implicates the pro-inflammatory process with production of IL-1 β in mild cognitive impairment and early AD (Agostinho et al., 2010). A proinflammatory signal typically follows from an infection leading to NF- κ B activation and synthesis of pro-IL-1 β . A second signal activates caspase-1, which cleaves pro-IL-1 β into its mature form, IL-1 β . Caspase-1 is derived from pro-caspase-1 following cleavage by a multiprotein complex called an inflammasome (He et al., 2010). The inflammasome contains three proteins, caspase-1, apoptosis-associated speck-like protein containing a caspase recruitment domain (ASC), and a nucleotide-binding oligomerization domain-like receptor (NLR). When an inflammatory response is needed, these three proteins will aggregate in order to cleave pro-caspase-1 and initiate the inflammatory response (He et al., 2010).

One specific inflammasome, NLRP-3, when activated produces IL-1 β in response to various fungal, viral, and/or bacterial infections including those caused by Cpn (He et al., 2010). *Chlamydia* utilize a type III secretion system to secrete virulence factors into the host cell cytosol to control intracellular reactions. These factors cause K⁺ efflux and formation of reactive oxygen species. This rise in reactive oxygen species is sufficient to initiate assembly of the NLRP-3 inflammasome (Abdul-Sater et al., 2009).

How Cpn infection affects autophagy and inflammasome gene expression in eukaryotic cells is important for understanding the role that infection plays in initiating acute damage and eventual chronic inflammatory responses resulting in AD pathogenesis.

Materials and Methods

Cell lines - Human SK-N-MC neuronal cells and THP1 monocytes obtained from the ATCC were used in these studies.

Infection with *Chlamydia pneumoniae* (Cpn)

The respiratory laboratory strain of Cpn, AR-39, at a multiplicity of infection of 1, was used for all infections for 24, 48, and 72hrs. For the neuronal cells, Cpn was added to a subconfluent monolayer followed by centrifugation at 800 rpm for 5 min and incubated for the allotted times. Monocytes were centrifuged, washed, and resuspended, followed by the addition of Cpn and incubated for the same time periods. Parallel uninfected control cells were grown under the same conditions for the times indicated.

RNA Isolation and First Strand Synthesis

RNA was extracted using the RNeasy Plus Mini kit from Qiagen, followed by cDNA production from RNA (1 μ g) using the RT² First Strand Kit from SABiosciences (Qiagen), all following manufacturer's directions.

Real Time - Polymerase Chain Reaction (RT-PCR)

RT-PCR for gene transcription used Autophagy (PAHS-084A) and Inflammasome (PAHS-097A) PCR Arrays from SABiosciences (Qiagen). Arrays were run on the ABI Prism 7000 Sequence Detection System from Applied Biosciences. The results were analyzed using software from SABiosciences (www.sabiosciences.com/pcrarraydataanalysis.php). Arrays were run for Cpn infected SK-N-MC neuronal cells, THP1 monocytes, and their uninfected controls at 24, 48, and 72hrs with each experiment performed in triplicate.

Data Analysis

For autophagy genes, ** significance of $p \leq 0.05$ and * significance of $p \leq 0.10$.

For inflammasome genes, only those greater than 4-fold change with significance of $p \leq 0.05$ are represented on the charts.

References

Abdul-Sater AA, Said-Sadier N, Ojcius DM, Yilmaz O, Kelly KA. (2009). Inflammasomes bridge signaling between pathogen identification and the immune response. *Drugs Today (Barc)*. 45 Suppl B:105-112.

Agostinho P, Cunha RA, Oliveira C. (2010). Neuroinflammation, oxidative stress and the pathogenesis of Alzheimer's disease. *Curr Pharm Des*. 16(25):2766-2778.

Akiyama H, Baegler S, Baum S, Brad B, Sauer J et al. (2000). Inflammation and Alzheimer's disease. *Neurobiol Aging*. 21(3):383-421.

Appelt DM, Roupas MR, Way DS, Bell MG, Albert EV, Hammond CJ, and Balin BJ. (2008). Inhibition of apoptosis in neuronal cells infected with *Chlamydia pneumoniae*. *BMC Neurosci*. 9:13.

Balin BJ, Gerard HC, Arking EJ, Appelt DM, Branigan FJ, Abrams JT, Whitlum-Hudson JA, and Hudson AP. (1998). Identification and localization of *Chlamydia pneumoniae* in the Alzheimer's brain. *Med. Microbiol. Immunol*. 187:23-42.

Duque L, Wirawan E, Vanden Berghe T, Vandendriessche P. (2009). Major cell death pathways at a glance. *Mol Biol Cell*. 20(11):3100-3104. Epub 2009 Sep 4.

Fischer SP, Vier J, Kirschnek S, Rios A, Hess S, Ying S, and Fischer G. (2004). *Chlamydia* inhibit host cell apoptosis by degradation of proapoptotic Bcl-2 only proteins. *J. Exp. Med*. 200, 905-16.

Funderburk SF, Marcelino BK, and Yue Z. (2010). Cell "Self-eating" (autophagy) mechanism in Alzheimer's disease. *Mount Sinai Journal of Medicine*. 77(1), 59-68.

Gerard HC, Dreses-Werringloer U, Wildt KS, Dekas S, Ozusut C, Balin BJ, Frey WH, 2nd, Bortdayo EZ, Whitlum-Hudson JA, Hudson AP (2006). *Chlamydia pneumoniae* in the Alzheimer's brain. *FEMS Immunol Med Microbiol*. 48(3):355-365.


Hammond CJ, Hallock LR, Howanski RJ, Appelt DM, Little CS, Balin BJ. (2010). Immunohistochemical detection of *Chlamydia pneumoniae* in the Alzheimer's disease brain. *BMC Neurosci*. 11:121.

He X, Mekasha S, Mavrogorgos N, Fitzgerald KA, Lien E, Ingalls RR (2010). Inflammation and fibrosis during *Chlamydia pneumoniae* infection is regulated by IL-1 and the NLRP3/ASC inflammasome. *J. Immunol*. 184(10):5743-5754.

Lee S, Sato Y, and Nixon RA. (2011). Lysosomal Proteolysis Inhibition Selectively Disrupts Axonal Transport of Degradative Organelles and Causes an Alzheimer's-Like Axonal Dystrophy. *J. Neurosci*. 31: 7817-30.

Nixon RA and Yang DS. (2011) Autophagy failure in Alzheimer's disease—locating the primary defect. *Neurobiol Dis*. 43(1):38-45.

Autophagy




Autophagy Gene Regulation

Gene Symbol	Gene Name
ATG3	ATG3 autophagy related 3 homolog (S. cerevisiae)
ATG4B	ATG4 autophagy related 4 homolog B (S. cerevisiae)
ATG4C	ATG4 autophagy related 4 homolog C (S. cerevisiae)
BECN1	Becn1
DRAM1	DNA-damage regulated autophagy modulator 1
GABARAP	GABA(A) receptor-associated protein
MAP1LC3A	Microtubule-associated protein 1 light chain 3 alpha
MAP1LC3B	Microtubule-associated protein 1 light chain 3 beta

Co-Regulation of Autophagy, Apoptosis, and the Cell Cycle


Gene Symbol	Gene Name
AKT1	V-akt murine thymoma viral oncogene homolog 1
APP	Amyloid beta (A β) precursor protein
BAD	BCL2-associated agonist of cell death
BAX	BCL2-associated X protein
BCL2L1	B-cell CLL/lymphoma 2-like 1
CASP3	Caspase 3, apoptosis-related cysteine peptidase
CASP8	Caspase 8, apoptosis-related cysteine peptidase
EIF2AK3	Eukaryotic translation initiation factor 2-alpha kinase 3
FAS	Fas (TNF receptor superfamily, member 6)
IFNA2	Interferon 2
IFNA4	Interferon 4
IFNG	Interferon, gamma
PIK3CG	Phosphoinositide-3-kinase, catalytic, gamma polypeptide
TP53	Tumor protein p53

- Genes encoding for co-regulation of autophagy, apoptosis and the cell cycle
- significantly deregulated in Cpn-infected neuronal cells
 - BCL2L1, FAS, PIK3CG
 - Changed in AD
 - Up-regulated significantly upon initiation of infection
 - APP, TP53
 - Changed in AD
 - Down-regulated over length of infection




Inflammasome Gene Regulation

PCR array genes with at least a 4-fold.




Inflammasome Gene Regulation

Cpn Infected THP1 Monocytes




Inflammasome Gene Regulation

Cpn Infected THP1 Monocytes



Inflammasome Gene Regulation

Cpn Infected THP1 Monocytes



- Genes encoding for inflammasomes are up-regulated in Cpn-infected THP1 monocytes
- Up-regulation of inflammasome genes may lead to pro-inflammatory cytokine and chemokine increase
- Three inflammasome genes (AIM2, NAI1, and NLR4) were significantly changed

- IL6, NOD2, CCL7, and CCL2 are up-regulated in infected monocytes
- Up-regulated in AD
- Enormous up-regulation of CCL2 ultimately leads to
 - recruitment of monocytes, dendritic cells, and T cells
 - activation of the innate immune response

Our study provides a mechanistic foundation for infection as an initiator and propagator in the pathogenesis of Alzheimer's disease