



**Nota científica**  
(*Short communication*)

**FIRST REPORT OF *DIGITONTHOPHAGUS GAZELLA* (FABRICIUS, 1787) AND NEW RECORDS FOR *EUONITICELLUS INTERMEDIUS* (REICHE, 1849) (COLEOPTERA: SCARABAEIDAE LATREILLE, 1802) IN EL SALVADOR**

**PRIMER REGISTRO DE *DIGITONTHOPHAGUS GAZELLA* (FABRICIUS, 1787) Y NUEVO REGISTRO PARA *EUONITICELLUS INTERMEDIUS* (REICHE, 1849) (COLEOPTERA: SCARABAEIDAE LATREILLE, 1802) EN EL SALVADOR**

**JOSÉ D. PABLO-CEA,<sup>1,\*</sup> MARYORY A. VELADO-CANO,<sup>2</sup> RENÉ FUENTES,<sup>1</sup> MELVIN CRUZ,<sup>3</sup> & JORGE ARI NORIEGA<sup>4,5</sup>**

<sup>1</sup> Escuela de Biología, Universidad de El Salvador. <pcea\_coleoptera@hotmail.com>; <refumos@yahoo.com>.

<sup>2</sup> Independent researcher, San Salvador, El Salvador. <mcano\_balaenoptera@hotmail.com>.

<sup>3</sup> Independent researcher, Chalatenango, El Salvador. <melvinpresent@gmail.com>.

<sup>4</sup> Laboratorio de Zoología y Ecología Acuática-LAZOEA, Universidad de Los Andes, Bogotá, Colombia.  
<jnorieg@hotmail.com>.

<sup>5</sup> Department of Biogeography and Global Change, National Museum of Natural Science (CSIC), Madrid, Spain.

\* Corresponding autor: <pcea\_coleoptera@hotmail.com>.

Recibido: 13/02/2017; aceptado: 30/08/2017.

Editor responsable: Cuauhtémoc Deloya.

**Pablo-Cea J. D., Velado-Cano M. A., Fuentes R., Cruz M., & Noriega J. A. (2017) First report of *Digitonthophagus gazella* (Fabricius, 1787) and new records for *Euoniticellus intermedius* (Reiche, 1849) (Coleoptera: Scarabaeidae Latreille, 1802) in El Salvador. *Acta Zoológica Mexicana* (n.s), 33(3), 527-531.**

**ABSTRACT.** The presence of the beetle *Digitonthophagus gazella* (Fabricius, 1787) is reported for the first time. In addition, the range of distribution of *Euoniticellus intermedius* (Reiche, 1849) in El Salvador is expanded, based on information obtained from collections made over twelve years and the current distribution map of *E. intermedius* in the territory is presented. In addition, the current situation of both species in the country and Central America is briefly discussed.

The introduction, dispersion and colonization process carried out by the African invasive dung species *Digitonthophagus gazella* (Fabricius, 1787) and the Afro-tropical *Euoniticellus intermedius* (Reiche, 1849), has been widely studied in Mexico (Rivera-Cervantes & García-Real, 1991; Kohlmann, 1994; Morales *et al.*, 2004), Central America (Maes *et al.* 1997; Solís & Kohlmann, 2012; Rivera & Schlein, 2014; Solís *et al.*, 2015) and South America (Noriega, 2002; Noriega *et al.*, 2006; Rivera &

**Pablo-Cea J. D., Velado-Cano M. A., Fuentes R., Cruz M. y Noriega J. A. (2017) Primer registro de *Digitonthophagus gazella* (Fabricius, 1787) y nuevo registro para *Euoniticellus intermedius* (Reiche, 1849) (Coleoptera: Scarabaeidae Latreille, 1802) en El Salvador. *Acta Zoológica Mexicana* (n.s), 33(3), 527-531.**

**RESUMEN.** Se reporta por primera vez la presencia del escarabajo *Digitonthophagus gazella* (Fabricius, 1787). Además, se amplía el rango de distribución de *Euoniticellus intermedius* (Reiche, 1849) en El Salvador, a partir de información obtenida de colectas realizadas a lo largo de doce años y se presenta el mapa de distribución actual de *E. intermedius* en el territorio. Adicionalmente, se discute brevemente la situación actual de ambas especies en el país y Centroamérica.

Wolff, 2007; Vidaurre *et al.*, 2008; Álvarez *et al.*, 2009; Noriega *et al.*, 2010; 2017) since their initial liberation in the United States of America (Fincher *et al.*, 1983; Wood & Kaufman, 2008). Recently, the taxonomy of the genus *Digitonthophagus* has been clarified (Génier & Moretto, 2017), showing that the genus had several species unknown to science and that the identity of the invasive species in the American continent must continue being *D. gazella* (Génier & Davis, 2017). In this note, we report

for the first time the presence of *D. gazella* in El Salvador, Central America and we extend the geographic distribution range of *E. intermedius* in the country.


Cara Sucia is a town located in the municipality of San Francisco Menéndez, Ahuachapán Department, in the Southwest of El Salvador. Cara Sucia presents high mercantile activity due mainly to its strategical location between the highway to La Hachadura (at the Guatemala and El Salvador border) and the urban center of the Sonsonate Department, which facilitates commercial trade between both regions. We collected the specimens on two cattle pastures on the outskirts of Cara Sucia (grassland #1: 13°47'06.1" N - 90°02'30.5" W, 15 m a.s.l. and grassland #2: 13°46'09.2" N - 90°03'25.6" W, 7 m a.s.l.) (Fig. 1), using 20 pitfall traps baited with cow dung on July 2016, as part of the project: "Dung Removal Rates in Grasslands around the World", coordinated by the last author.

We captured 128 beetles, of which 65 were *D. gazella* (Fig. 2a), providing the first record for this species in the country, and 17 of them were *E. intermedius* (Fig. 2b), extending this species' geographical range in El Salvador. Additionally, we gathered all the records of *E. intermedius* made by the authors during the past twelve years and a map of the current countrywide distribution of the species is presented (Fig. 1). In the traps, 10 individuals of *Copris*


*lugubris* Boheman, 1858 and 36 individuals of *Ataenius sculptor* Harold, 1868 were also captured.

First evidence for the arrival of *D. gazella* in Central America comes from Northern Guatemala in 1987 (Kohlmann, 1994). The first recorders of the species in Mexico (Barbero & López-Guerrero, 1992; Kohlmann, 1994) provide estimations for their average dispersal distance. For instance, Barbero & López-Guerrero (1992) estimated that *D. gazella* was able to travel between 16.5 and 800 km/y (177.81 km/y on average) from the South of the USA and the North of Mexico to Central America. Kohlmann (1994) estimated the average distance that beetles traveled, both, in the Atlantic (103 km/y) and in the Pacific regions (220 km/y). His observations suggested that the beetles seemed to be moving faster in the Pacific region, probably due to more favorable climatic conditions to the dung beetle (Kohlmann, 1994).

In the Pacific coast, the Southernmost distribution area of the species, reported by Kohlmann (1994) was Oaxaca (Mexico), where individuals of the species were collected in 1989, approximately 815 km away from the Ahuchapán Department in El Salvador. Following Kohlmann's estimations (220 km/y), if *D. gazella* had travelled at a steady pace, the species would have arrived in El Salvador by late 1993.



**Figure 1.** Map of the localities registered for current distribution of *Eouniticellus intermedius* in El Salvador. 2003: Tonacatepeque, San Salvador; 2005: El Majahual, La Libertad; 2008: Laguna Caldera, La Libertad; 2012: Colima, Cuscatlán; 2014: Tehuacán, San Vicente; 2016: Cara Sucia, Ahuachapán.



**Figure 2.** a) *Digithonthophagus gazella* and b) *Euoniticellus intermedius* collected in 2016 at Cara Sucia, El Salvador.

In the period between 1997 and 1999, numerous dung beetle surveys were carried out in several zones of the country, using pitfall traps baited with horse and cow dung in a variety of cattle pastures, including areas near to the current collection site, where the presence of the species was not recorded (Horgan, 2001; Horgan, 2002; Horgan & Fuentes, 2005). Fourteen years later, in 2013 a survey was carried out in the National Park (NP) El Impossible, a forest area located at less than 20 km from the current collection site, without any evidence of the species' presence (Pablo-Cea *et al.*, 2016). It is important to note that the altitudinal difference between these two regions can reach almost 900 m in some areas; however it doesn't exceed the altitudinal range limit reported for the species in America (Vidaurre *et al.*, 2008; Álvarez *et al.*, 2009; Noriega *et al.*, 2011). It is also important to mention that nor horse neither cow dung was used as bait in the NP El Impossible, however, the first author examined horse dung present around the area, during the whole period of the survey (June to November 2013) without successfully registering the presence the species.

It is very complicated to estimate the exact moment of arrival of this species in the country, since there hadn't been an active national monitoring program in past years. The entrance of *D. gazella* in El Salvador could be related to the cattle trade between Cara Sucia and adjacent areas in Guatemala (M. Cruz, personal observation), coming in from the neighboring country due to anthropogenic action.

*Euoniticellus intermedius* was collected for the first time by Fuentes in 2003 (Solís & Kohlmann, 2012) in Tona-catepeque, San Salvador ( $13^{\circ}48'7.95''$  N -  $89^{\circ}7'51.74''$  W, 525 m a.s.l.). In 2005, a specimen was found in El Majahual Beach in La Libertad ( $13^{\circ}29'19.81''$  N -  $89^{\circ}21'44.23''$  W, 5 m a.s.l.) (Fuentes, unpublished data). In 2008, specimens were collected near Laguna Caldera, also in La Libertad ( $13^{\circ}50'26.28''$  N -  $89^{\circ}21'26.17''$  W, 450 m a.s.l.) (Fuentes, 2008). In 2012, Pablo-Cea and Vellado-Cano (unpublished data) collected 112 specimens in pastures around the Natural Protected Area of Colima in Cuscatlán ( $14^{\circ}02'44.4''$  N -  $89^{\circ}07'39.57''$  W, 300 m a.s.l.), where the species displayed dominance (in number of individuals) over the native species populations, and even a few specimens were collected inside the forest areas (an uncommon habitat for this species that prefers open areas). In 2014, Fuentes (unpublished data) collected specimens in pastures near the eco-touristic park Tehuacán in San Vicente ( $13^{\circ}33'25.55''$  N -  $88^{\circ}47'1.01''$  W, 378 m a.s.l.). Tehuacán is considered the easternmost point of the species' distribution in the country, while the specimens collected in Cara Sucia represent the first record of the species' presence in the Western region of the country. Despite of that, we consider the west region, the one where *E. intermedius* got access to El Salvador, due to its location near Guatemala, where the species was collected in 2002, a year before Fuentes captured the first specimens in El Salvador (Solís & Kohlmann, 2012).

Of both invasive species, *E. intermedius* shows a higher adaptability and colonization potential. It has invaded faster into the national territory, presumably facilitated by its higher plasticity that has allowed it to adapt to the conditions and maintain relatively stable populations, during the dry season (Fuentes, 2008). As a result, the species was already found in the central El Salvador in 2002, it has colonized several areas of the country (from 0 to 525 m) and several more countries in Central America (Solís & Kohlmann, 2012; Rivera & Schlein, 2014; Solís *et. al.*, 2015). On the other hand, *D. gazella* is restricted to the Southwestern area of El Salvador. Its exact date of arrival has not been confirmed and its presence has not been reported yet in Belize, Honduras, Costa Rica, nor Panama.

To conclude, we stress the importance of tracking geographical distributions of invasive species in El Salvador, where is common to find cattle pastures surrounding natural protected areas, which can facilitate the potential entrance of foreign species into the small and isolated forest patches which still remain. This creates a potential risk for dispersing invasive species to outcompete and even displace the local native species (as seen for example, in the surroundings of Colima). Therefore, we recommend the creation and establishment of a National Monitoring Network to carry out more extensive surveys in an effort to cover more areas of the country, and to follow the current and future distribution of these and other potential invasive species occurring in El Salvador.

**ACKNOWLEDGMENTS.** We are in debt with Mr. José Domingo Cruz Franco for allowing us access to his cattle pastures to collect beetles and for all the logistic support he gave us to that end. We thank David Morris and Javier Santos for kindly checking the English version of the manuscript, and to Berth Kohlmann and an anonymous reviewer for invaluable comments to make this manuscript better. JAN has been supported by a Colombian COLCIENCIAS PhD scholarship.

## LITERATURE CITED

- Álvarez, M. C., Damborsky, M. P., Bar, M. E. & Ocampo, F. C. (2009). Registros y distribución de la especie afroasiática *Digitonthophagus gazella* (Coleoptera: Scarabaeidae: Scarabaeinae) en Argentina. *Revista de la Sociedad Entomológica Argentina*, 68(3-4), 373-376.
- Barbero, E. & López-Guerrero, Y. (1992). Some considerations on the dispersal power of *Digitonthophagus gazella* (Fabricius, 1787) in the New World (Coleoptera: Scarabaeidae Scarabaeinae). *Tropical Zoology*, 5, 115-120.
- Fincher, G. T., Stewart, T. B. & Hunter III, J. S. (1983). The 1981 distribution of *Onthophagus gazella* Fabricius from releases in Texas and *Onthophagus taurus* Schreber from an unknown release in Florida (Coleoptera: Scarabaeidae). *The Coleopterists Bulletin*, 37, 159-163.
- Fuentes, R. (2008). Abundancia y estacionalidad del escarabajo *Euoniticellus intermedius* (Reiche, 1849) Coleoptera: Scarabaeidae, Tonacatepeque, San Salvador, El Salvador. XXXV Congreso So-colen: Sociedad Colombiana de Entomología. Universidad Nacio-nal de Colombia.
- Génier, F. & Davis, A. (2017). *Digitonthophagus gazella* auctorum: an unfortunate case of mistaken identity for a widely introduced species (Coleoptera: Scarabaeidae: Scarabaeinae: Onthophagini). *Zootaxa*, 4221(4), 497-500.
- Génier, F. & Moretto, P. (2017). *Digitonthophagus* Balthasar, 1959: taxonomy, systematics, and morphological phylogeny of the genus revealing an African species complex (Coleoptera: Scarabaeidae: Scarabaeinae). *Zootaxa*, 4248 (1), 001-110.
- Horgan, F. (2001). Burial of bovine dung by coprophagous beetles (Coleoptera: Scarabaeidae) from horse and cow grazing sites in El Salvador. *European Journal of Soil Biology*, 37, 103-111.
- Horgan, F. (2002). Shady field boundaries and the colonization of dung by coprophagous beetles in Central American pastures. *Agriculture, Ecosystems and Environment*, 91, 25-36.
- Horgan, F. & Fuentes, R. (2005). Asymmetrical competition between Neotropical dung beetles and its consequences for assem-blage structure. *Ecological Entomology*, 30, 182-193.
- Kohlmann, B. (1994). A preliminary study of the invasion and dispersal of *Digitonthophagus gazella* (Fabricius, 1787) in Mexico (Coleoptera; Scarabaeidae: Scarabaeinae). *Acta Zoológica Mexicana* (n.s.) 61, 35-42.
- Morales, C. J., Ruiz, R. & Delgado, L. (2004). Primer registro de *Euoniticellus intermedius* (Reiche, 1849) y datos nuevos de dis-tribución de *Digitonthophagus gazella* (Fabricius, 1787) (Coleop-tera: Scarabaeidae) e *Hybosorus illigeri* Reiche, 1853 (Coleoptera: Hybosoridae) para el estado de Chiapas. *Dugesiana*, 11, 21-23.
- Noriega, J. A. (2002). First report of the presence of the genus *Digi-tonthophagus* (Coleoptera: Scarabaeidae) in Colombia. *Caldasia*, 24, 213-215.
- Noriega, J. A., Solís, C., Quintero, I., Pérez, L. G., García, H. G. & Ospino, D. A. (2006). Registro continental de *Digitonthophagus gazella* (Coleoptera: Scarabaeidae) en Colombia. *Caldasia*, 28, 379- 381.
- Noriega, J. A., Horgan, F., Larsen, T. & Valencia, G. (2010). Re-cords of an invasive dung beetle species, *Digitonthophagus gazella* (Fabricius, 1787) (Coleoptera: Scarabaeidae), in Peru. *Acta Zoológica Mexicana* (n.s.), 26, 451-456.
- Noriega, J. A., Moreno, J. & Otavo, S. (2011). Quince años del arri-bo del escarabajo coprófago *Digitonthophagus gazella* (Fabricius, 1787) (Coleoptera: Scarabaeidae) a Colombia: proceso de inva-sión y posibles efectos de su establecimiento. *Biota Colombiana*, 12, 35-43.
- Noriega, J. A., Delgado, O., Blanco, J. I., Gámez, J. & Clavijo, J. (2017). Introduction, establishment, and invasion of *Digiton-thophagus gazella* (Fabricius, 1787) (Coleoptera: Scarabaeinae) in the savannas of Venezuela and Colombia. *Natural Resources*, 8, 370-381.
- Pablo-Cea, J. D., Velado-Cano, M. A. & Solís, A. (2016). Primer reporte de cuatro especies de escarabajos coprófagos (Coleoptera:



- Scarabaeinae) en El Salvador. *Acta Zoológica Mexicana (n.s.)* 32, 218-221.
- Rivera-Cervantes, L. E. & García-Real, E.** (1991). New locality records for *Onthophagus gazella* Fabricius (Coleoptera: Scarabaeidae) in Jalisco, Mexico. *Coleoperists Bulletin*, 45, 370.
- Rivera, J. D. & Schlein, O.** (2014). Primer registro de *Euoniticellus intermedius* (Reiche, 1849) y *Sulcophanaeus noctis cupricollis* (Nevinson, 1891) (Coleoptera: Scarabaeidae) para Honduras. *Entomotropica*, 29, 187-190.
- Rivera, C. & Wolff, M.** (2007). *Digitonthophagus gazella* (Coleoptera: Scarabaeidae): distribución en América y dos nuevos registros para Colombia. *Revista Colombiana de Entomología*, 33, 190-192.
- Solís, A. & Kohlmann, B.** (2012). A checklist and distribution atlas of the Scarabaeinae (Coleoptera: Scarabaeidae) of Costa Rica. *Zootaxa*, 3482, 1-32.
- Solís, A., Cambra, R. A. & González, M.** (2015). Primer registro para Panamá de la especie invasiva *Euoniticellus intermedius* (Reiche, 1849) (Coleoptera: Scarabaeidae, Scarabaeinae). *Tecnociencia*, 17, N° 2.
- Vidaurre, T., Noriega, J. A. & Ledezma, M. J.** (2008). First report on the distribution of *Digitonthophagus gazella* (Fabricius, 1787) (Coleoptera: Scarabaeidae) in Bolivia. *Acta Zoológica Mexicana (n.s.)* 24, 217-220.
- Wood, L. & Kaufman, P.** (2008). *Euoniticellus intermedius* (Coleoptera: Scarabaeidae: Scarabaeinae: Tribe Coprini): its presence and relative abundance in cattle pastures in northcentral Florida. *Florida Entomologist*, 91, 128-130.