

Philadelphia College of Osteopathic Medicine
DigitalCommons@PCOM

PCOM Scholarly Papers

2010

Dopamine D1 receptor-mediated enhancement of NMDA receptor trafficking requires rapid PKC-dependent synaptic insertion in the prefrontal neurons

Yan-Chun Li

Gang Liu

Jian-Li Hu

Wen-Jun Gao

Yueqiao Huang

Philadelphia College of Osteopathic Medicine, yueqiaohu@pcom.edu

Follow this and additional works at: http://digitalcommons.pcom.edu/scholarly_papers

 Part of the [Neuroscience and Neurobiology Commons](#)

Recommended Citation

Li, Yan-Chun; Liu, Gang; Hu, Jian-Li; Gao, Wen-Jun; and Huang, Yueqiao, "Dopamine D1 receptor-mediated enhancement of NMDA receptor trafficking requires rapid PKC-dependent synaptic insertion in the prefrontal neurons" (2010). *PCOM Scholarly Papers*. Paper 1131.

http://digitalcommons.pcom.edu/scholarly_papers/1131

This Article is brought to you for free and open access by DigitalCommons@PCOM. It has been accepted for inclusion in PCOM Scholarly Papers by an authorized administrator of DigitalCommons@PCOM. For more information, please contact library@pcom.edu.

Dopamine D₁ receptor-mediated enhancement of NMDA receptor trafficking requires rapid PKC-dependent synaptic insertion in the prefrontal neurons

Yan-Chun Li,¹ Gang Liu,¹ Jian-Li Hu,¹ Wen-Jun Gao and Yue-Qiao Huang

Department of Neurobiology and Anatomy, Drexel University College of Medicine, Philadelphia, Pennsylvania, USA

Abstract

Understanding the interaction between dopamine and glutamate, particularly the interaction of dopamine and NMDA receptors, may enable a more rational approach to the treatment of schizophrenia, drug addiction, and other psychiatric disorders. We show that, in prefrontal cortical neurons, dopamine D₁-induced enhancement of NMDA receptor function depends on rapid insertion of new NMDA receptor 2B subunits on the synaptic surface. Protein kinase A (PKA) inhibitor, but not protein kinase C (PKC) inhibitor, completely blocked dopamine D₁ agonist SKF-81297-induced increase of the total expression of NMDA receptors. Furthermore, SKF-81297 failed to alter the surface expression and synaptic

insertion of NMDA receptors in the presence of PKA inhibitor, phospholipase C inhibitor, PKC inhibitor, or Src family kinase inhibitor. Our data suggest that D₁-mediated enhancement of NMDA current depends on the NMDA receptor trafficking through rapid synaptic insertion and both PKA and PKC signaling pathways play important roles in the regulatory process. Although both PKA and PKC mediate the D₁-induced enhancement of NMDA receptors, the phospholipase C-PKC-Src pathway is only required for surface expression and new synaptic insertion of NMDA receptors.

Keywords: D₁ receptors, dopamine, glutamate, NMDA receptors, prefrontal cortex, psychiatric disorders.

J. Neurochem. (2010) **114**, 62–73.

The prefrontal cortex (PFC) receives extensive dopaminergic inputs and is particularly impaired in schizophrenia (Goldman-Rakic *et al.* 2004). At the subcellular level, dendritic spines enriched with D₁ and NMDA receptors in PFC pyramidal neurons are common targets of both glutamatergic and dopaminergic axons and thus providing the cellular basis for dopamine (DA)-glutamate interactions (Cepeda and Levine 2006). DA functions are mediated by D₁ class (D₁/D₅) and D₂ class (D₂, D₃ and D₄) G-protein coupled receptors (Missale *et al.* 1998). Previous studies indicated that NMDA receptor-mediated current is differentially regulated by dopamine receptors, with enhancement by D₁ (Cepeda *et al.* 1998; Seamans *et al.* 2001; Flores-Hernandez *et al.* 2002) and suppression by D₂ (Zheng *et al.* 1999; Li *et al.* 2009). The D₁-mediated enhancement of NMDA receptors was observed in various brain regions (Chergui and Lacey 1999; Wang and O'Donnell 2001; Chen *et al.* 2004; Hallett *et al.* 2006; Gao and Wolf 2008; Tong and Gibb 2008; Varela *et al.* 2009). Despite the overwhelming evidence regarding D₁ and NMDA receptors in synaptic functions, the molecular mechanisms underlying the D₁

modulation of NMDA receptors remain elusive, particularly in prefrontal neurons.

It is widely accepted that D₁ receptors are positively coupled to Gs to regulate NMDA receptors via cAMP-protein kinase A (PKA)-DARPP-32 (dopamine- and cyclic AMP-regulated phosphoprotein of 32 kDa) cascade (Blank *et al.* 1997; Cepeda *et al.* 1998; Snyder *et al.* 1998;

Received January 20, 2010; revised manuscript received March 23, 2010; accepted March 24, 2010.

Address correspondence and reprint requests to Wen-Jun Gao or Yue-Qiao Huang, Department of Neurobiology and Anatomy, Drexel University College of Medicine, 2900 W. Queen Lane, Philadelphia, PA 19129, USA. E-mail: wgao@drexelmed.edu or yueqiao.huang@drexelmed.edu

¹These authors contributed equally to this study.

Abbreviations used: AMPA, α -amino-3-hydroxy-5-4-propionate; DA, dopamine; DARPP-32, dopamine- and cyclic AMP-regulated phosphoprotein of 32 kDa; EPSC, excitatory postsynaptic current; NBQX, 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[f]quinoxaline-2,3-dione; NR, NMDA receptor; PFC, prefrontal cortex; PKA, protein kinase A; PKC, protein kinase C; PLC, phospholipase C; PP2, 4-amino-5-(4-chlorophenyl)-7-(t-butyl)pyrazolo[3,4-d]pyrimidine; TTX, tetrodotoxin.

Greengard 2001; Flores-Hernandez *et al.* 2002; Seamans and Yang 2004). However, emerging evidence indicates that D₁ can also activate other types of G proteins, thereby activating Ca²⁺ and protein kinase C (PKC) (Chergui and Lacey 1999; Chen *et al.* 2004; Neve *et al.* 2004). In addition, dynamic trafficking of NMDA receptors to and from the synaptic sites is increasingly appreciated as a key determinant in synaptic transmission and plasticity (Wenthold *et al.* 2003; Nong *et al.* 2004), as well as in pathologic states such as schizophrenia (Lau and Zukin 2007). NMDA receptor subunits can be directly phosphorylated by serine/threonine kinases PKA and PKC (Scott *et al.* 2003; Chen and Roche 2007) or Src family kinases (Salter and Kalia 2004). Therefore, an important question is how NMDA receptor trafficking is regulated by DA and what are the roles of PKA and PKC in the D₁-mediated regulation of NMDA receptor trafficking. In this study, we hypothesized that D₁ activation enhances NMDA receptor trafficking through rapid synaptic insertion of NMDA receptors, which might be differentially regulated by PKA and PKC. We tested the effects of PKA and PKC inhibitors on D₁-induced trafficking processes of NMDA receptors, including total expression, surface expression, synaptic insertion, and synaptic function. We found that although both PKA and PKC mediate the D₁-induced NMDA receptor trafficking, the phospholipase C (PLC)-PKC-Src pathway is only required for the surface expression and synaptic insertion of NMDA receptors.

Experimental procedures

Detailed experimental protocols can be found in the Appendix S1.

Electrophysiology in PFC slices

Rats at P12–30 were deeply anesthetized and the brains were sliced at 300 μm (Gao *et al.* 2001). Whole-cell recordings were conducted in medial PFC and the recordings were conducted at ~35°C. To record NMDAR mEPSCs, cells were held at –60 mV in Mg²⁺-free solution with perfusion of tetrodotoxin (TTX, 0.5 μM), 2,3-dihydroxy-6-nitro-7-sulfamoyl-benzo[f]quinoxaline-2,3-dione (NBQX, 20 μM), and picrotoxin (50–100 μM, see Figure S1). The NMDA receptor-mediated excitatory postsynaptic currents (EPSCs) were evoked by stimulating layer 2/3 in the presence of Mg²⁺ (1 mM), picrotoxin (50–100 μM) and NBQX (20 μM) at +40 mV. In some cells, NMDA (100 μM) was puff-applied from a glass pipette (~2 MΩ) and NMDA currents were recorded at –60 mV under same conditions. The electric signals were amplified and acquired with pCLAMP 9.2. A typical NMDA mEPSC with 20–80% rise time 7 ms and 63% decay time 240 ms was selected to create a sample template for detecting mEPSCs. The frequency (event number) and amplitude of individual mEPSC events were automatically detected with Clampfit by using the template. The amplitudes of the evoked EPSCs were measured by averaging 30 traces from the onset to peak of EPSC.

Neuronal cultures and treatment schedules

Neurons at 16–18 days *in vitro* were treated for 10 min with SKF-81297 (1 μM). For inhibitor and antagonist experiments, cells were pre-incubated in the following drugs (Sigma, St. Louis, MO, USA) and then co-applied with SKF-81297 at the indicated times and concentrations: KT5720 (20 min, 5 μM), Go6983 (10 min, 20 μM), SCH-23390 (10 min, 10 μM), U73122 (10 min, 10 μM), 4-amino-5-(4-chlorophenyl)-7-(*t*-butyl)pyrazolo[3,4-*d*]pyrimidine (PP2) (10 min, 10 μM) and inactive structural analogue PP3 (10 min, 10 μM).

Immunocytochemical analysis

(i) *Immunostaining of total protein*: Immunocytochemical analysis was performed according to established protocols (Glynn and McAllister 2006). Briefly, coverslips were fixed with methanol for 10 min followed by permeabilization for 5 min in 0.25% Triton X-100. Primary antibodies were visualized with FITC- or Cy3-conjugated secondary antibodies (1 : 500–1 : 1000). The specificity of all of these antibodies has been demonstrated previously and was rechecked (see Figure S2). (ii) *Surface labeling*: We labeled surface NMDA receptor (NR) 2B using an NR2B *N*-terminal antibody (Zymed Laboratories Inc., South San Francisco, CA, USA). The labeling was performed as previously described (Lu *et al.* 2001). Live cultures were incubated with antibody for 30 min. Cells were then fixed for 15 min, blocked with 5% donkey serum for 60 min, and incubated with secondary antibody conjugated to FITC for 60 min under non-permeant conditions. (iii) *Receptor insertion assay*: The rate of NR2B insertion is determined by pre-blocking existing surface receptors with primary antibody and non-conjugated secondary antibody, bringing cells to 20–22°C to allow receptor insertion, and then detecting newly inserted receptors with a second round of immunostaining (Lu *et al.* 2001).

Image analysis

All groups were analyzed simultaneously using cells from the same culture preparation. For each experimental group, cells from at least three different wells were used, and approximately five cells from each well were analyzed. Processes located about 1 soma diameter from the soma were selected for analysis. Immunocytochemical quantification was performed using ImageJ software (Glynn and McAllister 2006).

Results

D₁ activation potentiates NMDA receptor-mediated mEPSCs in PFC pyramidal neurons

We first recorded NMDA mEPSCs from layer V pyramidal neurons in medial PFC in the presence of the sodium channel blocker TTX (0.5–1 μM), α-amino-3-hydroxy-5-methylisoxazole-4-propionate (AMPA) receptor antagonist NBQX (20 μM), and GABA_A receptor antagonist picrotoxin (50–100 μM) under conditions of zero-Mg²⁺ with membrane potential held at –60 mV. After baseline recording for 5–10 min (Fig. 1), we tested the effect of dopamine D₁ agonist SKF-81297 on the mEPSCs. Bath application of SKF-81297 (1 μM) resulted in a significant potentiation of mEPSCs. As




Fig. 1 D₁ activation enhances NMDA receptor-mediated synaptic responses. (a) An example of NMDA mEPSC recorded at -60 mV in zero-Mg²⁺ solution. The NMDA mEPSCs were clearly detectable under the control condition. (b) The NMDAR mEPSCs from the same cell in panel (a) were significantly increased by D₁ agonist SKF-81297 (1 μM). (c) Time course showing the significant increase of NMDA mEPSC numbers ($n = 6$, $p < 0.01$). Insets denote the average traces in the marked periods. (d) Summary graphs show that the amplitudes

of NMDAR mEPSCs were significantly increased during and after washing out of SKF-81297 ($n = 6$, $*p > 0.05$). (e) SKF-81297-induced changes in mEPSC frequency and amplitude were prevented by pre-treatment with D₁ antagonist SCH-23390 (10 μM, $n = 6$, $p > 0.05$). (f) Bath application of specific NR2B antagonist ifenprodil (3 μM) effectively blocked the potentiation of NMDA mEPSCs induced by SKF-81297 ($n = 6$; $**p < 0.01$).

shown in Fig. 1(a–d), both frequency and amplitude of NMDAR mEPSCs were significantly increased during and after washing out of SKF-81297 ($n = 6$, $p < 0.05$ for both). The NMDA mEPSCs were effectively blocked by NMDA receptor antagonist (2R)-amino-5-phosphonovaleric acid; (2R)-amino-5-phosphonopentanoate (APV, 50 μM, data not shown). Under conditions of zero-Mg²⁺, both frequency and amplitude of mEPSCs appeared to be stable, without significance, in the absence of SKF-81297 ($n = 4$; $p > 0.05$, see Figure S1). To confirm the D₁ action, D₁ antagonist SCH-23390 (10 μM) was administered for 5 min and then SCH-23390 and SKF-81297 were co-applied for 10 min. SCH-23390 itself exhibited no significant effects on the NMDA mEPSCs (data not shown). Under this condition,

SCH-23390 effectively blocked the potentiation of NMDA mEPSCs caused by SKF-81297 in both frequency and amplitude ($n = 6$; $p > 0.05$, Fig. 1e). In another set of experiments, we recorded the AMPA receptor-mediated mEPSCs at -60 mV in the presence of (2R)-amino-5-phosphonovaleric acid; (2R)-amino-5-phosphonopentanoate (APV), TTX, and picrotoxin. Under this condition, we found the neither frequency nor amplitude of the AMPA mEPSCs was changed by D₁ agonist SKF-81297 (1 μM) ($n = 6$, $p > 0.05$, Figure S3).

Next, we determined which NMDA receptor subunits are involved in D₁-mediated regulation of NMDA mEPSCs. Bath application of specific NR2B antagonist ifenprodil (3 μM), which preferentially inhibits NR1/

NR2B channels ($IC_{50} = 0.34 \mu\text{M}$) (Williams 1993; Tovar and Westbrook 1999), effectively blocked the potentiation of NMDA mEPSCs caused by SKF-81297 (Fig. 1f, $n = 6$; $p < 0.05$). These data suggest that D₁-mediated potentiation of NMDA mEPSCs mainly depends on NR1/NR2B, although NR1/NR2A cannot be excluded. Taken together, these data indicate that D₁ activation potentiates NMDA mEPSCs in PFC and the D₁ agonist is likely to initiate an intracellular signal pathway to strengthen synaptic transmission.

D₁ activation enhances NMDA-induced current and NMDA receptor-mediated EPSCs in PFC slices

Either pre- or post-synaptic mechanisms could be responsible for the potentiation of NMDA mEPSCs by SKF-81297. To differentiate between these possibilities, we directly examined SKF-81297 effect on NMDA-induced current. NMDA ($100 \mu\text{M}$) was applied every 90 s for 2 s through a glass pipette ($\sim 2 \text{ M}\Omega$) to elicit NMDA current in the presence of Mg^{2+} (1 mM), TTX ($0.5 \mu\text{M}$), NBQX ($20 \mu\text{M}$), and picrotoxin ($50 \mu\text{M}$) at -60 mV . After application of SKF-81297 ($1 \mu\text{M}$), NMDA-induced current was gradually increased (Fig. 2a). The maximal amplitude of NMDA current increased by $50.4 \pm 11.41\%$ in SKF-81297 ($n = 5$; $p < 0.01$; Fig. 2b). Furthermore, we examined the EPSC response to paired pulses, a measure that is exquisitely sensitive to transmitter release. As shown in Fig. 2(c), NMDA EPSCs were elicited by stimulating intracortical fibers in layer 2/3 with two successive stimuli of identical strength at an interval of 100 ms (10 Hz) before and after application of SKF-81297. Although the amplitudes of both the first and the second EPSCs were increased by SKF-81297 (Fig. 2d), the ratio of paired pulses was slightly decreased without significant difference ($n = 7$, $p > 0.05$; Fig. 2e). In addition, no significant difference in amplitude variability was found, as shown in analysis of coefficient of variation (Fig. 2f). These results suggest that the effect of D₁ on NMDA mEPSCs is very likely attributable to the changes of post-synaptic NMDA receptors, although altered pre-synaptic glutamate release cannot be completely excluded.

D₁ regulation of NMDA mEPSCs involves both PKA and PKC signaling pathways

D₁ activation-induced enhancement of NMDA responses can be mediated by a number of redundant and cooperative signaling cascades (Seamans and Yang 2004; Cepeda and Levine 2006). The most prominent action is through the activation of D₁ receptor-mediated intracellular cAMP/PKA/DARPP-32 cascade (Blank *et al.* 1997; Cepeda *et al.* 1998; Snyder *et al.* 1998; Flores-Hernandez *et al.* 2002). Previous studies have also shown that PKC plays critical roles in the D₁-NMDA interactions (Chergui and Lacey 1999; Chen *et al.* 2004). In order to explore the signaling pathway




Fig. 2 SKF-81297 potentiates the mEPSC through effects on the NMDA receptors in the post-synaptic site. (a and b) Representative traces of the inward current (at -60 mV) induced by puff-applied NMDA ($100 \mu\text{M}$, every 2 s for 90 s) in the presence of TTX ($0.5 \mu\text{M}$), NBQX ($20 \mu\text{M}$) and picrotoxin ($100 \mu\text{M}$). The NMDA-induced currents were significantly potentiated by bath-applied SKF-81297 ($1 \mu\text{M}$), with maximum amplitude increased by $50.4 \pm 11.4\%$ ($n = 5$, $**p < 0.01$). (c) Sample traces showing the effects of D₁ on NMDA EPSCs evoked by stimulating layer 2/3 with a paired-pulse stimulation (inter-stimulus interval = 100 ms or 10 Hz) in the presence of picrotoxin ($100 \mu\text{M}$) and NBQX ($20 \mu\text{M}$) with membrane potential held at $+40 \text{ mV}$. After 5 min stable baseline recording, SKF-81297 ($1 \mu\text{M}$) was bath-applied to the PFC slices. (d) Summary histogram showing that the amplitude of the first NMDA EPSC was significantly increased by SKF-81297 application ($n = 7$, $*p < 0.05$). (e and f) The paired-pulse ratio and coefficient of variation of the NMDAR EPSCs were unaltered by SKF-81297 ($p > 0.05$ for both).

involved in the D₁-mediated enhancement of NMDA currents, we first bath-applied PKA or PKC inhibitor to examine its effect in NMDA mEPSCs. After baseline recording, we pre-treated with PKA inhibitor KT5720 ($5 \mu\text{M}$) or H89 ($20 \mu\text{M}$), or PKC inhibitor Go6983 ($20 \mu\text{M}$), respectively, for 5 min, and then administered one of these drugs with SKF-81297 ($1 \mu\text{M}$) for 10 min. As shown in Fig. 3(a), the D₁-mediated effects on the frequency of NMDA mEPSCs were partially but nonetheless significantly blocked by either PKA or PKC inhibitor ($p < 0.05$ vs. control and $p < 0.01$ vs. SKF only). On the other hand, these drugs not only completely blocked the effects of SKF-81297 ($p < 0.01$ for all) but also induced slight decrease of mEPSC amplitudes in PKA inhibitor KT5720 (but not H89) and PKC inhibitor Go6983. The




Fig. 3 D₁ agonist-induced increase of NMDA-mEPSCs was attenuated by PKA inhibitor and PKC inhibitor. (a) The SKF-81297-induced potentiation of both NMDA mEPSC frequency and amplitude was significantly decreased by co-application of PKA inhibitor H89 (20 μ M), KT5720 (5 μ M), or PKC inhibitor Go6983 (20 μ M) with SKF-81297 (1 μ M) ($n = 6$, * $p < 0.05$, ** $p < 0.01$). Dialysis with PKA inhibitory

peptide PKI6–22 also significantly blocked the SKF-induced changes in the NMDA mEPSC frequency and amplitude ($n = 6$, $p < 0.05$). (b) Neither frequency nor amplitude of NMDA mEPSCs was affected by PKA inhibitor H89, KT5720, or PKC inhibitor Go6983 administered alone without SKF ($n = 6$, $p > 0.05$).

reason for this discrepancy is unknown. The mEPSC frequency changes do not follow the same pattern with the changes in amplitudes, but were consistent with several previous studies that have also reported a slight decrease of EPSC amplitude under similar conditions (Huang and Kandel 1995; Chergui and Lacey 1999; Smith *et al.* 2005). To further investigate the role of PKA in the enhancement of NMDA currents, we included a membrane-impermeant PKA inhibitor peptide fragment (PKI6–22) amide (20 μ M) in the recording pipette in one set of experiments. Dialysis with PKI6–22 significantly blocked the SKF-induced changes in the NMDA mEPSC frequency and amplitude ($n = 6$, $p < 0.05$, Fig. 3a). This result was consistent with a previous study (Smith *et al.* 2005) but differed from another report (Chen *et al.* 2004). In addition, neither frequency nor amplitude of NMDA mEPSCs was affected by PKC inhibitor or PKA inhibitor administered alone without SKF ($n = 6$, $p > 0.05$, Fig. 3b). These data suggested that not only PKA, PKC is also involved in the D₁-induced regulation of NMDA currents, consistent with previous reports (Neve *et al.* 2004; Seamans and Yang 2004; Gu *et al.* 2007). It is therefore possible that PKA and PKC coordinately mediate the D₁ effects on NMDA receptors (Scott *et al.* 2003; Chen and Roche 2007).

D₁-mediated increase of dendritic localization of NMDA receptors depends on PKA but not PKC signaling

To directly address the role of D₁ activation on NMDA receptor trafficking, we treated cultured PFC neurons with or without D₁ agonist SKF-81297 and measured the cluster number and fluorescence intensity of NR2B subunits. Because efforts to identify antibodies that are capable of labeling surface NR1 and NR2A subunits were unsuccessful (Hallett *et al.* 2006) (Li *et al.*, unpublished observations) and

NR2B subunits are mainly regulated by D₁ activation (see Fig. 1f) (Gao and Wolf 2008), we focused our effort on NR2B. Treatment with SKF-81297 (1 μ M) for 10 min led to significant increase of dendritic localization of NR2B subunits (Fig. 4(a–l) and m) (cluster numbers: 39.1 ± 1.09 in control vs. 65.4 ± 1.07 in SKF-81297, $n = 20$, $p < 0.05$). D₁ antagonist SCH-23390 (10 μ M) blocked the effect of SKF-81297 on NMDA receptors (cluster number: 32.44 ± 1.32 , $n = 20$; $p < 0.01$ compared with the SKF-81297 group). D₁ activation by SKF-81297, however, exhibited no alterations in cluster number or fluorescence intensity of postsynaptic density-95 (PSD-95), a scaffold post-synaptic protein closely associated with NMDA receptors (see Figure S4). Interestingly, inclusion of PKA inhibitor KT5720 (5 μ M), but not PKC inhibitor Go6983 (20 μ M), effectively blocked the D₁ effect on total NR2B expression (Fig. 4m, cluster numbers: SKF + PKA inhibitor, 42.7 ± 6.1 , $n = 20$, $p > 0.05$; SKF + PKC inhibitor, 54.2 ± 9.8 , $n = 20$, $p < 0.05$). These data suggest that PKA, but not PKC, mediated the D₁ activation-induced increase of total NR2B subunits at the neuronal dendrites.

Both PKA and PKC mediate D₁-induced increase of surface expression of NMDA receptors

To directly measure the effect of D₁ activation on the surface expression of NMDA receptors, neurons treated with SKF-81297 were labeled with an antibody against the extracellular N-terminal region of NR2B. Using the surface labeling method (Hallett *et al.* 2006), control neurons stained under non-permeabilizing conditions showed surface staining for endogenous NR2B in a punctate pattern that was expressed on the dendrites (Fig. 5a,b,a' and b'). Treatment with SKF-81297 increased the NR2B surface labeling on PFC neuronal dendrites, measured by either the NR2B cluster number




Fig. 4 D₁ activation increases total NR2B clusters in permeabilizing conditions. (a–l), PFC neurons at 16 days *in vitro* (DIV 16) were treated with vehicle (a, d, g, j), SKF-81297 (1 μM) (b, e, h, k) or SKF-81297 in the presence of SCH (10 μM) (c, f, i, l), respectively, and double labeled for endogenous NR2B (green) and PSD-95 (red) (scale bar = 5 μm). (m) Quantification of total NR2B immunofluo-

rescence staining. SKF-81297 significantly increased total NR2B puncta number and fluorescence intensity compared with control ($n = 20$, $*p < 0.05$, $**p < 0.01$). PKA inhibitor KT5720 (5 μM), but not PKC inhibitor Go6983 (20 μM), completely blocked D₁ effect on NR2B cluster numbers and fluorescence intensity ($n = 20$, $**p < 0.01$).

(control: 15.2 ± 0.6 , $n = 12$; SKF: 20.6 ± 0.9 , $n = 12$, $p < 0.01$) or fluorescence intensity ($p < 0.01$) (Fig. 5c). We further tested the signaling pathways required in the increased NR2B cell surface labeling induced by D₁ activation. Previous studies suggested that PKA mediates GluR1 trafficking/surface expression (Snyder *et al.* 2005; Gao *et al.* 2006; Man *et al.* 2007; Serulle *et al.* 2007) but not NMDA receptors (Gao and Wolf 2008). In contrast, long-term potentiation leads to rapid surface expression of NMDARs but not AMPA receptors in a PKC- and Src-family-dependent manner (Grosshans *et al.* 2002). We therefore wondered whether PKA and PKC play different

roles in the D₁-mediated trafficking of NMDA receptors. Consistent with a role of cAMP/PKA in D₁ receptor signaling, inclusion of PKA inhibitor KT5720 (5 μM) effectively blocked the effect of SKF-81297 (1 μM) on surface NR2B expression (cluster numbers: control 16.6 ± 1.2 ; SKF 22.4 ± 1.8 ; SKF + SCH 17.6 ± 2.1 ; SKF + PKA inhibitor 17.8 ± 0.5 ; $n = 12$; $p > 0.05$ for all comparisons, Fig. 5c). In the cortical and hippocampal neurons, the PKC pathway is known to regulate NMDA receptor function (Lan *et al.* 2001; Grosshans *et al.* 2002; Salter and Kalia 2004; Lin *et al.* 2006), although it is unknown whether this pathway also plays a role in D₁ modulation of NMDA




Fig. 5 D₁-mediated increase in cell surface expression of NR2B is mediated by both PKA and PLC-PKC-Src pathways. (a and b, a' and b') Neurons treated with various chemicals were stained with anti-N-terminal NR2B under non-permeabilized conditions to label cell surface NR2B (scale bar = 10 μm). (c) Quantitative analysis from 40-μm dendrites of immunostained neurons showing a significant increase in surface NR2B clusters and fluorescence intensity in SKF-81297-treated neurons ($n = 12$, $*p < 0.05$, $**p < 0.01$). The significant increase in surface NR2B clusters and fluorescence intensity in SKF-81297-treated neurons was blocked by D₁ antagonist SCH-23390 (10 μM), PKA inhibitor KT5720 (5 μM), and PKC inhibitor Go6983 (20 μM), respectively ($n = 12$, $p > 0.05$). (d) In contrast, neither PKA nor PKC inhibitor administered alone affected cell surface NR2B expression ($n = 6$ for each treatment, $p > 0.05$). (e) D₁-mediated increase in NR2B surface expression was blocked by PLC inhibitor (U73122 10 μM, $n = 12$, $p > 0.05$) and Src kinase inhibitor (PP2, 10 μM, $n = 12$, $p > 0.05$) but not by inactive structural analogue PP3 (10 μM, $n = 12$, $**p < 0.01$).

receptors because D₁ signaling is traditionally coupled to cAMP/PKA, not PKC. When PKC inhibitor Go6983 (20 μM) was included in the SKF-81297 treatment, D₁ activation no longer enhanced the surface labeling of NR2B clusters (17.9 ± 1.1 , $n = 12$, $p > 0.05$; Fig. 5c). Neither PKA inhibitor KT5720 nor PKC inhibitor Go6983 administered alone affected the cell surface expression of NR2B (Fig. 5d). PLC can activate PKC, which in turn activates Src in hippocampal neurons, so we further tested the requirement of PLC and Src family kinases. In these experiments, the cluster numbers increased from 17.1 ± 0.8 in control to 23.7 ± 0.8 in SKF-81297 ($n = 12$; Fig. 5e). Both the PLC inhibitor U73122 (10 μM) and Src family kinase inhibitor PP2 (10 μM) blocked the D₁ effect (SKF + PLC inhibitor, 17.7 ± 0.7 , $p > 0.05$; SKF + PP2, 16.8 ± 0.9 , $p > 0.05$, $n = 12$). Conversely, the inactive structural analogue PP3 (10 μM) did not block the D₁ effect (22.5 ± 1.1 , $n = 12$, $p < 0.01$).

Both PKA and PKC mediate D₁-induced increase of new insertion of NMDA receptors

An increase in surface expression of NMDA receptors could be because of increased insertion or decreased internalization of NMDA receptors or both. To directly address whether D₁ activation affects the rate of NMDA receptor insertion, we selectively labeled newly inserted NR2B using a pre-blocking method (see Appendix S1) (Lu *et al.* 2001). Neurons treated with SKF-81297 (1 μM) showed significantly higher levels of newly inserted NR2B than control cultures, demonstrating that D₁ receptor stimulation increased the rate of NR2B insertion (Fig. 6). Using this approach, we examined the role of PKA and PKC signaling in D₁ activation-induced insertion of NR2B receptors. We observed that both PKA and PKC inhibitors (5 μM KT5720 and 20 μM Go6983, respectively) significantly blocked the D₁ effect on NR2B insertion ($n = 12$, cluster numbers: control, 10.3 ± 0.4 ; SKF, 14.4 ± 0.6 , $p < 0.01$; SKF + PKA inhibitor, 10.7 ± 0.2 , $p > 0.05$; SKF




Fig. 6 D₁-mediated NR2B insertion to plasma membrane is PLC-PKC-Src pathway dependent. (a–f) NR2B newly inserted into cell membrane under various conditions (scale bar = 10 μm). D₁ enhancement of NR2B insertion into cell membrane appeared to depend on PKA (d) and PKC (f). (g) Summary histogram showing that D₁ agonist SKF-81297 significantly increased NR2B insertion into cell membrane ($n = 12$, $**p < 0.01$). PKA inhibitor KT5720 (5 μM) and

PKC inhibitor (Go6983, 20 μM) blocked the effects of SKF-81297 on NR2B expression ($n = 12$ in each condition, $p < 0.05$; g). (h) D₁ receptor-mediated NR2B insertion was also obstructed by coapplication of PLC inhibitor U73122 (10 μM, $n = 12$, $p > 0.05$) and Src inhibitor PP2 (10 μM, $n = 12$, $p > 0.05$) but not by inactive analogue PP3 (10 μM, $n = 12$, $*p < 0.05$, $**p < 0.01$).

+ PKC inhibitor, 11.4 ± 0.4 , $p > 0.05$) (Fig. 6g). Interestingly, neither PKA inhibitor (11.0 ± 0.4 , $n = 12$) nor PKC inhibitor Go6983 (11.2 ± 0.5 , $n = 12$) used alone affected NR2B insertion at the basal level (Fig. 6g). Furthermore, PLC inhibitor U73122 (10 μM) or Src family kinase inhibitor PP2 (10 μM), but not its inactive analogue PP3 (10 μM), abolished the enhancement of NR2B insertion induced by D₁ activation ($n = 12$, cluster numbers: control, 11.5 ± 0.5 ; SKF, 16.1 ± 0.6 , $p < 0.01$; SKF + PLC inhibitor, 12.4 ± 0.8 , $p > 0.05$; SKF + PP2, 12.3 ± 0.5 , $p > 0.05$; SKF + PP3, 15.9 ± 0.3 , $p < 0.01$) (Fig. 6h). This finding suggests that the D₁-mediated synaptic insertion of NR2B subunits requires both PKA and PLC-PKC-Src pathways and the trafficking of NR2B is responsible to the D₁ enhancement of NMDA receptor function.

The newly inserted NMDA receptors induced by D₁ activation are localized on synaptic site and are functional NMDA receptors need to be on the cell surface to receive the glutamate binding. Our data, as well as others (Hallett *et al.* 2006; Gao and Wolf 2008), indicated that D₁ receptor activation leads to rapid trafficking of NMDA receptor subunits and increases surface expression of both NR1 and NR2B subunits in the dendrites of cultured neurons. It remains, however, unknown whether the increased surface receptors are functional or not because increase of NMDA receptors on the cell surface can be caused by increased receptors on synaptic sites or extrasynaptic sites. Only an increase in NMDA receptors on synaptic sites can strengthen the function of newly inserted receptors. Therefore, D₁-mediated potentiation of NMDA mEPSCs could be because

of NMDA receptors undergoing translocation from an intracellular location to the synapses. We tested this assumption by using MK-801, an activity-dependent and irreversible NMDA receptor antagonist (Rosenmund *et al.* 1993). Previous studies showed that MK-801 could only block the synaptic NMDA receptors that opened in response to synaptically released glutamate and once the receptors are blocked, they become dysfunctional and the channel properties are irreversible (Tovar and Westbrook 2002). We took the advantage of these properties to examine whether D₁ strengthens NMDA current by translocating NMDA receptors to synapses. The NMDAR-mediated EPSCs were evoked by stimulating layer 2/3 with a single pulse (0.1 ms, 10–100 μ A, 0.1 Hz) in the presence of picrotoxin (50–100 μ M) and NBQX (20 μ M) at +40 mV. After 5 min stable recording, MK-801 (20 μ M) was bath-applied to the PFC slices. As shown in Fig. 7(a), the NMDA EPSCs were progressively and completely blocked after \sim 10 min. The NMDA EPSCs were at an almost undetectable level in the control group even 20 min after washout of the unbound MK-801 in the absence of stimulation. In contrast, when SKF-81297 (1 μ M, 10 min) was bath-applied during the washing period, a clear recovery of NMDA EPSCs in response to the 0.1-Hz stimulation was observed (Fig. 7a). The amplitude recovery of NMDA EPSCs in the SKF-81297-treated neurons was significantly greater than that in controls (6.54 \pm 6.89% in control vs. 43.4 \pm 2.55% in SKF-81297, $n = 6$, $p < 0.01$, Fig. 7b). Consistently, the recovery of NMDA EPSCs induced by SKF-81297 was blocked by co-application of SKF with either PKA inhibitor KT5720 or NR2B inhibitor ifenprodil (recovered only by 17.1 \pm 2.16% and 15.1 \pm 3.62%, respectively, $n = 6$ for each drug and $p < 0.01$ for both, Fig. 7b). These data further suggest that the newly inserted NR2B-containing NMDA receptors induced by D₁ activation are likely to be translocated from intracellular or extrasynaptic sites to the synaptic membrane where they become functional.

Discussion

This study has identified that rapid NMDA trafficking and synaptic insertion underlie the D₁-induced enhancement of NMDA receptors. Importantly, we show that both PKA and PKC signaling pathways play important roles in the regulatory process. However, it appears that, although both PKA and PKC mediate the D₁ activation-induced enhancement of NMDA receptors, the PLC-PKC-Src pathway is only required for surface expression and new synaptic insertion of NMDA receptors.

We found that D₁ receptor activation enhanced both frequency and amplitude of NMDA mEPSCs in PFC pyramidal neurons. Although the frequency of mEPSCs was previously interpreted as being of pre-synaptic origin, the change of mEPSC frequency could be directly attribut-




Fig. 7 D₁ promotes the recovery of NMDAR currents by translocating intracellular NMDA receptors to the synapse. (a) The NMDA EPSCs were evoked by stimulating layer 2/3 in the presence of picrotoxin (100 μ M) and NBQX (20 μ M) at +40 mV. After 5 min stable recording, MK-801 (20 μ M) was bath-applied to PFC slices for 10 min to progressively block the NMDA EPSCs. The stimulation was stopped for 20 min to wash out the residual MK-801 after the EPSCs were completely and irreversibly blocked. The NMDA EPSCs were at an almost undetectable level under control condition but were significantly increased when SKF-81297 (1 μ M) was bath-applied for 10 min during the washout period. Co-application SKF-81297 (1 μ M) with either PKA inhibitor KT5720 (5 μ M) or NR2B antagonist ifenprodil (3 μ M) blocked the recovery of NMDA EPSCs induced by SKF-81297. (b) Summary histogram showing that SKF-81297 significantly promoted the recovery of NMDA EPSCs ($n = 6$, $**p < 0.01$) and this effect was blocked by either PKA inhibitor or NR2B antagonist.

able to alterations of post-synaptic receptors (Smith *et al.* 2005). In fact, after washing D₁ agonist, the potentiated mEPSCs did not return to baseline and, instead, were even further facilitated. One possibility is that SKF-81297 primed the intracellular signal pathway(s) to potentiate receptor trafficking, and once it was triggered, it would not depend on agonist itself (Seamans *et al.* 2001). This assumption, however, depends on the D₁ action on post-synaptic receptors. Indeed, D₁ agonist SKF-81297 enhanced puff

NMDA-induced current and evoked NMDA EPSCs, but did not change the paired-pulse ratio and coefficient of variation of the evoked NMDA-EPSCs (Zheng *et al.* 1999; Chen *et al.* 2004), indicating a direct post-synaptic effect. The increased total NR2B subunits and clusters of both NR1 and NR2B in the neuronal dendrites also provided direct evidence of post-synaptic receptor change. Furthermore, the AMPA mEPSCs were unchanged by low concentration of SKF-81297 (1 μ M) under our recording conditions. At relative high concentrations, dopamine through D₁ activation actually produced a decrease in AMPA EPSCs, indicating a reduction of glutamate release at pre-synaptic site (Nicola and Malenka 1997; Gao *et al.* 2001; Seamans *et al.* 2001; Ding *et al.* 2003). Although it remains unclear how pre-synaptic reduction of glutamate release would reconcile with post-synaptic trafficking of AMPA and NMDA receptors, it is proposed that by increasing the NMDA EPSCs, yet slightly reducing release, D₁ activation may selectively enhance sustained synaptic inputs (Seamans *et al.* 2001).

NMDAR trafficking presents a dynamic and powerful mechanism for the regulation of synaptic transmission and plasticity (Wenthold *et al.* 2003; Lau and Zukin 2007). Previous studies indicated that the D₁-induced increase of NR2B expressions is possibly initiated by phosphorylation of existing receptors (Dunah and Standaert 2001; Hallett *et al.* 2006; Gao and Wolf 2008; Tong and Gibb 2008). In the PFC neurons, D₄ activation decreased NR1 surface expression (Wang *et al.* 2003), whereas D₁ receptors facilitated the trafficking of NMDA receptors (Chen *et al.* 2004; Gao and Wolf 2008). Our data provided novel evidence that D₁-induced enhancement of NMDA receptor function depends on rapid insertion of NMDA receptors to the synaptic surface. Most importantly, we found that the D₁ receptor-mediated increase in NMDA receptor trafficking depends on both PKA and PKC/Src but their roles appear to be different in the regulatory process.

Although both serine kinases PKA and PKC and tyrosine kinase Src/Fyn can regulate NMDA receptor trafficking (Lan *et al.* 2001; Scott *et al.* 2003; Salter and Kalia 2004; Chen and Roche 2007), it is unknown how PKA and PKC regulate the trafficking of NMDA receptors induced by D₁ activation. In this study, we defined a signaling cascade from PLC to PKC to Src family kinase that is important for the trafficking of NMDA receptors after D₁ activation. The requirement of PLC/PKC/Src signaling cascade in the enhanced trafficking of NMDA receptors after D₁ activation is intriguing because D₁ receptor is well known to activate the classical D₁/cAMP/PKA/DARPP-32 pathway (Blank *et al.* 1997; Cepeda *et al.* 1998; Snyder *et al.* 1998; Flores-Hernandez *et al.* 2002). In addition, although our results showed that PKA inhibitor blocked the D₁-induced changes of surface expression and synaptic insertion of NR2B subunits, its role in these trafficking processes could be not be confirmed or excluded because PKA inhibitor also blocked the D₁-induced changes

in total NR2B expression. As PKC inhibition did not block the D₁-induced increase in total NR2B at dendritic spines, but nonetheless blocked the surface insertion of NMDA receptors, we speculate a two step process: first is the increase of NMDA receptors underneath the dendritic spines (which is dependent on PKA) followed by the insertion of NMDA receptors into the spines (which is dependent on PKC). The first step could involve local protein synthesis, and/or PKA-dependent NMDA receptor re-distribution (Crump *et al.* 2001; Smith *et al.* 2005). This assumption, however, needs further exploration. Indeed, it has been shown that although PKA mediates the effect of D₁ stimulation on the surface expression of AMPA receptors (Mangiavacchi and Wolf 2004; Smith *et al.* 2005; Sun *et al.* 2005, 2008; Gao *et al.* 2006), PKA activation was not sufficient to alter the surface expression of NMDA receptors in the cultured PFC neurons (Chen *et al.* 2004; Gao and Wolf 2008). In contrast, our data suggest that the D₁/PLC/PKC pathway may mediate the delivery of NMDA receptors, likely involved in surface expression and new insertion of NMDA receptors only, in consistence with previous studies (Chergui and Lacey 1999; Chen *et al.* 2004; Gao and Wolf 2008). Therefore, it is possible that PKA activation targets NMDA receptors to synapses (Crump *et al.* 2001), whereas PKC activation delivers NMDA receptors to the neuronal surface (Lan *et al.* 2001; Fong *et al.* 2002; Grosshans *et al.* 2002).

Furthermore, we have provided evidence that the process of new synaptic insertion of NR2B subunits on the cell surface triggered by D₁ stimulation appeared to be very rapid. Indeed, a recent study showed that subunit composition of synaptic NMDA receptors was extremely fast at neonatal synapses (Bellone and Nicoll 2007), as rapid as that seen with AMPA receptors (Smith *et al.* 2005; Bellone and Nicoll 2007). This rapid potentiation of NMDA receptor trafficking in the dendrites of prefrontal neurons by D₁ activation at very low (presumably optimal) concentration will have immediate effects on the integrative capacity of the synapse and can lead to important functional outcomes (Bellone and Nicoll 2007), particularly for working memory and other cognitive functions in the PFC. Indeed, D₁ antagonists have deleterious effects on the performance of working memory tasks in the primate PFC (Williams and Goldman-Rakic 1995). The lack of D₁ receptor sequestration in response to agonists because of its heteromerization with NMDA receptors would prolong dopamine's action on synaptic strength (Fiorentini and Missale 2004) and cognitive functions. In addition, both dopaminergic dysfunction and glutamatergic hypofunction are important for the development and symptomatology of some mental diseases such as schizophrenia and drug addiction. The novel NMDA trafficking pathway identified in the D₁ and NMDA receptor interaction is therefore important for understanding the interaction between dopamine and glutamate in these psychiatric disorders.

Acknowledgements

Supported by NARSAD Young Investigator Awards (W.J.G. and Y.Q.H.), NIH R21MH232307 and R01MH232395 (W.J.G.), as well as Drexel University College of Medicine. We thank Drs. I. Fischer, P. Baas, and G. Gallo for comments.

Conflict of interest

The authors claim no conflict of interest.

Supporting information

Additional Supporting information may be found in the online version of this article:

Appendix S1. Supplementary Materials and Methods.

Figure S1. Control experiments for recording of NMDA mEPSCs under zero-Mg²⁺ condition.

Figure S2. Non-specificity because of secondary antibodies is negligible in immunostaining processes.

Figure S3. The AMPA receptor-mediated currents were not affected by D₁ agonist.

Figure S4. D₁ activation increases NMDA receptor NR1 clusters but not PSD-95 in total staining.

As a service to our authors and readers, this journal provides supporting information supplied by the authors. Such materials are peer-reviewed and may be re-organized for online delivery, but are not copy-edited or typeset. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.

References

- Bellone C. and Nicoll R. A. (2007) Rapid bidirectional switching of synaptic NMDA receptors. *Neuron* **55**, 779–785.
- Blank T., Nijholt I., Teichert U., Kugler H., Behrsing H., Fienberg A., Greengard P. and Spiess J. (1997) The phosphoprotein DARPP-32 mediates cAMP-dependent potentiation of striatal N-methyl-D-aspartate responses. *Proc. Natl Acad. Sci. USA* **94**, 14859–14864.
- Cepeda C. and Levine M. S. (2006) Where do you think you are going? The NMDA-D1 receptor trap. *Sci. STKE* **2006**, pe20–pe24.
- Cepeda C., Colwell C. S., Itri J. N., Chandler S. H. and Levine M. S. (1998) Dopaminergic modulation of NMDA-induced whole cell currents in neostriatal neurons in slices: contribution of calcium conductances. *J. Neurophysiol.* **79**, 82–94.
- Chen B. S. and Roche K. W. (2007) Regulation of NMDA receptors by phosphorylation. *Neuropharmacology* **53**, 362–368.
- Chen G., Greengard P. and Yan Z. (2004) Potentiation of NMDA receptor currents by dopamine D1 receptors in prefrontal cortex. *Proc. Natl Acad. Sci. USA* **101**, 2596–2600.
- Chergui K. and Lacey M. G. (1999) Modulation by dopamine D1-like receptors of synaptic transmission and NMDA receptors in rat nucleus accumbens is attenuated by the protein kinase C inhibitor Ro 32-0432. *Neuropharmacology* **38**, 223–231.
- Crump F. T., Dillman K. S. and Craig A. M. (2001) cAMP-dependent protein kinase mediates activity-regulated synaptic targeting of NMDA receptors. *J. Neurosci.* **21**, 5079–5088.
- Ding L., Perkel D. J. and Farries M. A. (2003) Presynaptic depression of glutamatergic synaptic transmission by D1-like dopamine receptor activation in the avian basal ganglia. *J. Neurosci.* **23**, 6086–6095.
- Dunah A. W. and Standaert D. G. (2001) Dopamine D1 receptor-dependent trafficking of striatal NMDA glutamate receptors to the postsynaptic membrane. *J. Neurosci.* **21**, 5546–5558.
- Fiorentini C. and Missale C. (2004) Oligomeric assembly of dopamine D1 and glutamate NMDA receptors: molecular mechanisms and functional implications. *Biochem. Soc. Trans.* **32**, 1025–1028.
- Flores-Hernandez J., Cepeda C., Hernandez-Echeagaray E., Calvert C. R., Jokel E. S., Fienberg A. A., Greengard P. and Levine M. S. (2002) Dopamine enhancement of NMDA currents in dissociated medium-sized striatal neurons: role of D1 receptors and DARPP-32. *J. Neurophysiol.* **88**, 3010–3020.
- Fong D. K., Rao A., Crump F. T. and Craig A. M. (2002) Rapid synaptic remodeling by protein kinase C: reciprocal translocation of NMDA receptors and calcium/calmodulin-dependent kinase II. *J. Neurosci.* **22**, 2153–2164.
- Gao C. and Wolf M. E. (2008) Dopamine receptors regulate NMDA receptor surface expression in prefrontal cortex neurons. *J. Neurochem.* **106**, 2489–2501.
- Gao W. J., Krimer L. S. and Goldman-Rakic P. S. (2001) Presynaptic regulation of recurrent excitation by D1 receptors in prefrontal circuits. *Proc. Natl Acad. Sci. USA* **98**, 295–300.
- Gao C., Sun X. and Wolf M. E. (2006) Activation of D1 dopamine receptors increases surface expression of AMPA receptors and facilitates their synaptic incorporation in cultured hippocampal neurons. *J. Neurochem.* **98**, 1664–1677.
- Glynn M. W. and McAllister A. K. (2006) Immunocytochemistry and quantification of protein colocalization in cultured neurons. *Nat. Protoc.* **1**, 1287–1296.
- Goldman-Rakic P. S., Castner S. A., Svensson T. H., Siever L. J. and Williams G. V. (2004) Targeting the dopamine D1 receptor in schizophrenia: insights for cognitive dysfunction. *Psychopharmacology (Berl)* **174**, 3–16.
- Greengard P. (2001) The neurobiology of slow synaptic transmission. *Science* **294**, 1024–1030.
- Grosshans D. R., Clayton D. A., Coultrap S. J. and Browning M. D. (2002) LTP leads to rapid surface expression of NMDA but not AMPA receptors in adult rat CA1. *Nat. Neurosci.* **5**, 27–33.
- Gu W. H., Yang S., Shi W. X., Jin G. Z. and Zhen X. C. (2007) Requirement of PSD-95 for dopamine D1 receptor modulating glutamate NR1a/NR2B receptor function. *Acta Pharmacol. Sin.* **28**, 756–762.
- Hallett P. J., Spoelgen R., Hyman B. T., Standaert D. G. and Dunah A. W. (2006) Dopamine D1 activation potentiates striatal NMDA receptors by tyrosine phosphorylation-dependent subunit trafficking. *J. Neurosci.* **26**, 4690–4700.
- Huang Y. and Kandel E. R. (1995) D1/D5 receptor agonists induce a protein synthesis-dependent late potentiation in the CA1 region of the hippocampus. *Proc. Natl Acad. Sci. USA* **92**, 2446–2450.
- Lan J. Y., Skeberdis V. A., Jover T., Grooms S. Y., Lin Y., Araneda R. C., Zheng X., Bennett M. V. L. and Zukin R. S. (2001) Protein kinase C modulates NMDA receptor trafficking and gating. *Nat. Neurosci.* **4**, 382–390.
- Lau C. G. and Zukin R. S. (2007) NMDA receptor trafficking in synaptic plasticity and neuropsychiatric disorders. *Nat. Rev. Neurosci.* **8**, 413–426.
- Li Y. C., Xi D., Roman J., Huang Y. Q. and Gao W. J. (2009) Activation of glycogen synthase kinase-3 beta is required for hyperdopamine and D2 receptor-mediated inhibition of synaptic NMDA receptor function in the rat prefrontal cortex. *J. Neurosci.* **29**, 15551–15563.
- Lin Y., Jover-Mengual T., Wong J., Bennett M. V. and Zukin R. S. (2006) PSD-95 and PKC converge in regulating NMDA receptor trafficking and gating. *Proc. Natl Acad. Sci. USA* **103**, 19902–19907.
- Lu W., Man H., Ju W., Trimble W. S., MacDonald J. F. and Wang Y. T. (2001) Activation of synaptic NMDA receptors induces membrane

- insertion of new AMPA receptors and LTP in cultured hippocampal neurons. *Neuron* **29**, 243–254.
- Man H. Y., Sekine-Aizawa Y. and Haganir R. L. (2007) Regulation of {alpha}-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor trafficking through PKA phosphorylation of the Glu receptor 1 subunit. *Proc. Natl Acad. Sci. USA* **104**, 3579–3584.
- Mangiavacchi S. and Wolf M. E. (2004) D1 dopamine receptor stimulation increases the rate of AMPA receptor insertion onto the surface of cultured nucleus accumbens neurons through a pathway dependent on protein kinase A. *J. Neurochem.* **88**, 1261–1271.
- Missale C., Nash S. R., Robinson S. W., Jaber M. and Caron M. G. (1998) Dopamine receptors: from structure to function. *Physiol. Rev.* **78**, 189–225.
- Neve K. A., Seamans J. K. and Trantham-Davidson H. (2004) Dopamine receptor signaling. *J. Recept. Signal Transduct. Res.* **24**, 165–205.
- Nicola S. M. and Malenka R. C. (1997) Dopamine depresses excitatory and inhibitory synaptic transmission by distinct mechanisms in the nucleus accumbens. *J. Neurosci.* **17**, 5697–5710.
- Nong Y., Huang Y. Q. and Salter M. W. (2004) NMDA receptors are movin' in. *Curr. Opin. Neurobiol.* **14**, 353–361.
- Rosenmund C., Clements J. D. and Westbrook G. L. (1993) Nonuniform probability of glutamate release at a hippocampal synapse. *Science* **262**, 754–757.
- Salter M. W. and Kalia L. V. (2004) Src kinases: a hub for NMDA receptor regulation. *Nat. Rev. Neurosci.* **5**, 317–328.
- Scott D. B., Blanpied T. A. and Ehlers M. D. (2003) Coordinated PKA and PKC phosphorylation suppresses RXR-mediated ER retention and regulates the surface delivery of NMDA receptors. *Neuropharmacology* **45**, 755–767.
- Seamans J. K. and Yang C. R. (2004) The principal features and mechanisms of dopamine modulation in the prefrontal cortex. *Prog. Neurobiol.* **74**, 1–58.
- Seamans J. K., Durstewitz D., Christie B. R., Stevens C. F. and Sejnowski T. J. (2001) Dopamine D1/D5 receptor modulation of excitatory synaptic inputs to layer V prefrontal cortex neurons. *Proc. Natl Acad. Sci. USA* **98**, 301–306.
- Serulle Y., Zhang S., Ninan I., Puzzo D., McCarthy M., Khatri L., Arancio O. and Ziff E. B. (2007) A GluR1-cGKII interaction regulates AMPA receptor trafficking. *Neuron* **56**, 670–688.
- Smith W. B., Starck S. R., Roberts R. W., Schuman E. M., Smith W. B., Starck S. R., Roberts R. W. and Schuman E. M. (2005) Dopaminergic stimulation of local protein synthesis enhances surface expression of GluR1 and synaptic transmission in hippocampal neurons. *Neuron* **45**, 765–779.
- Snyder G. L., Fienberg A. A., Haganir R. L. and Greengard P. (1998) A dopamine/D1 receptor/protein kinase A/dopamine- and cAMP-regulated phosphoprotein (Mr 32 kDa)/protein phosphatase-1 pathway regulates dephosphorylation of the NMDA receptor. *J. Neurosci.* **18**, 10297–10303.
- Snyder E. M., Colledge M., Crozier R. A., Chen W. S., Scott J. D. and Bear M. F. (2005) Role for A kinase-anchoring proteins (AKAPS) in glutamate receptor trafficking and long term synaptic depression. *J. Biol. Chem.* **280**, 16962–16968.
- Sun X., Zhao Y. and Wolf M. E. (2005) Dopamine receptor stimulation modulates AMPA receptor synaptic insertion in prefrontal cortex neurons. *J. Neurosci.* **25**, 7342–7351.
- Sun X., Milovanovic M., Zhao Y. and Wolf M. E. (2008) Acute and chronic dopamine receptor stimulation modulates AMPA receptor trafficking in nucleus accumbens neurons cocultured with prefrontal cortex neurons. *J. Neurosci.* **28**, 4216–4230.
- Tong H. and Gibb A. J. (2008) Dopamine D1 receptor inhibition of NMDA receptor currents mediated by tyrosine kinase-dependent receptor trafficking in neonatal rat striatum. *J. Physiol.* **586**, 4693–4707.
- Tovar K. R. and Westbrook G. L. (1999) The incorporation of NMDA receptors with a distinct subunit composition at nascent hippocampal synapses in vitro. *J. Neurosci.* **19**, 4180–4188.
- Tovar K. R. and Westbrook G. L. (2002) Mobile NMDA receptors at hippocampal synapses. *Neuron* **34**, 255–264.
- Varela J. A., Hirsch S. J., Chapman D., Leverich L. S. and Greene R. W. (2009) D1/D5 modulation of synaptic NMDA receptor currents. *J. Neurosci.* **29**, 3109–3119.
- Wang J. and O'Donnell P. (2001) D1 dopamine receptors potentiate nmda-mediated excitability increase in layer V prefrontal cortical pyramidal neurons. *Cereb. Cortex* **11**, 452–462.
- Wang X., Zhong P., Gu Z. and Yan Z. (2003) Regulation of NMDA receptors by dopamine D4 signaling in prefrontal cortex. *J. Neurosci.* **23**, 9852–9861.
- Wentholt R. J., Prybylowski K., Standley S., Sans N. and Petralia R. S. (2003) Trafficking of NMDA receptors. *Annu. Rev. Pharmacol. Toxicol.* **43**, 335–358.
- Williams K. (1993) Ifenprodil discriminates subtypes of the N-methyl-D-aspartate receptor: selectivity and mechanisms at recombinant heteromeric receptors. *Mol. Pharmacol.* **44**, 851–859.
- Williams G. V. and Goldman-Rakic P. S. (1995) Modulation of memory fields by dopamine D1 receptors in prefrontal cortex. *Nature* **376**, 572–575.
- Zheng P., Zhang X. X., Bunney B. S. and Shi W. X. (1999) Opposite modulation of cortical N-methyl-D-aspartate receptor-mediated responses by low and high concentrations of dopamine. *Neuroscience* **91**, 527–535.