

12-6-2017

Oct4 Is Crucial for Transdifferentiation of Hepatocytes to Biliary Epithelial Cells in an in Vitro Organoid Culture Model

Mboya Doffou

George Adams

William C Bowen

Shirish Paranjpe

Harish S Parihar

See next page for additional authors

Follow this and additional works at: https://digitalcommons.pcom.edu/scholarly_papers

 Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Doffou, Mboya; Adams, George; Bowen, William C; Paranjpe, Shirish; Parihar, Harish S; Nguyen, Huy; Michalopoulos, George K; and Bhave, Vishakha, "Oct4 Is Crucial for Transdifferentiation of Hepatocytes to Biliary Epithelial Cells in an in Vitro Organoid Culture Model" (2017). *PCOM Scholarly Papers*. 1896.

https://digitalcommons.pcom.edu/scholarly_papers/1896

This Article is brought to you for free and open access by DigitalCommons@PCOM. It has been accepted for inclusion in PCOM Scholarly Papers by an authorized administrator of DigitalCommons@PCOM. For more information, please contact library@pcom.edu.

Authors

Mboya Doffou, George Adams, William C Bowen, Shirish Paranjpe, Harish S Parihar, Huy Nguyen, George K Michalopoulos, and Vishakha Bhave

Oct4 is crucial for transdifferentiation of hepatocytes to biliary epithelial cells in an in vitro organoid culture model.

Mboya Doffou^{1*}, George Adams^{1*}, William C. Bowen², Shirish Paranjpe², Harish S. Parihar³,
Huy Nguyen¹, George K. Michalopoulos², and Vishakha Bhave¹

Department of ¹Pharmaceutical Sciences and ³Pharmacy Practice, Philadelphia College of
Osteopathic Medicine School of Pharmacy, 625 Old Peachtree Road NW, Suwanee, GA 30024

²Department of Pathology, University of Pittsburgh School of Medicine, 200 Lothrop Street,
Pittsburgh, PA 15261

*Contributed equally to the manuscript

Number of pages: text (16), figures (8)

Abstract word count: 204

Running title: Oct4 mediates hepatocyte to biliary transdifferentiation

Key words: Cholangiocyte differentiation, Organoid culture, Signaling pathways, Oval cell,
Cholestatic liver diseases

Corresponding author

Vishakha Bhave

625 Old Peachtree Road NW

Suwanee, GA 30024

Phone: 678 407 7363, Fax: 678 407 7347, email: vishakhabh@pcom.edu

ABSTRACT

Background: Hepatocyte to biliary transdifferentiation has been documented in various models of bile duct injury. In this process, mature hepatocytes transform into mature biliary epithelial cells by acquiring biliary phenotypic markers. Several signaling pathways including PI3 kinase, Notch, Hes1, Sox9, and Hippo are shown to be involved in the process. However, if Oct4 is involved in hepatocyte to biliary transdifferentiation is unknown. Methods: We investigated the role of Oct4 in hepatocyte to biliary transdifferentiation utilizing an in vitro organoid culture system as a model of transdifferentiation. Oct4 was inhibited using adenovirus containing Oct4 shRNA. Hepatocyte specific HNF-4 α and biliary specific HNF-1 β & CK19 expression were assessed to gauge the extent of transdifferentiation. Results: Oct4 was induced during hepatocyte to biliary transdifferentiation. Oct4 inhibition significantly downregulated the appearance of biliary cells from hepatocytes. This was accompanied by a significant downregulation of signaling pathways including Notch, Sox9, and Hippo. Conclusion: Our findings suggest that Oct4 is crucial for hepatocyte to biliary transdifferentiation and maturation and that it acts upstream of Notch, Sox9, and Hippo signaling in this model. This finding identifies new signaling through Oct4 in plasticity between hepatocytes and biliary epithelial cells, which can be potentially utilized to identify new strategies in chronic biliary diseases.

INTRODUCTION

Transdifferentiation, also known as lineage reprogramming, is a process where one mature somatic cell transforms into another mature somatic cell without undergoing an intermediate pluripotent state¹. Hepatocyte to biliary transdifferentiation (HBT) is activated when there is extensive loss of functional biliary cells, and the resident biliary cells are unable/insufficient to proliferate to compensate for the loss. Under such situations, hepatocytes can transdifferentiate to biliary cells and help regenerate the bile ducts²⁻⁴. There is now growing evidence for this pathway, and current and future studies in this area are geared toward identifying the mechanisms and drivers that bring about this change. In recent years, many efforts have been aimed at generating pluripotent stem cells from somatic cells by inducing high expression of a combination of transcription (reprogramming) factors including Sox2, Octamer binding protein 4 (Oct4), Nanog, Klf4, and cMyc⁵⁻⁷. While these reprogramming factors are expressed in stem cells, their expression in adult somatic cells with high potential for clonal expansion such as hepatocytes has been less explored. Our previous study has documented that adult rat liver expresses some of these reprogramming factors in hepatocytes and biliary cells during liver regeneration and hepatocyte proliferation⁸. We further demonstrated that inhibition of these reprogramming factors affects the survival and proliferation of hepatocytes in culture. Oct4 has been previously shown to be involved with cellular reprogramming in mammalian embryogenesis^{9,10}. High expression of Oct4 is also associated with dedifferentiation of somatic cells in prostate¹¹ and breast¹² tumorigenesis. However, whether Oct4 is involved in HBT (a process involving somatic cell reprogramming) is not known and is the objective of the present study. We chose the in vitro organoid culture model of HBT for this purpose due to ease of mechanistic intervention. Primary hepatocytes, when cultured in roller bottles with chemically

defined Hepatocyte Growth Medium (HGM)¹³, transdifferentiate into biliary epithelial cells over a period of 21 days^{14,15}. In this model, the top/surface layer of hepatocytes facing the culture medium starts transdifferentiating into biliary epithelial cells between days 6-10¹⁴. Previous studies have established that it is the hepatocytes that transdifferentiate into biliary epithelial cells both in vivo and in this model, by using the DPPIV chimeric rat liver in which DPPIV was only expressed in hepatocytes^{4,14}. To investigate the role of Oct4 in HBT, we inhibited Oct4 in this system using adenovirus containing shRNA for Oct4. Various signaling pathways that have previously been shown to participate in transdifferentiation were also assessed in this model. Notch signaling is involved with liver regeneration^{16,17} and biliary proliferation¹⁸. Recent studies have also identified the role of Notch signaling¹⁹ and Hippo pathway²⁰ in in vivo models of HBT. Hence we looked at the expression of Notch and Yes associated protein 1 (Yap1, Hippo pathway), in our model before and after Oct4 inhibition. Sekiya et al have reported that Hes family transcription factor 1 (Hes1) is essential for conversion of hepatocytes into biliary lineage cells at the onset of intrahepatic colangiocarcinoma²¹. Thus we also looked at Hes1 levels before and after Oct4 inhibition. Other studies have identified 'liver progenitor cells' that have differentiated into hepatocytes or biliary cells in different cell injury models. These progenitor cells are identified based on expression of markers such as Lgr5²², Afp²³, Sox9^{24,25}, and OV6²⁶, among others. Whether hepatocytes that transdifferentiate to biliary cells express any of these progenitor markers is not known and was also investigated in the present study. Farnesoid X Receptor (Fxr) acts as a bile acid sensor²⁷, and upon binding with bile acids and activation, translocates to the nucleus and induces or inhibits the expression of a variety of genes involved in metabolism and lipid homeostasis²⁸. Fxr-dependent bile acid signaling is also required for normal liver regeneration²⁹. Rat and human primary hepatocyte cultures are known to release

bile acids in media³⁰. In the hepatocyte organoid culture (transdifferentiation model), the surface/top layer of hepatocytes (that gets maximum exposure to bile acids in the media) is the one that undergoes transdifferentiation to biliary epithelial cells. To assess if Fxr signaling is altered in hepatocytes during HBT, Fxr levels were also investigated in this model.

MATERIALS AND METHODS

Organoid cultures

Male Fisher 344 rats (150–200 g) were purchased from Charles River Laboratories (Frederick, MD). Animals were allowed food and water ad libitum. All animals received humane care according to the criteria outlined in the Guide for the Care and Use of Laboratory Animals prepared by the National Academy of Sciences and our institute. Hepatocytes were isolated by adaptation of the calcium two step collagenase perfusion technique as previously described^{13,31}. Freshly isolated hepatocytes were added to roller bottles (850 cm² surface) obtained from Falcon (Franklin Lakes, NJ). Each bottle contained 210,000,000 freshly isolated hepatocytes in 250 ml of Hepatocyte Growth Medium (HGM) medium1 supplemented with HGF (20 ng/ml) and EGF (10 ng/ml). The bottles were rotated at a rate of 2.5 rotations per minute and kept in an incubator maintained at 37°C, saturated humidity, and 5% CO₂. The composition of HGM is described previously¹⁵. Organoid culture tissue was harvested on days 3, 6, 10, 15, and 21 for immunohistochemistry as well as mRNA and protein extraction for further analysis.

Total RNA extraction

Total RNA was extracted from plated cells according to the manufacturer's protocol using the RNA Bee reagent (Invitrogen, Carlsbad, USA). The isolated RNA was treated with Turbo DNA-free (Ambion, Austin, TX) according to the manufacturer's instructions. RNA was quantified by

spectrophotometry at 260 nm, and purity was assessed by optical density 260/280 ratio. The RNA was stored at -80°C .

cDNA synthesis

Five micrograms RNA per sample was reverse transcribed using random hexamer to cDNA by using SuperScriptIII reverse transcriptase (Invitrogen, CA) according to the manufacturer's protocol. A no reverse transcriptase (RT) control was also included.

Real time RTPCR

Gene-specific primers used for rat were obtained from SuperArray Biosciences Corporation as follows: Gapdh (PPR06557B), Oct4 (PPR59727A), HNF-4 α (PPR49773A), HNF-1 β (PPR45380A), CK19 (PPR44322A), Sox9 (PPR53329A), Afp (PPR44288A), Notch1 (PPR47971A), Notch2 (PPR51608A), Hes1 (PPR46895C), Yap1 (PPR54845A). Expression levels were determined by quantitative Reverse Transcription Polymerase Chain Reaction (qRT-PCR) using SYBR green, and levels were normalized relative to expression of Gapdh in each sample. Fold change in gene expression was calculated by using the $2^{-\Delta\Delta\text{Ct}}$ method³². Reverse transcribed samples were amplified in parallel on an ABI prism 7000 SDS instrument (Applied Biosystems, Foster city, CA). qRT-PCR for each sample was performed in triplicate in a 20 μl reaction with 50 ng of cDNA, 5 picomoles of each primer, and 1X SYBR green PCR mix (Qiagen # 330510). The standard conditions for real time PCR were as follows: 2 minutes at 50°C , 10 minutes at 95°C followed by 40 cycles of 15 seconds denaturation at 95°C , and elongation at 60°C for 45 seconds. A dissociation curve analysis was performed at the end of every run. A no RT and a no template control were also included in every run. PCR products were resolved on 2% agarose gels and visualized with GelRed™ Nucleic Acid Gel Stain.

Oct4 inhibition studies

Adenovirus containing shRNA for Oct4 was made at the University of Pittsburgh virus core facility. Short hairpin RNA (shRNA) targeting rat Oct4 was cloned in the BamH1-kpn1 site of adenovirus shuttle vector under the control of the human H1 promoter with enhanced green fluorescent protein (eGFP) marker. Appropriate scrambled shRNA negative controls were also included. We confirmed authenticity of the clones by digesting with Sfi 1 enzyme and sequencing (M13 Forward). Organoid cultures were treated with either adenovirus containing scramble shRNA (AD-Scr, control) or Oct4 shRNA (AD-Oct4). Knock down efficiency of target genes was confirmed by transfecting primary mouse hepatocytes using standard protocols. MOI of 5 was determined to be optimal for infection of primary hepatocytes based on pilot studies. Tissue was collected on day 21.

Rat Oct4 target sequence used was as follows:

Forward oligo:

5' GATCC GAACCGTGTGAGGTGGAAC **TCAAGAG** GTTCCACCTCACACGGTTC **TT**
TTTT GGTAC 3'

Reverse oligo:

5' C AAAAAA GAACCGTGTGAGGTGGAAC **CTCTTGA** GTTCCACCTCACACGGTTC
G 3'

Oct4 ELISA

Protein levels in nuclear extracts were assessed by harvesting cells from day 21 organoid cultures treated with AD-Scr and AD-Oct4. Nuclear extracts were prepared using NE-PER

Nuclear and cytoplasmic extraction kit according to manufacturer's protocol (Pierce

Biotechnology Cat. # 78833, Rockford, IL). For Oct4 ELISA, 50 µg of total protein was used according to manufacturer's protocol (MyBiosource, MBS2501619, San Diego, CA).

Immunohistochemistry on organoid culture tissues

Paraffin-embedded tissue sections (4 µm thick) were used for H&E, as well as immunohistochemical staining. Antigen retrieval was achieved by heating the slides in the microwave at high power in 1×citrate buffer for 10 minutes. The tissue sections were blocked in blue blocker for 20 minutes followed by incubation with primary antibody overnight at 4°C. The primary antibody was then linked to biotinylated secondary antibody followed by routine avidin biotin complex method. Diaminobenzidine was used as the chromogen, which resulted in a brown reaction product. Primary antibodies used were as follows: Oct4 (ab27985, 1:100), HNF-4α (R&D H1415, 1:500), HNF-1β (SC7411, 1:250).

Statistical evaluation. The statistical analysis was performed using SPSS v22 (IBM corp., Chicago, IL, USA). The data were analyzed using repeated measures one way analysis of variance (RMANOVA) for time course studies and independent Student's t-test for Oct4 inhibition studies. The results are expressed as means ± SEM obtained from three independent experiments. $p \leq 0.05$ was considered as statistically significant and $p \leq 0.001$ as highly significant. This is denoted as * $p \leq 0.05$, ** $p \leq 0.01$, and *** $p \leq 0.001$.

RESULTS

Organization of organoid cultures

H&E stain: The organoid cultures were started with freshly isolated hepatocytes following liver perfusion. Tissue samples collected over a time course of 21 days exhibited the following development (Fig.1): At day 3, the hepatocytes formed small islands in a random fashion. Over

time, the hepatocytes aligned themselves on top of each other to form a multilayered tissue. By day 6, a more distinct pattern of cells appeared as some hepatocytes aligned themselves to form the top layer. On day 10 and 15, the culture looked more defined with the hepatocyte nuclei of the top layer appearing smaller and cuboidal (potentially undergoing HBT) as compared to the bigger and rounder hepatocyte nuclei in the intermediate layer (thick arrows). By day 21, the organoid culture was complete with visibly distinct nuclei of hepatocytes and biliary epithelial cells. It consisted of multilayered cells arranged in a three dimensional fashion with the top layer of biliary epithelial cells facing the media (Fig.1, thin arrows), an intermediate layer of hepatocytes (Fig.1, thick arrows) and connective tissue, and a basal layer of endothelial cells attached to the inner surface of the bottle. The endothelial cells in this system are most likely from a small number of contaminating cells during liver perfusion and isolation of primary hepatocytes¹⁵. These observations are consistent with previous study establishing this model for HBT^{14,15}.

Appearance of biliary cells: Biliary specific Hepatocyte Nuclear Factor1 β (HNF-1 β) was used to assess the appearance of biliary epithelial cells in this model. As expected, we did not see any biliary cells at the beginning of hepatocyte culture at day 3 (Fig.2). By day 6, some nuclei stained positive for this biliary marker. The HNF-1 β positive cells were scattered throughout the organoid culture and did not form a distinct layer. Note that many of these nuclei that stained positive for HNF-1 β were rounder and bigger (resembling hepatocyte nuclei) rather than cuboidal and smaller (resembling biliary nuclei) indicating the acquisition of biliary markers by hepatocytes (Fig.2 D6, thin arrow) on their path to becoming biliary cells. These were the cells that would potentially transdifferentiate and mature to a biliary phenotype as the culture progressed. This changed by day 10, when a majority of these positive nuclei had aligned

themselves at the surface layer position and the positive nuclei were smaller and cuboidal as compared to the bigger and round hepatocyte nuclei in the intermediate layer. By day 15, a distinct surface layer of HNF-1 β positive nuclei was visible (thin arrows). By day 21, the biliary layer was very well defined (thin arrows). All the nuclei in the intermediate layer were negative for HNF-1 β indicating their commitment to hepatocytic phenotype (Fig.2, thick arrows).

Oct 4 is induced during transdifferentiation

To determine if Oct4 is induced in hepatocytes during transdifferentiation, Oct4 expression was measured at mRNA and protein level over a time course in organoid cultures (Fig.3).

Immunohistochemical staining indicated numerous Oct4 positive nuclei at day 6 (Fig.3A) scattered throughout the tissue. By day 10 and 15, the staining pattern became more distinct with nuclei in the surface layer (potential biliary cells, thin arrows) staining stronger than nuclei in the intermediate layer (that remain hepatocytes). Some nuclei remained negative for Oct4 indicating their commitment to hepatocytic phenotype. By day 21, the newly formed biliary cells that underwent transdifferentiation stained strongly for Oct4 (thin arrows), whereas the hepatocyte nuclei in the intermediate layer that did not transdifferentiate were negative (thick arrows). Oct4 was also significantly induced as early as 6 days in cultures when compared to day 3 (Fig.3B-C), and remained elevated through day 21 in cultures. Oct4 expression also exhibited two waves of induction, first at day 6 and second at day 15.

Biliary markers and precursors are induced during HBT

Biliary specific nuclear factor HNF-1 β was significantly induced starting at day 6 and remained elevated through day 21 in cultures (Fig.4A). This corroborates with generation of biliary phenotype in this model (Fig.2). Sox9, which is shown to be a biliary precursor/marker increased significantly on day 6 as compared to day 3 and remained elevated through day 21 in cultures

(Fig.4B). Biphenotypic marker Afp expression was significantly induced at day 10 in culture as compared to day 3 and remained induced through day 21 (Fig.4C). LGR5 and OV6 were assessed but were not detectable in our model. Bile acid receptor Fxr was induced significantly on day 10 and came back to day 3 levels by day 15 (Fig.4D).

Oct4 inhibition studies

Oct4 was inhibited using adenovirus containing shRNA for Oct4 (AD-Oct4) or scramble controls (AD-Scr) as per Fig.5A. Transfection of hepatocyte cultures led to 70% transfection as assessed by eGFP through day 14 in culture without significant toxicity (Fig.5B). Tissue samples from culture were collected at day 21 for all assessments following OCT4 inhibition. We observed the following: Oct4 expression: Oct4 mRNA (Fig.5C) and protein (ELISA) (Fig.5D) on day 21 was significantly lower in AD-Oct4 group as compared to AD-Scr group indicating that Oct4 was successfully inhibited using our experimental protocol. Biliary specific markers HNF-1 β and CK19 were significantly downregulated in the AD-Oct4 group as compared to AD-Scr controls (Fig.5E, G). At the same time, there were no significant differences in hepatocyte specific marker HNF-4 α expression between the AD-Scr and AD-Oct4 group, suggesting that Oct4 inhibition did not negatively affect the hepatocytes (Fig.5F). Decreased biliary marker expression was further corroborated by HNF-1 β staining pattern. While AD-Scr group exhibited abundant HNF-1 β positive biliary cells at the surface layer of cultures (Fig.6A-C, thin arrows), such cells were very scarce in the AD-Oct4 group (Fig. 6D-F). Moreover, the staining intensity was also lower in Oct4 inhibited group. We did observe, however, that the cells at the surface of the organoid culture in AD-Oct4 group, even though mostly negative for HNF1 β , were smaller than hepatocytes with an intermediate morphology (Fig.6D-F, thick arrows). Most of these HNF-1 β negative cells turned out to be positive for hepatocyte specific HNF-4 α (Fig. 7C-D),

suggesting that these cells were not able to complete their transition from hepatocyte to biliary phenotype. On the contrary, surface layer cells of AD-Scr cultures were negative for HNF-4 α staining (Fig. 7 A-B, thick arrows) suggesting their successful transition to biliary lineage.

Hepatocytes and biliary cells were counted in tissue sections stained for HNF-1 β before and after Oct4 inhibition in organoid culture. The following criteria were used for identification of cells: Hepatocytes: HNF-1 β negative (blue) big round nuclei. Biliary cells: HNF-1 β positive (brown) small cuboidal shaped nuclei. One ribbon of organoid culture tissue was considered as one unit. Number of hepatocytes and biliary cells in each unit were counted. Over 100 such ribbons were assessed from AD-Scr and AD-Oct4 treatment groups. Following Oct4 inhibition, there was a significant decrease in biliary to hepatocyte cell ratio as compared to AD-Scr controls (Fig.7E). The total number of hepatocytes between the two groups was comparable (Fig.7F) indicating that Oct4 inhibition did not affect the viability of hepatocytes that might have led to decreased biliary cells.

Previous studies have associated Notch signaling with generation of biliary phenotype. For this reason, we assessed the levels of Notch1 and Notch2 in our model. Both of these were significantly downregulated in AD-Oct4 group as compared to AD-Scr controls (Fig.8A-B, F). Sox9 (precursor for biliary phenotype) was significantly downregulated in Oct4 inhibited group as compared to control (Fig.8C, F). Yap1 signaling was also downregulated in Oct4 inhibited group (Fig.8D, F). Hes1, previously shown to be required for conversion of hepatocyte to biliary phenotype, was also downregulated by Oct4 inhibition (Fig.8E, F).

DISCUSSION

Our results indicate that Oct4 plays an important role in HBT. In our study, Oct4 expression is induced starting at day 6 in this HBT model. This is also the time point when some cells start expressing biliary marker HNF-1 β in this model¹⁴ (Fig.2, D6). This induction is seen in many hepatocytes in anticipation of transdifferentiation. However, only the surface cells that actually undergo HBT retain OCT4 staining till the end, whereas cells in intermediate layer that remain hepatocytes start losing Oct4 induction (Figure 3). Time course studies in organoid cultures show that induction of Oct4 is accompanied by induction of bi-phenotypic marker Afp at day 10 (Fig.4C). This is suggestive of the phenotypic transformation in hepatocytes during HBT and the presence of an intermediate state. Pathways previously shown to be involved in HBT such as Yap1, Sox9, and Hes1 are upregulated in this study, indicating their involvement in HBT process and endorsing previous studies reporting their role in HBT. Following Oct4 inhibition, these genes were significantly downregulated, suggesting that they are downstream of Oct4 induction in hepatocytes. This resulted in decreased number of biliary cells (Fig. 6, 7E) at the end of the study as assessed by biliary specific markers HNF-1 β and CK19 (Fig.5E, G). In our study, the first wave of Oct4 induction is seen around day 6, followed by a much stronger expression again at day 15 (Fig.3B-C). This could be attributed to previous studies showing that too little or too much Oct4 expression leads to differentiation of cells³³. It is possible that the second wave of Oct4 induction at day 15 in our study could drive differentiation and maturation of new biliary cells derived from hepatocytes. The expression of bile acid receptor Fxr in this model is interesting. Fxr receptor expression was significantly induced at day 10 in organoid culture (Fig.4D). This could be in response to high extracellular bile acid levels in the media generated by hepatocytes. Since increase in Fxr provides negative feedback for bile synthesis, this finding

is not surprising in this study utilizing organoid culture model, which lacks biliary cells at the beginning. Moreover, on day 15 onwards, following HBT and appearance of biliary cells, Fxr levels came back to normal, which further corroborates the role of Fxr signaling in intracellular bile production especially in the presence/absence of biliary cells. Based on our data, it appears that Oct4 is a driver for plasticity between hepatocytes and biliary cells in the in vitro organoid culture model of HBT. From the morphology of the cells in Fig.6 D-F, it appears that early steps towards full biliary transdifferentiation are taken in this process, but the expression of Oct4 is required for other signaling pathways (Sox9, Notch, etc.) to be activated for the fully mature HNF-1 β and CK19 expressing biliary cells to emerge. This is further confirmed by HNF-4 α positive cells in the surface layer of Oct4 inhibited cultures (Fig.7C-D) as compared to controls (Fig.7A-B). These cells although smaller and cuboidal and similar in shape to controls are negative for HNF-1 β staining. Moreover, CK19, a marker for mature biliary cells is also significantly downregulated in these cells (Fig.5G). Further studies are required to elucidate the extracellular signals that drive hepatocytes to undergo phenotypic transformation into biliary cells by upregulating Oct4. Whether increased extracellular bile levels can initiate this signaling in hepatocytes is an interesting avenue for future direction.

ACKNOWLEDGEMENTS

This work was supported by 1) University of Pittsburgh Pathology Post-doctoral Research Training Grant, 2) American Association of Colleges of Pharmacy New Investigator Award and 3) Center for Chronic Disorders of Aging grant, awarded to Vishakha Bhawe

REFERENCES

1. Michalopoulos GK. The liver is a peculiar organ when it comes to stem cells. *Am J Pathol.* 2014;184(5):1263-7.
2. Michalopoulos GK, Barua L, Bowen WC. Transdifferentiation of rat hepatocytes into biliary cells after bile duct ligation and toxic biliary injury. *Hepatology* 2005;41(3):535-44.
3. Sekiya S, Suzuki A. Hepatocytes, rather than cholangiocytes, can be the major source of primitive ductules in the chronically injured mouse liver. *Am J Pathol.* 2014;184(5):1468-78.
4. Chen YH, Chen HL, Chien CS, Wu SH, Ho YT, Yu CH, Chang MH. Contribution of Mature Hepatocytes to Biliary Regeneration in Rats with Acute and Chronic Biliary Injury. *PLoS One* 2015;10(8):e0134327.
5. Park IH, Zhao R, West JA, Yabuuchi A, Huo H, Ince TA, Lerou PH, Lensch MW, Daley GQ. Reprogramming of human somatic cells to pluripotency with defined factors. *Nature* 2008;451(7175):141-6.
6. Sullivan GJ, Hay DC, Park IH, Fletcher J, Hannoun Z, Payne CM, Dalgetty D, Black JR, Ross JA, Samuel K and others. Generation of functional human hepatic endoderm from human induced pluripotent stem cells. *Hepatology* 2010;51(1):329-35.
7. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007;131(5):861-72.
8. Bhave VS, Paranjpe S, Bowen WC, Donthamsetty S, Bell AW, Khillan JS, Michalopoulos GK. Genes inducing iPS phenotype play a role in hepatocyte survival and proliferation in vitro and liver regeneration in vivo. *Hepatology* 2011;54(4):1360-70.
9. Nichols J, Zevnik B, Anastassiadis K, Niwa H, Klewe-Nebenius D, Chambers I, Scholer H, Smith A. Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. *Cell* 1998;95(3):379-91.
10. Zeineddine D, Hammoud AA, Mortada M, Boeuf H. The Oct4 protein: more than a magic stemness marker. *Am J Stem Cells* 2014;3(2):74-82.
11. Ben-Porath I, Thomson MW, Carey VJ, Ge R, Bell GW, Regev A, Weinberg RA. An embryonic stem cell-like gene expression signature in poorly differentiated aggressive human tumors. *Nat Genet.* 2008;40(5):499-507.
12. Hu T, Liu S, Breiter DR, Wang F, Tang Y, Sun S. Octamer 4 small interfering RNA results in cancer stem cell-like cell apoptosis. *Cancer Res.* 2008;68(16):6533-40.
13. Block GD, Locker J, Bowen WC, Petersen BE, Katyal S, Strom SC, Riley T, Howard TA, Michalopoulos GK. Population expansion, clonal growth, and specific differentiation patterns in primary cultures of hepatocytes induced by HGF/SF, EGF and TGF alpha in a chemically defined (HGM) medium. *J Cell Biol.* 1996;132(6):1133-49.
14. Michalopoulos GK, Bowen WC, Mule K, Lopez-Talavera JC, Mars W. Hepatocytes undergo phenotypic transformation to biliary epithelium in organoid cultures. *Hepatology* 2002;36(2):278-83.
15. Michalopoulos GK, Bowen WC, Mule K, Stolz DB. Histological organization in hepatocyte organoid cultures. *Am J Pathol.* 2001;159(5):1877-87.
16. Geisler F, Strazzabosco M. Emerging roles of Notch signaling in liver disease. *Hepatology* 2015;61(1):382-92.
17. Kohler C, Bell AW, Bowen WC, Monga SP, Fleig W, Michalopoulos GK. Expression of Notch-1 and its ligand Jagged-1 in rat liver during liver regeneration. *Hepatology* 2004;39(4):1056-65.
18. Loomes KM, Russo P, Ryan M, Nelson A, Underkoffler L, Glover C, Fu H, Gridley T, Kaestner KH, Oakey RJ. Bile duct proliferation in liver-specific Jag1 conditional knockout mice: effects of gene dosage. *Hepatology* 2007;45(2):323-30.

19. Yanger K, Zong Y, Maggs LR, Shapira SN, Maddipati R, Aiello NM, Thung SN, Wells RG, Greenbaum LE, Stanger BZ. Robust cellular reprogramming occurs spontaneously during liver regeneration. *Genes Dev.* 2013;27(7):719-24.
20. Yimlamai D, Christodoulou C, Galli GG, Yanger K, Pepe-Mooney B, Gurung B, Shrestha K, Cahan P, Stanger BZ, Camargo FD. Hippo Pathway Activity Influences Liver Cell Fate. *Cell* 2014;157(6):1324-1338.
21. Sekiya S, Suzuki A. Intrahepatic cholangiocarcinoma can arise from Notch-mediated conversion of hepatocytes. *J Clin Invest.* 2012;122(11):3914-8.
22. Huch M, Dorrell C, Boj SF, van Es JH, Li VS, van de Wetering M, Sato T, Hamer K, Sasaki N, Finegold MJ and others. In vitro expansion of single Lgr5+ liver stem cells induced by Wnt-driven regeneration. *Nature* 2013;494(7436):247-50.
23. Li J, Xin J, Zhang L, Wu J, Jiang L, Zhou Q, Li J, Guo J, Cao H, Li L. Human hepatic progenitor cells express hematopoietic cell markers CD45 and CD109. *Int J Med Sci.* 2014;11(1):65-79.
24. Carpentier R, Suner RE, van Hul N, Kopp JL, Beaudry JB, Cordi S, Antoniou A, Raynaud P, Lepreux S, Jacquemin P and others. Embryonic ductal plate cells give rise to cholangiocytes, periportal hepatocytes, and adult liver progenitor cells. *Gastroenterology* 2011;141(4):1432-8, 1438 e1-4.
25. Furuyama K, Kawaguchi Y, Akiyama H, Horiguchi M, Kodama S, Kuhara T, Hosokawa S, Elbahrawy A, Soeda T, Koizumi M and others. Continuous cell supply from a Sox9-expressing progenitor zone in adult liver, exocrine pancreas and intestine. *Nat Genet.* 2011;43(1):34-41.
26. Oh SH, Hatch HM, Petersen BE. Hepatic oval 'stem' cell in liver regeneration. *Semin Cell Dev Biol.* 2002;13(6):405-9.
27. Karpen SJ. Exercising the nuclear option to treat cholestasis: CAR and PXR ligands. *Hepatology* 2005;42(2):266-9.
28. Claudel T, Staels B, Kuipers F. The Farnesoid X receptor: a molecular link between bile acid and lipid and glucose metabolism. *Arterioscler Thromb Vasc Biol.* 2005;25(10):2020-30.
29. Huang W, Ma K, Zhang J, Qatanani M, Cuvillier J, Liu J, Dong B, Huang X, Moore DD. Nuclear receptor-dependent bile acid signaling is required for normal liver regeneration. *Science* 2006;312(5771):233-6.
30. Ellis E, Goodwin B, Abrahamsson A, Liddle C, Mode A, Rudling M, Bjorkhem I, Einarsson C. Bile acid synthesis in primary cultures of rat and human hepatocytes. *Hepatology* 1998;27(2):615-20.
31. Seglen PO. Preparation of isolated rat liver cells. *Methods Cell Biol.* 1976;13:29-83.
32. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-Delta Delta C(T)} Method. *Methods* 2001;25(4):402-8.
33. Niwa H, Miyazaki J, Smith AG. Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat Genet.* 2000;24(4):372-6.

FIGURE LEGENDS

Figure 1: Organization of cells in organoid culture.

Representative photomicrographs of H&E stained tissue samples obtained over a time course of 21 days. Biliary cells with smaller cuboidal nuclei appear at the surface layer (thin arrows), whereas hepatocytes with bigger round nuclei form the intermediate later (thick arrows). Original magnification x 200.

Figure 2: Appearance of biliary cells form hepatocytes over a time course of 21 days in organoid culture.

Representative photomicrographs of biliary specific marker HNF-1 β in organoid culture tissue samples. –ve represents ‘no primary antibody’ control. Thin arrows indicate HNF-1 β positive nuclei undergoing/undergone HBT. Thick arrows indicate HNF-1 β negative nuclei that retain hepatocytic phenotype. Original magnification x 200, insert x 400.

Figure 3: Oct4 expression during HBT over a time course of 21 days in organoid culture.

a) Representative photomicrographs of Oct4 immunohistochemistry in organoid culture tissue samples. –ve represents ‘no primary antibody’ control. Thin arrows indicate Oct4 positive nuclei undergoing HBT. Thick arrows indicate Oct4 negative nuclei that retain hepatocytic phenotype. Original magnification x 200. b) Representative blots of PCR product and c) mRNA levels assessed by qRT-PCR and expressed as fold change relative to Gapdh. Significantly different from day 3 (D3) time point, **p \leq 0.01, ***p \leq 0.001.

Figure 4: Genes involved with HBT in organoid culture over a time course of 21 days.

mRNA levels of a) HNF-1 β , b) Sox9, c) Afp, and d) Fxr as assessed by qRT-PCR and expressed as fold change relative to Gapdh. Significantly different from day 3 (D3) time point, * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$.

Figure 5: Oct4 inhibition in organoid cultures

A) Time line for Oct4 inhibition using adenovirus in organoid culture. B) eGFP in hepatocytes infected with adenovirus containing shRNA for Oct4, MOI 5, >70 % infection observed at day 14 in culture. Original magnification x 100. C-G: analyzed from D21 samples treated with shRNA for Oct4 (AD-Oct4) or shRNA for scramble control (AD-Scr). C) mRNA levels of Oct4 assessed by qRT-PCR and expressed as fold change relative to GAPDH. D) Oct4 protein assessed by ELISA. mRNA levels as assessed by qRT-PCR and expressed as fold change relative to GAPDH of E) biliary specific marker HNF-1 β , F) hepatocyte specific marker HNF-4 α , and G) biliary specific marker CK19. Significantly different from AD-Scr control, * $p \leq 0.05$, ** $p \leq 0.01$.

Figure 6: Appearance of biliary cells from hepatocytes following Oct4 inhibition.

Representative photomicrographs of biliary specific marker HNF-1 β following Oct4 inhibition in organoid culture tissue samples on day 21. (A-C): treated with AD-Scr. (D-F): treated with AD-Oct4. Thin arrows indicate HNF-1 β positive biliary cells. Thick arrows indicate lack of HNF-1 β positive biliary cells. Original magnification x 200 (A, B, D, E) and x 400 (C & F).

Figure 7: Effect of Oct4 inhibition on hepatocytes

Representative photomicrographs of hepatocyte specific HNF-4 α following Oct4 inhibition in organoid culture tissue samples on day 21. (A-B): treated with AD-Scr. (C-D): treated with AD-Oct4. Thick arrows indicate HNF-4 α negative cells. Thin arrows indicate HNF-4 α positive cells. Original magnification x 400. Cell counts of hepatocytes and biliary epithelial cells following Oct4 inhibition. E) Biliary to hepatocyte ratio and F) number of hepatocytes. **Significantly different from AD-Scr control, $p \leq 0.01$.

Figure 8: Expression of genes associated with HBT following Oct4 inhibition.

mRNA levels of a) Notch1, b) Notch2, c) Sox9, d) YAP1, and e) Hes1 as assessed by qRT-PCR and expressed as fold change relative to Gapdh on day 21. f) Representative blots of PCR product following Oct4 inhibition. Significantly different from AD-Scr control, * $p \leq 0.05$, ** $p \leq 0.01$.




Figure 1




Figure 2




Figure 3

Gene Expression: 11




Figure 4

Gene Expression




Figure 5




Figure 6




Figure 7




Figure 8

Gene Expression: The J