

Diversity and Abundance of Bird Species in Mole National Park, Damongo, Ghana

Agyei-Ohemeng, James¹ Danquah, Emmanuel² Adu Yeboah, Bright¹
1. University of Energy and Natural Resources, Sunyani
2. Kwame Nkrumah University of Science and Technology, Kumasi

Abstract

A survey of birds in Mole National Park in the Northern Region of Ghana between February and May 2016 to identify and determine the diversity and abundance of bird species. Using Point Count Method, all birds were observed at fixed locations using an Opticron Polarex 8×40 field binocular and identification of birds' species were confirmed by Birds of Ghana and recorded vocal replay of birds. The Gamin GPS device was used to take the coordinate and location of all stations. 366 of individual birds consisting of 59 species were recorded, 45 were identified by sighting and 14 by vocalization, all belonging to 29 families. Species abundance indicated that Helmeted Guineafowl (*Numida meleagris*) (Relative Abundance (RA)= 4.90%), followed by Red-throated Bee-eater (*Merops hirundineus*) (RA =4.63%), Red-billed firefinch (*Lagonosticta Senegala*) (RA =4.36%), Little Weaver (*Ploceus luteolus*) (RA =4.09%), Senegal Parrot (*Piocephalus senegalus*) (RA =3.81%), Bar-breasted firefinch (*Lagonosticta rufopicta*) (RA =3.54%), White faced Whistling Duck (*Dendrocygna viduata*) (RA =3.27%) and Laughing Dove (*Sterptopelia senegalensis*) (RA =3.00%), whereas, the Bateleur Eagle (*Terathopius ecaudatus*) (0.54), African Dwarf Kingfisher (*Ceyx lecontei*) (0.27) and Klaas's Cuckoo (*Chrysococcyx caprius*) (0.27) had low relative species abundance. Estrildidae and Alcedinidae families were highest in abundance, (8.5%) of the total respectively, followed by Columbidae and Malaconotidae with the same relative abundance (6.8%), Ploceidae (5.1%). The practice of bird watching aimed at avifauna conservation can be encouraged in the fringe communities such as; Morgnori, Larabanga and Murugu, to document and protect bird species to improve community livelihood. Further investigations are required to compare avifauna within the off reserve and the protected areas (on reserve).

Keywords: Avifauna, Opticron, Polarex, Diversity, Abundance

1. Introduction

Biodiversity is formally defined by the Convention on Biological Diversity (CBD) as: "the variability among living organisms from all sources including, terrestrial, marine and other aquatic ecosystems and the ecological complexes of which they are part; this includes diversity within species, between species and of ecosystems" (UN 1992 Article 2).

Birds represent an important component of global biodiversity. Thanks to the interest, they stimulate with their behaviors, colors and songs, birds are the best known group of animals at the global level with the most extended time series of data available. The world's biodiversity is diminishing rapidly (Balmford *et al.*, 2003; Jenkins *et al.*, 2003), therefore at the 2002 World Summit on Sustainable Development, the nations of the world agreed to pursue more effective implementation of the objectives of the Convention on Biological Diversity (CBD) in order to achieve a significant reduction in the current rate of loss of biological diversity by 2010 (Secretariat of the Convention on Biological Diversity, 2003).

The study of avifaunal diversity is an essential ecological tool, which acts as an important indicator to evaluate different habitats both qualitatively and quantitatively. It also fulfills many ecological functions, which include disease regulation, biomass recycling, seed dispersal of fleshy fruits, and pollination (Helm, 2002).

Migratory birds also represent a component of biodiversity whose conservation is particularly challenging, given the need for an international approach to the relevant legislation for their protection. (Fernando, 2012).

Land-use change is another key driver of current and future biodiversity change in the tropics, and conversion of native forests to pastures, croplands, botanical gardens and other human dominated habitats are other primary causes of biodiversity loss (Luck and Daily, 2008; Sala *et al.*, 2000). According to Şekercioğlu *et al.* (2004, 2006) even though only 1.3% of bird species have gone extinct since 1500 AD, the global number of individual birds is estimated to have experienced a 20–25% reduction during the same period, indicating that avian populations and dependent ecosystem services are declining faster than species extinctions would indicate.

Although patterns of biodiversity loss have been explored extensively, their ecological implications have been extended to cover botanical gardens, (Wiafe and Faalong, 2012) and National Parks. The issue of resource conservation has grown along with human population growth and as such, numerous attempts and measures have been put in place to conserve the world's biodiversity. These measures include laws and policies and the creation of conducive environment to support the regeneration of species of conservation interest. This may take the form of establishment of Game, Resource and Strict Nature reserves, National Parks, Wildlife

Sanctuaries, Zoos (Cowlshaw and Dunbar, 2000)

People, worldwide, are rapidly degrading ecosystems, especially in the tropics, leading to a massive reduction in biodiversity (Laurance and Bierregaard, 1997; Vitousek *et al.*, 1997; Pimmand R, 2000 and Dirzoand R, 2003) indicate the effects of their change on biodiversity.

There are over 2,300 bird species in the African Region, a high proportion of which are endemic to the continent. Although birds present huge opportunities for nature tourism, it is also one of the reliable indicators of terrestrial biological richness (Bibby, 2000)

As West African forests are rapidly disappearing, the survivals of the birds of the Upper Guinea forests are becoming increasingly dependent on fewer areas. Despite a number of field studies conducted in the region in recent years, Demey *et al.*, 2004, 2005; Rainey *et al.*, 2005, De Laat, 2010 and Manu, 2011, the avifauna of the majority of these forest remain inadequately known. In West Africa, the decline has been on a large scale for the last 300-400 years especially for mammalian fauna (Decher *et al.*, 2000). Several reasons have been proposed for this decline. Prominent among them is the decline in original habitat, which has been observed at the global level to be the most common cause of population decline and species extinction (Stewart *et al.*, 1996; Mace *et al.*, 2000). It has been estimated that 36% of animal extinctions, where the cause is known, were due to habitat destruction Jenkins, 1992. Another estimate suggests that habitat destruction is responsible for the extinction of over 100 species daily Ehrlich *et al.*, 1991. The other major cause of extinctions is hunting pressure, which continues with increasing human populations in local communities around forests. Thus, there is now widespread agreement that many threats to species are anthropogenic in origin Stewart and Hutchings, 1996; Ghimire and Pimbert, 1997.

Ghana's forest and savanna lands still support a diverse array of plant and animal species, although much of the resource is severely depleted. Data on status of individual plant species are not readily available, but it is estimated that over 70% of the original 8.22 million hectares of closed forest in Ghana has been destroyed (IIED, 1992) and only about 10.9% to 11.8% (representing 15,800 to 17,200 square kilometers of forest cover) remains as intact forests. Deforestation rate in the country is estimated at 22,000 square kilometers per annum (Hawthorne and Mussah, 1993). At this rate, without adequate intervention there will be no intact forest left in the country within the next 100 years. The country's productive systems as well as the livelihood and very survival of Ghanaians are threatened by the severe environmental degradation associated with deforestation (e.g. soil erosion, local climate changes, instability of hydrological regimes and loss of biological diversity). Currently there are eight threatened bird species and 14 near-threatened species in Ghana i.e. species at risk and requiring monitoring (Collar and Stuart, 1985; Collar and Andrew, 1988). Four of the threatened species are restricted range species endemic to the Upper Guinea forest block, while two are Palearctic migrants, the Damara tern *Sterna balaerum* and the Roseate Tern *Sterna dougallii*. The Damara tern is recorded only rarely on the Ghana coast while the Roseate occurs in appreciable numbers.

In Ghana, there are about 760 species of birds Brown and Borrow, 2010). This contributes to the biodiversity of birds and populations of birds in the country. In Mole National Park there are about 350 species of birds Dosset and Dosset, 2008 that constitute about 40% of the overall Ghanaians bird population. Due to the large number of birds' species and a series of ornithological survey and studies conducted in the Park, the park is one of the Important Birds Areas in Ghana Bird Life International, 2005.

In the past decade a series of scientific research has been conducted in the park, including a few birds' surveys. The latter were conducted mainly by students from overseas in the summer rains of 1968 Harvey and Harrison 1970, and in July–August 1974 and 1975, and from October to December 1975 by Greig-Smith 1976, who drew up a first checklist of birds in tabulated form, taking into account earlier records, published or unpublished. Other accounts from that period include Genelly 1969, who identified some 90 species, July–October 1966, Maze 1971, 101 species in January, April and August 1967 and 1968, Sutton 1965, 1970, who visited in late March 1964 and December–January 1968–9, Taylor and Macdonald 1978, a brief April 1976 visit, Wink 1976 and Macdonald 1978a, b who added a few records of migrants.

Payne 1985, 2004 studied fire finches and indigo birds in October 1975, and his other records are included in Greig-Smith 1976, 1977a. Nevertheless, there has been no update on the diversity of birds in Mole National Park since 2008 by L. Dosset and R. Dosset.

Though bird species are often important mobile links Lundberg *et al.*, 2003, top consumers, and keystone species in some habitats Raffaelli, 2004; their value in the ecosystems cannot be over emphasized. In the face of structural developments in Mole National Park recently, it is important that bird watchers become keen to document the surviving species and know the distribution of existing bird species to add on to check on their status.

This study will provide information about bird species abundance, distribution and availability within the Mole Loop Game viewing road. The research will also add to the existing information on the previous checklist of birds of Mole National Park and to stimulate interest in bird watching, which will help Wildlife Division, Mole National Park, and researchers.

1.1 Biophysical setting

Mole National Park as well as its fringe communities works within several major vegetation types as in typical Guinea Savanna zone. The dominant vegetation is open woodlands, riverine forest and floodplain grasslands and swamps. The major occupations of the inhabitants are subsistence farming and hunting (Mole National Park Management Plan, 2005).

The average annual rainfall ranges from 1000mm to 1100 mm. More than 90% of the rainfalls in the rainy season start from April to October, with its peaks from July to September. The dry season lasts from November to March with a mean annual temperature of 28°C, varying from 26°C in December to 31°C in March. The average range from day to night is 13°C. It can be unpleasantly hot in March and April, with temperatures sometimes in the 40°C. (Mole National Park Management Plan, 2005)

1.2 Objectives

The objective of this project is to do a snap checklist of birds in Mole National Park by a quick survey to establish the diversity and abundance of bird species.

The specific objectives are:

1. To identify bird species along the Mole loop game viewing road in Mole National Park.
2. To estimate the density of birds in Mole loop game viewing loop road in Mole National Park.
3. To determine the abundance of bird's species along the Mole loop game viewing road in Mole National Park.

1.3 Methodology

1.3.1 Study Area

Mole National Park is one of six National Parks in Ghana and one of three established in the northern savanna zone. It is the largest protected area in Ghana and considered to be the most prestigious. In Ghana, Mole National Park is one of the prime sites for biodiversity conservation.

The Park is the largest national and most developed protected area for eco-tourism and nature conservation. Mole National Park is situated in the Northern Ghana between 09° 12' – 10° 06' N and 01° 25' – 02° 17' W and covers an area of 'Four thousand, Seven hundred and Fifty-five (4755) km². The elevation ranges from 120 to 490m above sea level (Mole National Park Management Plan, 2005) Figure 1. The park is about One hundred and Forty-six (146) km from Tamale the Regional capital and Twenty-six (26) km from Damongo the district capital. The Park shares boundary with four districts namely the West Gonja district, West Mamprusi district, Sawla-Tuna-Kalba district, and the Wa East district. There are thirty-three fringe communities around the park.




Figure 1: Map of Study Area

1.4 Experimental Procedure

The research was conducted in Mole National Park in the Northern Region of Ghana between February and May, 2016 using Point count method (Hamel *et al.*, 1996; Ralph *et al.*, 1993, 1995a, 1995b) along Mole loop game viewing road behind the park Headquarters of Mole National Park (MNP). Using Point-count method to monitor

bird populations, all birds observed at a fixed location were tallied at repeated observation periods. Information on the relative abundance of all bird species over time was recorded. The field work was carried out in the morning between 6:00 am to 9:00 am, for five (5) stations at regular interval of thirty (30) minutes each and five (5) minutes rest to a different station for a total distance of 9 km along the Mole loop game viewing road in Mole National Park.

An Opticron Polarex 8×40 field binocular was used to assist in the observation and identification of the bird species. The Gamin GPS device was used to take the coordinate and location of the stations. Nomenclature of birds was referenced in field book of birds; ‘Birds of Ghana’ by Borrow *et al.*, (2010) and vocal replay of birds were used.

1.5 Analysis of Data

All records were in a tabular form in the excel data sheet and the data were analyzed.

Given that;

1. Abundance: = $\frac{ni}{N}$;

Where ni = the number of individuals in the i^{th} species,

N = total number of individuals of species recorded during the study period.

2. Population density: = number of individuals/total area (km).

Bird diversity for each study site was calculated using Shannon Wiener diversity index (H) and Simpson indices (D) Shannon and Weaver, 1963, while the similarities in the five study sites were using Jaccard index (C_j) given that;

1. $H = - \sum [ni/N * \ln(ni/N)]$

2. $D = (\sum [ni * (ni - 1)]) / N(N - 1)$

3. $C_j = (|X \cap Y|) / (|X \cup Y|)$

Where; ni = number of individuals in the i^{th} species, N = total number of individuals of species recorded during the study period, $X \cap Y$ = number of species in both site and $X \cup Y$ = total number of species in either site Kangah-Kesse *et al.*, 2005.

One – way analysis of variance (ANOVA) was used to test for differences between sites in species richness and diversity values Simon and Okoth, 2016.

1.6 Result

A total of 366 of bird individuals consisting of 59 species, 29 families were recorded during this study. Using the point count method 59 species was recorded, out of which 45 species were recorded by sighting and 14 species were recorded by vocalization Figure 2.




Figure 2: The population of individual numbers and species of birds at different sites along the Mole loop game viewing road in the Mole National Park.

Analysis of variance was conducted to evaluate the hypothesis that, the diversity, abundance and distribution of bird species does not depend on different vegetation type. At alpha level of 0.05, the differences

of means of all populations at, were significant at $p < 0.05$ ($F = 1.39$, $DoF = 4$, $P = 0.02$) (Table 1).

Table 1: Analysis of Variance (ANOVA) between sites

	Sum of Squares	DoF	Mean Square	F value	P value
Between groups	47.30	4	11.82	1.39	0.024
Within groups	844.55	99	8.53		
Total	891.85	103			

Table 2: Species composition of birds along the Mole loop game viewing road in the Mole National Park

Study Site	GPS Coordinates	Number of Individuals	Number of Species
A	Lat. 9.2462 Lon. -1.8555	52	21
B	Lat. 9.2548 Lon. -1.8624	139	35
C	Lat. 9.2681 Lon. -1.8650	91	22
D	Lat. 9.2800 Lon. -1.8641	40	10
E	Lat. 9.2807 Lon. -1.8500	46	16

Species composition was recorded according to the GPS locations along the game viewing loop as in Table 2. The relative abundance of birds' species was presented in a histogram in figure 3 below.

The most abundant species in the park was Helmeted Guineafowl (*Numida meleagris*) (Relative Dominance (RD)= 4.90%), followed by Red-throated Bee-eater (*Merops hirundineus*) (RD =4.63%), Red-billed fire finch (*Lagonosticta Senegala*) (RD =4.36%), Little Weaver (*Ploceus lueteolus*) (RD =4.09%), Senegal Parrot (*Piocephalus senegalus*) (RD =3.81%), Bar-breasted fire finch (*Lagonosticta rufopicta*) (RD =3.54%), White faced Whistling Duck (*Dendrocygna viduata*) (RD =3.27%) and Laughing Dove (*Sterptopelia senegalensis*) (RD =3.00%). These species had relatively high abundant whereas, the Bateleur Eagle (*Terathopius ecaudatus*) (0.54), African Dwarf Kingfisher (*Ceyx lecontei*) (0.27) and Klaas's Cuckoo (*Chrysococcyx caprius*) (0.27) also had the lowest abundance.




Figure 3: Percentage Relative abundance of birds

The Shannon diversity index of bird species survey in the study site A was 2.82 and evenness of 0.93 (3.26, 1.69 at 95% Confidence Limit (CL)); in the study site B the diversity was 3.39 and evenness of 0.95 (4.99, 2.95 at 95% CL); in the study site C the diversity was 2.79 and evenness of 0.90 (5.67, 2.60 at 95% CL); in the study site D the diversity was 1.89 and evenness of 0.82 (6.17, 1.83 at 95% CL) and in the study site E the diversity was 2.57 and evenness of 0.93 (3.87, 1.88 at 95% CL) (Figure 3). The reverse Simpson index (i.e. 1-D)

yielded values of 0.95, 0.96, 0.93, 0.85 and 0.93 for the study site; ‘A’, ‘B’, ‘C’, ‘D’ and ‘E’ respectively, suggesting that the study site ‘B’ had the highest bird species diversity and the study site ‘C’ and ‘E’ had the same species diversity Table 3.

Table 3: Diversity and evenness of bird species at 95% Confident Level

Study Site	H	D	1 - D	E _H	CL	UL	LL
A	2.82	0.05	0.95	0.93	0.78	3.26	1.69
B	3.39	0.04	0.96	0.95	1.02	4.99	2.95
C	2.79	0.07	0.93	0.90	1.53	5.67	2.60
D	1.89	0.15	0.85	0.82	2.17	6.17	1.83
E	2.57	0.07	0.93	0.93	0.99	3.87	1.88

Legend

H – Shannon Wiener Biodiversity Index, D – Simpson Biodiversity Index, 1-D – Reverse Simpson Biodiversity Index, E_H – Shannon Species Evenness, CL – Confidence limit, UL – Upper Limit and LL – Lower Limit.

Jaccard index of similarity of bird species was calculated to support the hypotheses that there was similarity between the study sites, similarity between sites A, B was 0.27, similarity between site B, and C yielded a value of 0.19. Site D and E had a similarity of 0.15, the results show that the similarities between the study site was relatively low (‘1’ complete similarity; ‘0’ complete dissimilarity).

Both Estrildidae and Alcedinidae family had the highest species abundance, comprising 8.5% of the total species respectively, followed in order by Columbidae and Malaconotidae with the same family composition of 6.8%, Ploceidae (5.1%).

1.7 Discussion

A total of 366 of bird individuals consisting of 59 species in 29 families were recorded during this study. This indicates that the Mole loop game viewing road attracts diverse numbers of bird species for conservation. According to Bibby *et al.*, 2000 bird’s conspicuousness can vary with observer, weather, and time of the day.

In the Mole loop game viewing road, 93.22% of the species encountered were identified as savanna generals Kangah-Kesse, 2006, within the savanna general; 62.71% constitute pure savanna inhabitants and 30.51 encountered were savanna woodland inhabitants, whilst 6.77% of species encountered identified as forest inhabitants Grimes 1987. Most of the species encountered were residence and common within the study site, according to Dowsett-Lemaire & Dowsett, 2008, the near-lack of data for the season and the short duration of most visits mean that the status of many migrants and the duration of their stay remain incompletely documented.

The avifauna is typically Sudanian, and 36 of the 37 species considered endemic to the Sudanian (Sudan-Guinea) biome that occur generally in Ghana Fishpool and Evans, 2001 are represented in Mole National Park. It was observed that during the latter part of the dry season in March Dowsett-Lemaire and Dowsett 2008, shrub land and the riverine forest with the study site ‘B’ had the highest density of birds. This could be associated with greater openness in the habitat which supports shrubs that provides food and cover for different bird species Askin *et al.*, 2012; Shochast *et al.*, 2010.

The present study of bird diversity in the ‘Samole loop game viewing road’ indicates that the bird species by varieties are very high as compared to other Game Viewing Loop in the park in numbers, since birds are highly diverse and conspicuous species of the ecosystem and birds as sentinel to environmental stresses Newton, 1995; Navarro and Benítez, 1995; Blair, 1999; Hobson and Rempel, 2001; Turner, 2002.

Dowsett-Lemaire & Dowsett, 2008, explained in his discussion that the presence of some species within the study site was subjected to habitat changes. As poaching has been increasing over the years, large mammals were forced to concentrate in the better-protected area in the vicinity of the motel and park headquarters. Over-grazing was noticeable in the rains of 2004, both in the marsh and surrounding woodland, up to only 1 km north of the motel.

The species diversity index and evenness of habitats during the entire period revealed that the study site ‘B’ had the highest diversity and abundance of bird species as a result of the inaccessibility of people and large mammals, as compared to other study sites. Factors that promoted the high diversity include a wide variety of resources, high productivity and moderate levels of predation Miller, 2002.

The study site ‘C’ had a species richness of 91 individuals, Shannon’s diversity index (H) of 2.79 and evenness of 0.90 (5.67, 2.60 at 95% CL), whilst the study site ‘E’ had a species richness of 46 individuals, Shannon’s diversity index of 2.57 and evenness of 0.93 (3.87, 1.88 at 95% CL) means they different species richness and different diversity indices according to Shannon’s.

Further analysis made by using reverse Simpson’s index (1-D) proved statistically, that both site ‘C’ and ‘E’ had the same bird species diversity of 0.93 the findings were further supported by Tramer, 1967 that in a given community, bird diversity can be adequately described by merely counting the number of species present; i.e., their relative abundances can be disregarded. Therefore, we can conclude that the factors that regulate bird

species diversity do so by determining the number of species that can coexist in a given habitat. The result is that in most avian communities, the available species partition the breeding space in a characteristic way, and a series of samples from the same or from different localities will yield similar distributions of species diversity.

The distribution and abundance of many bird species are determined by the composition of the vegetation that forms a major element of their habitats. Bird species abundance depended significantly on vegetation characteristics, indicating that areas with high plant resources supported more birds. Thus, the relatively low abundance of birds in the highly disturbed sites to the other sites may be due to fewer resources provided by this vegetation type in within the sites. As vegetation changes along complex geographical and environmental gradients, a particular bird species may appear, increase or decrease in number and disappear as the habitat changes Lee and Rotenberry, 2005.

The most abundant species in the park was Helmeted Guineafowl, *Numida meleagris*, (Relative Abundance (RA)= 4.90%), followed by Red-throated Bee-eater, *Merops hirundineus*, (RA =4.63%), Red-billed fire finch, *Lagonosticta senegala*, (RA =4.36%), Little Weaver (*Ploceus lucteolus*) (RA =4.09%), Senegal Parrot, *Piocephalus senegalus*, (RA =3.81%), Bar-breasted fire finch, *Lagonosticta rufopicta*, (RA=3.54%), White faced Whistling Duck, *Dendrocygna viduata*, (RA =3.27%) and Laughing Dove, *Sterptopelia senegalensis*, (RA =3.00%). These species had relatively high abundant whereas, the Bateleur Eagle, *Terathopius ecaudatus*, (RA=0.54), African Dwarf Kingfisher, *Ceyx lecontei*, (RA=0.27).

The result indicates that most of the species from the family Estrildidae consisting of '*Lagonosticta rufopicta* Bar-breasted fire finch; *Uraeginthus bengalus* Red-check Cordon-bleu; *Estrilda melpoda* Orange-checked Waxbill; *Estrilda troglodytes* Black-rumped Waxbill and the family Ploceidae consisting of '*Ploceus lucteolus*, *Ploceus cucullatus* and *Plocepasser superciliosus*, were mostly dominant around study site 'C' with which the mole marsh is located. Dowsett-Lemaire and Dowsett (2008), also be attributed to the appearance of these species to the

presence of shorter grasses at the site and further explained that poaching has been increasing over the years, large mammals were forced to concentrate in the better-protected area in the vicinity of the motel and park Headquarters where over-grazing was highly notice both in the and the woodland causing habitat changes in area.

There were three (Hooded Vulture, Bateleur Eagle and African Dwarf Kingfisher) globally threatening birds recorded during the survey (Birdlife International, 2017). *Ceyx lecontei* African Dwarf Kingfisher, Cruse, (2002) recorded this rain forest species in Mole National Park and was later withdrawn by Cruse 2003 (Dowsett-Lemaire and Dowsett, 2008). During the survey the African Dwarf Kingfisher were encountered at study site 'E' which possess some riparian forest and small waterholes. These findings are consistent with the previous work of Kangah-Kesse *et al.*, 2005 who stated that, since the habitat choice of birds is highly specific and driven by plant communities, the presence of savanna species in the forest area suggest the trans-formation of forest vegetation into savanna.

1.8 Conclusion and Recommendations

The study established that the Mole loop game viewing road harbors a sizable number of bird species from forest and other vegetation zones such as thickets and savanna as primarily vegetation. Birds are well known indicator taxon due to their sensitivity to environmental perturbations, relevant to ecosystem functioning (e.g. pollination and seed dispersal) and relative ease in sampling. Consequently, bird diversity and abundance differed significantly among the vegetation types, and depended significantly on plant community diversity and structure of the vegetation.

The result indicates that the site with less human disturbance and other large mammal's interference had high species richness, diversity, composition, abundance and distribution and support many rare bird species, as compared to other sites, which are exposed to more Human-induced disturbance, deforestation and over-grazing by large mammals.

There was a positive relationship between the number of birds and the habitat types and the available within that habitat, i.e. as habitat type approaches the nature of forest, the bird numbers also increase. Therefore, disturbance from human and large mammals have negative effects on the diversity and distribution of avifauna.

Since avifauna species are organism without boundary, it is recommended that the practice of bird watching as a source of avifauna conservation should be encouraged in the fringe communities such as; Morgnori, Larabanga and Murugu, which would protect these bird species and improve community livelihood.

Further studies should be done within the fringe communities to compare avifauna survey, within off reserve and protected areas (on reserve)

To reduce the level of disturbance by tourist along the 'Samole Game Viewing' Loop the management of Mole National Park can build a game viewing post for tourists (bird watchers) along the loop.

Further studies can be done by comparing the population of Bird species both in the dry season and the raining season, to identify the appropriate season for bird watching and bird watchers.

Acknowledgement

We are very grateful to Zach a Wildlife tour guide in Mole National Park, who has been enskinned as the Bawena Chief for his enormous guidance throughout the study. We also appreciate the effort of the Park Administration for allowing us to do this work at minimal costs.

Reference

- Askins, R.A; Folsom-Okeefon, C. M and Hardly, M.C. (2012). Effect of vegetation, corridor width and Regional land use on Early Successional Birds on Powerline corridor *Plos one* 7 (2): 1-10.
- Balmford A, Moore JL, Brooks T, Burgess N, Hansen LA, (2001) Conservation conflicts across Africa, *Science* 291: 2616–2619.
- Bibby, C.J., Burgess, D. and Hill, D.A. (1992) *Bird Census Techniques*, Academic Press, London.
- Bibby C.J, Burgess ND, Hill and D.A, Mustoe S. (2000). *Bird Census Techniques* 2ndEd. Academic Press, London, pp.120.
- Birdlife International. (2000). *Threatened birds of the World*. Cambridge, UK: BirdLife Int. & Barcelona: Lynx.
- Birdlife International. (2004c). *Threatened birds of the world 2004* [CD-ROM]. Cambridge: Birdlife International.
- Birdlife International (2005). Birdlife Data Zone. [online.] Available at www.birdlife.org/datazone/index.html.
- Borrow, N. and Demey, R. (2001). *Helm Identification. Guides: Birds of Western Africa*, Christopher Helm Publishers.
- Bowles, I. A., R. E. Rice, R. A. Mittermeier, and G. A. B. da Fonseca. (1998). Logging and tropical forest conservation. *Science* 280:1899-1900.
- Brawn, J. D., Robinson, S. K., & Thompson III, F. R. (2001). The role of disturbance in the ecology and conservation of birds. *Annual review of Ecology and Systematics*, 251-276.
- Brooks TM, Pimm SL, Collar NJ. (1997). Deforestation predicts the number of threatened birds in insular Southeast Asia. *ConservBiol* 11: 382–394.
- Brooks TM, Pimm SL, Oyugi JO. (1999). Time lag between deforestation and bird extinction in tropical forest fragments. *ConservBiol* 13: 1140–1150.
- CBD. (2010b). *Global Biodiversity Outlook 3*. Secretariat of the Convention on Biological Diversity, Montreal.
- Clergeau P., Savard, J.-P. L., Mennechez G., Falardeau G. (1998). Bird abundance and diversity along an urban-rural gradient: a comparative study between two cities on different continents. *Condor* 100: 413–425.
- Collar, N. J. and Stuart, S. N. (1985). *Threatened birds of Africa and related islands: The ICBP/IUCN Red data Book, Part 1* (3rd ed.). Cambridge University Press, pp. 761.
- Connell, J.H. (1978). Diversity in tropical rain forests and coral reefs. *Science* 199, 1302–1310.
- Connor, E.F. & McCoy, E.D. (1979). The statistics and biology of the species. Area relationship. *Am. Nat.*, 113, 791.833.
- Cuarón, A. D. (2000). A global perspective on habitat disturbance and tropical rainforest mammals. *Conservation Biology* 14:1574-1579.
- Decher J., Kilpatrick C. W., Bahian K. L. (2000). *Small Mammal Zoogeography and Diversity in West African Forest Remnants*. Progress Report to the National Geographic Society Committee for Research and Exploration (Grant 6583–99).
- Dirzo, R. and Raven, P. H. (2003). Global state of biodiversity and loss. *Ann. Rev. Env. Res.* 28: 137-167.
- Dowsett-Lemaire F. and Dowsett R. J. (2004). *Ornithological surveys in the proposed Kyabobo N.P., Ghana (July-August 2004)*. Accra: GWD/IUCN Project 9786: 20 pp.
- Ehrlich P. R. and Wilson E. O. (1991). Biodiversity studies: science and policy. *Science* 253: 758–762.
- Ford, H.A., Paton, D.C. & Forde, N. (1979). Birds as pollinators of Australian plants. *New Zeal. J. Bot.* 17: 509-519.
- Ford, H.A. (1985). Nectar-feeding birds and bird pollination: why are they so prevalent in Australia yet absent from Europe? *Proc. Ecol. Soc. Austr.* 14: 153-158.
- Freemark, K.E. and Merriam, H.G. (1986). Importance of area and habitat heterogeneity to bird assemblages in temperate forest fragments. *Biol. Conserv*, 36, 115.141.
- Gaston, K. J., T. M. Blackburn, and K. K. Goldewijk. (2003). Habitat conversion and global avian biodiversity loss. *Proceedings of the Royal Society of London Series B-Biological Sciences* 270:1293-1300.
- Gaston, K.J. (2004). Macroecology and people. *Basic and applied ecology* 5:303-307.
- Gaston KJ, Spicer JI. (2004). *Biodiversity: an introduction*. 2nd Edition. Blackwell Publication.
- Genelly R.E. (1969). Birds of the Mole game reserve, Ghana. *Nigerian Field* 34: 171-182.
- Greig-Smith P.W. (1976). The composition and habitat preferences of the avifauna of Mole National Park, Ghana. *Bull. Nigerian Orn. Soc.* 12: 49-66.
- Greig-Smith P.W. (1977a). Bird migration at Mole National Park, Ghana. *Bull. Nigerian Orn. Soc.* 13: 3-14.

- Haila, Y., Hanski, I.K. and Raivio, S. (1987). Breeding bird distribution in fragmented coniferous taiga in southern Finland. *Ornis Fenn.*, 64, 90-106.
- Harvey W.G. and Harrison I.D. (1970). The birds of the Mole game reserve. *Bull. Nigerian Orn. Soc.* 7: 43-52, 63-75.
- Herrera, C.M. and Pellmyr, O. (2002). *Plant-Animal Interactions: an Evolutionary Approach*. Blackwell Science, Malden, Massachusetts.
- Hill, J. K., and K. C. Hamer. (2004). Determining impacts of habitat modification on diversity of tropical forest fauna: the importance of spatial scale. *Journal of Applied Ecology* 41:744-754.
- Hilty, S.L. (1980). Flowering and fruiting periodicity in a premontane rain forest in Pacific Colombia. *Biotropica* 12: 292-306.
- Hobson K.A., Rempel R., (2001). The Saskatchewan For Impacts Monitoring Scientific Advisory Board.
- Houston, D.C. and Cooper, J.E. (1975). The digestive tract of the Whiteback Griffon Vulture and its role in disease transmission among wild ungulates. *J. Wildl. Diseases* 11: 306-313.
- Houston, D.C. (1979). The adaptation of scavengers. Pp. 263-286 in: Sinclair, A.R.E. & Griffiths, N. eds. (1979). *Serengeti, Dynamics of an Ecosystem*. University of Chicago Press, Chicago, Illinois.
- Houston, D.C. (1983). The adaptive radiation of the griffon vultures. Pp. 360-363.
- Houston, D.C. (1986). Scavenging efficiency of Turkey Vultures in tropical forest. *Condor* 88: 318-323.
- Howe, H.F. and Smallwood, J. (1982). Ecology of seed dispersal. *Ann. Rev. Ecol. Syst.* 13: 201-228.
- Howe, H.F. (1984). Implications of seed dispersal by animals for tropical reserve management. *Biol. Conserv.* 30: 261-282.
- Howe, H.F., Schupp, E.W. and Westley, L.C. (1985). Early consequences of seed dispersal for a Neotropical tree (*Virola surinamensis*). *Ecology* 66: 781-791.
- Howe, H.F. (1989). Scatter- and clump-dispersal and seedling demography: hypothesis and implications. *Oecologia* 79: 417-426.
- International Institute for Environment and Development (IIED). (1992). Environmental Synopsis of Ghana O. D. A. Publication.
- Jankowski, D. (2012). The role of competition in structuring tropical bird. es. *Orn.*
- Jankowski, J. E; Ciecka A. L; Meyer, N. Y and Rabenold, K.N. (2009). Beta diversity along environmental gradient: implications of habitat specialization in tropical montane landscapes. *Journal of Animal ecology* 78:315-327.
- Jackson, D. A., P. R. Peres-Neto, and J. D. Olden. (2001). What controls who is where in freshwater fish communities: the roles of biotic, abiotic, and spatial factors. *Canadian Journal of Fisheries and Aquatic Sciences* 58:157-170.
- Jenkins, M. 2003. Prospects for biodiversity. *Science* 302:1175-1177.
- Kangah-Kesse L., Attuquayefio D., Owusu E., and Gbogbo F. (2005), Bird Species Diversity and Abundance in the Abiriw Sacred Grove in the Eastern Region of Ghana. *Department of Zoology, University of Ghana, Legon-Accra, Ghana and Ghana Wildlife Society, Accra, Ghana.*
- Karr, J.R. (1982). Avian extinction on Barro Colorado Island, Panama: a reassessment. *Am. Nat.*, 119, 220-239.
- Karr, J. R., and K. E. Freemark. (1983). Habitat selection and environmental gradients: dynamics in the "stable" tropics. *Ecology* 64:1481-1494.
- Laurance, W. F., and R. O. Bierregaard. (1997a). Preface: a crisis in the making. in W. F. Laurance and R. O. Bierregaard, editors. *Tropical forest remnants: ecology, management, and conservation of fragmented communities*. University of Chicago Press, Chicago.
- Laurance, W. F., and R. O. Bierregaard, editors. (1997b). *Tropical forest remnants: ecology, management, and conservation of fragmented communities*. University of Chicago Press, Chicago.
- Lee, P. and Rotenberry, J. T. (2005). Relationships between bird species in forested habitats of Eastern North America. *Journal of Biogeography*. 32: 1139-1150.
- Luck, G. W., G. C. Daily, and P. R. Ehrlich. (2003). Population diversity and ecosystem services. *Trends in Ecology and Evolution* 18:331-336.
- Macdonald M.A. (1978a). Records of Palaearctic migrants in Ghana. *Bull. Nigerian Orn. Soc.* 14: 66-70.
- Macdonald M.A. (1978b). Additions to local avifaunas. Mole game reserve, Ghana. *Bull. Nigerian Orn. Soc.* 14: 88.
- Macdonald M.A. (1979). Evidence for migration and other movements of African birds in Ghana. *Rev. Zool. Afr.* 93: 413-423.
- Manu SA. (2000). Effects of habitat fragmentation on the distribution of forest birds in southwestern Nigeria with particular reference to the Ibadan Malimbés and other Malimbés, PhD thesis. University of Oxford, p. 205.
- McKinney, M. I. and J. I. Lockwood. (2001). Biotic homogenization: a sequential and selective process. In: Lockwood, J. I. and M. I. McKinney (eds). *Biotic homogenization*. Kluwer Academic/Plenum, New York, NY, USA. Pp 1-17.

- Miller SA; (2002) JP Harley. *Zoology*. Published by McGraw-Hill Higher Education, (5thed.), pp. 80-81.
- Navarro A., Benitez H., (1995). El dominio del aire, La Ciencia desde México, (138), Fondo de Cultura Económica, A., Benitez, H, Mastering Air. 216. Science from Mexico, Fondo de Cultura Económica SEP- CONACYT 138.
- Newton I., (1995). *J Anim Ecol* 64; 675-696.
- Orians, G. H., R. Dirzo, and J. H. Cushman, editors. (1996a). Biodiversity and Ecosystem Processes in Tropical Forests. Springer-Verlag, Berlin.
- Orians, G. H., R. Dirzo, and J. H. Cushman. (1996b). Introduction. Pages 1-9 in G. H. Orians, R. Dirzo, and J. H. Cushman, editors. Biodiversity and Ecosystem Processes in Tropical Forests. Springer-Verlag, Berlin.
- Orians, G. H., R. Dirzo, and J. H. Cushman. (1996c). Synthesis. Pages 195-220 in G. H. Orians, R. Dirzo, and J. H. Cushman, editors. Biodiversity and Ecosystem Processes in Tropical Forests. Springer-Verlag, Berlin.
- Pickett, S. T. A, White, P. S., eds. (1985). The Ecology of Natural Disturbance and Patch Dynamics. San Diego: Academic. 472 pp.
- Pimm SL, Raven P. (2000). Biodiversity Extinction by numbers. *Nature* 403: 843–845.
- Redford, K. (1992). The empty forest. *BioScience* 42:412-422.
- Rittiboon, K and Karntanut, W. (2011). Distribution of resident birds in a protected tropical habitat in south Thailand. *Journal of Issaas* 17(2):95-103.
- Rodríguez-Estrella, R. (2007). Land use changes affect distributional patterns of desert birds in the Baja California peninsula, Mexico. *Diversity and Distribution* 13: 877–889.
- Sala OE, Chapin III FS, Armesto J. J, Berlow E, Bloomfield J, et al. (2000). Global biodiversity scenarios for the year 2100. *Science* 287: 1770–1774.
- Saunders D. A, Hobbs R. J and Margules CR. (1991) Biological consequences of ecosystem fragmentation: A review. *Conservation Biology* 5: 18–32
- Saunders D. A., Hobbs R. J. (1991). Nature conservation: the role of corridors. Surrey Beatty and Sons, Australia.
- Scharlemann J. P. W, Green RE, Balmford A. (2004) Land-use trends in endemic bird areas: Global expansion of agriculture in areas of high conservation value. *Global Change Biol* 10: 2046–2051.
- Schemske, D. W., and N. Brokaw. (1981). Treefalls and the distribution of understory birds in a tropical forest. *Ecology* 62:938-945.
- Schmitt K. & Adu-Nsiah M. (1993). *The vegetation of Mole National Park*. Accra: Forest Resource Management Project, GWD/IUCN Project 9786: 48 pp.
- Schwartz C. W. and Schwartz E.R. (1951). An ecological reconnaissance of the pheasants of Hawaii. *Auk*. 68: 281–314.
- Secretariat of the Convention of Biological Diversity. (2003). Handbook of the Convention on Biological Diversity (2nd edition). London: Earthscan. 937 p.
- Sekercioglu, C. H. (2002a). Effects of forestry practices on vegetation structure and bird community of Kibale National Park, Uganda. *Biol. Conservation*. 107: 229-240.
- Sekercioglu, C. H. (2002b). Forest fragmentation hits insectivorous birds hard. *Sci. World* 1: 62-64.
- Sekercioglu, C. H. (2002c). Impacts of birdwatching on human and avian communities. *Environ. Conservation*. 29: 282-289.
- Şekercioglu, Ç. H., Ehrlich, P. R., Daily, G. C., Aygen, D., Goehring, D. and Sandi, R. F. (2002). Disappearance of insectivorous birds from tropical forest fragments. *Proceedings of the National Academy of Sciences USA* 99: 263–267.
- Sekercioglu, C. H., G. C. Daily, and P. R. Ehrlich. (2004). Ecosystem consequences of bird declines. *Proceedings of the National Academy of Sciences of the United States of America* 101:18042-18047.
- Sekercioglu, C. H. (2006). Increasing awareness of avian ecological function. *Trends in Ecology and Evolution* 21: 464–471.
- Sekercioglu, C. H., Loarie, S. R., Oviedo, B. F., Ehrlich, P. R. and Daily, G. C. (2007). Persistence of forest birds in the Costa Rican agricultural countryside. *Conservation Biology* 21: 482–494.
- Shannon, C. E., and W. Weaver. 1963. The mathematical theory of communication. University of Illinois Press, Urbana.
- Shochat, E; Lerman, S and Fernández-Juricic, E. (2010). Birds in Urban Ecosystems: Population Dynamics, Community Structure, Biodiversity, and Conservation. *Urban Ecosystem Ecology* 55:75-86.
- Simon GS, Okoth EO. (2016). Species richness and abundance of birds in and around Nimule National Park, South Sudan. *Sci Lett*; 4(1):92-94
- Storch, D; Konvicka, M; Benes, J; Martinkova, J. and Gaston, J.K. (2003). Distribution patterns in butterflies and birds of the Czech Republic: separating effects of habitat and geographical position. *Journal of Biogeography* 30: 1195–1205.

- Sutton R.W.W. (1965). Notes on birds seen in Ghana in 1964. *Bull. Nigerian Orn. Soc.* 2: 55-62, 102-107.
- Sutton R.W.W. (1970). Bird records from Ghana in 1967 and 1968/1969. Parts 1 & 2. *Bull. Nigerian Orn. Soc.* 7: 53-56, 76-92.
- Taylor I.R. and Macdonald M.A. (1978). The status of some northern Guinea savanna birds in Mole National Park, Ghana. *Bull. Nigerian Orn. Soc.* 14: 4-8.
- Terborgh J. (1985). The role of ecotones in the distribution of Andean birds. *Ecol.*, 66: 1237-1246.
- Terborgh, J. (1995). Wildlife in managed tropical forests: a neotropical perspective. Pages 331-342 in A. E. Lugo and C. Lowe, editors. *Tropical forests: management and ecology*. Springer-Verlag, New York.
- Tilman, D. (1999). The ecological consequences of changes in biodiversity: a search for general principles. *Ecology* 80:1455-1474.
- Tramer, E. J. (1969). Bird Species Diversity: Components of Shannon's Formula. *Ecology*, Vol. 50, No. 5 *Department of Biology, Princeton University, Princeton, New Jersey*.
- Turner W.R., (2002). *Landscape Urban Plann* 65; 149-166.
- UNCED. (1992). *Agenda 21*. United Nations Conference on Environment and Development. <http://www.un.org/esa/sustdev/documents/agenda21/english/Agenda21.pdf>
- Wilbur, S.R. & Jackson, J.A. eds. (1983). *Vulture Biology and Management*. University of California Press, Berkeley, California.

Appendix 1: Species composition, abundance and density of birds along the Mole loop game viewing road in the Mole National Park.

Family	Common Name	Species	Number of Individual (ni)	Dominance	Density
Numididae	Helmeted Guineafowl	<i>Numida meleagris</i>	18	4.90	2.00
Meropidae	Red-throated Bee-eater	<i>Merops hirundineus</i>	17	4.63	1.89
Estrildidae	Red-billed firefinch	<i>Lagonosticta senegala</i>	16	4.36	1.78
Ploceidae	Little Weaver	<i>Ploceus luteolus</i>	15	4.09	1.67
Psittacidae	Senegal Parrot	<i>Piocephalus senegalus</i>	14	3.81	1.56
Estrildidae	Bare-breasted firefinch	<i>Lagonosticta rufopicta</i>	13	3.54	1.44
Anatidae	White faced Whistling Duck	<i>Dendrocygna viduata</i>	12	3.27	1.33
Ploceidae	Village Weaver	<i>Ploceus cucullastus</i>	11	3.00	1.22
Columbidae	Laughing Dove	<i>Sterptopelia senegalensis</i>	11	3.00	1.22
Estrildidae	Red-check Cordon-bleu	<i>Uraeginthus bengalus</i>	10	2.72	1.11
Columbidae	Vinaceous Dove	<i>Streptopelia vinacea</i>	9	2.45	1.00
Ardeidae	Cattle Egret	<i>Bubulcus ibis</i>	9	2.45	1.00
Phasianidae	Double-spurred Francolin	<i>Francolinus bicalcaratus</i>	8	2.18	0.89
Nectariniidae	Scarley-chested sunbird	<i>Chalcomitra senegalensis</i>	8	2.18	0.89
Hirundinidae	Wire-tailed Swallow	<i>Hirundo smithii</i>	8	2.18	0.89
Columbidae	Black-bill Wood Dove	<i>Turtur abyssinicus</i>	8	2.18	0.89
Passeridae	Grey headed Sparrow	<i>Passer griseus</i>	8	2.18	0.89
Bucerotidae	African Grey Hornbill	<i>Tockus nasutus</i>	7	1.91	0.78
Alcedinidae	Blue-breasted Kingfisher	<i>Halcyon malimbica</i>	7	1.91	0.78
Accipitridae	Hooded Vulture	<i>Necrosytres monachus</i>	7	1.91	0.78
Ploceidae	Chestnut-crown Sparrow Weaver	<i>Plocepasser superciliosus</i>	7	1.91	0.78
Malaconotidae	Northern Puffback	<i>Dryoscopus gambensis</i>	7	1.91	0.78
Alcedinidae	Grey Headed kingfisher	<i>Halcyon leucocephala</i>	7	1.91	0.78
Nectariniidae	Beautiful Sunbird	<i>Cinnyris pulchellus</i>	7	1.91	0.78
Timaliidae	Brown babbler	<i>Turdoides plebejus</i>	7	1.91	0.78
Estrildidae	Black-rumped Waxbill	<i>Estrilda troglodytes</i>	7	1.91	0.78
Sylviidae	Senegal Eremomela	<i>Eremomela pusilla</i>	6	1.63	0.67
Pycnonotidae	Common Garden Bulbul	<i>Pycnonotus barbatus</i>	6	1.63	0.67
Coraciidae	Broad-billed Roller	<i>Eurystomus glgucurus</i>	6	1.63	0.67
Estrildidae	Orange-checked Waxbill	<i>Estrilda melpoda</i>	6	1.63	0.67
Threskiornithidae	Hadada Ibis	<i>Bostrychia hagedash</i>	5	1.36	0.56
Muscicapidae	Northern Black flycatcher	<i>Melaenornis annamarulae</i>	5	1.36	0.56
Fringillidae	Yellow fronted canary	<i>Serinus mozambicus</i>	5	1.36	0.56
Charadriidae	African Wattle Lapwing	<i>Vanellus senegalus</i>	5	1.36	0.56
Musophagidae	Western Grey Plantain-eater	<i>Crinifer piscator</i>	4	1.09	0.44
Turdidae	African Thrush	<i>Turdus pelios</i>	4	1.09	0.44
Coraciidae	Blue-billed Roller	<i>Coracias abyssinicus</i>	4	1.09	0.44
Phasianidae	Stone Partridge	<i>Ptilopachus petrosus</i>	4	1.09	0.44

Family	Common Name	Species	Number of Individual (ni)	Dominance	Density
Malaconotidae	Yellow Crown Gonolek	<i>Laniarius barbarous</i>	4	1.09	0.44
Columbidae	Bruce's Green Pigeon	<i>Treron waalia</i>	4	1.09	0.44
Alcedinidae	African Pygmy kingfisher		3	0.82	0.33
Monarchidae	Africa Paradise flycatcher	<i>Terpsiphone viridis</i>	3	0.82	0.33
Platysteiridae	Senegal Batis	<i>Batis senegalensis</i>	3	0.82	0.33
Scopidae	Hamerkop	<i>Scopus umbretta</i>	3	0.82	0.33
Muscicapidae	Snowy-crowned Robin Chat	<i>Cossypha albicapilla</i>	3	0.82	0.33
Bucerotidae	Northern Red-billed Hornbill	<i>Tockus erythrorhynchus</i>	3	0.82	0.33
Ardeidae	Grey Heron	<i>Ardea cinerea</i>	2	0.54	0.22
Passeridae	Bush Petronia	<i>Petronia dentate</i>	2	0.54	0.22
Indicatoridae	Greater Honey Guide	<i>Indicator</i>	2	0.54	0.22
Malaconotidae	Black-crown Tchagra	<i>Tchagra senegalus</i>	2	0.54	0.22
Phoeniculidae	Green wood hoopoe	<i>Phoebastria purpureus</i>	2	0.54	0.22
Alcedinidae	Malackite Kingfisher	<i>Alcedo cristata</i>	2	0.54	0.22
Cisticolidae	Oriole Warbler	<i>Hypergerus atriceps</i>	2	0.54	0.22
Accipitridae	Bateleur Eagle	<i>Terathopus ecaudatus</i>	2	0.54	0.22
Cuculidae	Senegal Coucal	<i>Centropus senegalensis</i>	2	0.54	0.22
Sylviidae	Tawny-flanked Prinia	<i>Prinia subflava</i>	2	0.54	0.22
Alcedinidae	African Dwarf Kingfisher	<i>Ceyx lecontei</i>	1	0.27	0.11
Malaconotidae	Grey-headed bush shrike	<i>Malaconotus blanchoti</i>	1	0.27	0.11
Cuculidae	Klaas's Cuckoo	<i>Chrysococcyx caprius</i>	1	0.27	0.11
	Total Count		59		
	Total		367		

Appendix 2: Checklist of bird species with their status and habitat at the Mole loop game viewing road in Mole National Park

Family/ Species	Common Name	Site	Habitat	Status
		A B C D E		
Estrildidae				
<i>Lagonosticta rufopicta</i>	Bar-breasted firefinch	- 7 6 - -	S	c
<i>Lagonosticta senegala</i>	Red-billed firefinch	- 16 - - -	S	c
<i>Uraeginthus bengalus</i>	Red-check Cordon-bleu	- 10 - - -	S	c
<i>Estrilda melpoda</i>	Orange-checked Waxbill	- - 6 - -	S	c
<i>Estrilda troglodytes</i>	Black-rumped Waxbill	- - 7 - -	S	u
Total species =5 (8.5%)				
Alcedinidae				
<i>Ceyx pictus</i>	African Pygmy kingfisher	1 2 - - -	FC	f
<i>Halcyon malimbica</i>	Blue-breasted Kingfisher	1 3 - - 3	S	c
<i>Halcyon leucocephala</i>	Grey headed kingfisher	- - 3 2 2	S	c
<i>Ceyx lecontei</i>	African Dwarf Kingfisher	- - - - 1	F	r/s
<i>Alcedo cristata</i>	Malachite Kingfisher	- 2 - - -	S	c
Total species =5 (8.5%)				
Malaconotidae				
<i>Laniarius barbarous</i>	Yellow-crowned Gonolek	1 1 1 - 1	S	c
<i>Tchagra senegalus</i>	Black-crown Tchagra	- - - - 1	S	c
<i>Dryoscopus gambensis</i>	Northern Puffback	2 3 - 2 -	S	c
<i>Malaconotus blanchoti</i>	Grey-headed Bush Shrike	- - - - 1	SW	c
Total species =4 (6.8%)				
Columbidae				
<i>Turtur abyssinicus</i>	Black-bill Wood Dove	3 5 - - -	SW	c
<i>Streptopelia vinacea</i>	Vinaceous Dove	- 6 3 - -	S	c
<i>Treron waalia</i>	Bruce's Green Pigeon	- 2 2 - -	SW	c
<i>Sterptopelia senegalensis</i>	Laughing Dove	2 3 2 2 2	SW	c
Total species =4 (6.8%)				
Ploceidae				
<i>Ploceus lueteolus</i>	Little Weaver	- - 15 - -	SW	f/c
<i>Ploceus cucullastus</i>	Village Weaver	- - 11 - -	S	c

<i>Plocepasser superciliosus</i>	Chestnut Crown Sparrow Weaver	- 7 - - -	S	u/f
Total species =3 (5.1%)				
Sylviidae				
<i>Prinia subflava</i>	Tawny-flanked Prinia	- - - 2 -	S	c
<i>Eremomela pusilla</i>	Senegal Eremomela	2 2 2 - -	S	c
Total species =2 (3.4%)				
Cuculidae				
<i>Centropus senegalensis</i>	Senegal coucal	- 2 - - -	S	c
<i>Chrysococcyx caprius</i>	Klaas's Cuckoo	1 - - - -	F	f
Total species =2 (3.4%)				
Nectariniidae				
<i>Cinnyris pulchellus</i>	Beautiful sunbird	- - - - 7	SW	c
<i>Chalcomitra senegalensis</i>	Scarley-chested Sunbird	2 4 2 - -	S	c
Total species =2 (3.4%)				
Accipitridae				
<i>Necrosyrtes monachus</i>	Hooded Vulture	- 4 3 - -	S/F	c
<i>Terathopius ecaudatus</i>	Bateleur Eagle	- 2 - - -	S	u/f
Total species =2 (3.4%)				
Phasianidae				
<i>Francolinus bicalcaratus</i>	Double-spurred Francolin	3 5 - - -	SW	c
<i>Ptilopachus petrosus</i>	Stone Partridge	4 - - - -	S	f
Coraciidae				
<i>Coracias abyssinicus</i>	Blue-billed Roller	- 2 2 - -	SW	u/f
<i>Eurystomus glaucurus</i>	Broad-billed Roller	- 1 1 2 2	S	c
Ardeidae				
<i>Ardea cinerea</i>	Grey Heron	- 2 - - -	S	f/c
<i>Bubulcus ibis</i>	Cattle Egret	- - - 9 -	S	c
Muscicapidae				
<i>Melaenornis annamarulae</i>	Northern Black Flycatcher	- 3 - 2 -	S	f
<i>Cossypha albicapilla</i>	Snowy-crowned Robin Chat	1 2 - - -	S	f/c
Passeridae				
<i>Petronia dentate</i>	Bush Petronia	- - 2 - -	S	f/c
<i>Passer griseus</i>	Grey Headed Sparrow	- 8 - - -	S	c
Bucerotidae				
<i>Tockus nasutus</i>	African Grey Hornbill	2 1 2 1 1	SW	c
<i>Tockus erythrorhynchus</i>	Northern Red-billed Hornbill	1 - 2 - -	SW	u/f
Musophagidae				
<i>Crinifer piscator</i>	Western Grey Plantain-eater	- 4 - - -	SW	c
Total species =1 (1.7%)				
Pycnonotidae				
<i>Pycnonotus barbatus</i>	Common Garden bulbul	6 - - - -	S	c
Hirundinidae				
<i>Hirundo smithii</i>	Wire-tailed swallow	4 4 - - -	S	u/f
Indicatoridae				
<i>Indicator indicator</i>	Greater Honey Guide	- - 2 - -	SW	u/f
Numididae				
<i>Numida meleagris</i>	Helmeted Guinea fowl	8 - - 10 -	S	c
Phoeniculidae				
<i>Phoemicuslus purpureus</i>	Green Wood Hoopoe	- 1 2 - -	SW	c
Fringillidae				
<i>Serinus mozambicus</i>	Yellow-fronted Canary	1 4 - - -	SW	c
Timaliidae				
<i>Turdoides plebejus</i>	Brown Babblers	- - - - 7	SW	f
Charadriidae				

<i>Vanellus senegalus</i>	African Wattle Lapwing	1	4	-	-	-	S	f	
Threskiornithidae									
<i>Bostrychia hagedash</i>	Hadda Ibis	3	-	2	-	-	S	f	
Scopidae									
<i>Scopus umbretta</i>	Hamerkop	-	-	-	-	3	S	f	
Platysteiridae									
<i>Batis senegalensis</i>	Senegal Batis	-	-	-	-	3	S	f	
Turdidae									
<i>Turdus pelios</i>	African Thrush	-	4	-	-	-	SW	c/f	
Anatidae									
<i>Dendrocygna viduata</i>	White faced Whistling Duck	-	-	4	8	-	S	f	
Cisticolidae									
<i>Hypergerus atriceps</i>	Oriole Warbler	-	2	-	-	-	SW	u	
Meropidae									
<i>Merops hirundineus</i>	Red-throated Bee-eater	-	11	6	-	-	S	c	
Monarchidae									
<i>Terpsiphone viridis</i>	Africa Paradise flycatcher	-	3	-	-	-	SW	c	
Psittacidae									
<i>Piocephalus senegalus</i>	Senegal Parrot	3	7	4	-	-	SW	c	

Total 52 139 91 40 46

Legend: Habitat: S – Savanna, SW – Savanna Woodland, F – Forest, FC – Forest Clearings and CS – Coastal Savanna

Status: c – common, f – fairly common, u – uncommon, s – scarce and r – rare.