

## Helminth Parasites of Fruits and Vegetables Planted in Crop and Landscape Management Garden College of Agricultural Sciences, Ebonyi State University Abakaliki, Implication for Public Health.

UHUO COSMAS.A<sup>1\*</sup>; NWANCHOR KELECHI.<sup>2</sup>; UMENE DAVID.A<sup>1</sup>; OGBANSHI MOSES.E; NWELE DAVID.E<sup>1</sup>; IBIAM GIDEON.A<sup>4</sup>; & AZI SIMON.O<sup>4</sup>.

1. Department Of Applied Biology, Faculty Of Biological Sciences, Ebonyi State University, Abakaliki. Ebonyi State Nigeria Pmb 053 Abakaliki.
2. Department Of Crop And Landscape Management, Faculty Of Agric Sciences, Ebonyi State University, Abakaliki. Ebonyi State. Nigeria Pmb 053 Abakaliki.
3. Department Of Biochemistry, Faculty Of Biological Sciences, Ebonyi State University, Abakaliki Ebonyi State Nigeria Pmb 053 Abakaliki.
4. Department Of Medical Laboratory, Faculty Of Health Sciences Ebonyi State University Abakaliki. Ebonyi State Nigeria Pmb 053 Abakaliki.

Email of the corresponding author: [\\*coscusanas@gmail.com](mailto:coscusanas@gmail.com), coscusans@yahoo.com, Uhuo,Cosmas

### Abstract

The parasitological examination of soil samples extracted from the roots of pre-nursery crops of fruits and vegetables for the prevalence of soil transmitted helminths was conducted using Zinc-Sulphate floatation, centrifugation methods and microscopy respectively. Out of 1,400 soil samples and 1000 fruits examined 58% and 42% respectively were positive. The study revealed that soil transmitted parasites were implicated in their full developmental stage of adult worms, larvae and eggs respectively. These include *Capillaria*, *Strongyloides stercoralis*, *Trichuris trichuria*, *Hookworms*, and *Ascaris lumbricoides*. Their presence could be as a result of using animal dung/feeces as organic fertilizer, direct urination and open defecation on the farmland by students, staff, and other student farmers in the nursery. Thus contaminating the nursery and posing health problems to students, farm workers in the campus and consumers of the contaminated fruits and vegetables. However, infected students could be treated with antihelminthics while avoidance of the use of animal dung/feeces, open defecation and urination on the farmland will reduce the parasite presence. Students treatment on infection, can be achieved through public health awareness scheme and provision of toilet facilities and finally teach them how to wash their fruits and vegetables with salt or potassium iodide which are predisposing factors for infection.

**Keywords:** Helminths, Parasites, Soil, Fruits, Vegetables, EBSU.

### 1. Introduction

In Africa, more than half of the population do not have access to portable water and about two-thirds lack good sanitation means of excreta and urine disposal. Nigeria, like other developing countries is faced with the problem of inadequate disposal of excreta-related human waste discharged into the environment and water bodies. Agbolade *et al*, (2008). Thus, in rural farming communities of eastern Nigeria, promiscuous defecation on open fields and farm lands is still a common practice. CAS campus lack disposal facilities and their students engage on uncommon practice of indiscriminate disposal of excreta commonly called bush method. Excreta related diseases especially helminths that cause gastrointestinal problems in man rank high on the scale of disease burden in mixed populace that lack portable drinking water and good sanitation. WHO (1989) reported that inadequate sanitation, lack of access to clean potable water and poor domestic hygiene are the cause of 80% of all infectious diseases in the world and responsible for 10-25 million deaths each year, most of them under 5 years age group.

These are transmitted through faecal-oral route through faecally contaminated water, food and soil. Communities characterised by poor and no hygiene and a large proportion of children will generate excreta rich in enteric pathogens Wargovich, (2000). Helminth infections are of particular concern in developing countries and many of these parasitic worms have human hosts. To highlight the actual risk to public health affecting student farmers as well as consumers of crops grown on faecally polluted soil as a result of post harvest contamination with helminths eggs, ova and adult worms were studied. The environment of this study provided conducive atmosphere for the prevalence of these soil transmitted helminths. Montessoro *et al*, (2003) also reported that moist soil and average temperature favours the fecundity of soil transmitted helminths and could suggest that this quickens their reproduction. This was carried out in farm produce from crop and landscape management nursery Ebonyi State University Abakaliki to highlight the actual implication to public health affecting student farmers as well as consumers of crops grown on focally polluted soil as a result of postharvest contamination with helminth adult worm, eggs and larvae.

## 2. Methods

### Study Area:

Ebonyi State geographically lies between co-ordinate 70 31` and 80 30` N and between 50 40`, and 60 45` E. The climate of the area is more tropical and the vegetation characteristic is predominantly the rain forest with an average atmospheric temperature of 250-350C. There are two distinct seasons, the rainy and the dry seasons; the former stretches between April and October, while the latter occurs from November to March. The study was conducted in crop and landscape management garden CAS campus Ebonyi State University Abakaliki. The population here is made of student and staff farmers under the management of Kelechi Nwanchor. Sanitation facilities are either non-existent or grossly inadequate in this study area. Thus, defecation on open farm lands is a popular practice among students and staff due to lack of disposing facilities which ensures a predisposing factor to many parasitic diseases.

### Methodology:

1400 soil samples from farm lands were collected from Crop and landscape management garden, College of Agriculture campus Ebonyi State University Abakaliki as well as fruits and vegetables from student farmers were examined for helminths eggs, ova and adultworms using the zinc sulphate flotation technique and microscopy.

## 3. Result and Discussion:

Soil samples from farm lands of students and staff of Ebonyi state university college of agriculture sciences (CAS) as well as fruits and vegetables harvested directly from the farm were examined for helminth eggs and larvae using Zinc-Sulphate floatation, centrifugation methods and microscopy respectively, to highlight the public health significant of using untreated domestic animal dungs as fertilizer. All the fruits and vegetables were found to be contaminated.

Out of the 1,400 soil samples and 1000 fruits examined using zinc-sulphate flotation technique and microscopy, 58% and 42% respectively were positive. Eggs of *Ascaris lumbricoides* (15%); Hookworms (11.8%); *Capillaria* (4%) and larvae of *Strongyloides stercoralis* (1.2%) were recovered from soil. The samples contained *Ascaris lumbricoides* (30.4%); Hookworms (20%); *Capillaria* (9.6%) and *Stongyloides stercoralis* (1%) respectively. Mixed parasite infections were also observed. All the fruits were found to be contaminated with one or more species of helminthes adult worm, eggs, ova or larvae, with the fruits being more contaminated (52.9%) and (47.1%) respectively. *Solanum eathropicum* leaf recorded the highest contamination rate (23%) while *Telferia occidentalis* recorded the least contamination (8.3%). *Solanum eathropicum* and *Solanum melongina* fruits recorded the highest contamination rate (15%) while *Lycopesicum lycopesicum* recorded the least rate of contamination (5.6%) for the fruits. Helminth ova and larvae recorded were *Ascaris lumbricoide*, *Capillaria*, Hookworms, *Strongyloides stercoralis* with *Ascaris lumbricoides* being the most frequently encountered (60.3%). These parasites affect human and animals and are directly linked to faecal contamination which is also reported by Simonart *et al*, (2003). The recovery of helminthic eggs and larvae on the vegetables and fruits is indeed of great public health importance. This shows that hygienic level of the consumers and the behavioural attitude of farmers in application of untreated human and animal dung as organic manure, use of irrigation source which receives raw affluent from human or animal wastes lead to the transmission of zoonotic infection as vegetables harbour the parasites through these sources, Shrestha, (2010). The contamination might have resulted from rain splashes of contaminated soil during heavy rainfall which deposit contaminated soil surface on leaves of vegetables or fallen fruits, (Agbolade, *et al.*, 2008; WHO, 1980). It is not also wrong to say that heavy wind may carry dust particles on leaves and fruits. Some of the vegetables and fruits may be eaten raw or undercooked and might cause infection and disease. This is true that the consumption of raw or undercooked vegetables and fruits play a significant role in human nutrition, especially as source of vitamins (C, A, B<sub>6</sub>) thiamine, niacin, E), minerals and dietary fibre Quebedeaux and Bliss, (1988); Quebedeaux and Eisa, (1990); Wargovich, (2000).

Special care must therefore be taken when introducing sanitation facilities that potentially increases contact between householders and excreta which may contain viable *Ascaris* ova. The risk of handling waste must be clearly identified and methods of reducing risk must be instituted wherever possible. This is because large number of students were found promiscuous in the farm yards using vegetations as cover during the exercise, this could be blamed on the non provision of toilet facilities by the campus management. Helminth infections are of particular concern in developing countries and many of these parasitic worms have human hosts. *Ascaris lumbricoides* is one of the most significant human pathogens in sanitary waste, particularly in developing communities. Its importance derives from the fact that it has ova which are extremely persistent in the environment outside the host. Shrestha *et al*,(2010); Gaspard and Schwarzbrod (1995), reported that an important source of exposure for humans to *Ascaris* ova exists in regions were excreta are used as soil conditioners or

fertilizers, so that both the person handling the waste and those consuming unprocessed crops grown in these soils are at risk of infection.

### Conclusion

Investigations in this study showed that in this campus, large amount of untreated human waste is discharged into the soil daily leading to the seeding of the soil with pathogenic organisms including geohelminth adult worm, eggs and larvae. Sanitation particularly, proper sewage treatment remains a major factor in poor rural farming systems of developing areas like College of Agriculture Ebonyi State university.

**Figure 1: Larva of *Strongyloides stercoralis***



### References

- Agbolade, O.M., Akinboye, D.O., Fajebe, O.T., Abolade, O, A. & Adebambo, A.A. (2008). Human Urinary Schistosomiasis Transmission Foci and Period in an endemic town of Ijebu North, Southwest, Nigeria. *Tropical Biomedical* **21**:15-22.
- Ayres RM, Stott R, Lee DL, Mara DD, & Silva SA (1992) Contamination of lettuces with nematode eggs by spray irrigation with treated and untreated wastewater, *Water Science and Technology* **26**: 1615-1623.
- Faust EC, Beaver PC, & Jung RC (1975) *Animal Agents and Vectors of Human Disease* (Fourth Edition): 237-244, Henry Kimpton Publisher, London.
- Gaspard PG, & Schwartzbrod J (1995) Helminth eggs in wastewater: quantification Helminth parasite eggs in Nepal. Southeast Asian. *Journal of Tropical Medicine Public Health*,**14**(2):115-123
- Quebedeaux & Esia, HM. (1990). Horticulture and human health. Contributions of fruits and vegetables. Production. 2nd International Symposium Horticulture and Human Health. *Horticulture of Science*, **25**:1473-1532.
- Quebedeaux, B. & Bliss, FA. (1988). *Horticulture and Human Health*. Contributions of fruits and vegetables Production. 1st International Symposium of Horticulture and Human Health. Practice hall, Englewood, N.J. 1338 pp.
- Schonning C, & Stenstrom TA (2004) Guidelines on the Safe use of Urine and Faeces in Ecological Sanitation Systems. EcoSan Res Programme, Stockholm Environmental Institute, **32**(1): 388-393.
- Shrestha, B. (2010). Intestinal Parasitic Infection in Health School Children of Lalitpur District. *Journal of Nepal Medical Association*, **41**:266-270.
- Simonart T, Roussel S, & Gireaudot-Liepman MF (2003) Desk study on European standard for the enumeration of viable helminth ova in sludge, soil and solid waste, Horizontal WP3-5-Draft Report on Helminth ova. Techniques, *Water Science and Technology* **31**: 443- 446.
- Wargovich, M.J. (2000). Anticancer properties of fruits and vegetables. *Horticulture of Science* **35**:573- 575.
- WHO Scientific Group (1989). Health Guidelines for the Use of Wastewater in Agriculture and Aquaculture. World Health Organization. (1980). *Parasite related diarrhoeas*. Scientific Working Group Series No 500. World Health Organization. Geneva, 54.

### Acknowledgement

The authors are highly delighted to acknowledge the effort of Nwanchor Kelechi and staff of crops and landscape management for giving us the research tools to carry out this study. The principal researcher Uhoo Cosmas is highly appreciated for his major contributions.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

## CALL FOR PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/Journals/>

The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

## IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

