

Genetic Variation of Coconut Tall (*Cocos nucifera* L., Arecaceae) in Bali, Indonesia Based on Microsatellite DNA.

Eniek Kriswiyanti*, I Gede Rai Maya Temaja**, I Made Sudana**
 and I Gusti Ngurah Alit Susanta Wirya**

*Post Graduate Student of Agriculture Faculty and *Department of Biology Mathc and Natural Science Faculty*,
Udayana University Campus Bukit Jimbaran, Kuta Bali Indonesia ** *Faculty of Agriculture, Udayana*
University Campus Sudirman, Denpasar, Bali Indonesia

Corresponding author, Email : eniek_kriswiyanti@yahoo.co.id

The research was funded by Director General of Higher Education, Ministry of Education and Culture of Republic Indonesia (grant of Doctorate Student) and Udayana University, Bali Indonesia

Abstract

The coconut has important roles for economics, traditional medicine and culture, especially for Hindu's ceremonial purpose in Bali Island (Indonesia). Each coconut cultivar has unique characteristics. The aimed of this research was to determine genetic variation of coconut tall (*Cocos nucifera* L., Arecaceae) in Bali, based on DNA microsatellite. Six pairs microsatellite markers were used to determine heterozygosity. The results showed that total of 80 alleles were detected by microsatellite with an average of 13.33 alleles per locus, there were 12 alleles on microsatellite locus CnCirA3, 12 alleles on locus CnCirC3, 16 alleles on locus CNZ05, 14 alleles on CNZ09, 17 alleles on CNZ21, and 9 alleles on microsatellite primer pairs CNZ51. The mean values of gene diversity (He) and observed heterozygosity (Ho) were 0.8835 and 0.5421, respectively. The highest heterozygosity was on *bulan tall* coconuts cultivar (0.816), the lowest heterozygosity was on *bluluk tall* coconuts (0.35).

Keywords: unique characteristic, microsatellite, allele, heterozygosity, coconut

1.Introduction

Coconut (*Cocos nucifera* L.) is one of the primary sources of income for most Indonesian farmers. Coconut fruit is processed into various products like vegetable oil, raw material for food, industry, medicine, cosmetic and oleo chemicals. Total land area plantation of coconut is about 3.7 million hectares, with 97% of overall production from farmers (Novariant, et al., 2005). In Bali, beside of its economic value, parts of coconut tree (leaves, fruit, young flower) also used for ceremonial and medicinal purposes, which called "nyuh madan". Morphologically, some coconut plants differ in some characters, which is very unique and specific to its individual plant. Based on the unique characters, there were more than twenty unique coconuts identified that differ individually. Some of them are *ancak* which has branches on its trunk; the *be julit* has plicated lamina leaves, the *bingin* 's root grow from nodes of stem, the *Bojog* 's fruit husk is colored like the hair of long tailed monkey. Differences in fruit color such as white, green, yellow and red were used to distinguish between the *bulan*, *gadang*, *gading* and *surya* coconuts respectively. Inflorescentia spicata was characteristic of *bluluk*, while the *udang* and *mulung* were characterized by red mesocarpium. The *Rangda* coconut has petiole and the apex of the stem were twisted. The *sudamala* was characterized by many kinds of unique characters, which some of them are doubled spatha, and flat apex of male spikelet (Kriswiyanti, 2012).

To study its genetic diversity, simple sequence repeats (SSR) microsatellite markers were used. This technique had been used successfully by many scientists to characterize the genetic diversity of the coconut population (Perera et al. 2001; Manimekalai and Nagarajan 2006; Kumar et al., 2011). The microsatellite analysis of the 9 coconut populations with 8 primers revealed a total of 37 alleles (Devakumar et.al., 2010). From the used 26 SSR markers, detected a total of 188 alleles with an average of 7.23 per locus (Liu, et.al., 2011). In this research, genetic variation of *tall coconut* (*Cocos nucifera* L., Arecaceae) was determined based on DNA microsatellite, using alleles size on each individual of tall coconut. The alleles data from each accession can be applied to predict natural breeding for unique coconut conservation purpose.

Materials and Methods

Plant material used: Leaf samples were collected from Klungkung, Bangli, Karangasem, Gianyar, Tabanan, Badung, Jembrana residence and Denpasar, Bali province, Indonesia. Of the 58 samples from twenty cultivars

were ‘tall coconut’ category (*nyuh anak, biasa, bulan, bingin, gadang, gading, kebo, srogsogan, mulung, rangda, bluluk, sudamala, surya, udang, kapas, be julit, bojog, macan, naga, pudak*). DNA extraction and detection of microsatellite polymorphisms were analysed at Forensics and Primata Laboratory Udayana University, Bali Indonesia.

DNA extraction: DNA was extracted from fresh coconut young leaves using a CTAB based protocol modified from Doyle and Doyle (1987). The primer sequences and associated information are given in Table 1 (Pandin, et.al., 2008).

SSR analysis

Polymerase Chain Reactions (PCR) assay and gel analysis : DNA was amplified in 13 µL reactions containing 2 µL sample, 3.5 µL H₂O, 6.5 µL Mastermix (Qiagen), and 1 µL primer. The PCR programmed for 30 cycles of 45 seconds each at 94°C, 1 min at the different annealing temperatures standardized for the individual SSR locus, and extension for 1 min 30 seconds at 72°C. The first cycle was preceded by a 5 min denaturation at 94°C and the last cycle ended with 5 min extension at 72° C. Reaction products were separated on 6% polyacrylamide (denatured) and visualized with silver nitrate staining (Tegelstorm, 1984).The alleles were scored based on the size of each PCR amplified fragment by electrophoresis all samples in a single gel. Allele size were determined by semilog plotting of distance migration of amplicon on PAGE (Hutchinson, 2001). Diversity values based on allele frequencies were calculated for each microsatellite locus and coconut cultivar using Nei’s methods (1987).

Table .1 Details of microsatellites primer used in the analysis

Primer	Urutan Basa	
	Forward (5’-3’)	Riverse (5’ – 3’)
CNZ 05	CTTATCCAAATCGTCACAGAG	AGGAGAAGCCAGGAAAGATTT
CNZ 09	ATCTACCAGTGTGGTCTCTC	ACCAGGAAAAAGAGCGGAGAA
CNZ 21	ATGTTTTAGCTTCACCATGAA	TCAAGTTCAAGAAGACCTTTG
CNZ51	CTTTAGGGAAAAAGGACTGAG	ATCCATGAGCTGAGCTTGAAC
CnCir A3	AATCTAAATCTACGAAAGCA	AATAATGTGAAAAAGCAAAG
CnCir C3	AGAAAGCTGAGAGGGAGGATT	GTGGGGCATGAAAAGTAAC

Results

From 58 extracted DNA samples, some sample were successfully amplified on each primer used. For examples locus CNZ09 on figure 1, C and D were not successfully amplified.




Figure 1. An example of allelic polymorphisme at microsatellite locus CNZ09 in 12 individuals coconut tall from some village in Bali island . A.*Be Julit tall* Babung, B *Be Julit tall* Buruan, C *Be Julit tall* Tuwed D. *Be Julit tall* Tamblang, E.*Biasa tall* Pejeng, F.*Bingin tall* Jelekungkang, G.*Bingin tall* Babung, H. *Biasa tall* Babung, I.*Ancak tall* Pikat, J.*Surya tall* Jelekungkang, K.*Gadang tall* Babung, L.*Rangda tall* Babung

Genetic Variation (Heterozygosity)

Total number of investigated alleles, number alleles, heterozygosity, and variation alleles length of twenty tall coconut cultivars for each SSR locus were showed in Table1. The combination of six SSR loci generated a total of 80 alleles, with a mean of 13.33 allele per locus and ranging from nine (CNZ51) to seventeen alleles (CNZ21).

Gene diversity range from 0.8561 for the loci CNZ51 to 0.9204 for the loci CNZ21, with an overall mean 0.8835. The loci CnCirC3 and CNZ05 presented the lowest (0.45) and the highest value (0.7619) respectively, for observed heterozygosity, with a mean value of 0.5421.

Table 1. Number of alleles per locus, heterozygosity and variation in allele length of Bali tall coconut, using 6 SSR loci

Primer	Number of amplified samples	Number of allele	Gene diversity/ Expected Heterozygosity (He±SE)	Observed Heterozygosity (Ho)
CnCirA3	32	12	0.8922±0.0001	0.4687
CnCirC3	40	12	0.862±0.0005	0.45
CNZ05	42	16	0.9093±0.0012	0.7619
CNZ09	47	14	0.8612±0.0003	0.4893
CNZ21	42	17	0.9204±0.0002	0.5714
CNZ51	43	9	0.8561 ±0.0005	0.5116
Mean±SD		13.333±2.9	0.8835±0.027	0.5421

Heterozygosity among the coconut cultivars was showed in Table 2. The expected heterozygosity range 0.35 (Bluluk tall) to 0.8166 (Bulan tall), with a mean of 0.5941.

The highest heterozygosity was in Bulan tall (0.776) and the lowest value in Pudak tall and Kapas tall (0.33).

Table 2. Description and heterozygosity of tall coconuts from Bali, Indonesia

Coconut Name	Description	Expected Heterozygosity (He) ± SD		Observed Heterozygosity (Ho)	Number of sample
Ancak tall	branched trunk	0.6458	0.2186	0.5766	3
Bejulut tall	plicated lamina of young leaves	0.5989	0.163	0.6833	3
Biasa tall	Ordinary tall	0.4835	0.351	0.264	3
Bingin tall	the root growth from nodes trunk	0.6331	0.259	0.6067	3
Bojog tall	fruit husk colored like the hair of long tailed monkey	0.6043	0.177	0.4716	3
Bulan tall	White fruit	0.8166	0.154	0.776	3
Gadang tall	Green fruit with sweet water	0.6777	0.333	0.6067	3
Gading tall	Golden yellow fruit	0.8042	0.2106	0.526	3
Srogsogan tall	Damage endosperm = kopyor	0.4525	0.225	0.36	3
Mulung tall	Green fruit with red mesocarp	0.5805	0.2061	0.608	3
Rangda tall	petiole and the apex of the stem were twisted	0.4199	0.363	0.358	3
Bluluk tall	Inflorescentia spicata	0.35	0.418	0.415	4
Sudamala tall	Double spatha, reduction of some male spikelet , tip of primordial leaves like hook	0.6833	0.178	0.553	3
Surya tall	Red fruit and tip lamina (old)	0.7137	0.208	0.555	3
Udang tall	Brown fruit with red mesocarp	0.6102	0.131	0.34	3
Kapas tall	Edible white husk (immature)	0.4166	0.52	0.333	2
Kebo tall	Small nut with thick husk	0.6498	0.392	0.5	3
Macan tall	Black spot on fruit skin	0.6748	0.186	0.416	3
Naga tall	Green fruit colour, angled of stem	0.6833	0.2185	0.4666	3
Pudak tall	Brown fruit colour, spatha>2	0.3832	0.525	0.33	2
Mean		0.5941		0.4872	

Discussion

The alleles size of several locus from tall coconut that explored during this research differed to that found in other studies of coconut tree population using same SSR markers. Alleles size on locus CnCirA3 range 210-280 bp, with highest frequency on 255bp allele (0,203). Rajesh *et.al.*,(2008) and Ribeiro et al (2010) found 228-240bp alleles sizes. The alleles sizes on the CNZ21 locus that range 180-300bp with highest frequency on 240 bp (0,207), on locus CNZ51 range 110-230 bp with the highest frequency 0,221 on 140 and 160 bp. Pandin et al (2009) found 270bp on loci CNZ21 and 110bp on loci CNZ5 .

Total alleles were found in this research were 80 alleles, with range 9-17 alleles and mean values was 13,3. This result had more variation than others. Perera et al. (2001) found 56 alleles sizes range 3-10 on 33 samples coconut in Srilangka used 8 loci. Thirty seven alleles were found on tall and dwarf from nine population from Agatti and Kavaratti Island, Lakshadweep, India (Devakumar *et.al.*, 2010). Other research that used 8 microsatellites loci from 14 accession from Kidu India coconut population, found 28 alleles with mean two alleles per locus (Kumar *et al.*, 2011). Overall, genetic diversity in tall coconuts in Bali was very high (Table 1). Mean values of gene diversity (H_e) and observed heterozygosity (H_o) were 0.8835 and 0.5421, respectively. The highest expected heterozygosity was on loci CNZ21 (0.9204), and the lowest gene diversity on CNZ51 (0.8561). Genetic variation found in this result higher than others research, e.g research coconut population in Brazil, Phillipina, India, Sri Langka (Perera et al, 2003; Devakumar *et al.*, 2010; Liu, *at al.*,2011; Xiao, *et al.*, 2013).

Genetic variation on the tall coconut cultivars range value 0.35-0.8166 (Table 2), high expected heterozygosity on bulan tall (0.816), and lowest on bluluk tall (0.35). The estimate of heterozygosity was high (0.590) for Laccadive Micro Tall and Laccadive Small Tall was lowest (0,240) (Devakumar, *et.al* 2010). Genetic diversity of 10 coconut accessions from six location in Hainan province, China; expected heterozygosity of Haikou Green Tall accession was significantly higher (0.4753) than other tall type (Liu *et al.*, 2011)

Higher heterozygosity was determined by number of alleles variation and frequency. Higher diversity in tall coconuts in Bali, Indonesia was caused by germplasm variation or natural mutation.

References

- Devakumar, K; V.Niral, B.A Jerard, C Jayabose, R Chandramohan and P M Jacob. (2010). Microsatellite Analysis of Distint Coconut Accsseions from Agatti and Kavaratti Island Lakshadweep, India. *Scientia Hort* 125 (2010) : 309-315
- Doyle J.J. and Doyle J.L. (1987). A Rapid DNA Isolation Procedure for Small Amount of Leaf Tissue. *Phytochem. Bull.* 19 : 11-15.
- Hutchinson, F.(2001). DNA Band Size Semi-Log Plotting: Estimation Size of DNA Band by Semi-Log. Cancer Research Center : Science Education Patnership.
- Kriswiyanti, E.(2012). Diversity of Coconut Cultivar (*Cocos nucifera* L., Arecaceae) in Bali Island, Indonesia. Research report, Department of Biology Udayana University
- Kumar, S. P; Manimekalai, R. and Kumari B.D.R. 2011. Microsatellite Marker Based Characterization of South Pasific Coconut (*Cocos nucifera* L.) Accessions. *Int. J. Plant Breed. Genet.*, 5: 34-43.
- Liu, X; Tang,H; Li, D and Hou, L. (2011). Genetic Diversity of Coconut Cultivars in China by Microsatellite (SSR).*Moleculer Plant Breeding* 2 (2):
- Manimekalai, R and Nagarajan, P. (2010) : SSR and ISSR Markers Based Population on Genetic Structure of Coconut (*Cocos nucifera* L.) Germplasm Accssion Indian J. of Plant Genetic Resources, 23 (1) : 77-82.
- Nei, M.(1987). *Moleculer Evolusionary Genetic*. Columbia University Press, New York, USA
- Noel, K K J., Edmond, K.K., Konan, K J L and Eugene, K.K. (2011). Microsatellite gene diversity within Philippines dwarf coconut palm (*Cocos nucifera* L.) resources at Port-Bout d'Ivoire. *Scientific research Essays* 6 (28): 5986-5992.
- Novarianto, H., Akuba, R.H., Mashud, N., Tenda, E and Kumaunang, J.(2005). Status of Coconut Genetic Resoures Research in Indonesia. In *Coconut Genetic Resources*. Batugal, P., Ramanatha Rao, V. and Oliver, J.

(eds). International Plant Genetic Resources Institute – Regional Office for Asia, the Pacific and Oceania (IPGRI-APO), Serdang, Selangor D.E, Malaysia, 608-617.

Pandin, D.S., Hartana, A., Aswidinnoor, H. and Setiawan, A. (2008). Parentage analysis of Mapanget Tall Coconut No.32 (DMT-32) population via gene flow based on Micro-satellite Markers (SSR). *Littri Journal*, 14 (4) : 131-140.

Pandin, D.S.(2010). DNA Marker in Coconut (*Cocos nucifera* L.) Breeding Programm. *Perspektif* 9 (1): 21-35.

Perera, L; Russel, R., Provan, J.and Powell, W. (2001). Levels and Distribution of Genetic Diversity of Coconut (*Cocos nucifera* L., var *Typica* from *typica*) from Sri Lanka by Microsatellite Markers. *Euphytica* 122 : 381-389.

Perera L., Russell J.R., Provan J., and Powell W., 2003, Studying genetic relationships among coconut varieties/populations using microsatellitemarkers, *Euphytica*, 132(1): 121-128.

Sambrook, J., Fritsch, and Maniatis, T. (1989). *Molecular Cloning. A Laboratory Manual* Second Edition. Cold Spring Harbor Laboratory Press.

Tegelstöm H. (1986). Mitochondrial DNA in Natural Population : an Improved Routine for Screening of Genetic Variation Base on Sensitive Silver Staining. *Electrophoresis*. 7: 226-229

Xiai, Y., Lou, Y., Yang, Y., Fan, H., Xia, W., Mason, A.S., Zhao, S., Sager, R., and Qiao, F. (2013). Development of Microsatellite Markers in *Cocos nucifera* and their application in evaluating the level of Genetic Diversity of *Cocos nucifera* *POJ*: 6(3):193-200.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Recent conferences: <http://www.iiste.org/conference/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

