

Investigation on the Effect of Chemical and Biological Control of Bacterial Soft Root Disease of Potato in Storage.

Abeer H. Makhoul⁽¹⁾ and rehab abdeen⁽²⁾.

1. Faculty of Agriculture, Minufiya University,

2. Health science program, King Khalid University, Saudi Arabia.

hyabeer@yahoo.com,* rabdeen@kku.edu.sa

Abstract

Potato soft rot, caused by *Erwinia carotovora* subsp. *carotovora*, greatly affect potato tuber quality in storage of Egypt and indicated that it can affect all potato cultivars. *Erwinia carotovora* subsp. *carotovora* was isolated from the infected potato tuber and was identified by pathological, morphological and biochemical studies. One antimicrobial chemical compound chitosans (CS) with concentration (1, 3, 5 %) combine with three biocontrol agents (*Bacillus subtilis*, *Pseudomonas fluorescens* and *Trichoderma viridi*) each. These treatments were screened out in vitro and in storage against the growth of *Erwinia carotovora* subsp. *carotovora*. All bio-control agents combined with Cs reduced the bacterial soft rot disease to various degrees. The stronger antagonistic activity against *Erwinia carotovora* was found in treatment Cs 5% with biocontrol agents (*T. viridi*, *P. fluorescens* and *B. subtilis* respectively). All treatments reduced the soft rot infection to 20-week storage with two types of potatoes (*Spunta* and *Cara*) varieties by (91, 86, 83.6 and 77.3% respectively in *Spunta* c.v.) and (88.6, 86.4 and 79.8% in *Cara* c.v.). The lowest antagonistic activity against *Erwinia carotovora* was found in treatment CS 1% with biocontrol agents (*T. viridi*, *P. fluorescens* and *B. subtilis* respectively) varieties by (64.2, 58.6 and 43.7% respectively) compared with the treatments of biocontrol agents individually.

Keywords: Potato - *Erwinia carotovora*- Chitosan- *T. viridi* - *P. fluorescens*- *Bacillus subtilis*

1. Introduction

Potato (*Solanum tuberosum*) is a worldwide cultivated tuber-bearing plant which is the fourth main food crop in the world after rice (*Oryza sativa*), maize (*Zea mays*) and wheat (*Triticum aestivum*), (Douches et al., 1996). Potato does not require special growth conditions; it has been for a long time a major field crop in temperate regions, and increasingly in warmer regions (Haverkort, 1990). In Egypt, potato crop has an important position among all vegetable crops, where about 20% of total area devoted for vegetable production is cultivated with potato. In addition, the total cultivation of potatoes reached 197,250 feddans which produce 2,039,350 tons of tubers with an average yield of 10.34 tons/feddans (Douches et al., 1996). This crop is economically important to Egypt and disturbance in its production affects severely its local and export impact.

Harmful bacteria or pathogens can affect agricultural crops and produce which results in economical losses (Bathily et al., 2010). *Erwinia carotovora*, a plant pathogen, causes bacterial soft rot in vegetables such as potatoes, cabbages, carrots and many other plants (Altin, N. & Bora, T., 2001). Soft rot is a serious situation in wet and warm areas after harvest. When vegetables are harvested or transported to storage, small cuts and bruises might be inflicted on them. This allows *Erwinia carotovora* to enter the vegetable and infect it with soft rot, causing it to deteriorate over the next few days. *E. carotovora* produce extracellular pectinase and breaks down plant tissue by making them osmotically fragile. Signs of soft rot can be seen when the vegetable becomes slimy, mushy and has a darkened area. It also gives off a foul odour. This causes most of the infected crops to be unsuitable for consumption, leading to economical losses as they have to be discarded.

Biological-control treatments consisting of living microorganisms or abiotic products can provide disease protection essentially through one or more of the following: (i) production of antibiotics or other molecules that are deleterious to the pathogen's development, (ii) competition with the pathogen for nutrients and space, or (iii) induced plant resistance. It has more specific effect on the pathogen and has limited impact on the environment (Sigeo, 1993). The strategy for biological control of tubers involves the use of antagonistic microorganisms. The bacteria (*Bacillus subtilis* and *Pseudomonas fluorescens*) and the fungi *Trichoderma viridi*. *Bacillus subtilis* shows biological activity against phytopathogenic bacteria by producing peptide antibiotics (Backman et al., 1997) and reducing the soft rot decaying stored potato tubers (Klopper et al., 2004). Soil fluorescent and non-fluorescent *Pseudomonas* spp. have shown biological control of soft rot disease of potato by producing a variety

of secondary antibacterial metabolites including siderophores, antibiotics and surfactants (Compant et al., 2005). *Trichoderma Viride* is the potential antagonistic fungus which prevents the crops from diseases and able to suppress more than 60 species of pathogens (Abd-El-Khair & Karima, 2007)

In recent years there has been an increasing interest in finding alternatives to bactericides and fungicides. Chemical-control treatments considered as safe, with negligible risk to human health and environment among these strategies, Chitosan (CS) is one of the most abundant biological polysaccharide derived from nature and is obtained by deacetylation of chitin present in crustaceans. Chitosan is composed of linked 2-deoxy-2-amino-D-glucopyranose units and β (1 \rightarrow 4) linked 2-deoxy-2-acetamido-D-glucopyranose units Figure (1). In spite of its abundance in nature, the commercial utilization of chitosan has been developed only over the last 2 decades; it has emerged as a new biomaterial for food (Rhoades & Roller, 2000), pharmaceutical (Illum, 1998 and Rouget, 1859), medical (Tharanathan & Kittur, 2003 and Utaida et al., 2003), textile (Takai et al., 2002), agricultural (Dodane & Vilivalam, 1998), and other industries, as well as for waste water purification (Babel & Kurniawan, 2003).




Figure (1): structure of chitosan.

Some satisfactory results have been reported using natural compounds such as chitosan (Muzzarelli, 1983). Chitosan and its derivatives have attracted much attention as antimicrobial agents against fungi, bacteria, viruses and as elicitors of plant defense mechanisms (Chirkov, 2002, Helander et al., 2001 and O'Byrne & Booth, 2002). In fact, a number of commercial applications of chitosan benefit from its antimicrobial activity, including its use in food preservation (Rhoades & Roller, 2000), manufacture of wound dressings (Tharanathan & Kittur, 2003) and antimicrobial-finished textiles (Takai et al., 2002). The lack of understanding of how this industrially valuable biopolymer exerts its antibacterial activities led us to our more systematic study of its mechanism of action. Taking these antecedents into account, the aim of the study reported here was to test the effect of bio-control agents (*Bacillus subtilis*, *Pseudomonas fluorescens* and *Trichoderma viridi*) individually, biochemical on *Erwinia carotovora* subsp. Followed by testing the effect of combining biochemical and bio-control agent. Then study the effect of this treatment on protein, starch and total sugar content and also define dry matter and specific graphite.

1.1. MATERIALS AND METHODS

1.1.1. Materials

Chitosan (CS) with the deacetylation degree of 70 % with medium molecular weight (Mw) was purchased from Aldrich chemicals.

1.1.2. Potato tuber source

The local farmers provided the fresh mature tubers of potato variety Cara and Spunta. Apparently healthy and uniformly sized tubers were selected for the experiments. These selected tubers were washed under running tap water and stored in the laboratory conditions prior to the onset of the experiments.

1.1.3. Isolation and purification of the causal organism

Erwinia carotovora subsp. *carotovora* was isolated from infected potato tubers, and identified as the principal pathogen of potato soft rot in storage. *E. carotovora* cultures were maintained on potato dextrose agar (PDA) slants at 4°C. The spores suspension was prepared from 3-day-old cultures grown at 30°C in nutrient broth medium. After the incubation period the medium was centrifuged at 3000 rpm for 20 min and the pellet was

collected in sterile distilled water (which was considered as bacterial spore and cell suspension) and diluted to a concentration of 6×10^6 spores/ml (Dodane & Vilivalam, 1998)

1.1.4. Identification of the causal organism

Morphological tests shape of cell and Motility. Physiological and Biochemical tests Cavity formation on crystal violet pectate (CVP) medium and fluorescence on King's B medium were performed as described earlier (Kelman, 1974 and Fang, 1998). The tests used in this study were Gram staining (Gerhardt, 1981), growth in 1-5% NaCl and at 37°C performed as described in Bergey's Manual of Systematic Bacteriology (Garrity et al., 1984), Catalase and Nitrate test (Schaad et al., 2001), Strach hydrolysis test (Cowan, 1974), V.P test, Methyl red reaction and H₂S production (Bergey's, 1984) and acid production from carbohydrates utilized as source of carbon (Schaad et al., 2001) glucose, maltose, sucrose, lactose, inositol and α -methyl glucoside, amylolysis, utilization of citrate.

1.1.5. Isolation and identification of Biocontrol agents

Six samples from the rhizospheric soils were taken (up to 10 cm depth). Samples were placed in sterile polyethylene bags, closed tightly and stored in the refrigerator at 4°C until use. Samples of each rhizospheric soil were first mixed, suspended in sterile distilled water (1 g in 100 ml) and shaken on rotatory shaker (200 rev/min, 30 min). All treated samples were serially diluted up to 10⁶ and spread (0.1 ml) over the surface of nutrient agar (Difco, USA) and soil extract agar (Barakate et al., 2002). When the bacterial colony appeared on the medium, representative isolates were picked for this study. Pure cultures of biocontrol agent strains were identified using the morphological and physiological characteristics according to the methods of (Schaad et al., 2001, Lelliot & Stead, 1987 and Klement et al., 1990).

1.1.6. In Vitro Antimicrobial Activity on Petri Dishes

Antagonistic activity of the bacterial isolates was tested in vitro using plate chloroform method (Wakimoto et al., 1986 and Furuya et al., 1997). Briefly, one loop full of 1-2-day-old probable antagonistic bacterial colony (*Bacillus subtilis* and *Pseudomonas fluorescens*) grown in yeast peptone dextrose agar YPDA medium was transferred to the center of a petri dish containing 20 mL of YPDA with four replicates of each bacteria. The plates were incubated at 30°C for 2 to 3 days. When the bacteria colonies were formed as several millimeters in diameter, the plate was turned upside down. A sheet of filter paper was placed in the petri dish lid with 0.5 mL of chloroform. The dish was kept at room temperature for 2 h. After complete evaporation of chloroform, 5 mL suspension of pathogenic bacteria (ca. 10⁸ cfu/mL) was overlaid on each plate. One plate of the soft rot bacterial strain, *E. carotovora* subsp. *Carotovora* was used as an indicator bacterium. The plate thus prepared was incubated at 30°C for 2 days. When an inhibition zone appeared, its diameter was measured to evaluate the antibacterial activity of the probable antagonistic bacteria (Furuya et al., 1997). The bacterial isolates which showed antagonistic effect against indicator bacteria were selected for the further studies. But antagonistic activity of the fungal isolate *T. viridi* was tested in vitro using disc diffusion method (Furuya et al., 1997). A disc of 5 mm in diameter from *T. viridi* (from 7 days old culture) was placed on the surface of OMA plates seeded with potato scab pathogen. The plates were incubated at 28 °C for 48 h. The inhibition zone around the discs indicated the antagonistic interaction.

1.1.7. Preparation of different concentration of chemical control chitosan.

Under magnetic stirring 1 g of chitosan mixed with 500 ml of 10 % acetic anhydride the mixture was allowed to stir in room temperature for 24 hrs to form a homogenous solution. The same procedure was repeated with 3g and 5 g to prepare 3% and 5% respectively.

1.1.8. Tuber inoculation

Prior to inoculation, the tubers were surface sterilized in 10.0% sodium hypochlorite solution for three minutes followed by washing in five changes of sterile tap water. Tubers were allowed to dry at room temperature (about 30°C). After drying, the tubers were wounded (except control) by punching 5 holes about 5 mm deep using 2 mm thick nails. Next the tubers were inoculated by submerging in the bacterial suspension for ten minutes. After the inoculation the tubers were allowed to dry at room temperature for thirty minutes prior to the control agent.

1.1.9. Biological and chemical Control of Soft Rot Disease under Storage Conditions.

To evaluate the effectiveness of the selected antagonistic bacteria, fungi and chitosan in reducing soft rot infection in storage potatoes Table 1, fresh tubers of each two potato varieties, Spinta and Cara with six replicates of each treatment were dipped in suspensions of antagonistic bacterium, fungi and chitosan for 30 min and air-dried separately. The treated potato tubers were inoculated with soft rot bacteria *E. carotovora* subsp. *Carotovora* by spraying with inoculum suspensions (10^6 – 10^7 cfu/mL).

Table (1): Seventeen treatments were tested on potato plants.

1. Control (non sterilized tuber).

2. Tuber infestation with *E. carotovora*

3. Tuber infestation with CS (1%)	6. Tuber infestation with <i>Bacillus subtilis</i> .	9. Tuber infestation with <i>E. carotovora</i> + <i>Bacillus subtilis</i> + CS (1 %)
		10. Tuber infestation with <i>E. carotovora</i> + <i>Bacillus subtilis</i> + CS (3 %)
		11. Tuber infestation with <i>E. carotovora</i> + <i>Bacillus subtilis</i> + CS (5 %)
4. Tuber infestation with CS (3%)	7. Tuber infestation with <i>Pseudomonas fluorescens</i>	12. Tuber infestation with <i>E. carotovora</i> + <i>Pseudomonas fluorescens</i> + CS (1 %)
		13. Tuber infestation with <i>E. carotovora</i> + <i>Pseudomonas fluorescens</i> + CS (3 %)
		14. Tuber infestation with <i>E. carotovora</i> + <i>Pseudomonas fluorescens</i> + CS (5 %)
5. Tuber infestation with CS (5%)	8. Tuber infestation with <i>Trichoderma viridi</i>	15. Tuber infestation with <i>E. carotovora</i> + <i>Trichoderma viridi</i> + CS (1 %)
		16. Tuber infestation with <i>E. carotovora</i> + <i>Trichoderma viridi</i> + CS (3 %)
		17. Tuber infestation with <i>E. carotovora</i> + <i>Trichoderma viridi</i> + CS (5 %)

Inoculated potato tubers were stored separately at room temperature. Data on soft rot incidence was recorded after 20 weeks of inoculation. Number and weight of soft rot infected tubers were recorded and expressed in percentage using the following formula (Abd-El-Khair and Karima, 2007):

Infection % = number of infected tubers / Total number of tubers × 100.

Loss of weight % = Initial weight – weight after discarding the infected sample / Initial weight × 100

Percentage of disease reduction (PDR) was calculated according to the following formula (Hajhamed et al., 2007):

$$PDR = \frac{Ack - Atr}{Ack} \times 100, \quad (2)$$

Where Ack and Atr represent the severity of the disease in control and treated samples, respectively. Total sugar (Dubois et al., 1956), protein content (Thomes & Dutcher, 1924), specific gravity (S.g) (Nissen, 1955), starch content (Simmonds, 1977) and dry weight percentage of tubers (A.O.A.C. 1990).

1.1.10. Data analysis

The collected data were subjected to the analysis of variance and the mean compared either using least significant difference (LSD) or plotting standard error of -means at 5% level of probability (Gomez & Gomez, 1984).

1.2. Results and Discussion

1.2.1. Isolation and Identification of the causal organism

The isolated bacteria identified as *Erwinia carotovora* subsp. *Carotovora* according to its morphological and biochemical characters (Table 2). The bacterium was rod shaped with rounded ends, convex, creamy white colonies cells appeared singly and also in pairs. Abd El- Khair and Karima found similar types of colonies (Abd-El-Khair and Karima, 2007). The isolated bacteria were Gram negative. No results were obtained with Starch hydrolysis, Nitrate reduction and V.P test. Positive results were obtained with methyl red reaction, indole formation and bacteria were grew under anaerobic condition. The production of acid from Fructose, Galactose and Glucose were positive results but it was negative from Arabinose, Lactose and Maltose.

(Table 2). Morphological, physiological and biochemical characters of the pathogenic bacterial isolate *E. carotovora* obtained from naturally infected potato tubers.

Character	Bacterial isolate Od23
Shape of cell	Rod
Motility	+
Gram Reaction	-
Spore forming	+
Growth in 1-5 NaCl	+
Catalase production	+
Yellow pigment	+
Starch hydrolysis	-
Nitrate reduction	-
V.P test	-
Methyl red reaction	+
Indole formation	+
Aerobiosis	+
H ₂ S production	±
Carbon source utilization	
Arabinose	-
Fructose	+
Galactose	+
Glucose	+
Lactose	-
Mannitol	+
Maltose	-

(+) = Positive reaction; (-) = Negative reaction; (±) = Weekly reaction

1.2.2. Characterization and Identification of o biocontrol agents.

Two biocontrol bacteria , *Bacillus subtilis* and *Pseudomonas fluorescens* were identified according to their morphological, cultural and physiological characteristic as stated in Bergey's Manual of Systematic Bacteriology, *Bacillus subtilis* was rod ,long, grame positive , sporulating , motile, negative in Pigmentation and soft rot . The tested isolate *Bacillus subtilis* resulted positive reactions with fructose, sucrose, glucose, galactose, maltose, dextrose, glycerol, starch hydrolysis, gelatin liquefaction, V.P. Test and Catalase activity. Negative reactions with mannitol, arabinose, lactose, menthol, raffinose and Indole formation. *Pseudomonas fluorescens* was rod, short, grame negative, non-sporulating and motile, negative in soft rot. the tested isolate *Pseudomonas fluorescens* resulted positive reactions with fructose, glucose, maltose , glycerol and Catalase activity . Negative reactions with Mannitol, sucrose,arabinose, , galactose, lactose, dextrose, menthol, raffinose, starch hydrolysis ,gelatin liquefaction , Indole formation and V.P. Test (Table 3).

On the basis of the above physiological, biochemical, and carbon sources utilization test results, the antagonistic bacterial were identified as a member of the genus *Bacillus subtilis* and *Pseudomonas fluorescens*. The fungal biocontrol agent *Trichoderma viridi* was identified according to its morphological, cultural characteristic. Conidial wart is conspicuous and Conidiophores are Irregularly branched; branches usually not paired on PDA after incubation at optimum temperature for 48 h (Gams % Bisset, 1998 and Rifai, 1969).

(Table 3). Characterization and identification of two bacterial strains *Pseudomonas fluorescens* and *Bacillus subtilis*.

Character	<i>Pseudomonas fluorescens</i>	<i>Bacillus subtilis</i>
Shape of cells	Rod	Rod
Size	Short	Long
Gram's staining	-	+
Sporulation	-	+
Motility	+	+
Pigmentation	+	-
Soft rot	-	-
Growth in 1- 5%	+	+
Utilization of sugar:		
Mannitol	-	-
Fructose	A	A
Sucrose	-	A
Arabinose	-	-
Glucose	A	A
Galactose	-	A
Lactose	-	-
Maltose	A	A
Dextrose	-	A
Glycerol	A	A
Menthol	-	-
Raffinose	-	-
Physiological characteristics		
Starch hydrolysis	-	+
Gelatin liquefaction	-	+
Indole formation	-	-
V.P. Test	-	+
Catalase activity	+	+
Oxidase reaction	-	-

(+): Positive reaction or growth, (-): Negative reaction or no growth, (A): Acid

1.2.3. In Vitro antimicrobial and chemical activity on petri dishes.

All biocontrol agents, *Bacillus subtilis*, *Pseudomonas fluorescens*, *Trichoderma viridi* and chemical control Cs by concentration 1, 3, 5% were significantly superior in inhibiting the growth of *Erwinia carotovora* subsp. *Carotovora* (Table 4). *Trichoderma viridi* showed highest inhibition zone (26.3 mm) after 48 hours of

incubation followed by *Bacillus subtilis* with inhibition zone (23.3 mm) and *Pseudomonas fluorescens* with inhibition zone (22.0 mm) after 48 hours of incubation (Table 4). Long et al. reported that the genus *Bacillus* and *Pseudomonas* have antagonistic activity against various plant pathogenic bacteria including soft rot bacterium *E. carotovora* subsp. *carotovora* in vitro (Long et al., 2003). Abd El-Khair and Karima reported that *Trichoderma* has antagonistic activity against *E. carotovora* subsp. *carotovora* in vitro (Abd-El-Khair and Karima, 2007). Cs showed highly inhibition zone with concentration 5% (28.6 mm) followed by Chitosan with concentration 3% (25.3 mm) and Chitosan with concentration 1% (22.9 mm) after 48 hours of incubation (Raju et al., 2006 and Hadwiger & Lee, 2013).

Table (4): Comparative efficiency of biocontrol agents and Chitosan against *Erwiniacarotovorasubsp. Carotovora*.

Antagonism	Inhibition zone in mm
<i>Bacillus subtilis</i>	23.3
<i>Pseudomonas fluorescens</i>	22.0
<i>Trichodermavirdi</i>	26.3
Chitosan 1%	22.9
Chitosan3%	25.3
Chitosan5%	28.6

1.2.4. Biological and chemical Control of Soft Rot Disease under Storage Conditions.


All tested bio control agents have significantly reduced severity of bacterial soft rot disease of potato individually or combined with Cs compared with the control. The highest severity of disease completely decreased by the treatment Cs 5% with biocontrol agents (*T. viridi*, *B. subtilis* and *P. fluorescens* respectively) each after 20-week storage potatoes of two Spunta and Cara varieties by (91.8, 83.6 and 80.3% respectively in Spunta c.v) and (88.6, 84.1 and 79.3% respectively in Cara c.v) as show in Figures (2,3 and 4). Followed by treatment chitosan 3% with biocontrol agents (*T. viridi*, *P. fluorescens* and *B. subtilis* respectively in Spunta c.v) by (88.6, 80.4 and 76% respectively in Spuntac.v) and (85, 80.1 and 77.3% respectively in Cara c.v) and treatment chitosan 1% with biocontrol agents (*T. viridi*, *P. fluorescens* and *B. subtilis* respectively) each by (75.3,71.3 and 66.4% respectively in Spunta c.v) and (69.2, 65.5 and 62.4% respectively in Cara c.v) compared with *T. viridi*, *B. subtilis* and *P. fluorescens* individually as show in Figure (5).



Figure(2): Effect of Chitosan 5% and *T. viridi* on potato tubers (Spunta and Cara c.v). A: Healthy tuber (Untreated control); B: Tuber infected with *E. carotovorasubsp. Carotovora*; C: Tuber infected with *E. carotovora subsp. Carotovora + Chitosan 5%+ T. viridi*



Figure (3): Effect of Chitosan 5% and *B. subtilis* on potato tubers (Spunta and Carac.v). A: Healthy tuber (Untreated control).; B:Tuber infected with *E.carotovorasubsp. Carotovora*; C: Tuber infected with *E. carotovora subsp. Carotovora + Chitosan 5%+ B. subtilis*.



Figure(4): Effect of Chitosan 5% and *P. fluorescens* on potato tubers (Spunta and Cara c.v). A: Healthy tuber (Untreated control).; B:Tuber infected with *E.carotovorasubsp. Carotovora*; C: Tuber infected with *E.carotovorasubsp. Carotovora + Chitosan 5%+ P. fluorescens*




Figure (5): Effect of biocontrol agent and chitosan on the severity of potato tuber soft rot disease. Bars indicate the standard error.

Total sugar, protein content, starch content, specific gravity (S.g) of tubers were completely increased by the treatment Cs 1%, Cs 3% and the highest result with Cs 5% each with biocontrol agents (T. viridi, B. subtilis and P. fluorescens respectively) of two Spunta and Cara varieties compared with T. viridi, P. fluorescens and B. subtilis individually and control. The results are shown in Figures (6, 7, 8 and 9). And percentages losing of weight were decreasing by the pathogen as shown in figure (10).




Figure (6): Effect of biocontrol agents and chitosan on Total sugar of tubers. Bars indicate the standard error.




Figure (7): Effect of biocontrol agents and chitosan on protein content of tubers. Bars indicate the standard error.




Figure (8): Effect of biocontrol agents and chitosan on starch content of tubers. Bars indicate the standard error.




Figure (9): Effect of biocontrol agents and chitosan on specific gravity of tubers. Bars indicate the standard error.




Figure (10): Effect of biocontrol agents and chitosan on Losing weight % of tubers. Bars indicate the standard error.

The results of the study demonstrated that the identified antagonistic bacterial (*T. viridi*, *B. subtilis* and *P. fluorescens*) combined with Cs 5% can significantly inhibit the growth of soft rot bacteria *in vitro* and in storage. *Trichoderma viride* was reported by several workers as the best antagonists for growth inhibition of several plant pathogen. It has various mechanisms of bio control include antibiosis, parasitism, inducing host-plant resistance, and competition confrontation with fungi (Howell, 2003), fungal cell wall degrading enzymes (Sharma et al., 2009 and Chutrakul et al., 2008). Fluorescent pseudomonads are aggressive rhizosphere colonizers and produce a wide range of antimicrobial compounds (Gross, 1988). Certain fluorescent pseudomonads can protect plants from diseases caused by root pathogens and often this biocontrol effect involves antimicrobial compounds such as siderophores (Bakker et al., 1987), hydrogen cyanide (Voisard et al., 1989) and antibiotics (Raaijmakers et al., 2002) like phenazine-1-carboxylate, pyrroluteorin and 2,4-diacetyl phoroglucinol. *Bacillus* sp. was considered one

of the best antagonistic bacteria for limiting growth of many pathogens demonstrated strong antagonistic actions against the soft rot bacterial pathogen which was previously identified as the most virulent strain of *Erwinia carotovora* subsp. *carotovora* both in *invitro* and under storage conditions. Many researchers previously exploited *Bacillus* sp. to control soft rot bacteria in various plants (Abd-El-Khair and Karima, 2007 and Sharga & Lyon, 1998).

Conclusion:

The three treatment of biocontrol agents combined with chitosan 5% were showed the highest level of antibiosis against pathogenic organism (*Erwinia carotovora*) the causal organism of potato soft rot, and achieve a significant control of disease on the tested varieties Cara and Spunta.

Reference:

- 1) Douches, D.S., Maas D., Jastrzebski, K. & Chase, R.W. (1996). Assessment of potato breeding progress in the USA over the last century. *Crop Science*, 36, 1544–52.
- 2) Haverkort, A.J.(1990). Ecology of potato cropping systems in relation to latitude and altitude. *Agricultural Systems*, 32, 251–72
- 3) Bathily, H., Babana, A. H., Samaké, F. (2010). *Bacillus pumilus* a new pathogen on potato tubers in Mali. *Afr. J. Microbiol. Res.*, 4(20), 2067 – 2071.
- 4) Altin, N., Bora, T. (2001). Biological control studies by fluorescent *Pseudomonas* against to *Erwinia carotovora* subsp. *Carotovora* caused soft rot on potato. Ninth Turkish Phytopathology Congress, 3–8 September 2001, Tekirdag, Turkey, pp. 104–110
- 5) Sigeo, D.C. (1993). *Bactenol plant pathology: cell and molecular aspects*. Cambridge: Cambridge University Press.
- 6) Backman, P. A., Wilson, M. and Murphy, J. F. (1997). Bacteria for biological control of plant diseases. In: *Environmentally Safe Approaches to Crop Disease Control*. Eds. N. A. Rechcigl and J. E. Rechcigl. Boca Rota: CRC Press, 95-109.
- 7) Kloepper, J. W., Ryn, C. M. & Zhang, S. (2004). Induced systemic resistance and promotion of plant growth by *Bacillus* sp. *Phytopathology*, 94, 1259-1266.
- 8) Compant, S., Duffy, B., Nowak, J., Clement, C. and Barka, E. A. (2005). Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanisms of action and future prospects. *Applied and Environmental Microbiology*, 71, 4951-4959.
- 9) Abd-El-Khair, H., Karima, H.E.H.(2007). Application of some bactericides and bioagents for controlling the soft rot disease in potato. *Research Journal Agricultural Biological Science*. 3(5):463–473
- 10) Rhoades, J., & Roller, S.(2000). Antimicrobial actions of degraded and native chitosan against spoilage organisms in laboratory media and foods. *Appl. Environ. Microbiol.* 66, 80-86.
- 11) Illum, L. (1998). Chitosan and its use as a pharmaceutical excipient. *Pharm. Res.* 15, 1326-1331.
- 12) Rouget, M. C. (1859). Des substances amylicées dans les tissus des animaux, spécialement des Articulés (chitine). *Comp. Rend.* 48,792-795.
- 13) Tharanathan, R. N., & F. S. Kittur. (2003). Chitin—the undisputed biomolecule of great potential. *Crit. Rev. Food Sci. Nutr.* 43,61-87.
- 14) Utaida, S., P. M., Dunman, D., Macapagal, E., Murphy, S. J., Projan, V. K., Singh, R. K. , Jayaswal, & Wilkinson. B. J. (2003). Genome-wide transcriptional profiling of the response of *Staphylococcus aureus* to cell-wall-active antibiotics reveals a cell-wall-stress stimulon. *Microbiology* 149:2719-2732.
- 15) Takai, K., Ohtsuka T., Senda, Y., Nakao, M., Yamamoto, K., Matsuoka, J. & Hirai. Y. (2002). Antibacterial properties of antimicrobial-finished textile products. *Microbiol. Immunol.* 46:75-81.
- 16) Dodane, V., & Vilivalam. V. D. (1998). Pharmaceutical applications of chitosan. *Pharm. Sci. Technol. Today* 1, 246-253.
- 17) Babel, S., & Kurniawan. T. A. (2003). Low-cost adsorbents for heavy metals uptake from contaminated water: a review. *J. Hazard. Mater.* 97, 219-243.

- 18) Muzzarelli R.A.A. (1983). Chitin and its derivatives: New trend of applied research. *Carbohydrate Polymer*, 3, 53–75.
- 19) Chirkov, S. N. (2002). The antiviral activity of chitosan. *Appl. Biochem. Microbiol.* 38,1-8. (In Russian.).
- 20) Helander, I. M., Nurmiäho-Lassila, E.-L., Ahvenainen, R., Rhoades, J., & Roller. S. (2001). Chitosan disrupts the barrier properties of the outer membrane of Gram-negative bacteria. *Int. J. Food Microbiol.* 71:235-244.
- 21) O'Byrne, C. P., & Booth. I. R. (2002). Osmoregulation and its importance to food-borne microorganisms. *Int. J. Food Microbiol.* 74, 203-216.
- 22) Dodane, V., & Vilivalam. V. D. (1998). Pharmaceutical applications of chitosan. *Pharm. Sci. Technol. Today.* 1, 246-253.
- 23) Kelman, A. (1974). Evaluation of selective media for isolation of soft-rot bacteria from soil and plant tissue. *Phytopathology.* 64, 468-475
- 24) Fang Z., 1998. Researching methods of plant pathogens. China Agriculture Press, Beijing, China.
- 25) Gerhardt, P. (1981). *Manual of methods of General Bacteriology.* American Society of Microbiology, Washington DC, USA, 127 p.
- 26) Garrity, G.M., Bell, J.A., Lilburn, T.G. (1984). *Bergey's Manual of Systematic Bacteriology.* 8th Ed. Science Press, Beijing, China.
- 27) Schaad, N.W., Jones, J.B., Chun, W. (2001). *Laboratory guide for identification of plant pathogenic bacteria.* Third Ed, APS Press, St. Paul, Minnesota, USA., p. 373
- 28) Cowan, S. T. (1974). *Cowan and Steel's Manual for the Identification of Medical Bacteria,* 2nd ed. Cambridge University Press, Cambridge, p. 67-83.
- 29) *Bergey's Manual of Systemic Bacteriology* (1984). Soc. Amer. Jmct. Williams and Wilkins Company. Baltimore, USA
- 30) Schaad, W., Jones, J.B. & Chun, W. (2001). *Laboratory guide for identification of plant pathogenic bacteria.* Third Ed, APS Press, St. Paul, Minnesota, USA., p. 373
- 31) Barakate, M., Ouhdouch, Y., Oufdou, Kh., Beaulieu, C. (2002). Characterization of rhizospheric soil streptomycetes from Moroccan habitats and their antimicrobial activities. *World J. Microbiol. Biotechnol.*, 18, 49–54.
- 32) Lelliot, R.A. & Stead, D.E. (1987). *Methods for the Diagnosis of Bacterial Diseases of Plants.* In: *Methods in Plant Pathology,* Preece, T.F. (Ed.). Blackwell Science, Oxford, pp: 2- 216
- 33) Klement, Z., Rudolph, K. & Sands, D.C. (1990). *Methods in Phytobacteriology.* Akademiai Kiado, Budapest, P. 568.
- 34) Wakimoto, K., Hirayae, K., Tsuchiya, K., Kushima, Y., Furuya, N., & Matsuyama, N. (1986). Production of antibiotics by plant pathogenic pseudomonads. *Annual Phytopathology Society of Japan*, 52, 835–842.
- 35) Furuya, N., Yamasaki, S., Nishioka, M., Shirasshi, I., Ityama, K., & Matsuyama, N. (1997). Antibacterial activities of pseudomonads against plantpathogenic organisms and efficacy of *Pseudomonas aeruginosa* ATCC7700 against bacterial wilt of tomato. *Annual Phytopathology Society of Japan*, 63, 417–424.
- 36) Hajhamed, A., Sayed, W. M. A. E., Yazied, A. A. E., \$ Ghaffar, N.Y. A. E. (2007). Suppression of bacterial soft rot disease of potato, *Egypt Journal Phytopathology*, 35 (2), 69– 80, 2007.
- 37) Dubois, M., Gilles, K.A., Hamilton, J.K., Robers, P.A., & Smith F. (1956). Colorimetric method for determination of sugars and related substances. *Anal. Chem.* 28, 350-356.

- 38) Thomes, W. & Dutcher, R. A. (1924). The colorimetric determination of CHO plants by Picric acid reduction methods. *J. Amer. Chem. Soc.*, 46, 7-12.
- 39) Nissen, M. (1955). The weight of potatoes in water. *Amer. Potato. J.* 32(9), 332- 339.
- 40) Simmonds, N. W. (1977). Relation between specific gravity, dry matter content and starch content of potatoes. *Potato Res.* 20, 137-140.
- 41) A.O.A.C. (1990). Official method of analysis of the association of official agriculture methods. 15th ed., Published by Association of official Analytical Chemists, Arlington, Virginia, USA.
- 42) Gomez, K.A. & Gomez, A.A. (1984). *Statistical Procedures for Agriculture Research*. 2nd Edn., Wiley-IEEE, New York, P. 680.
- 43) Gams, W., Bisset, J. (1998). Morphology and identification of *Trichoderma*. In: Kubicek cp, Harman GE (Eds *Trichoderma and Gliocladium*, vol1. Taulor and Francis, London, UK, p. 3-33
- 44) Rifai, M.A. (1969). A revision of the genus *Trichoderma*. Mycological paper No. 116. Faculty of Pure Science, University of Sheffield. England, p. 56
- 45) Long, H. H., Furuya, N., Kurose, D., Takeshita, M. and Takanami, Y. (2003). Isolation of endophytic bacteria from *Solanum* sp. And their antibacterial activity against plant pathogenic bacteria. *Journal of the Faculty of Agriculture, Kyushu University*, 48(1-2), 21-28
- 46) Raju, M. R. B., Vijai, P. and Jalali, I. (2006). Antagonistic rhizospheric bacteria for management of bacterial soft rot of radish seed crop. *Annual Plant Protection Science*, 14(2), 393-395.
- 47) Hadwiger, Lee A. (2013). "Multiple effects of chitosan on plant systems: Solid science or hype". *Plant Science* 208: 42–9. doi:10.1016/j.plantsci.2013.03.007
- 48) Howell, C.R. (2003). Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: the history and evolution of current concepts, *Plant Disease*, 87, 4–10.
- 49) Sharma, K., Kumar, M., & Misra, R.M. (2009). Morphological, biochemical and molecular characterization of *Trichoderma harzianum* isolates for their efficacy as biocontrol agents. *Journal of Phytopathology*, 157, 51-56.
- 50) Chutrakul, C., Alcocer, M., Bailey, K and Peberdy, J.F. (2008). The production and characterization of Trichotox in peptaibols by *Trichoderma asperellum*. *Chemistry and Biodiversity*. 5, 1694- 1706.
- 51) Gross, D.C. (1988). Maximising rhizosphere populations of fluorescent pseudomonads on potatoes and their effects on *Erwinia carotovora*. *Amer. Potato J.*, 65, 697-710.
- 52) Bakker, P.A.H.M., Bakker, A.W., Marugg, J.D., Weisbeek, P.J., & Schippers, B. (1987). Bioassays for studying the role of siderophores in potato growth stimulation by *Pseudomonas* spp. in short potato rotations. *Soil Biol. Biochem.*, 19, 443-449.
- 53) Voisard, C., Keel, C., Haas, D., Deëfago, G. (1989). Cyanide production by *Pseudomonas fluorescens* helps suppress black root rot of tobacco under gnotobiotic conditions. *EMBO J.*, 8, 351-358.
- 54) Raaijmakers. J.M., Vlami, M. & de Souza, J.T. (2002). Antibiotic production by bacterial Biocontrol agents. *Antonie Van Leeuwenhoek*, 81, 537-547.
- 55) Sharga, B.M. & Lyon, G.D. (1998). *Bacillus subtilis* BS 107 as an antagonist of potato blackleg and soft rot bacteria. *Canadian Journal of Microbiology*. 44(8), 777–783.

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:
<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Recent conferences: <http://www.iiste.org/conference/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

